

LAGUERRE-TYPE BELL POLYNOMIALS

P. NATALINI AND P. E. RICCI

Received 14 October 2005; Revised 4 May 2006; Accepted 9 May 2006

We develop an extension of the classical Bell polynomials introducing the Laguerre-type version of this well-known mathematical tool. The Laguerre-type Bell polynomials are useful in order to compute the n th Laguerre-type derivatives of a composite function. Incidentally, we generalize a result considered by L. Carlitz in order to obtain explicit relationships between Bessel functions and generalized hypergeometric functions.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

The Bell polynomials [1] appear in different frameworks. They are often used in combinatorial analysis [20], and even in statistics [14], although without explicit references. Moreover these polynomials have been applied even in many other contexts, such as the Blissard problem (see [20, page 46]), the representation of Lucas polynomials of the first and second kinds [4, 9], the representation formulas of Newton sum rules for polynomials' zeros [12, 13], the recurrence relations for a class of Freud-type polynomials [3], the representation of symmetric functions of a countable set of numbers, generalizing the classical algebraic Newton-Girard formulas [15]. Consequently they were also used [6] in order to find reduction formulas for the *orthogonal invariants* of a strictly positive compact operator, deriving in a simple way the so-called Robert formulas [21].

Some generalized forms of Bell polynomials already appeared in literature (see, e.g., [11, 17, 19]). A generalization of the Bell polynomials suitable for the differentiation of multivariable composite functions can also be found in [18]. Lastly, in [2], the so-called multidimensional Bell polynomials of higher order were introduced, which are suitable for representing the derivative of a composite function of several (say m) variables $f(\varphi^{(1)}(t), \varphi^{(2)}(t), \dots, \varphi^{(m)}(t))$, where $\varphi^{(i)}(t) = \phi^{(i,1)}(\phi^{(i,2)}(\dots\phi^{(i,r_i)}(t)))$, ($i=1, 2, \dots, m$).

In this article we find explicit representation formulas for the n th Laguerre-type derivatives of a composite function. The case of the first Laguerre derivative DxD , $D := d/dx$ is essentially related to an article by Carlitz [5], originated by a preceding paper by Lardner [16] in which the powers $(DxD)^n$ of this derivative appear.

2. Recalling the Bell polynomials

We recall that the Bell polynomials are a classical mathematical tool for representing the n th derivative of a composite function. In fact by considering the composite function $\Phi(t) := f(g(t))$ of functions $x = g(t)$ and $y = f(x)$ defined in suitable intervals of the real axis and n times differentiable with respect to the relevant independent variables and by using the following notations:

$$\begin{aligned} \Phi_h &:= D_t^h \Phi(t), & f_h &:= D_x^h f(x)|_{x=g(t)}, & g_h &:= D_t^h g(t), \\ ([f, g]_n) &:= (f_1, g_1; f_2, g_2; \dots; f_n, g_n), \end{aligned} \tag{2.1}$$

they are defined as follows:

$$Y_n([f, g]_n) := \Phi_n. \tag{2.2}$$

For example one has

$$\begin{aligned} Y_1([f, g]_1) &= f_1 g_1, \\ Y_2([f, g]_2) &= f_1 g_2 + f_2 g_1^2, \\ Y_3([f, g]_3) &= f_1 g_3 + f_2 (3g_2 g_1) + f_3 g_1^3. \end{aligned} \tag{2.3}$$

Further examples can be found in [20, page 49].

Inductively, we can write

$$Y_n([f, g]_n) = \sum_{k=1}^n A_{n,k}(g_1, g_2, \dots, g_n) f_k, \tag{2.4}$$

where the coefficient $A_{n,k}$, for any $k = 1, \dots, n$, is a polynomial in g_1, g_2, \dots, g_n , homogeneous of degree k and *isobaric* of weight n (i.e., it is a linear combination of monomials $g_1^{k_1} g_2^{k_2} \dots g_n^{k_n}$ whose weight is constantly given by $k_1 + 2k_2 + \dots + nk_n = n$).

For them the following result holds true.

PROPOSITION 2.1. *The Bell polynomials satisfy the recurrence relation*

$$\begin{aligned} Y_0([f, g]_0) &:= f_1, \\ Y_{n+1}([f, g]_{n+1}) &= \sum_{k=0}^n \binom{n}{k} Y_{n-k}([f, g]_{n-k}) g_{k+1}, \end{aligned} \tag{2.5}$$

where

$$([f_1, g]_{n-k}) := (f_2, g_1; f_3, g_2; \dots; f_{n-k+1}, g_{n-k}). \quad (2.6)$$

An explicit expression for the Bell polynomials is also given by the Faà di Bruno formula [10]:

$$\Phi_n = Y_n([f, g]_n) = \sum_{\pi(n)} \frac{n!}{j_1! j_2! \dots j_n!} f_j \left[\frac{g_1}{1!} \right]^{j_1} \left[\frac{g_2}{2!} \right]^{j_2} \dots \left[\frac{g_n}{n!} \right]^{j_n}, \quad (2.7)$$

where the sum runs over all partitions $\pi(n)$ of the integer n , that is, $n = j_1 + 2j_2 + \dots + nj_n$. In (2.7) j_h denotes the number of parts of size h , and $j = j_1 + j_2 + \dots + j_n$ denotes the number of parts of the considered partition. A proof of the Faà di Bruno formula can be found in [20]. In [22] the proof is based on the *umbral calculus* (see [23] and the references therein).

3. Laguerre-type derivatives

The Laguerre-type derivatives were introduced in [7, 8] in connection with a differential isomorphism denoted by the symbol $\mathcal{T} := \mathcal{T}_x$, acting onto the space $\mathcal{A} := \mathcal{A}_x$ of analytic functions of the x variable by means of the correspondence

$$D := \frac{d}{dx} \longrightarrow \hat{D}_L := Dx D; \quad x \cdot \longrightarrow \hat{D}_x^{-1}, \quad (3.1)$$

where

$$\begin{aligned} \hat{D}_x^{-1} f(x) &:= \int_0^x f(\xi) d\xi, \\ \hat{D}_x^{-n} f(x) &:= \frac{1}{(n-1)!} \int_0^x (x-\xi)^{n-1} f(\xi) d\xi, \end{aligned} \quad (3.2)$$

so that

$$\mathcal{T}_x(x^n) = \hat{D}_x^{-n}(1) := \frac{1}{(n-1)!} \int_0^x (x-\xi)^{n-1} d\xi = \frac{x^n}{n!}. \quad (3.3)$$

According to this isomorphism, the exponential operator e^x is transformed into the first Laguerre-type exponential $e_1(x) := \sum_{k=0}^{\infty} x^k / (k!)^2$ which is an eigenfunction of the *Laguerre derivative operator* $D_L := Dx D$. We have, in fact,

$$\mathcal{T}_x(e^x) = \sum_{k=0}^{\infty} \frac{\mathcal{T}_x(x^k)}{k!} = \sum_{k=0}^{\infty} \frac{x^k}{(k!)^2} = e_1(x), \quad (3.4)$$

$$\hat{D}_L e_1(ax) = a e_1(ax), \quad \forall a \in \mathbb{C}.$$

4 Laguerre-type Bell polynomials

This result can be generalized by considering the r Laguerre-type exponential $e_r(x) := \sum_{k=0}^{\infty} x^k / (k!)^{r+1}$, the r th Laguerre-type derivative operator $D_{rL} := Dx Dx D \cdots Dx D$ (containing $r + 1$ ordinary derivatives), and the iterated isomorphism \mathcal{T}^r , since

$$\mathcal{T}_x^r(e^x) = \sum_{k=0}^{\infty} \frac{\mathcal{T}_x(x^k)}{(k!)^r} = \sum_{k=0}^{\infty} \frac{x^k}{(k!)^{r+1}} = e_r(x), \quad (3.5)$$

$$\widehat{D}_{rL} e_r(ax) = a e_r(ax), \quad \forall a \in \mathbb{C}.$$

Remark 3.1. The above results show that, for every positive integer r , we can define a Laguerre-type exponential function $e_r(x)$, satisfying an eigenfunction property, which is an analog of the elementary property of the exponential. This function reduces to the exponential function when $r = 0$, so that we can put by definition

$$e_0(x) := e^x, \quad \widehat{D}_{0L} := D. \quad (3.6)$$

Obviously, $\widehat{D}_{1L} := \widehat{D}_L$.

For this reason we will refer to such functions as L -exponential functions, or shortly L -exponentials.

4. Laguerre-type Bell polynomials

The problem of constructing Bell polynomials can be extended in the natural way to the case of the Laguerre-type derivatives.

To this aim, by using notations in (2.1), we introduce the following definition.

Definition 4.1. The n th Laguerre-type Bell polynomial, denoted by ${}_{rL}Y_n(x; [f, g]_n)$, represents the n th r Laguerre-type derivative of the composite function $f(g(t))$.

We will show that ${}_{rL}Y_n$ can be expressed as a polynomial in the independent variable x , depending on $f_1, g_1; f_2, g_2; \dots; f_n, g_n$ in terms of the classical Bell polynomials.

We start noting that, according to a general result due to Viskov [24], the Laguerre derivative satisfy

$$(D_L)^n = (Dx D)^n = D^n x^n D^n, \quad (4.1)$$

and furthermore, for any order r , it turns out that

$$(D_{rL})^n = (Dx Dx \cdots Dx D)^n = D^n x^n D^n x^n \cdots D^n x^n D^n. \quad (4.2)$$

According to the above equations, the proof of Carlitz [5] can be reduced to a simple application of the Leibnitz rule, since

$$\begin{aligned} (Dx D)^n &= D^n (x^n D^n) = \sum_{k=0}^n \binom{n}{k} D^{n-k} x^n D^{n+k} \\ &= \sum_{k=0}^n \left[\binom{n}{k} \right]^2 (n-k)! x^k D^{n+k} = \sum_{k=0}^n \frac{n!}{k!} \binom{n}{k} x^k D^{n+k}. \end{aligned} \quad (4.3)$$

Therefore, the following representation formula for the Laguerre-type Bell polynomials, denoted by ${}_L Y_n$, holds true.

THEOREM 4.2. *The ${}_L Y_n$ polynomials are expressed in terms of the ordinary Bell polynomials according to the equation*

$${}_L Y_n(x; [f, g]_n) = \sum_{k=0}^n \frac{n!}{k!} \binom{n}{k} x^k Y_{n+k}([f, g]_{n+k}). \quad (4.4)$$

The above results can be easily generalized, since

$$\begin{aligned} (D_{2L})^n &= (DxDxD)^n = D^n x^n (D^n x^n D^n) \\ &= \sum_{k_1=0}^n \sum_{k_2=0}^n \frac{n!}{k_1!} \frac{(n+k_1)!}{(k_1+k_2)!} \binom{n}{k_1} \binom{n}{k_2} x^{k_1+k_2} D^{n+k_1+k_2}. \end{aligned} \quad (4.5)$$

5. The general case

The following result follows by induction.

THEOREM 5.1. *The powers of the r th Laguerre-type derivative operator $D_{rL} := DxDxD \cdots DxD$ (containing $r + 1$ ordinary derivatives) can be expanded in the form*

$$\begin{aligned} (D_{rL})^n &= (DxDxD \cdots DxD)^n = D^n x^n D^n x^n \cdots D^n x^n D^n \\ &= \sum_{k_1=0}^n \sum_{k_2=0}^n \cdots \sum_{k_r=0}^n \frac{n!}{k_1!} \frac{(n+k_1)!}{(k_1+k_2)!} \cdots \frac{(n+k_1+k_2+\cdots+k_{r-1})!}{(k_1+k_2+\cdots+k_r)!} \\ &\quad \times \binom{n}{k_1} \binom{n}{k_2} \cdots \binom{n}{k_r} x^{k_1+k_2+\cdots+k_r} D^{n+k_1+k_2+\cdots+k_r}. \end{aligned} \quad (5.1)$$

Therefore, for the r th Laguerre-type Bell polynomials denoted by ${}_{rL} Y_n$, the following result holds true.

THEOREM 5.2. *The ${}_{rL} Y_n$ polynomials are expressed in terms of the ordinary Bell polynomials according to the equation*

$$\begin{aligned} {}_{rL} Y_n(x; [f, g]_n) &= \sum_{k_1=0}^n \sum_{k_2=0}^n \cdots \sum_{k_r=0}^n \frac{n!}{k_1!} \frac{(n+k_1)!}{(k_1+k_2)!} \cdots \frac{(n+k_1+k_2+\cdots+k_{r-1})!}{(k_1+k_2+\cdots+k_r)!} \\ &\quad \times \binom{n}{k_1} \binom{n}{k_2} \cdots \binom{n}{k_r} x^{k_1+k_2+\cdots+k_r} Y_{n+k_1+k_2+\cdots+k_r}([f, g]_{n+k_1+k_2+\cdots+k_r}). \end{aligned} \quad (5.2)$$

Acknowledgments

This paper was concluded in the framework of the Italian National Group for Scientific Computation (GNCS). Useful discussions with Professor Dr. Y. Ben Cheikh are gratefully recognized.

References

- [1] E. T. Bell, *Exponential polynomials*, Annals of Mathematics. Second Series **35** (1934), no. 2, 258–277.
- [2] A. Bernardini, P. Natalini, and P. E. Ricci, *Multidimensional Bell polynomials of higher order*, Computers & Mathematics with Applications **50** (2005), no. 10–12, 1697–1708.
- [3] A. Bernardini and P. E. Ricci, *Bell polynomials and differential equations of Freud-type polynomials*, Mathematical and Computer Modelling **36** (2002), no. 9–10, 1115–1119.
- [4] M. Bruschi and P. E. Ricci, *I polinomi di Lucas e di Tchebycheff in più variabili [Lucas and Čebyšev polynomials in several variables]*, Rendiconti di Matematica. Serie VI **13** (1980), no. 4, 507–529 (1981).
- [5] L. Carlitz, *Some reduction formulas for generalized hypergeometric functions*, SIAM Journal on Mathematical Analysis **1** (1970), 243–245.
- [6] C. Cassisa and P. E. Ricci, *Orthogonal invariants and the Bell polynomials*, Rendiconti di Matematica e delle sue Applicazioni. Serie VII **20** (2000), 293–303.
- [7] G. Dattoli and P. E. Ricci, *Laguerre-type exponentials, and the relevant L-circular and L-hyperbolic functions*, Georgian Mathematical Journal **10** (2003), no. 3, 481–494.
- [8] G. Dattoli and A. Torre, *Operatorial methods and two variable Laguerre polynomials*, Atti della Accademia delle Scienze di Torino. Classe di Scienze Fisiche, Matematiche e Naturali **132** (1998), 3–9.
- [9] A. Di Cave and P. E. Ricci, *Sui polinomi di Bell ed i numeri di Fibonacci e di Bernoulli [On Bell polynomials and Fibonacci and Bernoulli numbers]*, Le Matematiche **35** (1980), no. 1–2, 84–95.
- [10] F. Faà di Bruno, *Théorie des formes binaires*, Brero, Turin, 1876.
- [11] D. Fujiwara, *Generalized Bell polynomials*, Sūgaku (Mathematics) **42** (1990), no. 1, 89–90.
- [12] T. Isoni, P. Natalini, and P. E. Ricci, *Symbolic computation of Newton sum rules for the zeros of orthogonal polynomials*, Advanced Special Functions and Integration Methods (Melfi, 2000), Proc. Melfi Sch. Adv. Top. Math. Phys., vol. 2, Aracne, Rome, 2001, pp. 97–112.
- [13] ———, *Symbolic computation of Newton sum rules for the zeros of polynomial eigenfunctions of linear differential operators*, Numerical Algorithms **28** (2001), no. 1–4, 215–227, special volume in memory of W. Gross.
- [14] M. G. Kendall and A. Stuart, *The Advanced Theory of Statistics*, Griffin, London, 1958.
- [15] A. Kurosh, *Cours d'Algèbre Supérieure*, Éditions Mir, Moscow, 1971.
- [16] T. J. Lardner, *Relations between ${}_0F_3$ and Bessel functions*, SIAM Review **11** (1969), 69–72.
- [17] P. Natalini and P. E. Ricci, *An extension of the Bell polynomials*, Computers & Mathematics with Applications **47** (2004), no. 4–5, 719–725.
- [18] S. Noschese and P. E. Ricci, *Differentiation of multivariable composite functions and Bell polynomials*, Journal of Computational Analysis and Applications **5** (2003), no. 3, 333–340.
- [19] P. N. Rai and S. N. Singh, *Generalization of Bell polynomials and related operational formula*, Vijnana Parishad Anusandhan Patrika **25** (1982), no. 3, 251–258 (Hindi).
- [20] J. Riordan, *An Introduction to Combinatorial Analysis*, Wiley Publications in Mathematical Statistics, John Wiley & Sons, New York, 1958.
- [21] D. Robert, *Invariants orthogonaux pour certaines classes d'opérateurs*, Journal de Mathématiques Pures et Appliquées. Neuvième Série **52** (1973), 81–114.
- [22] S. M. Roman, *The formula of Faà di Bruno*, The American Mathematical Monthly **87** (1980), no. 10, 805–809.

- [23] S. M. Roman and G.-C. Rota, *The umbral calculus*, Advances in Mathematics **27** (1978), no. 2, 95–188.
- [24] O. V. Viskov, *A commutative-like noncommutative identity*, Acta Scientiarum Mathematicarum (Szeged) **59** (1994), no. 3-4, 585–590.

P. Natalini: Dipartimento di Matematica, Università di Roma Tre, Largo S. Leonardo Murialdo 1, 00146 Roma, Italy

E-mail address: natalini@mat.uniroma3.it

P. E. Ricci: Dipartimento di Matematica, Università degli Studi di Roma “La Sapienza,”

P. le Aldo Moro 2, 00185 Roma, Italy

E-mail address: paoloemilio.ricci@uniroma1.it