

EIGENVALUES OF BOUNDARY VALUE PROBLEMS FOR HIGHER ORDER DIFFERENTIAL EQUATIONS

PATRICIA J. Y. WONG

Division of Mathematics, Nanyang Technological University, Singapore

RAVI P. AGARWAL

Department of Mathematics, National University of Singapore, Singapore

(Received 23 October 1995)

We shall consider the boundary value problem

$$\begin{aligned}y^{(n)} + \lambda Q(t, y, y', \dots, y^{(n-2)}) &= \lambda P(t, y, y', \dots, y^{(n-1)}), \quad n \geq 2, \quad t \in (0, 1), \\y^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 3, \\ \alpha y^{(n-2)}(0) - \beta y^{(n-1)}(0) &= 0, \\ \gamma y^{(n-2)}(1) + \delta y^{(n-1)}(1) &= 0,\end{aligned}$$

where $\lambda > 0$, α , β , γ and δ are constants satisfying $\alpha\gamma + \alpha\delta + \beta\gamma > 0$, $\beta, \delta \geq 0$, $\beta + \alpha > 0$ and $\delta + \gamma > 0$ to characterize the values of λ so that it has a positive solution. For the special case $\lambda = 1$, sufficient conditions are also established for the existence of positive solutions.

AMS No.: 34B15

KEYWORDS: Eigenvalues, positive solutions, differential equations

1. INTRODUCTION

In this paper we shall consider the n th order differential equation

$$y^{(n)} + \lambda Q(t, y, y', \dots, y^{(n-2)}) = \lambda P(t, y, y', \dots, y^{(n-1)}), \quad t \in (0, 1), \quad (1.1)$$

together with the boundary conditions

$$y^{(i)}(0) = 0, \quad 0 \leq i \leq n - 3, \quad (1.2)$$

$$\alpha y^{(n-2)}(0) - \beta y^{(n-1)}(0) = 0, \quad (1.3)$$

$$\gamma y^{(n-2)}(1) + \delta y^{(n-1)}(1) = 0, \quad (1.4)$$

where $n \geq 2$, $\lambda > 0$, α , β , γ and δ are constants so that

$$\rho = \alpha\gamma + \alpha\delta + \beta\gamma > 0, \tag{1.5}$$

and

$$\beta \geq 0, \quad \delta \geq 0, \quad \beta + \alpha > 0, \quad \delta + \gamma > 0. \tag{1.6}$$

We remark that condition (1.6) allows α and γ to be negative.

Further, we assume that there exist continuous functions $f: [0, \infty) \rightarrow (0, \infty)$ and $p, p_1, q, q_1: (0, 1) \rightarrow \mathfrak{R}$ such that

(A1) f is nondecreasing;

(A2) for $u \in [0, \infty)$,

$$q(t) \leq \frac{Q(t, u, u_1, \dots, u_{n-2})}{f(u)} \leq q_1(t), \quad p(t) \leq \frac{P(t, u, u_1, \dots, u_{n-1})}{f(u)} \leq p_1(t);$$

(A3) $q(t) - p_1(t)$ is nonnegative and is not identically zero on any subinterval of $(0, 1)$;

(A4) $\int_0^1 (\beta + \alpha t)[\delta + \gamma(1 - t)][q_1(t) - p(t)]dt < \infty$.

We shall establish upper and lower bounds for λ so that the boundary value problem (1.1)–(1.4) has a positive solution. By a *positive solution* y of (1.1)–(1.4), we mean $y \in C^{(n)}(0, 1) \cap C^{(n-1)}[0, 1]$, y satisfies (1.1) on $(0, 1)$, y fulfills (1.2)–(1.4), and y is nonnegative on $[0, 1]$, positive on some subinterval of $[0, 1]$. If, for a particular λ the boundary value problem (1.1)–(1.4) has a positive solution y , then we shall call λ an *eigenvalue* and y a corresponding *eigenfunction* of (1.1)–(1.4). Throughout, we shall let

$$E = \{\lambda > 0 \mid (1.1)–(1.4) \text{ has a positive solution}\}.$$

We note that E is the set of eigenvalues of (1.1)–(1.4).

Next, for the special case $\lambda = 1$, we shall investigate the existence of positive solutions for the boundary value problem (1.1)–(1.4), assuming further that

(A5) f is either superlinear or sublinear.

To be precise, we introduce the notations

$$f_0 = \lim_{u \rightarrow 0} \frac{f(u)}{u}, \quad f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}.$$

The function f is said to be *superlinear* if $f_0 = 0, f_\infty = \infty$, and f is *sublinear* provided $f_0 = \infty, f_\infty = 0$.

The motivation for the present work stems from many recent investigations. In fact, when $n = 2$ the boundary value problem (1.1)–(1.4) arises in various physical phenomena such as gas diffusion through porous media [3], thermal self ignition of a chemically active mixture of gases in a vessel [7,18,21], catalysis theory [10], chemically reacting systems

[25], as well as adiabatic tubular reactor processes [9]. For the present work we refer particularly to [6,8,15–17]. In all these papers, eigenvalue characterizations for particular cases of (1.1)–(1.4) are discussed. For example in [16], Fink, Gatica and Hernandez deal with the boundary value problem

$$\begin{aligned}y'' + \lambda q(t)f(y) &= 0, \quad t \in (0, 1), \\y(0) &= y(1) = 0.\end{aligned}\tag{1.7}$$

Their results are extended in [17] to systems of second order boundary value problems. In [6] and [15], the authors tackle a different boundary value problem

$$\begin{aligned}y'' + \frac{N-1}{t}y' + \lambda q(t)f(y) &= 0, \quad t \in (0, 1), \\y'(0) &= y(1) = 0.\end{aligned}\tag{1.8}$$

Recently, Chyan and Henderson [8] have studied a more general problem than (1.7), namely,

$$\begin{aligned}y^{(n)} + \lambda q(t)f(y) &= 0, \quad t \in (0, 1), \\y^{(i)}(0) &= y^{(n-2)}(1) = 0, \quad 0 \leq i \leq n-2.\end{aligned}\tag{1.9}$$

Our results not only generalize and extend the known eigenvalue theorems for (1.7)–(1.9), but also complement the discrete analog studied by Wong and Agarwal [29], as well as include several other known criteria offered in [1].

In the special case that $\lambda = 1$, applications of (1.1)–(1.4) and its discrete version have been made to singular boundary value problems by Agarwal and Wong [2,27]. Other particular cases of (1.1)–(1.4) and their discrete analogs have also been the subject matter of several recent publications on singular boundary value problems, e.g., see [1,11,20,22,23 and the references cited therein]. Further, in the particular case that $n = 2$, (1.1)–(1.4) arises in applications involving nonlinear elliptic problems in annular regions, for this we refer to [4,5,19,26]. In all these applications, it is frequent that only solutions that are positive are useful. We are particularly motivated by the work of [12–14], and our result is a generalization and extension of theirs. We further remark that other than the differential equation (1.1) considered is more general, the conditions on the coefficients in the boundary conditions are also weakened, and we allow some coefficients to be negative. Our work also complements naturally the discrete problem considered in [28].

The plan of this paper is as follows: In Section 2 we shall state a fixed point theorem due to Krasnosel'skii [24], and present some properties of a Green's function which will be used later. In Section 3 we define an appropriate Banach space and cone so that the set E can be characterized. Finally, in Section 4 we consider the special case $\lambda = 1$ and apply the fixed point theorem from [24] to yield a positive solution for (1.1)–(1.4).

2. PRELIMINARIES

THEOREM 2.1. [24] *Let B be a Banach space, and let $C \subset B$ be a cone in B . Assume Ω_1, Ω_2 are open subsets of B with $0 \in \Omega_1, \bar{\Omega}_1 \subset \Omega_2$, and let*

$$S : C \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow C$$

be a completely continuous operator such that, either

- (a) $\|Sy\| \leq \|y\|, y \in C \cap \partial\Omega_1$, and $\|Sy\| \geq \|y\|, y \in C \cap \partial\Omega_2$, or
- (b) $\|Sy\| \geq \|y\|, y \in C \cap \partial\Omega_1$, and $\|Sy\| \leq \|y\|, y \in C \cap \partial\Omega_2$.

Then, S has a fixed point in $C \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

To obtain a solution for (1.1)–(1.4), we need a mapping whose kernel $g(t, s)$ is the Green’s function of the boundary value problem

$$\begin{aligned} -y^{(n)} &= 0, \\ y^{(i)}(0) &= 0, 0 \leq i \leq n - 3, \\ \alpha y^{(n-2)}(0) - \beta y^{(n-1)}(0) &= 0, \\ \gamma y^{(n-2)}(1) + \delta y^{(n-1)}(1) &= 0. \end{aligned}$$

It can be verified that

$$G(t, s) = \frac{\partial^{n-2}}{\partial t^{n-2}} g(t, s)$$

is the Green’s function of the boundary value problem

$$\begin{aligned} -w'' &= 0, \\ \alpha w(0) - \beta w'(0) &= 0, \\ \gamma w(1) + \delta w'(1) &= 0. \end{aligned}$$

Further, we have [2]

$$G(t, s) = \frac{1}{\rho} \begin{cases} (\beta + \alpha s)[\delta + \gamma(1 - t)], & 0 \leq s \leq t \\ (\beta + \alpha t)[\delta + \gamma(1 - s)], & t \leq s \leq 1 \end{cases} \tag{2.1}$$

We observe that the conditions (1.5) and (1.6) imply that $G(t, s)$ is nonnegative on $[0, 1] \times [0, 1]$, and positive on $(0, 1) \times (0, 1)$.

LEMMA 2.1. For $(t, s) \in [1/4, 3/4] \times [0, 1]$, we have

$$G(t, s) \geq K G(s, s), \tag{2.2}$$

where $0 < K < 1$ is given by

$$K = \min \left\{ \frac{4\beta + \alpha}{4(\beta + \alpha)}, \frac{4\delta + \gamma}{4(\delta + \gamma)}, \frac{4\beta + 3\alpha}{4\beta + \alpha}, \frac{4\delta + 3\gamma}{4\delta + \gamma} \right\}. \tag{2.3}$$

Proof. For $0 \leq s \leq t$, using (2.1) the inequality (2.2) reduces to

$$\delta + \gamma(1 - t) \geq K[\delta + \gamma(1 - s)]. \tag{2.4}$$

In order that (2.4) holds, it is sufficient that K satisfies

$$\min_{t \in [\frac{1}{4}, \frac{3}{4}]} [\delta + \gamma(1 - t)] \geq K \max_{s \in [0, \frac{3}{4}]} [\delta + \gamma(1 - s)]. \tag{2.5}$$

If $\gamma \geq 0$, then (2.5) gives

$$\delta + \frac{1}{4}\gamma \geq K(\delta + \gamma), \quad \text{or} \quad K \leq \frac{4\delta + \gamma}{4(\delta + \gamma)}. \tag{2.6}$$

If $\gamma < 0$, then it follows from (2.5) that

$$\delta + \frac{3}{4}\gamma \geq K(\delta + \frac{1}{4}\gamma), \quad \text{or} \quad K \leq \frac{4\delta + 3\gamma}{4\delta + \gamma}. \tag{2.7}$$

Next, for $t \leq s \leq 1$ the inequality (2.2) becomes

$$\beta + \alpha t \geq K(\beta + \alpha s).$$

Again, it suffices to find K such that

$$\min_{t \in [\frac{1}{4}, \frac{3}{4}]} (\beta + \alpha t) \geq K \max_{s \in [\frac{1}{4}, 1]} (\beta + \alpha s). \tag{2.8}$$

If $\alpha \geq 0$, then from (2.8) we obtain

$$\beta + \frac{1}{4}\alpha \geq K(\beta + \alpha), \quad \text{or} \quad K \leq \frac{4\beta + \alpha}{4(\beta + \alpha)}. \tag{2.9}$$

If $\alpha < 0$, then (2.8) yields

$$\beta + \frac{3}{4}\alpha \geq K \left(\beta + \frac{1}{4}\alpha \right), \quad \text{or} \quad K \leq \frac{4\beta + 3\alpha}{4\beta + \alpha}. \quad (2.10)$$

Taking into account (2.6), (2.7), (2.9) and (2.10), we immediately get (2.3).

LEMMA 2.2. For $(t, s) \in [0, 1] \times [0, 1]$, we have

$$G(t, s) \leq L G(s, s), \quad (2.11)$$

where $L \geq 1$ is given by

$$L = \max \left\{ 1, \frac{\beta}{\beta + \alpha}, \frac{\delta}{\delta + \gamma} \right\}. \quad (2.12)$$

Proof. For $0 \leq s \leq t$, using (2.1) the inequality (2.11) can be written as

$$\delta + \gamma(1 - t) \leq L[\delta + \gamma(1 - s)]. \quad (2.13)$$

In the case that $\gamma \geq 0$, it is clear that we may take $L = 1$ in (2.13). If $\gamma < 0$, in order that (2.13) holds it is sufficient that L satisfies

$$\max_{t \in [0,1]} [\delta + \gamma(1 - t)] \leq L \min_{s \in [0,1]} [\delta + \gamma(1 - s)],$$

which provides

$$L \geq \frac{\delta}{\delta + \gamma}. \quad (2.14)$$

Next, for $t \leq s \leq 1$, using (2.1) the inequality (2.11) reduces to

$$\beta + \alpha t \leq L(\beta + \alpha s). \quad (2.15)$$

If $\alpha \geq 0$, then we may take $L = 1$ in (2.15).

If $\alpha < 0$, then as before it suffices to have

$$\max_{t \in [0,1]} (\beta + \alpha t) \leq L \min_{s \in [0,1]} (\beta + \alpha s),$$

which yields

$$L \geq \frac{\beta}{\beta + \alpha}. \quad (2.16)$$

The expression (2.12) is immediate from (2.14) and (2.16).

LEMMA 2.3. For $0 \leq i \leq n - 2$ and $(t, s) \in [0, 1] \times [0, 1]$, we have

$$0 \leq \frac{\partial^i}{\partial t^i} g(t, s) \leq L G(s, s) \frac{t^{n-2-i}}{(n-2-i)!}. \tag{2.17}$$

In particular, for $(t, s) \in [0, 1] \times [0, 1]$,

$$0 \leq g(t, s) \leq \frac{L}{(n-2)!} G(s, s). \tag{2.18}$$

Proof. We note that for $0 \leq i \leq n - 3$,

$$\frac{\partial^i}{\partial t^i} g(t, s) = \int_0^t \frac{\partial^{i+1}}{\partial t^{i+1}} g(\tau, s) d\tau. \tag{2.19}$$

Hence, we find

$$\frac{\partial^{n-3}}{\partial t^{n-3}} g(t, s) = \int_0^t G(\tau, s) d\tau \leq \int_0^t L G(s, s) d\tau = L G(s, s) t, \tag{2.20}$$

where we have used Lemma 2.2. It follows from (2.19) and (2.20) that

$$\frac{\partial^{n-4}}{\partial t^{n-4}} g(t, s) = \int_0^t \frac{\partial^{n-3}}{\partial t^{n-3}} g(\tau, s) d\tau \leq \int_0^t L G(s, s) \tau d\tau = L G(s, s) \frac{t^2}{2!}.$$

Continuing the process, we obtain inequality (2.17) from which (2.18) is immediate.

We shall need the following notations later: Let

$$v(t) = q_1(t) - p(t), \quad \text{and} \quad u(t) = q(t) - p_1(t).$$

For a nonnegative y on $[0, 1]$, we denote

$$\theta = \int_0^1 G(s, s)v(s)f(y(s))ds, \quad \text{and} \quad \Gamma = \int_0^1 G(s, s)u(s)f(y(s))ds.$$

In view of (A2) and (A3), it is clear that $\theta \geq \Gamma > 0$. Further, we define the constant

$$\xi = \frac{K\Gamma}{L\theta}.$$

It is noted that $0 < \xi < 1$.

3. MAIN RESULTS

Let the Banach space

$$B = \{y \in C^{(n)}(0, 1) \cap C^{(n-1)}[0, 1] \mid y^{(i)}(0) = 0, 0 \leq i \leq n - 3\}$$

with norm $\|y\| = \sup_{t \in [0, 1]} |y^{(n-2)}(t)|$, and let

$$C = \{y \in B \mid y^{(n-2)}(t) \text{ is nonnegative and is not identically zero} \\ \text{on } [0, 1]; \min_{t \in [\frac{1}{4}, \frac{3}{4}]} y^{(n-2)}(t) \geq \xi \|y\|\}.$$

It is noted that C is a cone in B . Further, we let

$$C_M = \{y \in C \mid \|y\| \leq M\}.$$

LEMMA 3.1. *Let $y \in B$. For $0 \leq i \leq n - 2$, we have*

$$|y^{(i)}(t)| \leq \frac{t^{n-2-i}}{(n-2-i)!} \|y\|, t \in [0, 1]. \quad (3.1)$$

In particular,

$$|y(t)| \leq \frac{1}{(n-2)!} \|y\|, t \in [0, 1]. \quad (3.2)$$

Proof. For $y \in B$, we have

$$y^{(n-3)}(t) = \int_0^t y^{(n-2)}(s) ds, t \in [0, 1],$$

which implies

$$|y^{(n-3)}(t)| \leq t \|y\|, t \in [0, 1]. \quad (3.3)$$

Next, since

$$y^{(n-4)}(t) = \int_0^t y^{(n-3)}(s) ds, t \in [0, 1],$$

on using (3.3) we get

$$|y^{(n-4)}(t)| \leq \int_0^t s \|y\| ds = \frac{t^2}{2!} \|y\|, \quad t \in [0, 1].$$

Continuing in the same manner we obtain (3.1) and also (3.2).

LEMMA 3.2. *Let $y \in C$. For $0 \leq i \leq n - 2$, we have*

$$y^{(i)}(t) \geq 0, \quad t \in [0, 1], \tag{3.4}$$

and

$$y^{(i)}(t) \geq \left(t - \frac{1}{4}\right)^{n-2-i} \frac{\xi}{(n-2-i)!} \|y\|, \quad t \in \left[\frac{1}{4}, \frac{3}{4}\right]. \tag{3.5}$$

In particular,

$$y(t) \geq \frac{\xi}{4^{n-2}(n-2)!} \|y\|, \quad t \in \left[\frac{1}{2}, \frac{3}{4}\right]. \tag{3.6}$$

Proof. Inequality (3.4) is obvious from the fact that

$$y^{(i)}(t) = \int_0^t y^{(i+1)}(s) ds, \quad t \in [0, 1], \quad 0 \leq i \leq n - 3.$$

To prove (3.5), we note that

$$y^{(n-3)}(t) = \int_0^t y^{(n-2)}(s) ds \geq \int_{\frac{1}{4}}^t \xi \|y\| ds = \xi \|y\| \left(t - \frac{1}{4}\right), \quad t \in \left[\frac{1}{4}, \frac{3}{4}\right]. \tag{3.7}$$

Next, on using (3.7), we find

$$\begin{aligned} y^{(n-4)}(t) &= \int_0^t y^{(n-3)}(s) ds \geq \int_{\frac{1}{4}}^t \xi \|y\| \left(s - \frac{1}{4}\right) ds \\ &= \left(t - \frac{1}{4}\right)^2 \frac{\xi}{2!} \|y\|, \quad t \in \left[\frac{1}{4}, \frac{3}{4}\right]. \end{aligned}$$

Continuing the process we obtain (3.5). Inequality (3.6) is immediate from (3.5) by taking $i = 0$ and substituting $t = 1/2$ in the right side of (3.5).

Remark 3.1. If $y \in C$ is a solution of (1.1)–(1.4), then (3.4) and (3.6) imply that y is a positive solution of (1.1)–(1.4).

To obtain a positive solution of (1.1)–(1.4), we shall seek a fixed point of the operator λS in the cone C , where $S: C \rightarrow B$ is defined by

$$\begin{aligned}
 Sy(t) &= \int_0^1 g(t, s)[Q(s, y, y', \dots, y^{(n-2)}) \\
 &\quad - P(s, y, y', \dots, y^{(n-1)})]ds, \quad t \in [0, 1].
 \end{aligned} \tag{3.8}$$

In view of condition (A2) we get

$$Uy(t) \leq Sy(t) \leq Vy(t), \quad t \in [0, 1], \tag{3.9}$$

where

$$Uy(t) = \int_0^1 g(t, s)u(s)f(y(s))ds, \tag{3.10}$$

and

$$Vy(t) = \int_0^1 g(t, s)v(s)f(y(s))ds. \tag{3.11}$$

Since for $t \in [0, 1]$,

$$(Sy)^{(n-2)}(t) = \int_0^1 G(t, s)[Q(s, y, y', \dots, y^{(n-2)}) - P(s, y, y', \dots, y^{(n-1)})]ds,$$

$$(Uy)^{(n-2)}(t) = \int_0^1 G(t, s)u(s)f(y(s))ds,$$

and

$$(Vy)^{(n-2)}(t) = \int_0^1 G(t, s)v(s)f(y(s))ds,$$

using (A2) again gives

$$(Uy)^{(n-2)}(t) \leq (Sy)^{(n-2)}(t) \leq (Vy)^{(n-2)}(t), \quad t \in [0, 1]. \tag{3.12}$$

We shall now show that the operator S is compact on the cone C . Let us consider the case when $u(t)$ is unbounded in a deleted right neighborhood of 0 and also in a deleted left neighborhood of 1. Clearly, $v(t)$ is also unbounded near 0 and 1. For $m \in \{1, 2, 3, \dots\}$, define $u_m, v_m: [0, 1] \rightarrow \mathfrak{R}$ by

$$u_m(t) = \begin{cases} u\left(\frac{1}{m+1}\right), & 0 \leq t \leq \frac{1}{m+1} \\ u(t), & \frac{1}{m+1} \leq t \leq \frac{m}{m+1} \\ u\left(\frac{m}{m+1}\right), & \frac{m}{m+1} \leq t \leq 1, \end{cases} \tag{3.13}$$

$$v_m(t) = \begin{cases} v\left(\frac{1}{m+1}\right), & 0 \leq t \leq \frac{1}{m+1} \\ v(t), & \frac{1}{m+1} \leq t \leq \frac{m}{m+1} \\ v\left(\frac{m}{m+1}\right), & \frac{m}{m+1} \leq t \leq 1, \end{cases} \tag{3.14}$$

and the operators $U_m, V_m: C \rightarrow B$ by

$$U_m y(t) = \int_0^1 g(t, s) u_m(s) f(y(s)) ds,$$

$$V_m y(t) = \int_0^1 g(t, s) v_m(s) f(y(s)) ds.$$

It is standard that for each m , both U_m and V_m are compact operators on C . Let $M > 0$ and $y \in C_M$. Then, in view of (2.18) and (3.2), we find

$$\begin{aligned} & |V_m y(t) - V y(t)| \\ &= \int_0^1 g(t, s) |v_m(s) - v(s)| f(y(s)) ds \\ &= \int_0^{\frac{1}{m+1}} g(t, s) |v_m(s) - v(s)| f(y(s)) ds \\ &\quad + \int_{\frac{m}{m+1}}^1 g(t, s) |v_m(s) - v(s)| f(y(s)) ds \\ &\leq \frac{L}{(n-2)!} f\left(\frac{M}{(n-2)!}\right) \left[\int_0^{\frac{1}{m+1}} G(s, s) |v_m(s) - v(s)| ds \right. \\ &\quad \left. + \int_{\frac{m}{m+1}}^1 G(s, s) |v_m(s) - v(s)| f(y(s)) ds \right] \\ &= \frac{L}{(n-2)!} f\left(\frac{M}{(n-2)!}\right) \left[\int_0^{\frac{1}{m+1}} G(s, s) \left| v\left(\frac{1}{m+1}\right) - v(s) \right| ds \right. \\ &\quad \left. + \int_{\frac{m}{m+1}}^1 G(s, s) \left| v\left(\frac{m}{m+1}\right) - v(s) \right| f(y(s)) ds \right]. \end{aligned}$$

The integrability of $G(t, t)v(t)$ (condition (A4)) implies that V_m converges uniformly to V on C_M . Hence, V is compact on C . Similarly, we can verify that U_m converges uniformly to U on C_M and therefore U is compact on C . It follows from inequality (3.9) that the operator S is compact on C .

THEOREM 3.1. *There exists a $c > 0$ such that the interval $(0, c] \subseteq E$.*

Proof. Let $M > 0$ be given. Define

$$c = M \left\{ \frac{L}{\rho} f \left(\frac{M}{(n-2)!} \right) \int_0^1 (\beta + \alpha s) [\delta + \gamma(1-s)] v(s) ds \right\}^{-1}. \quad (3.15)$$

Let $y \in C_M$ and $0 < \lambda \leq c$. We shall prove that $\lambda Sy \in C_M$. For this, first we shall show that $\lambda Sy \in C$. From (3.12) and (A3), we find

$$(\lambda Sy)^{(n-2)}(t) \leq \lambda \int_0^1 G(t, s) u(s) f(y(s)) ds \geq 0, \quad t \in [0, 1]. \quad (3.16)$$

Further, it follows from (3.12) and Lemma 2.2 that

$$\begin{aligned} (Sy)^{(n-2)}(t) &\leq \int_0^1 G(t, s) v(s) f(y(s)) ds \\ &\leq L \int_0^1 G(s, s) v(s) f(y(s)) ds, \quad t \in [0, 1]. \end{aligned}$$

Therefore,

$$\|Sy\| \leq L \int_0^1 G(s, s) v(s) f(y(s)) ds = L\theta. \quad (3.17)$$

Now, on using (3.12), Lemma 2.1 and (3.17), we find for $t \in \left[\frac{1}{4}, \frac{3}{4} \right]$,

$$\begin{aligned} (\lambda Sy)^{(n-2)}(t) &\geq \lambda \int_0^1 G(t, s) u(s) f(y(s)) ds \\ &\geq \lambda K \int_0^1 G(s, s) u(s) f(y(s)) ds \\ &= \lambda K \Gamma \geq \lambda \xi \|Sy\| = \xi \|\lambda Sy\|. \end{aligned}$$

Hence,

$$\min_{t \in \left[\frac{1}{4}, \frac{3}{4} \right]} (\lambda Sy)^{(n-2)}(t) \geq \xi \|\lambda Sy\|. \quad (3.18)$$

It follows from (3.16) and (3.18) that $\lambda Sy \in C$.

Next, on using (3.12), Lemma 2.2, (3.2), (2.1) and (3.15) successively, we get

$$\begin{aligned} (\lambda Sy)^{(n-2)}(t) &\leq \lambda \int_0^1 G(t, s) v(s) f(y(s)) ds \\ &\leq L \lambda \int_0^1 G(s, s) v(s) f(y(s)) ds \\ &\leq L \lambda \int_0^1 G(s, s) v(s) f \left(\frac{M}{(n-2)!} \right) ds \end{aligned}$$

$$\begin{aligned}
 &= \frac{L\lambda}{\rho} \int_0^1 (\beta + \alpha s)[\delta + \gamma(1 - s)]v(s)f\left(\frac{M}{(n - 2)!}\right) ds \\
 &\leq M, t \in [0, 1],
 \end{aligned}$$

which implies

$$\|\lambda Sy\| \leq M.$$

Hence, $(\lambda S)(C_M) \subseteq C_M$. Also, the standard arguments yield that λS is completely continuous. By Schauder fixed point theorem, λS has a fixed point in C_M . Clearly, this fixed point is a positive solution of (1.1)–(1.4) and therefore λ is an eigenvalue of (1.1)–(1.4). Since $0 < \lambda \leq c$ is arbitrary, it follows immediately that $(0, c] \subseteq E$.

The next theorem makes use of the monotonicity and compactness of the operator S on the cone C . We refer to [16, Theorem 3.2] for its proof.

THEOREM 3.2. [16] *Suppose that $\lambda_0 \in E$. Then, for each $0 < \lambda < \lambda_0$, $\lambda \in E$.*

The following corollary is immediate from Theorem 3.2.

COROLLARY 3.1. *E is an interval.*

We shall establish conditions under which E is a bounded or unbounded interval. For this, we need the following results.

THEOREM 3.3. *Let λ be an eigenvalue of (1.1)–(1.4) and $y \in C$ be a corresponding eigenfunction.*

(a) Suppose that $\delta = \beta = 0$ and $\gamma = \alpha = 1$. If

$$y^{(n-1)}(0) = \nu \tag{3.19}$$

for some $\nu > 0$, then λ satisfies

$$a(\nu)\nu \left[f\left(\frac{\nu}{(n - 1)!}\right) \right]^{-1} \leq \lambda \leq a(u)\nu [f(0)]^{-1}, \tag{3.20}$$

where

$$a(z) = \left\{ \int_0^1 (1 - s)z(s)ds \right\}^{-1}. \tag{3.21}$$

(b) Suppose that $\delta > 0$, $\beta = 0$ and (3.19) holds for some $\nu > 0$.

(i) If $\gamma \geq 0$, then λ satisfies

$$b(\nu)\nu(\gamma + \delta) \left[f\left(\frac{\nu}{(n - 1)!}\right) \right]^{-1} \leq \lambda \leq b(u)\nu(\gamma + \delta)[f(0)]^{-1}, \tag{3.22}$$

where

$$b(z) = \left\{ \int_0^1 [\gamma(1-s) + \delta]z(s)ds \right\}^{-1}. \quad (3.23)$$

(ii) If $\gamma < 0$, then λ satisfies

$$c \left(u, v, f, 0, \frac{v}{(n-1)!} \right) v(\gamma + \delta) \leq \lambda \leq c \left(v, u, f, \frac{v}{(n-1)!}, 0 \right) v(\gamma + \delta), \quad (3.24)$$

where

$$c(z, w, f, \theta_1, \theta_2) = \left\{ \gamma f(\theta_1) \int_0^1 (1-s)z(s)ds + \delta f(\theta_2) \int_0^1 w(s)ds \right\}^{-1}. \quad (3.25)$$

(c) Suppose that $\delta = 0$ and $\beta > 0$. If

$$y^{(n-2)}(0) = \mu, \quad y^{(n-1)}(0) = \nu \quad (3.26)$$

for some $\mu, \nu > 0$ such that $\alpha\mu = \beta\nu$, then λ satisfies

$$a(\nu) (\mu + \nu) \left[f \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \right]^{-1} \leq \lambda \leq a(\mu) (\mu + \nu) [f(0)]^{-1}, \quad (3.27)$$

where $a(\cdot)$ is defined in (3.21).

(d) Suppose that $\delta > 0$, $\beta > 0$ and (3.26) holds for some $\mu, \nu > 0$ such that $\alpha\mu = \beta\nu$.

(i) If $\gamma \geq 0$, then λ satisfies

$$\begin{aligned} b(\nu) [\gamma(\mu + \nu) + \delta\nu] \left[f \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \right]^{-1} \\ \leq \lambda \leq b(\mu) [\gamma(\mu + \nu) + \delta\nu] [f(0)]^{-1}, \end{aligned} \quad (3.28)$$

where $b(\cdot)$ is defined in (3.23).

(ii) If $\gamma < 0$, then λ satisfies

$$\begin{aligned} c \left(u, v, f, 0, \frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) [\gamma(\mu + \nu) + \delta\nu] \\ \leq \lambda \leq c \left(v, u, f, \frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!}, 0 \right) [\gamma(\mu + \nu) + \delta\nu], \end{aligned} \quad (3.29)$$

where $c(\cdot, \cdot, \cdot, \cdot, \cdot)$ is defined in (3.25).

Proof. (a) In this case, the boundary conditions (1.2)–(1.4) reduce to

$$\begin{aligned} y^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 2, \\ y^{(n-2)}(1) &= 0. \end{aligned} \tag{3.30}$$

For $m \in \{1, 2, 3, \dots\}$, we define $f_m = f * \eta_m$, where η_m is a standard mollifier [8, 16] such that f_m is Lipschitz and converges uniformly to f .

For a fixed m , let λ_m be an eigenvalue and y_m , with $y_m^{(n-1)}(0) = \nu$, be a corresponding eigenfunction of the following boundary value problem

$$\begin{aligned} y_m^{(n)} + \lambda_m Q_m(t, y_m, y_m', \dots, y_m^{(n-2)}) &= \lambda_m P_m(t, y_m, y_m', \dots, y_m^{(n-1)}), \\ t &\in [0, 1], \end{aligned} \tag{3.31}$$

$$\begin{aligned} y_m^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 2, \\ y_m^{(n-2)}(1) &= 0, \end{aligned} \tag{3.32}$$

where Q_m and P_m converge uniformly to Q and P respectively, and

$$u_m(t) \leq \frac{Q_m(t, z, z_1, \dots, z_{n-2}) - P_m(t, z, z_1, \dots, z_{n-1})}{f_m(z)} \leq v_m(t) \tag{3.33}$$

(see (3.13) and (3.14) for the definitions of $u_m(t)$ and $v_m(t)$).

Clearly, y_m is the unique solution of the initial value problem (3.31),

$$\begin{aligned} y_m^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 2, \\ y_m^{(n-1)}(0) &= \nu. \end{aligned} \tag{3.34}$$

Since

$$\begin{aligned} y_m^{(n)}(t) &= \lambda_m [P_m(t, y_m, y_m', \dots, y_m^{(n-1)}) - Q_m(t, y_m, y_m', \dots, y_m^{(n-2)})] \\ &\leq -\lambda_m u_m(t) f_m(y_m(t)) \\ &\leq 0, \end{aligned}$$

we have $y_m^{(n-1)}$ is nonincreasing and hence

$$y_m^{(n-1)}(t) \leq y_m^{(n-1)}(0) = \nu, \quad t \in [0, 1]. \tag{3.35}$$

Using the initial conditions (3.34) and (3.35), we find for $t \in [0, 1]$,

$$y_m^{(n-2)}(t) = \int_0^t y_m^{(n-1)}(s) ds \leq \int_0^t v ds = vt.$$

This in turn leads to

$$y_m^{(n-3)}(t) = \int_0^t y_m^{(n-2)}(s) ds \leq \int_0^t v s ds = v \frac{t^2}{2}, t \in [0, 1].$$

Continuing the process we obtain for $t \in [0, 1]$,

$$y_m(t) \leq v \frac{t^{n-1}}{(n-1)!} \leq \frac{v}{(n-1)!}. \quad (3.36)$$

Now, from (3.31), (3.33) and (3.36) we get for $t \in [0, 1]$,

$$\lambda_m u_m(t) f_m(0) \leq -y_m^{(n)}(t) \leq \lambda_m v_m(t) f_m \left(\frac{v}{(n-1)!} \right). \quad (3.37)$$

An integration of (3.37) from 0 to t provides

$$\phi_1(t) \leq y_m^{(n-1)}(t) \leq \phi_2(t), t \in [0, 1], \quad (3.38)$$

where

$$\phi_1(t) = v - \lambda_m f_m \left(\frac{v}{(n-1)!} \right) \int_0^t v_m(s) ds,$$

and

$$\phi_2(t) = v - \lambda_m f_m(0) \int_0^t u_m(s) ds.$$

Again, we integrate (3.38) from 0 to t , and subsequently change the order of integration, to obtain

$$\phi_3(t) \leq y_m^{(n-2)}(t) \leq \phi_4(t), t \in [0, 1], \quad (3.39)$$

where

$$\phi_3(t) = vt - \lambda_m f_m \left(\frac{v}{(n-1)!} \right) \int_0^t (t-s) v_m(s) ds,$$

and

$$\phi_4(t) = vt - \lambda_m f_m(0) \int_0^t (t-s) u_m(s) ds.$$

From inequality (3.39), in order to have $y_m^{(n-2)}(1) = 0$ (see (3.32)), it is necessary that

$$\phi_3(1) \leq 0, \quad \text{and} \quad \phi_4(1) \geq 0,$$

or equivalently,

$$\lambda_m \geq a(v_m)v \left[f_m \left(\frac{v}{(n-1)!} \right) \right]^{-1}, \tag{3.40}$$

and

$$\lambda_m \leq a(u_m)v [f_m(0)]^{-1}. \tag{3.41}$$

A combination of (3.40) and (3.41) yields

$$a(v_m)v \left[f_m \left(\frac{v}{(n-1)!} \right) \right]^{-1} \leq \lambda_m \leq a(u_m)v [f_m(0)]^{-1}. \tag{3.42}$$

It follows from (3.38) that $\{y_m^{(n-1)}\}_{m=1}^\infty$ is a uniformly bounded sequence on $[0, 1]$. Using the initial conditions (3.34) and repeated integrations, we find that $\{y_m^{(i)}\}_{m=1}^\infty, 0 \leq i \leq n - 1$ is a uniformly bounded sequence. Thus, there exists a subsequence, which can be relabelled as $\{y_m\}_{m=1}^\infty$, that converges uniformly (in fact, in $C^{(n-1)}$ -norm) to some y on $[0, 1]$. We note that each $y_m(t)$ can be expressed as

$$\begin{aligned} y_m(t) &= \lambda_m \int_0^1 g(t, s)[Q_m(s, y_m, y_m', \dots, y_m^{(n-2)}) \\ &\quad - P_m(s, y_m, y_m', \dots, y_m^{(n-1)})]ds, \quad t \in [0, 1]. \end{aligned} \tag{3.43}$$

Since $\{\lambda_m\}$ is a bounded sequence (from (3.42)), there is a subsequence, which can be relabelled as $\{\lambda_m\}$, that converges to some λ . Then, letting $m \rightarrow \infty$ in (3.43) yields

$$\begin{aligned} y(t) &= \lambda \int_0^1 g(t, s)[Q(s, y, y', \dots, y^{(n-2)}) \\ &\quad - P(s, y, y', \dots, y^{(n-1)})]ds, \quad t \in [0, 1]. \end{aligned}$$

This means that y is an eigenfunction of (1.1)–(1.4) corresponding to the eigenvalue λ . Further, $y^{(n-1)}(0) = v$, and inequality (3.20) follows from (3.42) immediately.

(b) Here, the boundary conditions (1.2)–(1.4) reduce to

$$\begin{aligned} y^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 2, \\ \gamma y^{(n-2)}(1) + \delta y^{(n-1)}(1) &= 0. \end{aligned}$$

Using a similar technique as in case (a), for a fixed m we let λ_m be an eigenvalue and y_m , with $y_m^{(n-1)}(0) = \nu$, be a corresponding eigenfunction of the boundary value problem (3.31),

$$\begin{aligned} y_m^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 2, \\ \gamma y_m^{(n-2)}(1) + \delta y_m^{(n-1)}(1) &= 0. \end{aligned} \tag{3.44}$$

It is obvious that the eigenfunction y_m is the unique solution of the initial value problem (3.31), (3.34). As before we get the inequalities (3.38) and (3.39).

If $\gamma \geq 0$, then

$$\gamma \phi_3(t) + \delta \phi_1(t) \leq \gamma y_m^{(n-2)}(t) + \delta y_m^{(n-1)}(t) \leq \gamma \phi_4(t) + \delta \phi_2(t). \tag{3.45}$$

If $\gamma < 0$, then (3.38) and (3.39) lead to

$$\gamma \phi_4(t) + \delta \phi_1(t) \leq \gamma y_m^{(n-2)}(t) + \delta y_m^{(n-1)}(t) \leq \gamma \phi_3(t) + \delta \phi_2(t). \tag{3.46}$$

Since y_m satisfies $\gamma y_m^{(n-2)}(1) + \delta y_m^{(n-1)}(1) = 0$ (from (3.44)), in inequality (3.45) it is necessary that

$$\gamma \phi_3(1) + \delta \phi_1(1) \leq 0, \quad \text{and} \quad \gamma \phi_4(1) + \delta \phi_2(1) \geq 0,$$

which provide

$$b(v_m)\nu(\gamma + \delta) \left[f_m \left(\frac{\nu}{(n-1)!} \right) \right]^{-1} \leq \lambda_m \leq b(u_m)\nu(\gamma + \delta)[f_m(0)]^{-1}, \quad \gamma \geq 0. \tag{3.47}$$

Likewise, in inequality (3.46) we must have

$$\gamma \phi_4(1) + \delta \phi_1(1) \leq 0, \quad \text{and} \quad \gamma \phi_3(1) + \delta \phi_2(1) \geq 0,$$

which give

$$\begin{aligned} c \left(u_m, v_m, f_m, 0, \frac{\nu}{(n-1)!} \right) \nu(\gamma + \delta) &\leq \lambda_m \leq c \left(v_m, u_m, f_m, \frac{\nu}{(n-1)!}, 0 \right) \\ &\times \nu(\gamma + \delta), \quad \gamma < 0. \end{aligned} \tag{3.48}$$

Using a similar argument as in case (a), we get y_m converges to y (satisfying (3.19)), λ_m converges to λ , and y is a eigenfunction of (1.1)–(1.4) corresponding to the eigenvalue λ . Further, inequalities (3.22) and (3.24) follow from (3.47) and (3.48), respectively.

(c) In this case, the boundary conditions (1.2)–(1.4) become

$$\begin{aligned} y^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 3, \\ y^{(n-2)}(1) &= 0, \\ \alpha y^{(n-2)}(0) - \beta y^{(n-1)}(0) &= 0. \end{aligned}$$

For a fixed m , let λ_m be an eigenvalue and y_m , with $y_m^{(n-2)}(0) = \mu, y_m^{(n-1)}(0) = \nu, \alpha\mu = \beta\nu$, be a corresponding eigenfunction of the boundary value problem (3.31),

$$\begin{aligned} y_m^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 3, \\ y_m^{(n-2)}(1) &= 0, \\ \alpha y_m^{(n-2)}(0) - \beta y_m^{(n-1)}(0) &= 0. \end{aligned} \tag{3.49}$$

Clearly, the eigenfunction y_m is the unique solution of the differential equation (3.31), together with the initial conditions

$$\begin{aligned} y_m^{(i)}(0) &= 0, \quad 0 \leq i \leq n - 3, \\ y_m^{(n-2)}(0) &= \mu, \\ y_m^{(n-1)}(0) &= \nu. \end{aligned} \tag{3.50}$$

As in case (a), we see that $y_m^{(n-1)}$ is nonincreasing and hence (3.35) holds. In view of the initial conditions (3.50) and (3.35), we find

$$y_m^{(n-2)}(t) = \mu + \int_0^t y_m^{(n-1)}(s) ds \leq \mu + \int_0^t \nu ds = \mu + \nu t, \quad t \in [0, 1].$$

It follows that

$$y_m^{(n-3)}(t) = \int_0^t y_m^{(n-2)}(s) ds \leq \int_0^t (\mu + \nu s) ds = \mu t + \nu \frac{t^2}{2}, \quad t \in [0, 1].$$

Continuing the process we obtain for $t \in [0, 1]$,

$$y_m(t) \leq \mu \frac{t^{n-2}}{(n-2)!} + \nu \frac{t^{n-1}}{(n-1)!} \leq \frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!}. \tag{3.51}$$

Now, it follows from (3.31), (3.33) and (3.51) that for $t \in [0, 1]$,

$$\lambda_m u_m(t) f_m(0) \leq -y_m^{(n)}(t) \leq \lambda_m v_m(t) f_m \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right). \tag{3.52}$$

An integration of (3.52) from 0 to t gives

$$\phi_5(t) \leq y_m^{(n-1)}(t) \leq \phi_6(t), \quad t \in [0, 1], \quad (3.53)$$

where

$$\phi_5(t) = \nu - \lambda_m f_m \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \int_0^t \nu_m(s) ds,$$

and

$$\phi_6(t) = \nu - \lambda_m f_m(0) \int_0^t u_m(s) ds.$$

Once again, we integrate (3.53) from 0 to t , to get

$$\phi_7(t) \leq y_m^{(n-2)}(t) \leq \phi_8(t), \quad t \in [0, 1], \quad (3.54)$$

where

$$\phi_7(t) = \mu + \nu t - \lambda_m f_m \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \int_0^t (t-s) \nu_m(s) ds,$$

and

$$\phi_8(t) = \mu + \nu t - \lambda_m f_m(0) \int_0^t (t-s) u_m(s) ds.$$

Since y_m satisfies the boundary condition $y_m^{(n-2)}(1) = 0$ (see (3.49)), in inequality (3.54) we must have

$$\phi_7(1) \leq 0, \quad \text{and} \quad \phi_8(1) \geq 0,$$

or equivalently,

$$\lambda_m \geq a(\nu_m) (\mu + \nu) \left[f_m \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \right]^{-1}, \quad (3.55)$$

and

$$\lambda_m \leq a(u_m) (\mu + \nu) [f_m(0)]^{-1}. \quad (3.56)$$

Again, using a similar argument as in case (a), we find that y_m converges to y (satisfying (3.26)), λ_m converges to λ , and y is a eigenfunction of (1.1)–(1.4) corresponding to the eigenvalue λ . Further, inequality (3.27) follows immediately from (3.55) and (3.56). (d)

For a fixed m , let λ_m be an eigenvalue and y_m , with $y_m^{(n-2)}(0) = \mu, y_m^{(n-1)}(0) = \nu, \alpha\mu = \beta\nu$, be a corresponding eigenfunction of the boundary value problem (3.31),

$$y_m^{(i)}(0) = 0, 0 \leq i \leq n - 3,$$

$$\alpha y_m^{(n-2)}(0) - \beta y_m^{(n-1)}(0) = 0,$$

$$\gamma y_m^{(n-2)}(1) + \delta y_m^{(n-1)}(1) = 0.$$

It is obvious that the eigenfunction y_m is the unique solution of the initial value problem (3.31), (3.50). As in case (c), we get the inequalities (3.53) and (3.54) which lead to

$$\gamma\phi_7(t) + \delta\phi_5(t) \leq \gamma y_m^{(n-2)}(t) + \delta y_m^{(n-1)}(t) \leq \gamma\phi_8(t) + \delta\phi_6(t), \gamma \geq 0, \tag{3.57}$$

and

$$\gamma\phi_8(t) + \delta\phi_5(t) \leq \gamma y_m^{(n-2)}(t) + \delta y_m^{(n-1)}(t) \leq \gamma\phi_7(t) + \delta\phi_6(t), \gamma < 0. \tag{3.58}$$

Since y_m satisfies the boundary condition $\gamma y_m^{(n-2)}(1) + \delta y_m^{(n-1)}(1) = 0$, in inequality (3.57) it is necessary that

$$\gamma\phi_7(1) + \delta\phi_5(1) \leq 0, \quad \text{and} \quad \gamma\phi_8(1) + \delta\phi_6(1) \geq 0,$$

which reduce to

$$\begin{aligned} b(v_m) [\gamma(\mu + \nu) + \delta\nu] \left[f_m \left(\frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) \right]^{-1} \\ \leq \lambda_m \leq b(u_m) [\gamma(\mu + \nu) + \delta\nu] [f_m(0)]^{-1}, \gamma \geq 0. \end{aligned} \tag{3.59}$$

Likewise, in inequality (3.58) we must have

$$\gamma\phi_8(1) + \delta\phi_5(1) \leq 0, \quad \text{and} \quad \gamma\phi_7(1) + \delta\phi_6(1) \geq 0,$$

which provide

$$\begin{aligned} c \left(u_m, v_m, f_m, 0, \frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!} \right) [\gamma(\mu + \nu) + \delta\nu] \\ \leq \lambda_m \leq c \left(v_m, u_m, f_m, \frac{\mu}{(n-2)!} + \frac{\nu}{(n-1)!}, 0 \right) [\gamma(\mu + \nu) + \delta\nu], \\ \gamma < 0. \end{aligned} \tag{3.60}$$

Once again, using a similar argument as in case (a), we find that y_m converges to y (satisfying (3.26)), λ_m converges to λ , and y is a eigenfunction of (1.1)–(1.4) corresponding to the eigenvalue λ . Further, inequalities (3.28) and (3.29) follow immediately from (3.59) and (3.60) respectively.

THEOREM 3.4. *Let λ be an eigenvalue of (1.1)–(1.4) and $y \in C$ be a corresponding eigenfunction. Further, let $\eta = \|y\|$. Then,*

$$\lambda \geq \frac{\eta\rho}{L} \left\{ f\left(\frac{\eta}{(n-2)!}\right) \int_0^1 (\beta + \alpha s)[\delta + \gamma(1-s)]v(s)ds \right\}^{-1}. \tag{3.61}$$

Also, there exists a $c > 0$ such that

$$\lambda \leq \frac{\eta\rho}{f(c\eta)} \left\{ \int_{\frac{1}{4}}^{\frac{1}{2}} \left(\delta + \frac{\gamma}{2}\right) (\beta + \alpha s)u(s)ds \right\}^{-1}. \tag{3.62}$$

Proof. We observe that $y^{(n)}$ is nonpositive and hence $y^{(n-2)}$ is concave on $[0, 1]$. This, together with the fact that $y^{(n-2)}$ is nonnegative, implies the existence of a unique $t_0 \in (0, 1)$ such that

$$\eta = \|y\| = y^{(n-2)}(t_0).$$

To prove that (3.61) holds, we use (3.12), Lemma 2.2, (3.2) and (2.1) successively, to get

$$\begin{aligned} \eta &= y^{(n-2)}(t_0) = (\lambda S y)^{(n-2)}(t_0) \\ &\leq \lambda \int_0^1 G(t_0, s)v(s) f(y(s))ds \\ &\leq \lambda L \int_0^1 G(s, s)v(s) f(y(s))ds \\ &\leq \lambda L \int_0^1 G(s, s)v(s) f\left(\frac{\eta}{(n-2)!}\right) ds \\ &= \frac{\lambda L}{\rho} f\left(\frac{\eta}{(n-2)!}\right) \int_0^1 (\beta + \alpha s)[\delta + \gamma(1-s)]v(s)ds. \end{aligned}$$

The inequality (3.61) now follows immediately.

Next, to prove (3.62) we shall consider four cases.

Case 1 $\delta = \beta = 0, \gamma = \alpha = 1$

Here, $y^{(n-2)}(0) = y^{(n-2)}(1) = 0$. By the concavity of $y^{(n-2)}$, we find

$$\begin{aligned} y^{(n-2)}(t) &\geq \begin{cases} \frac{\eta}{t_0} t, & t \in [0, t_0] \\ \frac{\eta}{1-t_0} (1-t), & t \in [t_0, 1] \end{cases} \\ &\geq \eta t(1-t), t \in [0, 1]. \end{aligned} \tag{3.63}$$

Thus, on using (1.2) and (3.63) we get for $t \in [0, 1]$,

$$y^{(n-3)}(t) = \int_0^t y^{(n-2)}(s) ds \geq \int_0^t \eta s(1-s) ds = \eta \left(\frac{t^2}{2} - \frac{t^3}{3} \right).$$

Continuing the integration process we obtain

$$y(t) \geq \eta \psi(t), \quad t \in [0, 1], \tag{3.64}$$

where

$$\psi(t) = \frac{t^{n-1}}{(n-1)!} - 2 \frac{t^n}{n!}.$$

We note that

$$\psi'(t) = \frac{t^{n-2}}{(n-2)!} \left(1 - \frac{2t}{n-1} \right)$$

is nonnegative for $t \in I \equiv \left[0, \frac{n-1}{2} \right]$. Hence, in particular $\psi(t)$ is nondecreasing for $t \in J \equiv \left[\frac{1}{4}, \frac{1}{2} \right] \subseteq I$. It follows from (3.64) that

$$y(t) \geq c\eta, \quad t \in J, \tag{3.65}$$

where

$$c = \psi \left(\frac{1}{4} \right) = \frac{1}{4^{n-1}(n-1)!} - \frac{2}{4^n n!} > 0. \tag{3.66}$$

Now, in view of (3.12), (3.65) and (2.1), we find

$$\begin{aligned} \eta &\geq y^{(n-2)} \left(\frac{1}{2} \right) = (\lambda Sy)^{(n-2)} \left(\frac{1}{2} \right) \\ &\geq \lambda \int_0^1 G \left(\frac{1}{2}, s \right) u(s) f(y(s)) ds \\ &\geq \lambda \int_{1/4}^{1/2} G \left(\frac{1}{2}, s \right) u(s) f(y(s)) ds \\ &\geq \lambda \int_{1/4}^{1/2} G \left(\frac{1}{2}, s \right) u(s) f(c\eta) ds \\ &= \frac{\lambda}{\rho} f(c\eta) \int_{1/4}^{1/2} \left(\delta + \frac{\gamma}{2} \right) (\beta + \alpha s) u(s) ds \end{aligned}$$

from which (3.62) follows immediately.

Case 2 $\delta > 0, \beta = 0$

In this case, $y^{(n-2)}(0) = 0, y^{(n-2)}(1) \neq 0$. Hence, for $t \in [0, 1]$,

$$y^{(n-2)}(t) \geq y^{(n-2)}(1) t \geq y^{(n-2)}(1) t(1-t). \quad (3.67)$$

Using a similar technique as in Case 1, it follows from (3.67) and successive integrations that

$$y(t) \geq y^{(n-2)}(1) \psi(t), t \in [0, 1]. \quad (3.68)$$

This leads to (3.65), where

$$c = \frac{y^{(n-2)}(1)}{\eta} \left[\frac{1}{4^{n-1}(n-1)!} - \frac{2}{4^n n!} \right] > 0. \quad (3.69)$$

The rest of the proof is similar to that of Case 1.

Case 3 $\delta = 0, \beta > 0$

In this case, $y^{(n-2)}(0) \neq 0, y^{(n-2)}(1) = 0$. Thus, for $t \in [0, 1]$,

$$y^{(n-2)}(t) \geq y^{(n-2)}(0) (1-t) \geq y^{(n-2)}(0) t(1-t). \quad (3.70)$$

Again, as in Case 1 it follows from (3.70) and successive integrations that

$$y(t) \geq y^{(n-2)}(0) \psi(t), t \in [0, 1]. \quad (3.71)$$

Inequality (3.71) implies (3.65), where

$$c = \frac{y^{(n-2)}(0)}{\eta} \left[\frac{1}{4^{n-1}(n-1)!} - \frac{2}{4^n n!} \right] > 0. \quad (3.72)$$

The rest of the proof is similar to that of Case 1.

Case 4 $\delta > 0, \beta > 0$

Here, $y^{(n-2)}(0) \neq 0, y^{(n-2)}(1) \neq 0$. Let

$$m = \min\{y^{(n-2)}(0), y^{(n-2)}(1)\}.$$

Then,

$$y^{(n-2)}(t) \geq m \geq mt(1-t), t \in [0, 1]. \quad (3.73)$$

Once again, it follows from (3.73) and successive integrations that

$$y(t) \geq m \psi(t), t \in [0, 1], \tag{3.74}$$

which in turn leads to inequality (3.65), where

$$c = \frac{m}{\eta} \left[\frac{1}{4^{n-1}(n-1)!} - \frac{2}{4^n n!} \right] > 0. \tag{3.75}$$

The rest of the proof is similar to that of Case 1.

This completes the proof of the theorem.

THEOREM 3.5. *Let*

$$F_B = \left\{ f \mid \frac{u}{f(u)} \text{ is bounded for } u \in [0, \infty) \right\},$$

$$F_0 = \left\{ f \mid \lim_{u \rightarrow \infty} \frac{u}{f(u)} = 0 \right\},$$

$$F_\infty = \left\{ f \mid \lim_{u \rightarrow \infty} \frac{u}{f(u)} = \infty \right\}.$$

- (a) *If* $f \in F_B$, *then* $E = (0, c)$ *or* $(0, c]$ *for some* $c \in (0, \infty)$.
- (b) *If* $f \in F_0$, *then* $E = (0, c]$ *for some* $c \in (0, \infty)$.
- (c) *If* $f \in F_\infty$, *then* $E = (0, \infty)$.

Proof. (a) This is immediate from (3.62).

(b) Since $F_0 \subseteq F_B$, it follows from case (a) that $E = (0, c)$ or $(0, c]$ for some $c \in (0, \infty)$. In particular,

$$c = \sup E. \tag{3.76}$$

Let $\{\lambda_m\}_{m=1}^\infty$ be a monotonically increasing sequence in E which converges to c , and let $\{y_m\}_{m=1}^\infty$ in C be a corresponding sequence of eigenfunctions. Further, let $\eta_m = \|y_m\|$. Then, (3.61) implies that no subsequence of $\{\eta_m\}_{m=1}^\infty$ can diverge to infinity. Thus, there exists $M > 0$ such that $\eta_m \leq M$ for all m . In view of (3.2), we find that y_m is uniformly bounded. Hence, there is a subsequence of $\{y_m\}$, relabelled as the original sequence, which converges uniformly to some $y \in C$.

Noting that $\lambda_m S y_m = y_m$, we have

$$cSy_m = \frac{c}{\lambda_m} y_m. \quad (3.77)$$

Since $\{cSy_m\}_{m=1}^{\infty}$ is relatively compact, y_m converges to y and λ_m converges to c , it follows from (3.77) that

$$cSy = y,$$

i.e., $c \in E$. This completes the proof for case (b).

(c) This follows from Corollary 3.1 and (3.61).

Example 3.1. Consider the boundary value problem

$$\begin{aligned} y^{(3)} + \lambda \left\{ \phi(t, y, y') + \frac{2t}{[t(7-t-t^3)+5]^r} \right\} (y+5)^r \\ = \lambda \phi(t, y, y') (y+5)^r, \quad t \in (0, 1), \\ y(0) = 0, \\ -2y'(0) - 7y''(0) = 0, \\ 14y'(1) + y''(1) = 0, \end{aligned}$$

where $\lambda > 0$, $0 \leq r < 1$, and $\phi(t, y, y')$ is any function of t, y and y' .

Taking $f(y) = (y+5)^r$, we find

$$\frac{Q(t, y, y')}{f(y)} = \phi(t, y, y') + \frac{2t}{[t(7-t-t^3)+5]^r}$$

and

$$\frac{P(t, y, y', y'')}{f(y)} = \phi(t, y, y').$$

Hence, we may take

$$q(t) = \phi(t, y, y') + \frac{t}{[t(7-t-t^3)+5]^r},$$

$$q_1(t) = \phi(t, y, y') + \frac{2t}{[t(7-t-t^3)+5]^r}$$

and

$$p(t) = p_1(t) = \phi(t, y, y').$$

Since $f \in F_\infty$, by Theorem 3.5(c) the set $E = (0, \infty)$. As an example when $\lambda = 12$, the boundary value problem has a positive solution given by $y(t) = t(7 - t - t^3)$.

Example 3.2. Consider the boundary value problem

$$y'' + \lambda \left\{ \phi(t, y) + \frac{1}{[t(1-t) + 5]^r} \right\} (y + 3)^r = \lambda \phi(t, y) (y + 3)^r, t \in (0, 1),$$

$$y(0) - 2y'(0) = 0,$$

$$y(1) + 2y'(1) = 0,$$

where $\lambda > 0$, $0 \leq r < 3/2$, and $\phi(t, y)$ is any function of t and y .

Taking $f(y) = (y + 3)^r$, we find

$$\frac{Q(t, y)}{f(y)} = \phi(t, y) + \frac{1}{[t(1-t) + 5]^r}$$

and

$$\frac{P(t, y, y')}{f(y)} = \phi(t, y).$$

Hence, we may choose

$$q(t) = \phi(t, y) + \frac{1}{2[t(1-t) + 5]^r} \quad q_1(t) = \phi(t, y) + \frac{1}{[t(1-t) + 5]^r}$$

and

$$p(t) = p_1(t) = \phi(t, y).$$

Case 1 $0 \leq r < 1$

Since $f \in F_\infty$, by Theorem 3.5(c) the set $E = (0, \infty)$. For example when $\lambda = 2$, the boundary value problem has a positive solution given by $y(t) = t(1 - t) + 2$.

Case 2 $r = 1$

Since $f \in F_B$, by Theorem 3.5(a) the set E is an open or half-closed interval. Further, we note from Case 1 and Theorem 3.2 that E contains the interval $(0, 2]$.

Case 3 $1 < r < \frac{3}{2}$

Since $f \in F_0$, by Theorem 3.5(b) the set E is a half-closed interval. Again, it is noted that $(0, 2] \subseteq E$.

4. SPECIAL CASE: $\lambda = 1$

THEOREM 4.1. *Suppose that (A1)–(A5) hold. Then, (1.1)–(1.4) has a positive solution.*

Proof. To obtain a positive solution of (1.1)–(1.4), we shall seek a fixed point of the operator S (defined in (3.8)) in the cone C . We have seen that S is compact on the cone C . Further, we observe from the proof of Theorem 3.1 that S maps C into itself. Also, the standard arguments yield that S is completely continuous.

Case 1 Suppose that f is superlinear. Since $f_0 = 0$, we may choose $a_1 > 0$ such that $f(u) \leq \epsilon u$ for $0 < u \leq a_1$, where $\epsilon > 0$ satisfies

$$\frac{L\epsilon}{(n - 2)!} \int_0^1 G(s, s)v(s)ds \leq 1. \tag{4.1}$$

Let $y \in C$ be such that $\|y\| = a_1(n - 2)!$. Then, from (3.2) we have $|y(t)| \leq a_1, t \in [0, 1]$. Hence, applying (3.12), Lemma 2.2, (3.2) and (4.1) successively gives for $t \in [0, 1]$,

$$\begin{aligned} (Sy)^{(n-2)}(t) &\leq \int_0^1 G(t, s)v(s)f(y(s))ds \\ &\leq L \int_0^1 G(s, s)v(s)f(y(s))ds \\ &\leq L\epsilon \int_0^1 G(s, s)v(s)y(s)ds \\ &\leq L\epsilon \int_0^1 G(s, s)v(s) \frac{1}{(n - 2)!} \|y\| ds \leq \|y\|. \end{aligned}$$

Consequently,

$$\|Sy\| \leq \|y\|. \tag{4.2}$$

If we set

$$\Omega_1 = \{y \in B \mid \|y\| < a_1(n - 2)!\},$$

then (4.2) holds for $y \in C \cap \partial\Omega_1$.

Next, since $f_\infty = \infty$, we may choose $\bar{a}_2 > 0$ such that $f(u) \geq Mu$ for $u \geq \bar{a}_2$, where $M > 0$ satisfies

$$\frac{\xi M}{4^{n-2}(n - 2)!} \int_{1/2}^{3/4} G\left(\frac{1}{2}, s\right) u(s)ds \geq 1. \tag{4.3}$$

Let

$$a_2 = \max \left\{ 2a_1(n - 2)!, \frac{4^{n-2}(n - 2)!}{\xi} \bar{a}_2 \right\},$$

and let $y \in C$ be such that $\|y\| = a_2$. Then, from (3.6) we have

$$y(t) \geq \frac{\xi}{4^{n-2}(n-2)!} \|y\| \geq \frac{\xi}{4^{n-2}(n-2)!} \cdot \frac{4^{n-2}(n-2)!}{\xi} \bar{a}_2 = \bar{a}_2, t \in \left[\frac{1}{2}, \frac{3}{4} \right].$$

Hence, $f(y(t)) \geq My(t)$ for $t \in [\frac{1}{2}, \frac{3}{4}]$. In view of (3.12), (3.6) and (4.3), we find

$$\begin{aligned} (Sy)^{(n-2)} \left(\frac{1}{2} \right) &\geq \int_0^1 G \left(\frac{1}{2}, s \right) u(s) f(y(s)) ds \\ &\geq \int_{\frac{1}{2}}^{\frac{3}{4}} G \left(\frac{1}{2}, s \right) u(s) f(y(s)) ds \\ &\geq M \int_{\frac{1}{2}}^{\frac{3}{4}} G \left(\frac{1}{2}, s \right) u(s) y(s) ds \\ &\geq M \int_{\frac{1}{2}}^{\frac{3}{4}} G \left(\frac{1}{2}, s \right) u(s) \frac{\xi}{4^{n-2}(n-2)!} \|y\| ds \geq \|y\|. \end{aligned}$$

Therefore,

$$\|Sy\| \geq \|y\|. \tag{4.4}$$

If we set

$$\Omega_2 = \{y \in B \mid \|y\| < a_2\},$$

then (4.4) holds for $y \in C \cap \partial\Omega_2$.

In view of (4.2) and (4.4), it follows from Theorem 2.1 that S has a fixed point $y \in C \cap (\bar{\Omega}_2 \setminus \Omega_1)$, such that

$$a_1(n-2)! \leq \|y\| \leq a_2.$$

This y is a positive solution of (1.1)–(1.4).

Case 2 Suppose that f is sublinear. Since $f_0 = \infty$, there exists $a_3 > 0$ such that $f(u) \geq \bar{M}u$ for $0 < u \leq a_3$, where $\bar{M} > 0$ satisfies

$$\frac{\xi \bar{M}}{4^{n-2}(n-2)!} \int_{\frac{1}{2}}^{\frac{3}{4}} G \left(\frac{1}{2}, s \right) u(s) ds \geq 1. \tag{4.5}$$

Let $y \in C$ be such that $\|y\| = a_3(n-2)!$. Then, from (3.2) we have $|y(t)| \leq a_3, t \in [0, 1]$. Hence, on using (3.12), (3.6) and (4.5) successively, we get

$$\begin{aligned}
 (Sy)^{(n-2)}\left(\frac{1}{2}\right) &\geq \int_0^1 G\left(\frac{1}{2}, s\right) u(s)f(y(s))ds \\
 &\geq \int_{\frac{1}{2}}^{3/4} G\left(\frac{1}{2}, s\right) u(s)f(y(s))ds \\
 &\geq \bar{M} \int_{\frac{1}{2}}^{3/4} G\left(\frac{1}{2}, s\right) u(s)y(s)ds \\
 &\geq \bar{M} \int_{\frac{1}{2}}^{3/4} G\left(\frac{1}{2}, s\right) u(s) \frac{\xi}{4^{n-2}(n-2)!} \|y\| ds \geq \|y\|.
 \end{aligned}$$

from which the inequality (4.4) follows immediately. If we set

$$\Omega_1 = \{y \in B \mid \|y\| < a_3(n-2)!\},$$

then (4.4) holds for $y \in C \cap \partial\Omega_1$.

Next, in view of $f_\infty = 0$, we may choose $\bar{a}_4 > 0$ such that $f(u) \leq \bar{\epsilon} u$ for $u \geq \bar{a}_4$, where $\bar{\epsilon} > 0$ satisfies

$$\frac{L\bar{\epsilon}}{(n-2)!} \int_0^1 G(s, s)v(s)ds \leq 1. \tag{4.6}$$

There are two cases to consider, namely, f is bounded and f is unbounded.

Case (i) Suppose that f is bounded, i.e., $f(u) \leq R, u \in [0, \infty)$ for some $R > 0$. Let

$$a_4 = \max \left\{ 2a_3(n-2)!, \frac{LR}{(n-2)!} \int_0^1 G(s, s)v(s)ds \right\},$$

and let $y \in C$ be such that $\|y\| = a_4(n-2)!$. For $t \in [0, 1]$, from (3.12) and Lemma 2.2, we find

$$\begin{aligned}
 (Sy)^{(n-2)}(t) &\leq \int_0^1 G(t, s)v(s)f(y(s))ds \\
 &\leq R \int_0^1 G(t, s)v(s)ds \\
 &\leq LR \int_0^1 G(s, s)v(s)ds \\
 &\leq a_4(n-2)! = \|y\|.
 \end{aligned}$$

Hence, (4.2) holds.

Case (ii) Suppose that f is unbounded, i.e., there exists

$$a_4 > \max \{2a_3(n-2)!, \bar{a}_4\},$$

such that $f(u) \leq f(a_4)$ for $0 < u \leq a_4$. Let $y \in C$ be such that $\|y\| = a_4 (n - 2)!$. Then, from (3.2) we have $|y(t)| \leq a_4, t \in [0, 1]$. Hence, applying (3.12), Lemma 2.2 and (4.6) successively gives for $t \in [0, 1]$,

$$\begin{aligned} (Sy)^{(n-2)}(t) &\leq \int_0^1 G(t, s)v(s)f(y(s))ds \\ &\leq L \int_0^1 G(s, s)v(s)f(y(s))ds \\ &\leq L \int_0^1 G(s, s)v(s)f(a_4)ds \\ &\leq L \int_0^1 G(s, s)v(s)\bar{\epsilon} a_4 ds \\ &\leq a_4(n - 2)! = \|y\| \end{aligned}$$

from which (4.2) follows immediately.

In both Cases (i) and (ii), if we set

$$\Omega_2 = \{y \in B \mid \|y\| < a_4(n - 2)!\},$$

then (4.2) holds for $y \in C \cap \partial\Omega_2$.

Now that we have obtained (4.4) and (4.2), it follows from Theorem 2.1 that S has a fixed point $y \in C \cap (\bar{\Omega}_2 \setminus \Omega_1)$, such that

$$a_3(n - 2)! \leq \|y\| \leq a_4(n - 2)!.$$

This y is a positive solution of (1.1)–(1.4).

The proof of the theorem is complete.

The following two examples illustrate Theorem 4.1.

Example 4.1. Consider the boundary value problem

$$\begin{aligned} y^{(4)} + \left\{ \phi(t, y, y', y'') + \frac{24}{[t^2(9 - t - t^2) + 1]^r} \right\} (y + 1)^r \\ = \phi(t, y, y', y'') (y + 1)^r, t \in (0, 1), \\ y(0) = y'(0) = y''(1) = 0, \\ -y''(0) - 3y^{(3)}(0) = 0, \end{aligned}$$

where $0 \leq r < \frac{3}{4}$ and $\phi(t, y, y', y'')$ is any function of t, y, y' and y'' .

Taking $f(y) = (y + 1)^r$ (which is sublinear), we find

$$\frac{Q(t, y, y', y'')}{f(y)} = \phi(t, y, y', y'') + \frac{24}{[t^2(9 - t - t^2) + 1]^r}$$

and

$$\frac{P(t, y, y', y'', y^{(3)})}{f(y)} = \phi(t, y, y', y'').$$

Hence, we may choose

$$q(t) = \phi(t, y, y', y'') + \frac{2}{[t^2(9 - t - t^2) + 1]^r}$$

$$q_1(t) = \phi(t, y, y', y'') + \frac{24}{[t^2(9 - t - t^2) + 1]^r}$$

and

$$p(t) = p_1(t) = \phi(t, y, y', y'').$$

All the conditions of Theorem 4.1 are fulfilled and therefore the boundary value problem has a positive solution. One such solution is given by $y(t) = t^2(9 - t - t^2)$.

Example 4.2. Consider the boundary value problem

$$y'' + \left\{ \phi(t, y) + \frac{6t}{(10 - t - t^3)^r} \right\} (y + 7)^r = \phi(t, y) (y + 7)^r, t \in (0, 1),$$

$$-y(0) - 3y'(0) = 0,$$

$$4y(1) + y'(1) = 0,$$

where $0 \leq r < \frac{4}{3}$, $r \neq 1$, and $\phi(t, y)$ is any function of t and y .

Taking $f(y) = (y + 7)^r$ (which is superlinear if $r > 1$, and sublinear if $r < 1$), we find

$$\frac{Q(t, y)}{f(y)} = \phi(t, y) + \frac{6t}{(10 - t - t^3)^r}$$

and

$$\frac{P(t, y, y')}{f(y)} = \phi(t, y).$$

Hence, we may take

$$q(t) = \phi(t, y) + \frac{t}{(10 - t - t^3)^r}, \quad q_1(t) = \phi(t, y) + \frac{8t}{(10 - t - t^3)^r}$$

and

$$p(t) = p_1(t) = \phi(t, y).$$

Again, all the conditions of Theorem 4.1 are satisfied and so the boundary value problem has a positive solution. Indeed, $y(t) = 3 - t - t^3$ is one such solution.

References

1. R. P. Agarwal, *Difference Equations and Inequalities*, Marcel Dekker, New York, 1992.
2. R. P. Agarwal and P. J. Y. Wong, Existence of solutions for singular boundary value problems for higher order differential equations, *Rendiconti del Seminario Matematico e Fisico di Milano*, to appear.
3. D. Aronson, M. G. Crandall and L. A. Peletier, Stabilization of solutions of a degenerate nonlinear diffusion problem, *Nonlinear Analysis* 6(1982), 1001–1022.
4. C. Bandle, C. V. Coffman and M. Marcus, Nonlinear elliptic problems in annular domains, *J. Diff. Eqns.* 69(1987), 322–345.
5. C. Bandle and M. K. Kwong, Semilinear elliptic problems in annular domains, *J. Appl. Math. Phys.* 40(1989), 245–257.
6. N. P. Cac, A. M. Fink and J. A. Gatica, Nonnegative solutions of quasilinear elliptic boundary value problems with nonnegative coefficients, preprint.
7. Y. S. Choi and G. S. Ludford, An unexpected stability result of the near-extinction diffusion flame for non-unity Lewis numbers, *Q. J. Mech. Appl. Math.* 42, part 1(1989), 143–158.
8. C. J. Chyan and J. Henderson, Positive solutions for singular higher order nonlinear equations, *Diff. Eqns. Dyn. Sys.* 2(1994), 153–160.
9. D. S. Cohen, Multiple stable solutions of nonlinear boundary value problems arising in chemical reactor theory, *SIAM J. Appl. Math.* 20(1971), 1–13.
10. E. N. Dancer, On the structure of solutions of an equation in catalysis theory when a parameter is large, *J. Diff. Eqns.* 37(1980), 404–437.
11. P. W. Eloe and J. Henderson, Singular nonlinear boundary value problems for higher order ordinary differential equations, *Nonlinear Analysis* 17(1991), 1–10.
12. P. W. Eloe and J. Henderson, Positive solutions for higher order ordinary differential equations, preprint.
13. P. W. Eloe, J. Henderson and P. J. Y. Wong, Positive solutions for two-point boundary value problems, In *Dynamic Systems and Applications*, Vol 2, eds. G. S. Ladde and M. Sambandham, to appear.
14. L. H. Erbe and H. Wang, On the existence of positive solutions of ordinary differential equations, *Proc. Amer. Math. Soc.* 120(1994), 743–748.
15. A. M. Fink, *The radial Laplacian Gel'fand problem*, *Delay and Differential Equations* (Ames, IA, 1991), 93–98, World Sci. Publishing, River Edge, NJ, 1992.
16. A. M. Fink, J. A. Gatica and G. E. Hernandez, Eigenvalues of generalized Gel'fand models, *Nonlinear Analysis* 20(1993), 1453–1468.
17. A. M. Fink and J. A. Gatica, Positive solutions of systems of second order boundary value problems, *J. Math. Anal. Appl.*, to appear.
18. H. Fujita, On the nonlinear equations $\Delta u + e^u = 0$ and $\partial v / \partial t = \Delta v + e^v$, *Bull. Amer. Math. Soc.* 75(1969), 132–135.
19. X. Garaizar, Existence of positive radial solutions for semilinear elliptic problems in the annulus, *J. Diff. Eqns.* 70(1987), 69–72.

20. J. A. Gatica, V. Olikar and P. Waltman, Singular nonlinear boundary value problems for second-order ordinary differential equations, *J. Diff. Eqs.* 79(1989), 62–78.
21. I. M. Gel'fand, Some problems in the theory of quasilinear equations, *Uspehi Mat. Nauka* 14(1959), 87–158; English translation, *Trans. Amer. Math. Soc.* 29(1963), 295–381.
22. J. Henderson, Singular boundary value problems for difference equations, *Dynamic Systems and Applications* 1(1992), 271–282.
23. J. Henderson, Singular boundary value problems for higher order difference equations, In *Proceedings of the First World Congress on Nonlinear Analysts 1992*, ed. V. Lakshmikantham, Walter de Gruyter and Co., 1139–1150.
24. M. A. Krasnosel'skii, *Positive Solutions of Operator Equations*, Noordhoff, Groningen, 1964.
25. S. Parter, Solutions of differential equations arising in chemical reactor processes, *SIAM J. Appl. Math.* 26(1974), 687–716.
26. H. Wang, On the existence of positive solutions for semilinear elliptic equations in the annulus, *J. Diff. Eqns.* 109(1994), 1–7.
27. P. J. Y. Wong and R. P. Agarwal, On the existence of solutions of singular boundary value problems for higher order difference equations, *Nonlinear Analysis*, to appear.
28. P. J. Y. Wong and R. P. Agarwal, On the existence of positive solutions of higher order difference equations, *Topological Methods in Nonlinear Analysis* to appear.
29. P. J. Y. Wong and R. P. Agarwal, On the eigenvalues of boundary value problems for higher order difference equations, *Rocky Mountain J. Math.* to appear.

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie