

On Absolute Stability of Nonlinear Systems with Small Delays*

M.I. GIL^{a,†}, A. AILON^{b,‡} and B.H. AHN^b

^a*Department of Mathematics, Ben Gurion University of the Negev, P.O. Box 653, Beer-Sheva 84105, Israel;* ^b*Department of Mechatronics, Kwangju Institute of Science and Technology, Kwangju 506-303, Korea*

(Received 17 February 1998; Revised 31 March 1998)

Nonlinear nonautonomous retarded systems with separated autonomous linear parts and continuous nonlinear ones are considered. It is assumed that deviations of the argument are sufficiently small. Absolute stability conditions are derived. They are formulated in terms of eigenvalues of auxiliary matrices.

Keywords: Nonlinear differential–difference systems; Absolute stability

AMS (MOS) Subject Classification: 34 K 20

1 INTRODUCTION AND STATEMENT OF THE RESULT

Stability of nonlinear differential–difference systems has been discussed by many authors (see [1–3], and references given therein). The basic method for the stability analysis is the Lyapunov functionals one. By this method many very strong results are obtained. But finding Lyapunov's functionals is usually difficult. In [4] (see also [2, p. 111]) an explicit absolute stability criterion for a class of retarded systems was derived. It is formulated in the terms of the eigenvalues of the characteristic matrix-valued functions. That result was further developed in [5].

* This research was supported by the Israel Ministry of Science and Technology.

[†] Corresponding author. Tel.: 07-6461608/9. E-mail: gilmi@black.bgu.ac.il.

[‡] A. Ailon is on leave from the Departments of Electrical and Computer Engineering, Ben Gurion University of the Negev, Israel.

The present paper is devoted to a class of nonlinear nonautonomous differential–difference systems having sufficiently small deviations of the argument. We will obtain absolute exponential stability conditions in terms of eigenvalues of auxiliary matrices. These conditions allow us to avoid the analysis of quasipolynomials. There have been many stability conditions for nonlinear systems with sufficiently small delays in recent years usually in the category of “robust stability” for time-delay systems (e.g. [6,7], and references therein). Besides, only systems with constant discrete delays are considered. Below we will consider systems with arbitrary (time-dependent and discrete, in general) delays.

Let \mathbf{C}^n be a Euclidean space with the Euclidean norm $\|\cdot\|_{\mathbf{C}^n}$. Moreover, $C([a, b], \mathbf{C}^n)$ is the space of continuous functions defined on the real segment $[a, b]$ with values in \mathbf{C}^n and equipped with the norm

$$\|v\|_{C([a,b],\mathbf{C}^n)} := \sup_{a \leq t \leq b} \|v(t)\|_{\mathbf{C}^n} \quad (v \in C([a, b], \mathbf{C}^n)).$$

In addition, let $L^2([a, b], \mathbf{C}^n)$ be the space of n -vector-valued functions defined on $[a, b]$ with values in \mathbf{C}^n and equipped with the norm

$$\|v\|_{L^2([a,b],\mathbf{C}^n)} := \left[\int_a^b \|v(t)\|_{\mathbf{C}^n}^2 dt \right]^{1/2} \quad (v \in L^2([a, b], \mathbf{C}^n)).$$

Let $\eta < \infty$ be a positive constant. As usually, for all $t \geq 0$ and $x \in C([0, \infty), \mathbf{C}^n)$, x_t is defined by the relation

$$x_t(\theta) \equiv x(\theta + t), \quad -\eta \leq \theta \leq 0,$$

cf. [1, p. 36], [2, p. 19]. That is, for any $x \in C([0, \infty), \mathbf{C}^n)$, x_t is some function from $C([-\eta, \infty), \mathbf{C}^n)$.

Consider in \mathbf{C}^n the equation

$$\dot{x}(t) = \sum_{k=0}^m A_k x(t - h_k) + F(t, x_t) \quad (t \geq 0, \dot{x} = dx/dt), \quad (1.1)$$

where A_k ($k = 0, \dots, m$) are constant matrices,

$$0 = h_0 < h_1 < \dots < h_m = \eta$$

are numbers, and F continuously maps $[0, \infty) \times C([-\eta, 0], \mathbf{C}^n)$ into \mathbf{C}^n .

It is assumed that for every $u \in L^2([-\eta, \infty), \mathbf{C}^n) \cap C([-\eta, \infty), \mathbf{C}^n)$, the inequality

$$\|F(\cdot, u_t)\|_{L^2([0, \infty), \mathbf{C}^n)} \equiv \left[\int_0^\infty \|F(t, u_t(t))\|_{\mathbf{C}^n}^2 dt \right]^{1/2} \leq q \|u\|_{L^2([-\eta, \infty), \mathbf{C}^n)} \tag{1.2}$$

is fulfilled. For instance, let there exist constants $b_j > 0$ and $q_j \geq 0$ ($j = 1, \dots, m < \infty$), such that the relation

$$\|F(t, u_t)\|_{\mathbf{C}^n} \leq \sum_{k=1}^m q_k \|u(t - h_k(t))\|_{\mathbf{C}^n} \quad \text{for all } u \in L^2([-\eta, \infty), \mathbf{C}^n) \tag{1.3}$$

holds, where $h_j(t)$ ($j = 1, \dots, m$) are differentiable scalar-valued functions with the properties

$$1 - \dot{h}_j(t) \geq b_j > 0 \quad \text{and} \quad 0 \leq h_j(t) \leq \eta \quad (t \geq 0, j = 1, \dots, m). \tag{1.4}$$

Then by Proposition 8.1.2 of [8], condition (1.2) holds with

$$q = \sum_{j=1}^m q_j b_j^{-1/2}. \tag{1.5}$$

Take the initial condition

$$x(t) = \Phi(t) \quad \text{for } -\eta \leq t \leq 0, \tag{1.6}$$

where $\Phi(t)$ is a given continuous vector-valued function defined on $[-\eta, 0]$. A solution of (1.1) is an absolutely continuous vector-valued function $x : [-\eta, \infty) \rightarrow \mathbf{C}^n$ which satisfies that equation over $R_+ := [0, \infty)$ almost everywhere with a given initial condition (1.6). The solution existence is assumed.

DEFINITION 1.1 We will say that the zero solution of (1.1) is absolutely exponentially stable in the class of nonlinearities (1.2) if there exist positive constants N and ϵ independent of the specific form of the function F (but dependent on q) such that for any solution $x(t)$

of (1.1) with an initial function Φ , the inequality

$$\|x(t)\|_{C^n} \leq N \exp[-\epsilon t] \|\Phi\|_{C([- \eta, 0], C^n)} \quad (t \geq 0)$$

is fulfilled.

Let A be an $n \times n$ -matrix, and let $\lambda_k(A)$ ($k = 1, \dots, n$) denote the eigenvalues of A including with their multiplicities. The following quantity plays an essential role hereafter:

$$g(A) = (N^2(A) - \sum_{k=1}^n |\lambda_k(A)|^2)^{1/2},$$

where $N(A)$ is the Hilbert–Schmidt (Frobenius) norm of A , i.e. $N^2(A) = \text{Trace}(AA^*)$. The following properties of $g(A)$ are proved in [8, Sections 1.1 and 1.3]: $g^2(A) \leq N^2(A) - |\text{Trace } A^2|$,

$$\begin{aligned} g(A) &\leq \sqrt{1/2} N(A^* - A) \quad \text{and} \quad g(Ae^{i\tau} + zI) \\ &= g(A) \quad \text{for every } \tau \in \mathbf{R}, z \in \mathbf{C}. \end{aligned} \quad (1.7)$$

Here and below, I is the unit matrix. In [9, p. 185], $g(A)$ is called the deviation from normality of A , since $g(A) = 0$ if A is a normal matrix, i.e., if $AA^* = A^*A$. Further, set

$$\tilde{A} = \sum_{k=0}^m A_k \quad \text{and} \quad V_2 = \sum_{k=1}^m h_k \|A_k\|_{C^n}.$$

Everywhere below it is assumed that \tilde{A} is a Hurwitz matrix. Denote

$$\alpha(\tilde{A}) = \max_k \text{Re} \lambda_k(\tilde{A}), \quad \omega(\tilde{A}) = \max_k |\text{Im} \lambda_k(\tilde{A})|,$$

and

$$\Gamma(\tilde{A}) = \sum_{k=1}^{n-1} \frac{g^k(\tilde{A})}{(k!)^{1/2} |\alpha(\tilde{A})|^{k+1}}.$$

The aim of this paper is to prove the following.

THEOREM 1.2 *Let the condition*

$$[(\alpha^2(\tilde{A}) + \omega^2(\tilde{A}))^{1/2}V_2 + q]\Gamma(\tilde{A}) < 1 \tag{1.8}$$

hold. Then the zero solution of Eq. (1.1) is absolutely exponentially stable in the class of nonlinearities (1.2).

Remark 1.3 Since $V_2 = h_1\|A_1\|_{\mathbf{C}^n} + \dots + h_m\|A_m\|_{\mathbf{C}^n}$, if $q\Gamma(\tilde{A}) < 1$, then sufficiently small delays provide the stability.

If \tilde{A} is a normal (in particular, Hermitian) matrix. Then $g(\tilde{A}) = 0$, $\Gamma(\tilde{A}) = |\alpha(\tilde{A})|^{-1}$, and condition (1.8) takes the form

$$[(\alpha^2(\tilde{A}) + \omega^2(\tilde{A}))^{1/2}V_2 + q]|\alpha(\tilde{A})|^{-1} < 1 \tag{1.9}$$

Remark 1.4 Theorem 1.2 is exact, and condition (1.8) cannot be improved without additional restrictions.

In fact, let us consider the system

$$\dot{x}(t) = Ax(t) + F(t, x(t)), \tag{1.10}$$

where A is a constant Hurwitz matrix, and F continuously maps $[0, \infty) \times \mathbf{C}^n$ into \mathbf{C}^n with the property

$$\|F(t, h)\|_{\mathbf{C}^n} \leq q\|h\|_{\mathbf{C}^n} \quad (h \in \mathbf{C}^n; t \geq 0). \tag{1.11}$$

Under consideration $V_2 = 0$, and (1.8) takes the form $q\Gamma(A) < 1$. If A is normal, then the inequality

$$q|\alpha(A)|^{-1} < 1 \tag{1.12}$$

gives the absolute stability conditions of the zero solution of Eq. (1.10) in the class of nonlinearities (1.11). Take $F(t, h) = qh$ ($h \in \mathbf{C}$). Then for the asymptotic stability of Eq. (1.10) it is necessary and sufficient that $\alpha(A) + q < 0$. But this condition coincides with (1.12).

2 PROOF OF THEOREM 1.2

For brevity put $\|\cdot\|_{L^2([0, \infty), \mathbf{C}^n)} = \|\cdot\|_L$, and

$$\|y(i\omega)\|_{\tilde{L}} \equiv \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} \|y(i\omega)\|_{\mathbf{C}^n}^2 d\omega \right]^{1/2} \quad (y \in L^2(\mathbf{R}, \mathbf{C}^n)).$$

Denote

$$c_0(\tilde{A}) = \sup_{\omega \in \mathbf{R}} \| (i\omega I - \tilde{A})^{-1} \|_{\mathbf{C}^n} \quad \text{and} \quad c_1(\tilde{A}) = \sup_{\omega \in \mathbf{R}} \| \omega(i\omega I - \tilde{A})^{-1} \|_{\mathbf{C}^n}.$$

Let us consider the equation

$$\dot{x}(t) = \sum_{k=0}^m A_k x(t - h_k) + f(t) \quad (t \geq 0). \quad (2.1)$$

LEMMA 2.1 *Under the condition $c_1(\tilde{A})V_2 < 1$, let $f \in L^2([0, \infty), \mathbf{C}^n)$. Then the bound*

$$\|x\|_L \leq [1 - c_1(\tilde{A})V_2]^{-1} c_0(\tilde{A}) \|f\|_L + m_1 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \quad (2.2)$$

is valid with a constant m_1 independent of Φ for any solution $x(t)$ of (2.1) with a continuous initial function Φ .

Proof Rewrite (2.1) in the form

$$\dot{x}(t) = \tilde{A}x(t) + \sum_{k=1}^m A_k (x(t - h_k) - x(t)) + f(t) \quad (t \geq 0). \quad (2.3)$$

The Laplace transform of $x(t - h_k)$ is $\tilde{x}(\lambda)e^{-h_k\lambda} + v_k(\lambda)$ where $\tilde{x}(\lambda)$ is the Laplace transform of $x(t)$, and

$$v_k(\lambda) = \int_0^{h_k} e^{-t\lambda} x(t - h_k) dt = \int_0^{h_k} e^{-t\lambda} \Phi(t - h_k) dt.$$

Thus the application of the Laplace transformation to Eq. (2.3) gives

$$\begin{aligned} \lambda \tilde{x}(\lambda) - x(0) &= \tilde{A} \tilde{x}(\lambda) + \sum_{k=1}^m A_k [\tilde{x}(\lambda)(e^{-h_k\lambda} - 1) \\ &\quad + v_k(\lambda)] + \tilde{F}(\lambda) \quad (\lambda \in \mathbf{C}), \end{aligned}$$

where $\tilde{F}(\lambda)$ is the Laplace transform of the function $f(t)$. Hence,

$$\tilde{x}(\lambda) = (\lambda I - \tilde{A})^{-1} \left[\sum_{k=1}^m A_k (e^{-h_k\lambda} - 1)x(\lambda) + \tilde{F}(\lambda) \right] + e(\lambda) \quad (\lambda \in \mathbf{C}), \quad (2.4)$$

where

$$e(\lambda) = (\lambda I - \tilde{A})^{-1} \left(x(0) + \sum_{k=1}^m A_k v_k(\lambda) \right).$$

Since $|1 - e^{-hi\omega}| \leq h|\omega|$ ($\omega \in \mathbf{R}, h > 0$), we have

$$\begin{aligned} & \|(\tilde{A} - i\omega I)^{-1} \sum_{k=1}^m A_k (1 - e^{-hki\omega}) \tilde{x}(i\omega)\|_{\tilde{L}} \\ & \leq \|\omega(\tilde{A} - i\omega I)^{-1}\|_{C^n} \sum_{k=1}^m h_k \|A_k\|_{C^n} \|\tilde{x}(i\omega)\|_{\tilde{L}} \\ & \leq c_1(\tilde{A}) V_2 \|\tilde{x}(i\omega)\|_{\tilde{L}} \quad (\omega \in \mathbf{R}). \end{aligned} \tag{2.5}$$

Due to (2.4) and (2.5), it can be written

$$\|\tilde{x}(i\omega)\|_{\tilde{L}} \leq V_2 c_1(\tilde{A}) \|\tilde{x}(i\omega)\|_{\tilde{L}} + c_0(\tilde{A}) \|\tilde{F}\|_{\tilde{L}} + \|e(i\omega)\|_{\tilde{L}}.$$

Hence, taking into account that $c_1(\tilde{A}) V_2 < 1$, we get

$$\|\tilde{x}(i\omega)\|_{\tilde{L}} \leq (1 - V_2 c_1(\tilde{A}))^{-1} (c_0(\tilde{A}) \|\tilde{F}\|_{\tilde{L}} + \|e(i\omega)\|_{\tilde{L}}). \tag{2.6}$$

The simple calculations show that

$$\|v_k(i\omega)\|_{C^n} \leq h_k \|\Phi\|_{C([- \eta, 0], C^n)} \quad (\omega \in \mathbf{R})$$

and

$$\begin{aligned} \|e(i\omega)\|_{\tilde{L}} & \leq \|(i\omega I - \tilde{A})^{-1}\|_{\tilde{L}} \left(\|x(0)\|_{C^n} + \sum_{k=1}^m h_k \|A_k\|_{C^n} \sup_{\omega \in \mathbf{R}} \|v_k(i\omega)\|_{C^n} \right) \\ & = \|(i\omega I - \tilde{A})^{-1}\|_{\tilde{L}} \left(\|\Phi(0)\|_{C^n} + V_2 \|\Phi\|_{C([- \eta, 0], C^n)} \right). \end{aligned}$$

Hence,

$$\|e(i\omega)\|_{\tilde{L}} \leq \|(i\omega I - \tilde{A})^{-1}\|_{\tilde{L}} (1 + V_2) \|\Phi\|_{C([- \eta, 0], C^n)}.$$

Relation (2.6) yields

$$\|\tilde{x}(i\omega)\|_{\tilde{L}} \leq (1 - V_2 c_1(\tilde{A}))^{-1} c_0(\tilde{A}) \|\tilde{F}\|_{\tilde{L}} + m_1 \|\Phi\|_{C([- \eta, 0], C^n)}.$$

Now due to the Parseval equality, we arrive at the required result.

LEMMA 2.2 Under condition (1.2), let the inequality

$$c_1(\tilde{A})V_2 + c_0(\tilde{A})q < 1 \quad (2.7)$$

be fulfilled. Then for any solution $x(t)$ of (1.1) with a continuous initial function $\Phi(t)$, the bound

$$\|x\|_L \leq m_0 [1 - (c_1(\tilde{A})V_2 + qc_0(\tilde{A}))]^{-1} \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)}$$

is valid with a constant m_0 .

Proof Rewrite (1.1) in the form (2.1) with $f(t) = F(t, x(t))$, where x is the solution of (1.1). Combining the previous lemma with condition (1.2), we easily get

$$\begin{aligned} \|x\|_L &\leq [1 - c_1(\tilde{A})V_2]^{-1} c_0(\tilde{A}) \|f\|_L + m_1 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \\ &\leq [1 - c_1(\tilde{A})V_2]^{-1} c_0(\tilde{A})q (\|x\|_L + \|\Phi\|_{L^2([- \eta, 0], \mathbf{C}^n)}) \\ &\quad + m_1 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)}. \end{aligned}$$

Consequently,

$$\|x\|_L \leq [1 - c_1(\tilde{A})V_2]^{-1} c_0(\tilde{A})q \|x\|_L + m_2 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)}$$

with a constant m_2 . Condition (2.7) implies

$$[1 - c_1(\tilde{A})V_2]^{-1} c_0(\tilde{A})q < 1.$$

Thus,

$$\begin{aligned} \|x\|_L &\leq (1 - c_0(\tilde{A})q [1 - c_1(\tilde{A})V_2]^{-1}) m_2 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \\ &= (1 - c_0(\tilde{A})q - c_1(\tilde{A})V_2)^{-1} m_0 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)}. \end{aligned}$$

As claimed.

LEMMA 2.3 The relations $c_0(\tilde{A}) \leq \Gamma(\tilde{A})$ and

$$c_1(\tilde{A}) \leq \Gamma(\tilde{A}) [\alpha^2(\tilde{A}) + \omega^2(\tilde{A})]^{1/2}$$

are true.

Proof By Corollary 1.2.4 of [8], for any $n \times n$ -matrix A , we have the inequality

$$\|(A - I\lambda)^{-1}\|_{C^n} \leq \sum_{k=0}^{n-1} \frac{g^k(A)}{\sqrt{k!}\rho^{k+1}(A, \lambda)} \quad \text{for all regular } \lambda, \quad (2.8)$$

where $\rho(A, \lambda)$ is the distance between the spectrum of A and a complex point λ . Since \tilde{A} is a Hurwitz matrix,

$$\rho(\tilde{A}, i\omega) \geq |\alpha(\tilde{A})| \quad (\omega \in \mathbf{R}), \quad (2.9)$$

and therefore $c_0(\tilde{A}) \leq \Gamma(\tilde{A})$.

Further, we have

$$|\lambda_k(\tilde{A}) - i\omega|^2 \geq \alpha^2(\tilde{A}) + (\omega_k - \omega)^2$$

for any eigenvalue $\lambda_k(\tilde{A})$ of \tilde{A} with $\text{Im}\lambda_k(\tilde{A}) = \omega_k$. Hence simple calculations show that

$$\max_{\omega \in \mathbf{R}} \omega^2 |\lambda_k(\tilde{A}) - i\omega|^{-2} \leq \frac{\alpha^2(\tilde{A}) + \omega_k^2}{\alpha^2(\tilde{A})}.$$

Thus

$$\omega^2 \rho^{-2}(\tilde{A}, \omega) \leq \frac{\alpha^2(\tilde{A}) + \omega^2(\tilde{A})}{\alpha^2(\tilde{A})} \quad (\omega \in \mathbf{R}).$$

Therefore according to (2.8) and (2.9),

$$\begin{aligned} \|\omega(\tilde{A} - i\omega I)^{-1}\| &\leq \sum_{k=0}^{n-1} \frac{|\omega|g^k(\tilde{A})}{\sqrt{k!}\rho^{k+1}(\tilde{A}, \omega)} \\ &\leq (\alpha^2(\tilde{A}) + \omega^2(\tilde{A}))^{1/2} \sum_{k=0}^{n-1} \frac{g^k(\tilde{A})}{\sqrt{k!}|\alpha(\tilde{A})|^{k+1}} \\ &= (\alpha^2(\tilde{A}) + \omega^2(\tilde{A}))^{1/2} \Gamma(\tilde{A}) \quad (\omega \in \mathbf{R}). \end{aligned}$$

As claimed.

Lemmas 2.2 and 2.3 yield

COROLLARY 2.4 *Let conditions (1.2) and (1.8) hold. Then for any solution $x(t)$ of (1.1) with an initial function $\Phi(t)$, the bound*

$$\|x\|_L \leq m_0 [1 - \Gamma(\tilde{A})(V_2(\alpha^2(\tilde{A}) + \omega^2(\tilde{A}))^{1/2} + q)]^{-1} \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)}$$

($m_0 = \text{const.}$)

is valid.

Proof of Theorem 1.2 From (1.1) it follows the inequality

$$\begin{aligned} \|\dot{x}\|_L &\leq \sum_{k=0}^m \|A_k\|_{\mathbf{C}^n} \|x(t - h_k)\|_L + \|F\|_L \\ &\leq V_1 (\|x\|_L + \|x\|_{L^2([- \eta, 0], \mathbf{C}^n)}) + \|F\|_L, \end{aligned}$$

where

$$V_1 = \sum_{k=0}^m \|A_k\|_{\mathbf{C}^n}.$$

Due to (1.2) and (1.6) this gives

$$\|\dot{x}\|_L \leq (q + V_1)(\|x\|_L + \|\Phi\|_{L^2([- \eta, 0], \mathbf{C}^n)}).$$

Now Corollary 2.4 implies the relation

$$\|\dot{x}\|_L \leq N_1 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \quad (N_1 = \text{const.}).$$

By the trivial Lemma 8.4.5 of [8], we get

$$\|\dot{x}\|_{C([0, \infty), \mathbf{C}^n)} \leq (2\|x\|_L \|\dot{x}\|_L)^{1/2} \leq N_2 \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \quad (N_2 = \text{const.}).$$

(2.10)

Further, substitute in (1.1) the relation

$$x(t) = x_\epsilon(t) e^{-\epsilon t} \quad (2.11)$$

with a small enough $\epsilon > 0$. Applying our reasoning above, we can assert according to (2.10) that

$$\|x_\epsilon\|_{C([- \eta, \infty), \mathbf{C}^n)} \leq N_\epsilon \|\Phi\|_{C([- \eta, 0], \mathbf{C}^n)} \quad (N_\epsilon = \text{const.}).$$

Hence, (2.11) yields the absolute exponential stability.

3 EXAMPLE

Let us consider in \mathbf{C}^n the equation

$$\dot{x}(t) = A_0 x(t) + A_1 x(t - \eta) + f(t, x(t)), \int_0^\eta T(t, s)x(t - s) ds, \quad (3.1)$$

where the function $f: [0, \infty) \times \mathbf{C}^{2n} \rightarrow \mathbf{C}^n$ has the property

$$\|f(t, h, w)\|_{\mathbf{C}^n} \leq q_0 \|h\|_{\mathbf{C}^n} + q_1 \|w\|_{\mathbf{C}^n} \quad (h, w \in \mathbf{C}^n; t \geq 0), \quad (3.2)$$

and the matrix-valued function $T(t, s)$ defined on $[0, \infty) \times [0, \eta]$ satisfies the inequality

$$m_T \equiv \sup_{t \geq 0} \int_0^\eta \|T(t, s)\|_{\mathbf{C}^n} ds < \infty. \quad (3.3)$$

In the considered case

$$F(t, x_t) = f(t, x(t), \int_0^\eta T(t, s)x(t - s) ds).$$

Taking into account, that

$$\begin{aligned} & \int_0^\infty \left\| \int_0^\eta T(t, s)x(t - s) ds \right\|_{\mathbf{C}^n}^2 dt \\ & \leq \int_0^\infty \left(\int_0^\eta \|T(t, s)\|_{\mathbf{C}^n} \|x(t - s)\|_{\mathbf{C}^n} ds \right)^2 dt \\ & \leq \sup_{t \geq 0} \left(\int_0^\eta \|T(t, s)\|_{\mathbf{C}^n} ds \right)^2 \int_{-\eta}^\infty \|x(t)\|_{\mathbf{C}^n}^2 dt = m_T^2 \|x\|_{L^2([- \eta, 0], \mathbf{C}^n)}^2, \end{aligned}$$

we easily get the inequality

$$\left[\int_0^\infty \left\| f(t, u(t), \int_0^\eta T(t, s)u(t-s) ds) \right\|_{\mathbf{C}^n}^2 dt \right]^{1/2} \leq q_0 \|u\|_{L^2([0, \infty); \mathbf{C}^n)} + q_1 m_T \|u\|_{L^2([- \eta, \infty), \mathbf{C}^n)}$$

for any $u \in L^2([- \eta, \infty), \mathbf{C}^n) \cap C([- \eta, \infty), \mathbf{C}^n)$. Hence it follows that (1.2) holds with

$$F(t, u_t) = f\left(t, u(t), \int_0^\eta T(t, s)u(t-s) ds\right) \text{ and } q = q_0 + q_1 m_T. \tag{3.4}$$

Clearly, under consideration

$$\tilde{A} = A_0 + A_1, \quad V_2 = \eta \|A_1\|_{\mathbf{C}^n}. \tag{3.5}$$

Now Theorem 1.2 yields the following result. Let \tilde{A} , q and V_2 be defined by (3.4) and (3.5). Let condition (1.8) hold. Then the zero solution of Eq. (3.1) is absolutely exponentially stable in the class of nonlinearities satisfying conditions (3.2) and (3.3).

4 CONCLUDING REMARKS

We have derived the sufficient absolute stability conditions for Eq. (1.1) in the class of nonlinearities (1.2) in terms of the eigenvalues of the matrix \tilde{A} . These conditions are exact according to Remark 1.4. As it was mentioned, the stability conditions suggested in [6,7] can be applied to systems with constant discrete delays only, while Theorem 1.2 allows us to investigate systems containing time-variant and distributed delays (see relations (1.3) and (3.4) above). Consider now the equation

$$\dot{x} = A_0 x(t) + F(x(t-h)), \tag{4.1}$$

where A_0 is a Hurwitz matrix, $F: \mathbf{C}^n \rightarrow \mathbf{C}^n$ satisfies the inequality

$$\|F(z)\|_{\mathbf{C}^n} \leq L \|z\|_{\mathbf{C}^n} \quad (L = \text{const.}, z \in \mathbf{C}^n).$$

In the paper [10] among other very interesting results the following stability condition for Eq. (4.1) is established:

$$\Lambda(A_0) + L < 0, \quad (4.2)$$

where

$$\Lambda(A_0) = \lim_{h \rightarrow 0^+} (\|I + hA_0\| - 1)/h$$

with some norm $\|\cdot\|$ in \mathbf{C}^n . But if A_0 is not dissipative with respect to that norm, condition (4.2) cannot be applied. At that same time, as the example shows, Theorem 1.2 gives the stability conditions for (4.1) if L is sufficiently small.

References

- [1] J.K. Hale and S.M. Verduyn Lunel, *Introduction to Functional Differential Equations*. Springer-Verlag, Berlin, 1993.
- [2] V. Kolmanovskii and A. Myshkis, *Applied Theory of Functional Differential Equations*, Kluwer Academic Publishers, London, 1992.
- [3] V. Kolmanovskii and V. Nosov, *Stability of Functional Differential Equations*, Academic Press, New York, 1986.
- [4] M.I. Gil', On absolute stability of differential-delay systems, *IEEE Trans. Automatic Control* **39**(12) (1994), 2481–2484.
- [5] M.I. Gil', On absolute stability of neutral type functional differential systems, *Non-linear Studies* **3**(1) (1996), 59–72.
- [6] K.K. Shyu and J.J. Yan, Robust stability of uncertain time-delay systems, and its stabilization by variable structure control. *Int. Journal of Control* **57**(1) (1993), 237–246.
- [7] S.S. Wang, B.S. Chen and T.P. Lin, Robust stability of uncertain time-delay systems. *Int. Journal of Control* **46**(3) (1987), 963–976.
- [8] M.I. Gil', *Norm Estimations for Operator-valued Functions and Applications*. Marcel Dekker Inc., New York, 1995.
- [9] L. Collatz, *Functional Analysis and Numerical Mathematics*. Academic Press, New York, London, 1966.
- [10] V. Kolmanovskii, Applications of differential inequalities for stability of some functional differential equations. *Nonlinear Analysis, TMA* **25**(9–10) (1995), 1017–1028.

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be