


Journal of Inequalities in Pure and Applied Mathematics

L'HOSPITAL TYPE RULES FOR MONOTONICITY, WITH APPLICATIONS

IOSIF PINELIS

Department of Mathematical Sciences,
Michigan Technological University,
Houghton, MI 49931, USA
EMail: ipinelis@mtu.edu

©2000 Victoria University
ISSN (electronic): 1443-5756
010-01



volume 3, issue 1, article 5,
2002.

*Received 29 January, 2001;
accepted 11 August, 2001.*

Communicated by: F. Qi

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

Let f and g be differentiable functions on an interval (a, b) , and let the derivative g' be positive on (a, b) . The main result of the paper implies that, if $f(a+) = g(a+) = 0$ and $\frac{f'}{g'}$ is increasing on (a, b) , then $\frac{f}{g}$ is increasing on (a, b) .

2000 Mathematics Subject Classification: Primary: 26A48, 26D10; Secondary: 26D07, 60E15, 62B10, 94A17

Key words: L'Hospital's Rule, Monotonicity, Information inequalities

Contents

1	L'Hospital Type Rule for Monotonicity	3
2	Applications to Information Inequalities	7
	References	

L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

1. L'Hospital Type Rule for Monotonicity

Let $-\infty \leq a < b \leq \infty$. Let f and g be differentiable functions on the interval (a, b) . Assume also that the derivative g' is nonzero and does not change sign on (a, b) ; in other words, either $g' > 0$ everywhere on (a, b) or $g' < 0$ on (a, b) . The following statement reminds one of the l'Hospital rule for computing limits and turns out to be useful in a number of contexts.

Proposition 1.1. *Suppose that $f(a+) = g(a+) = 0$ or $f(b-) = g(b-) = 0$. (Then g is nonzero and does not change sign on (a, b) , since g' is so.)*


1. If $\frac{f'}{g'}$ is increasing on (a, b) , then $\left(\frac{f}{g}\right)' > 0$ on (a, b) .
2. If $\frac{f'}{g'}$ is decreasing on (a, b) , then $\left(\frac{f}{g}\right)' < 0$ on (a, b) .

Proof. Assume first that $f(a+) = g(a+) = 0$. Assume also that $\frac{f'}{g'}$ is increasing on (a, b) , as in part 1 of the proposition. Fix any $x \in (a, b)$ and consider the function

$$h_x(y) := f'(x)g(y) - g'(x)f(y), \quad y \in (a, b).$$

This function is differentiable and hence continuous on (a, b) . Moreover, for all $y \in (a, x)$,

$$\frac{d}{dy}h_x(y) = f'(x)g'(y) - g'(x)f'(y) = g'(x)g'(y) \left(\frac{f'(x)}{g'(x)} - \frac{f'(y)}{g'(y)} \right) > 0,$$



L'Hospital Type Rules for
Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 3 of 12

because g' is nonzero and does not change sign on (a, b) and $\frac{f'}{g'}$ is increasing on (a, b) . Hence, the function h_x is increasing on (a, x) ; moreover, being continuous, h_x is increasing on $(a, x]$. Note also that $h_x(a+) = 0$. It follows that $h_x(x) > 0$, and so,

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - g'(x)f(x)}{g(x)^2} = \frac{h_x(x)}{g(x)^2} > 0.$$


This proves part 1 of the proposition under the assumption that $f(a+) = g(a+) = 0$. The proof under the assumption that $f(b-) = g(b-) = 0$ is similar; alternatively, one may replace here $f(x)$ and $g(x)$ for all $x \in (a, b)$ by $f(a + b - x)$ and $g(a + b - x)$, respectively. Thus, part 1 is proved.

Part 2 follows from part 1: replace f by $-f$. □

Remark 1.1. *Instead of the requirement that f and g be differentiable on (a, b) , it would be enough to assume, for instance, only that f and g are continuous and both have finite right derivatives f'_+ and g'_+ (or finite left derivatives f'_- and g'_-) on (a, b) and then use $\frac{f'_+}{g'_+}$ (or, respectively, $\frac{f'_-}{g'_-}$) in place of $\frac{f'}{g'}$. In such a case, one would need to use the fact that, if a function h is continuous on (a, b) and $h'_+ > 0$ on (a, b) or $h'_- > 0$ on (a, b) , then h is increasing on (a, b) ; cf. e.g. Theorem 3.4.4 in [1].*

The following corollary is immediate from Proposition 1.1.

Corollary 1.2. *Suppose that $f(a+) = g(a+) = 0$ or $f(b-) = g(b-) = 0$.*



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit


Page 4 of 12

1. If $\frac{f'}{g'}$ is increasing on (a, b) , then $\frac{f}{g}$ is increasing on (a, b) .
2. If $\frac{f'}{g'}$ is decreasing on (a, b) , then $\frac{f}{g}$ is decreasing on (a, b) .

Remark 1.2. *The related result that any family of probability distributions with a monotone likelihood ratio is stochastically monotone is well known in statistics; see e.g. [2] for this and many other similar statements. For the case when f and g are probability tail functions, a proof of Corollary 1.2 may be found in [3]. In fact, essentially the same proof remains valid for the general setting, at least when the double integrals below exist and possess the usual properties; we are reproducing that proof now, for the readers' convenience: if $f(a+) = g(a+) = 0$, $\frac{f'}{g'}$ is increasing on (a, b) , g' does not change sign on (a, b) , and $a < x < y < b$, then*

$$\begin{aligned}
 (1.1) \quad f(x) \cdot (g(y) - g(x)) &= \iint_{\substack{u \in (a, x) \\ v \in (x, y)}} f'(u)g'(v) \, du \, dv \\
 &< \iint_{\substack{u \in (a, x) \\ v \in (x, y)}} g'(u)f'(v) \, du \, dv \\
 &= g(x) \cdot (f(y) - f(x)),
 \end{aligned}$$

whence $f(x)g(y) < g(x)f(y)$, and so, $\frac{f(x)}{g(x)} < \frac{f(y)}{g(y)}$; inequality (1.1) takes



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit


Page 5 of 12

place because $u < v$ implies $\frac{f'(u)}{g'(u)} < \frac{f'(v)}{g'(v)}$, and so, $f'(u)g'(v) < g'(u)f'(v)$.

The proof in the case when one has $f(b-) = g(b-) = 0$ instead of $f(a+) = g(a+) = 0$ is quite similar.

Ideas similar to the ones discussed above were also present, albeit implicitly, in [5].

Remark 1.3. Corollary 1.2 will hold if the terms “increasing” and “decreasing” are replaced everywhere by “non-decreasing” and “non-increasing”, respectively.



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

2. Applications to Information Inequalities

In this section, applications of the above l'Hospital type rule to information inequalities are given. Other applications, as well as extensions and refinements of this rule, will be given in a series of papers following this one: in [7], extensions to non-monotonic ratios of functions, with applications to certain probability inequalities arising in bioequivalence studies and to problems of convexity; in [6], applications to monotonicity of the relative error of a Padé approximation for the complementary error function; in [8], applications to probability inequalities for sums of bounded random variables.

With all these applications, apparently we have only “scratched the surface”. Yet, even the diversity of the cited results suggests that the monotonicity counterparts of the l'Hospital Rule may have as wide a range of application as the l'Hospital Rule itself.

Consider now the entropy function

$$H(p, q) := -p \ln p - q \ln q,$$


where $q := 1 - p$ and $p \in (0, 1)$. In effect, it is a function of one variable, say p .

Topsøe [9] proved the inequalities

$$(2.1) \quad \ln p \cdot \ln q \leq H(p, q) \leq \frac{\ln p \cdot \ln q}{\ln 2}$$

and

$$(2.2) \quad \ln 2 \cdot 4pq \leq H(p, q) \leq \ln 2 \cdot (4pq)^{1/\ln 4}$$



L'Hospital Type Rules for
Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12

for all $p \in (0, 1)$ and also showed that these bounds on the entropy are exact; namely, they are attained when $p \downarrow 0$ or $p = 1/2$. Topsøe also indicated promising applications of bounds (2.2) in statistics. He noticed that the bounds in (2.1) and (2.2), as well as their exactness, would naturally be obtained from the monotonicity properties stated below, using also the symmetry of the entropy function: $H(p, q) = H(q, p)$.

Conjecture 2.1. [9] *The ratio*

$$r(p) := \frac{\ln p \ln q}{H(p, q)}$$

is decreasing in $p \in (0, 1/2]$, from $r(0+) = 1$ to $r(1/2) = \ln 2$.

Conjecture 2.2. [9] *The ratio*


$$R(p) := \frac{\ln \left(\frac{H(p, q)}{\ln 2} \right)}{\ln(4pq)}$$

is decreasing on $(0, 1/2)$, from $R(0+) = 1$ to $R\left(\frac{1}{2}-\right) = \frac{1}{\ln 4}$.

We shall now prove these conjectures, based on Proposition 1.1 of the previous section.

Proof of Conjecture 2.1. On $(0, 1)$,

$$r = \frac{f}{g},$$



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12

where $f(p) := \ln p \ln q$ and $g(p) := H(p, q)$. Consider, for $p \in (0, 1)$,

$$r_1(p) := \frac{f'(p)}{g'(p)} = \frac{\frac{1}{p} \ln q - \frac{1}{q} \ln p}{\ln q - \ln p}; \quad r_2(p) := \frac{f''(p)}{g''(p)} = \frac{f_2(p)}{g_2(p)},$$

where

$$f_2(p) := -(pq)^2 f''(p) = p^2 \ln p + q^2 \ln q + 2pq$$

and $g_2(p) := -(pq)^2 g''(p) = pq;$

$$r_3(p) := \frac{f'_2(p)}{g'_2(p)} = \frac{2p \ln p - 2q \ln q + q - p}{q - p}; \quad r_4(p) := \frac{f''_2(p)}{g''_2(p)} = -1 - \ln pq.$$

Now we apply Proposition 1.1 repeatedly, four times. First, note that r_4 is decreasing on $(0, 1/2)$ and $f'_2(1/2) = g'_2(1/2) = 0$; hence, r_3 is decreasing on $(0, 1/2)$. This and $f_2(0+) = g_2(0+) = 0$ imply that r_2 is decreasing on $(0, 1/2)$. This and $f'(1/2) = g'(1/2) = 0$ imply that r_1 is decreasing on $(0, 1/2)$. Finally, this and $f(0+) = g(0+) = 0$ imply that r is decreasing on $(0, 1/2)$. \square

Proof of Conjecture 2.2. On $(0, 1/2)$,

$$R = \frac{F}{G},$$

where $F(p) := \ln \left(\frac{H(p, q)}{\ln 2} \right)$ and $G(p) := \ln(4pq)$. Next,

$$(2.3) \quad \frac{F'}{G'} = \frac{F_1}{G_1},$$



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit


Page 9 of 12

where $F_1(p) := \ln q - \ln p$ and $G_1(p) := \left(\frac{1}{p} - \frac{1}{q}\right) H(p, q)$. Further, $\frac{F'_1}{G'_1} = \frac{1}{2 - r_2}$, where r_2 is the same as in the proof of Conjecture 2.1, and r_2 is decreasing on $(0, 1/2)$, as was shown. In addition, $r_2 < 2$ on $(0, 1)$. Hence, $\frac{F'_1}{G'_1} = \frac{1}{2 - r_2}$ is decreasing on $(0, 1/2)$. Also, $F_1(1/2) = G_1(1/2) = 0$. Now

Proposition 1.1 implies that $\frac{F_1}{G_1}$ is decreasing on $(0, 1/2)$; hence, by (2.3), $\frac{F'}{G'}$ is decreasing on $(0, 1/2)$. It remains to notice that $F(1/2) = G(1/2) = 0$ and use once again Proposition 1.1. \square

It might seem surprising that these proofs uncover a connection between the two seemingly unrelated conjectures – via the ratio r_2 .

Concerning other proofs of Conjecture 2.1, see the final version of [9]. Concerning another conjecture by Topsøe [9], related to Conjecture 2.2, see [7].



L'Hospital Type Rules for Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 10 of 12

References

- [1] R. KANNAN AND C.K. KRUEGER, *Advanced Analysis on the Real Line*, Springer, New York, 1996.
- [2] J. KEILSON AND U. SUMITA, Uniform stochastic ordering and related inequalities, *Canad. J. Statist.*, **10** (1982), 181–198.
- [3] I. PINELIS, Extremal probabilistic problems and Hotelling's T^2 test under symmetry condition, *Preprint* (1991).
- [4] I. PINELIS, Extremal probabilistic problems and Hotelling's T^2 test under a symmetry condition, *Ann. Stat.*, **22** (1994), 357–368.
- [5] I. PINELIS, On the Yao-Iyer inequality in bioequivalence studies. *Math. Inequal. Appl.* (2001), 161–162.
- [6] I. PINELIS, Monotonicity Properties of the Relative Error of a Padé Approximation for Mills' Ratio, *J. Ineq. Pure & Appl. Math.*, **3**(2) (2002), Article 20. (http://jipam.vu.edu.au/v3n2/012_01.html).
- [7] I. PINELIS, L'Hospital type rules for oscillation, with applications, *J. Ineq. Pure & Appl. Math.*, **2**(3) (2001), Article 33. (http://jipam.vu.edu.au/v2n3/011_01.html).
- [8] I. PINELIS, L'Hospital type rules for monotonicity: an application to probability inequalities for sums of bounded random variables, *J. Ineq. Pure & Appl. Math.*, **3**(1) (2002), Article 7. (http://jipam.vu.edu.au/v3n1/013_01.html).



L'Hospital Type Rules for
Monotonicity, with Applications

Iosif Pinelis

Title Page

Contents




Go Back

Close

Quit

Page 11 of 12

- [9] F. TOPSØE, Bounds for entropy and divergence for distributions over a two-element set, *J. Ineq. Pure & Appl. Math.*, **2**(2) (2001), Article 25. (http://jipam.vu.edu.au/v2n2/044_00.html).



**L'Hospital Type Rules for
Monotonicity, with Applications**

Iosif Pinelis

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12