

COEFFICIENT INEQUALITIES FOR CERTAIN CLASSES OF RUSCHEWEYH TYPE ANALYTIC FUNCTIONS

S. LATHA

Department of Mathematics and Computer Science
Maharaja's College
University of Mysore
Mysore - 570 005, INDIA.
EMail: drlatha@gmail.com

Received: 18 January, 2007

Accepted: 5 May, 2008

Communicated by: **S.S. Dragomir**

2000 AMS Sub. Class.: 30C45.

Key words: Convolution, Ruscheweyh derivative, Uniformly starlike and Uniformly convex.

Abstract: A class of univalent functions which provides an interesting transition from starlike functions to convex functions is defined by making use of the Ruscheweyh derivative. Some coefficient inequalities for functions in these classes are discussed which generalize the coefficient inequalities considered by Owa, Polatoğlu and Yavuz.

Coefficient Inequalities

S. Latha

vol. 9, iss. 2, art. 52, 2008

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1 Introduction	3
2 Main Results	5



Coefficient Inequalities

S. Latha

vol. 9, iss. 2, art. 52, 2008

[Title Page](#)

[Contents](#)



Page **2** of 10


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

[Title Page](#)[Contents](#)

Page 3 of 10

[Go Back](#)[Full Screen](#)[Close](#)

1. Introduction

Let \mathcal{N} denote the class of functions of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disc $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$.

We designate $\mathcal{V}(\beta, b, \delta)$ as the subclass of \mathcal{N} consisting of functions f obeying the condition

$$(1.2) \quad \Re \left\{ 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1} f(z)}{D^{\delta} f(z)} \right\} > \beta$$

where, $b \neq 0$, $\delta > -1$, $0 \leq \beta < 1$ and $D^{\delta} f$ is the Rushceweyh derivative of f [5] given by,

$$(1.3) \quad D^{\delta} f(z) = \frac{z}{(1-z)^{1+\delta}} * f(z) = z + \sum_{n=2}^{\infty} a_n B_n(\delta) z^n,$$

where $*$ stands for the convolution or Hadamard product of two power series and

$$B_n(\delta) = \frac{(\delta+1)(\delta+2) \cdots (\delta+n-1)}{(n-1)!}.$$

This class is obtained by putting $k = 2$ and $\lambda = 0$ in the class $\mathcal{V}_k^{\lambda}(\beta, b, \delta)$ introduced by Latha and Nanjunda Rao [2]. The class $\mathcal{V}_k^{\lambda}(\beta, b, \delta)$ is of special interest for it contains many well known as well as new classes of analytic univalent functions studied in literature. It provides a transition from starlike functions to convex functions. More specifically, $\mathcal{V}_2^0(\beta, 2, 0)$ is the family of starlike functions of order β and $\mathcal{V}_2^0(\beta, 1, 1)$ is the class of convex functions of order β . Shams, Kulkarni

and Jahangiri [6] introduced the subclass $\mathcal{SD}(\alpha, \beta)$ of \mathcal{N} consisting of functions f satisfying

$$(1.4) \quad \Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \alpha \left| \frac{zf'(z)}{f(z)} - 1 \right| + \beta$$

for some $\alpha \geq 0$, $0 \leq \beta < 1$ and $z \in \mathcal{U}$.

The class $\mathcal{KD}(\alpha, \beta)$, another subclass of \mathcal{N} , is defined as the set of all functions f obeying

$$(1.5) \quad \Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha \left| \frac{zf''(z)}{f'(z)} - 1 \right| + \beta$$

for some $\alpha \geq 0$, $0 \leq \beta < 1$ and $z \in \mathcal{U}$.

We introduce the class $\mathcal{VD}(\alpha, \beta, b, \delta)$ as the subclass of \mathcal{N} consisting of functions f which satisfy

$$\Re \left\{ 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} \right\} > \alpha \left| \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} - 1 \right| + \beta$$

where, $b \neq 0$, $\alpha \geq 0$, and $0 \leq \beta < 1$.

For the parametric values $b = 2$, $\delta = 0$ and $b = \delta = 1$ we obtain the classes $\mathcal{SD}(\alpha, \beta)$ and $\mathcal{KD}(\alpha, \beta)$ respectively.



[Title Page](#)

[Contents](#)




Page 4 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

[Title Page](#)[Contents](#)

Page 5 of 10

[Go Back](#)[Full Screen](#)[Close](#)

2. Main Results

We prove some coefficient inequalities for functions in the class $\mathcal{VD}(\alpha, \beta, b, \delta)$.

Theorem 2.1. *If $f(z) \in \mathcal{VD}(\alpha, \beta, b, \delta)$ with $0 \leq \alpha \leq \beta$, or, $\alpha > \frac{1+\beta}{2}$, then $f(z) \in \mathcal{V}\left(\frac{\beta-\alpha}{1-\alpha}, b, \delta\right)$.*

Proof. Since $\Re\{\omega\} \leq |\omega|$ for any complex number ω , $f(z) \in \mathcal{VD}(\alpha, \beta, b, \delta)$ implies that

$$(2.1) \quad \Re \left\{ 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} \right\} > \alpha \left| \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} - \frac{2}{b} \right| + \beta.$$

Equivalently,

$$(2.2) \quad \Re \left\{ 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} \right\} > \frac{\beta - \alpha}{1 - \alpha}, \quad (z \in \mathcal{U}).$$

If $0 \leq \alpha \leq \beta$, we have, $0 \leq \frac{\beta-\alpha}{1-\alpha} < 1$, and if $\alpha > \frac{1+\beta}{2}$, then we have $-1 < \frac{\alpha-\beta}{\alpha-1} \leq 0$. \square

Corollary 2.2. *For the parametric values $b = 2$ and $\delta = 0$, we get Theorem 2.1 in [3] which reads as:*


If $f(z) \in \mathcal{SD}(\alpha, \beta)$ with $0 \leq \alpha \leq \beta$, or, $\alpha > \frac{1+\beta}{2}$, then $f(z) \in \mathcal{S}^\left(\frac{\beta-\alpha}{1-\alpha}\right)$.*

Corollary 2.3. *The parametric values $b = \delta = 1$, yield the Corollary 2.2 in [3] stated as:*

If $f(z) \in \mathcal{KD}(\alpha, \beta)$ with $0 \leq \alpha \leq \beta$, or, $\alpha > \frac{1+\beta}{2}$, then $f(z) \in \mathcal{K}\left(\frac{\beta-\alpha}{1-\alpha}\right)$.

Theorem 2.4. *If $f(z) \in \mathcal{VD}(\alpha, \beta, b, \delta)$ then,*

$$(2.3) \quad |a_2| \leq \frac{b(1-\beta)}{|1-\alpha|}$$



[Title Page](#)

[Contents](#)



Page 6 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

and

$$(2.4) \quad |a_n| \leq \frac{b(1-\beta)(\delta+1)}{(n-1)|1-\alpha|B_n(\delta)} \prod_{j=1}^{n-2} \left(1 + \frac{b(\delta+1)(1-\beta)}{j|1-\alpha|} \right), \quad (n \geq 3).$$

Proof. We note that for $f(z) \in \mathcal{VD}(\alpha, \beta, b, \delta)$,

$$\Re \left\{ 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} \right\} > \frac{\beta - \alpha}{1 - \alpha}, \quad (z \in \mathcal{U}).$$

We define the function $p(z)$ by

$$(2.5) \quad p(z) = \frac{(1-\alpha) \left[1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} \right] - (\beta - \alpha)}{(1-\beta)}, \quad (z \in \mathcal{U}).$$

Then, $p(z)$ is analytic in \mathcal{U} with $p(0) = 1$ and $\Re\{p(z)\} > 0$ and $z \in \mathcal{U}$.

Let $p(z) = 1 + p_1z + p_1z^2 + \dots$. We have

$$(2.6) \quad 1 - \frac{2}{b} + \frac{2}{b} \frac{D^{\delta+1}f(z)}{D^\delta f(z)} = 1 + \left(\frac{1-\beta}{1-\alpha} \right) \sum_{n=1}^{\infty} p_n z^n.$$

That is,

$$2(D^{\delta+1}f(z) - D^\delta f(z)) = bD^\delta f(z) \left(\frac{1-\beta}{1-\alpha} \sum_{n=1}^{\infty} p_n z^n \right).$$

which implies that

$$\begin{aligned} & \frac{2B_n(\delta)(n-1)a_n}{(\delta+1)} \\ &= \frac{b(1-\beta)}{(1-\alpha)} [p_{n-1} + B_2(\delta) + a_2p_{n-2} + B_3(\delta)a_3p_{n-3} + \dots + B_{n-1}(\delta)a_{n-1}p_1]. \end{aligned}$$

[Title Page](#)[Contents](#)

Page 7 of 10

[Go Back](#)[Full Screen](#)[Close](#)

Applying the coefficient estimates $|p_n| \leq 2$ for Carathéodory functions [1], we obtain,

$$(2.7) \quad |a_n| \leq \frac{b(1-\beta)(\delta+1)}{|1-\alpha|(n-1)B_n(\delta)} \times [1 + B_2(\delta)|a_2| + B_3(\delta)|a_3| + \cdots + B_{n-1}(\delta)|a_{n-1}|].$$

For $n = 2$, $|a_2| \leq \frac{b(1-\beta)}{|1-\alpha|}$, which proves (2.3).

For $n = 3$,

$$|a_3| \leq \frac{b(1-\beta)(\delta+1)}{2|1-\alpha|B_3(\delta)} \left[1 + \frac{b(1-\beta)(\delta+1)}{|1-\alpha|} \right].$$


Therefore (2.4) holds for $n = 3$.

Assume that (2.4) is true for $n = k$.

Consider,

$$\begin{aligned} |a_{k+1}| &\leq \frac{b(1-\beta)(\delta+1)}{kB_{k+1}(\delta)} \left\{ \left(1 + \frac{b(1-\beta)(\delta+1)}{|1-\alpha|} \right) \right. \\ &\quad + \frac{b(1-\beta)(\delta+1)}{|1-\alpha|B_2(\delta)} \left(1 + \frac{b(1-\beta)(\delta+1)}{|1-\alpha|} \right) \\ &\quad \left. + \cdots + \frac{b(1-\beta)(\delta+1)}{(k-1)!|1-\alpha|B_k(\delta)} \prod_{j=1}^{k-2} \left(1 + \frac{b(1-\beta)(\delta+1)}{j(|1-\alpha|)} \right) \right\} \\ &= \frac{b(1-\beta)(\delta+1)}{kB_{k+1}(\delta)} \prod_{j=1}^{k-1} \left[1 + \frac{b(1-\beta)(\delta+1)}{j(|1-\alpha|)} \right]. \end{aligned}$$

Therefore, the result is true for $n = k + 1$. Using mathematical induction, (2.4) holds true for all $n \geq 3$. \square



[Title Page](#)

[Contents](#)



Page 8 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

Corollary 2.5. *The parametric values $b = 2$ and $\delta = 0$ yield Theorem 2.3 in [3] which states that:*

If $f(z) \in \mathcal{SD}(\alpha, \beta)$, then

$$(2.8) \quad |a_2| \leq \frac{2(1-\beta)}{|1-\alpha|}$$

and

$$(2.9) \quad |a_n| \leq \frac{2(1-\beta)}{(n-1)|1-\alpha|} \prod_{j=1}^{n-2} \left(1 + \frac{2(1-\beta)}{j|1-\alpha|} \right), \quad (n \geq 3).$$

Corollary 2.6. *Putting $\alpha = 0$ in Corollary 2.5, we get*

$$(2.10) \quad |a_n| \leq \frac{\prod_{j=1}^n (j-2\beta)}{(n-1)!}, \quad (n \geq 2),$$

a result by Robertson [4].

Corollary 2.7. *For the parametric values $b = \delta = 1$ we obtain Corollary 2.5 in [3] given by:*

If $f(z) \in \mathcal{KD}(\alpha, \beta)$ then,

$$(2.11) \quad |a_2| \leq \frac{(1-\beta)}{|1-\alpha|}$$

and

$$(2.12) \quad |a_n| \leq \frac{2(1-\beta)}{n(n-1)|1-\alpha|} \prod_{j=1}^{n-2} \left(1 + \frac{2(1-\beta)}{j|1-\alpha|} \right), \quad (n \geq 3).$$

Corollary 2.8. Letting $\alpha = 0$ in Corollary 2.7, we get the inequality by Robertson [4] given by:

$$(2.13) \quad |a_n| \leq \frac{\prod_{j=1}^n (j - 2\beta)}{n!}, \quad (n \geq 2).$$



Coefficient Inequalities

S. Latha

vol. 9, iss. 2, art. 52, 2008

Title Page

Contents



Page 9 of 10

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] C. CARATHÉODORY, Über den variabilitätsbereich der Fourier'schen konstanten von positiven harmonischen funktionen, *Rend. Circ. Palermo.*, **32** (1911), 193–217.
- [2] S. LATHA AND S. NANJUNDA RAO, Convex combinations of n analytic functions in generalized Ruscheweyh class, *Int. J. Math, Educ. Sci. Technology.*, **25**(6) (1994), 791–795.
- [3] S. OWA, Y. POLATOĞLU AND E. YAVUZ, Coefficient inequalities fo classes of uniformly starlike and convex functions, *J. Ineq. in Pure and Appl. Math.*, **7**(5) (2006), Art. 160. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=778>].
- [4] M.S. ROBERTSON, On the theory of univalent functions, *Ann. Math.*, **37** (1936), 374–408.
- [5] S. RUSCHEWEYH, A new criteria for univalent function, *Proc. Amer. Math. Soc.*, **49**(1) (1975), 109–115.
- [6] S. SHAMS, S.R. KULKARNI AND J. M. JAHANGIRI, Classes of uniformly starlike convx functions, *Internat. J. Math. and Math. Sci.*, **55** (2004), 2959–2961.



Coefficient Inequalities

S. Latha

vol. 9, iss. 2, art. 52, 2008

Title Page

Contents



Page 10 of 10

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756