

Journal of Inequalities in Pure and Applied Mathematics

ON CERTAIN NEW INTEGRAL INEQUALITIES AND THEIR APPLICATIONS

S.S. DRAGOMIR AND YOUNG-HO KIM

School of Communications and Informatics
Victoria University of Technology
PO Box 14428, Melbourne City MC
8001 Victoria, AUSTRALIA.

*E*Mail: sever.dragomir@vu.edu.au

*U*RL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>


Department of Applied Mathematics
Changwon National University
Changwon 641-773, Korea.

*E*Mail: yhkim@sarim.changwon.ac.kr

©2000 School of Communications and Informatics, Victoria University of Technology

ISSN (electronic): 1443-5756

034-02



volume 3, issue 4, article 65,
2002.

Received 8 April, 2001;
accepted 15 June, 2002.

Communicated by: D. Bainov

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

The aim of the present paper is to establish some variants integral inequalities in two independent variables. These integral inequalities given here can be applied as tools in the boundedness and uniqueness of certain partial differential equations.

2000 Mathematics Subject Classification: 26D10, 26D15.

Key words: Integral inequality, Two independent variables, Partial differential equations, Nondecreasing, Nonincreasing.

Contents

1	Introduction	3
2	Main Results	4
3	Some Applications	14
	References	

On Certain New Integral Inequalities and their Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 2 of 17


1. Introduction

The integral inequalities involving functions of one and more than one independent variables which provide explicit bounds on unknown functions play a fundamental role in the development of the theory of differential equations (see [1]–[11]). In recent year, Pachpatte [11] discovered some new integral inequalities involving functions of two independent variables. These inequalities are applied to study the boundedness and uniqueness of the solutions of following terminal value problem for the hyperbolic partial differential equation (1.1) with the condition (1.2).

$$(1.1) \quad u_{xy}(x, y) = h(x, y, u(x, y)) + r(x, y),$$

$$(1.2) \quad u(x, \infty) = \sigma_{\infty}(x), u(\infty, y) = \tau_{\infty}(y), u(\infty, \infty) = d.$$

Our main objective here, motivated by Pachpatte's inequalities [11], is to establish additional new integral inequalities involving functions of two independent variables which can be used in the analysis of certain classes of partial differential equations.



**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 3 of 17

2. Main Results

Throughout the paper, all the functions which appear in the inequalities are assumed to be real-valued and all the integrals are involved in existence on the domains of their definitions. We shall introduce some notation, \mathbb{R} denotes the set of real numbers and $\mathbb{R}_+ = [0, \infty)$ is the given subset of \mathbb{R} . The first order partial derivatives of a functions $z(x, y)$ defined for $x, y \in \mathbb{R}$ with respect to x and y are denoted by $z_x(x, y)$ and $z_y(x, y)$ respectively.

We need the inequalities in the following Lemma 2.1 and Lemma 2.2, which are given in the book [1].

Lemma 2.1. *Let g be a monotone continuous function in an interval I , containing a point u_0 , which vanishes in I . Let u and k be continuous functions in an interval $J = [\alpha, \beta]$ such that $u(J) \subset I$, and suppose that k is of fixed sign in J . Let $a \in I$.*

(i) *Assume that g is nondecreasing and k is nonnegative. If*


$$u(t) \leq a + \int_{\alpha}^t k(s)g(u(s)) ds, \quad t \in J,$$

then

$$u(t) \leq G^{-1} \left(G(a) + \int_{\alpha}^t k(s)ds \right), \quad \alpha \leq t \leq \beta_1,$$

where $G(u) = \int_{u_0}^u dx/g(x)$, $u \in I$, and $\beta_1 = \min(v_1, v_2)$, with

$$v_1 = \sup \left\{ v \in J : a + \int_{\alpha}^t k(s)g(u(s)) ds \in I, \alpha \leq t \leq v \right\},$$



On Certain New Integral
Inequalities and their
Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 4 of 17

$$v_2 = \sup \left\{ v \in J : G(a) + \int_{\alpha}^t k(s) ds \in G(I), \alpha \leq t \leq v \right\}.$$

(ii) Assume that $J = (\alpha, \beta]$. If

$$u(t) \leq a + \int_t^{\beta} k(s)g(u(s)) ds, \quad t \in J,$$

then

$$u(t) \leq G^{-1} \left(G(a) + \int_t^{\beta} k(s) ds \right), \quad \alpha_1 < t \leq \beta,$$

where $\alpha_1 = \max(\mu_1, \mu_2)$, with

$$\mu_1 = \sup \left\{ \mu_1 \in J : a + \int_t^{\beta} k(s)g(u(s)) ds \in I, \mu \leq t \leq \beta \right\},$$


$$\mu_2 = \sup \left\{ \mu \in J : G(a) + \int_t^{\beta} k(s) ds \in G(I), \mu \leq t \leq \beta \right\}.$$

The proofs of the inequalities in (i), (ii) can be completed as in [1, p. 40–42]. Here we omit the details.

Let $u(x, y)$, $a(x, y)$, $b(x, y)$ be nonnegative continuous functions defined for $x, y \in \mathbb{R}_+$.

Lemma 2.2. (i) Assume that $a(x, y)$ is nondecreasing in x and nonincreasing in y for $x, y \in \mathbb{R}_+$. If

$$u(x, y) \leq a(x, y) + \int_0^x \int_y^{\infty} b(s, t)u(s, t) dt ds$$



On Certain New Integral
Inequalities and their
Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents




Go Back

Close

Quit

Page 5 of 17



Title Page

Contents



Go Back

Close

Quit

Page 6 of 17

for all $x, y \in \mathbb{R}_+$, then

$$u(x, y) \leq a(x, y) \exp \left(\int_0^x \int_y^\infty b(s, t) dt ds \right).$$

(ii) Assume that $a(x, y)$ is nonincreasing in each of the variables $x, y \in \mathbb{R}_+$.

If

$$u(x, y) \leq a(x, y) + \int_x^\infty \int_y^\infty b(s, t) u(s, t) dt ds$$

for all $x, y \in \mathbb{R}_+$, then

$$u(x, y) \leq a(x, y) \exp \left(\int_x^\infty \int_y^\infty b(s, t) dt ds \right).$$

The proofs of the inequalities in (i), (ii) can be completed as in [1, p. 109-110]. Here we omit the details.


To establish our results, we require the class of functions S as defined in [2]. A function $g : [0, \infty) \rightarrow [0, \infty)$ is said to belong to the class S if

(i) $g(u)$ is positive, nondecreasing and continuous for $u \geq 0$,

(ii) $(1/v)g(u) \leq g(u/v)$, $u > 0, v \geq 1$.

Theorem 2.3. Let $u(x, y), a(x, y), b(x, y), c(x, y), d(x, y)$ be nonnegative continuous functions defined for $x, y \in \mathbb{R}_+$, let $g \in S$. Define a function $z(x, y)$ by

$$z(x, y) = a(x, y) + c(x, y) \int_0^x \int_y^\infty d(s, t) u(s, t) dt ds$$



with $z(x, y)$ is nondecreasing in x and $z(x, y) \geq 1$ for $x, y \in \mathbb{R}_+$. If

$$(2.1) \quad u(x, y) \leq z(x, y) + \int_{\alpha}^x b(s, y)g(u(s, y)) ds,$$

for $\alpha, x, y \in \mathbb{R}_+$ and $\alpha \leq x$, then

$$(2.2) \quad u(x, y) \leq p(x, y) \left[a(x, y) + c(x, y)e(x, y) \right. \\ \left. \times \exp \left(\int_0^x \int_y^{\infty} d(s, t)p(s, t)c(s, t) dt ds \right) \right],$$

for $x, y \in \mathbb{R}_+$, where

$$(2.3) \quad p(x, y) = G^{-1} \left(G(1) + \int_{\alpha}^x b(s, y) ds \right),$$

$$(2.4) \quad e(x, y) = \int_0^x \int_y^{\infty} d(s, t)p(s, t)a(s, t) dt ds,$$

$$(2.5) \quad G(u) = \int_{u_0}^u \frac{ds}{g(s)}, \quad u \geq u_0 > 0,$$

G^{-1} is the inverse function of G , and

$$G(1) + \int_{\alpha}^x b(s, y) ds \in \text{Dom} (G^{-1}).$$

Proof. Let $z(x, y)$ is a nonnegative, continuous, nondecreasing and let $g \in S$. Then (2.1) can be restated as

$$(2.6) \quad \frac{u(x, y)}{z(x, y)} \leq 1 + \int_{\alpha}^x b(s, y) \frac{1}{z(x, y)} g(u(s, y)) ds.$$

**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents




Go Back

Close

Quit

Page 7 of 17



Title Page

Contents



Go Back

Close

Quit

Page 8 of 17

Define a function $w(x, y)$ by the right side of (2.6), then $[u(x, y)/z(x, y)] \leq w(x, y)$ and

$$(2.7) \quad w(x, y) \leq 1 + \int_{\alpha}^x b(s, y)g(w(s, y)) ds.$$

Treating $y, y \in \mathbb{R}_+$ fixed in (2.7) and using (i) of Lemma 2.1 to (2.7), we get

$$(2.8) \quad w(x, y) \leq G^{-1} \left(G(1) + \int_{\alpha}^x b(s, y) ds \right).$$

Using (2.8) in $[u(x, y)/z(x, y)] \leq w(x, y)$, we obtain

$$u(x, y) \leq z(x, y)p(x, y),$$

where $p(x, y)$ is defined by (2.3). From the definition of $z(x, y)$ we have

$$(2.9) \quad u(x, y) \leq p(x, y) (a(x, y) + c(x, y)v(x, y)),$$

where $v(x, y)$ is defined by

$$v(x, y) = \int_0^x \int_y^{\infty} d(s, t)u(s, t) dt ds.$$

From (2.9) we get

$$\begin{aligned} v(x, y) &\leq \int_0^x \int_y^{\infty} d(s, t)p(s, t) (a(s, t) + c(s, t)v(s, t)) dt ds \\ &= e(x, y) + \int_0^x \int_y^{\infty} d(s, t)p(s, t)c(s, t)v(s, t) dt ds, \end{aligned}$$

where $e(x, y)$ is defined by (2.4). Clearly, $e(x, y)$ is nonnegative, continuous, nondecreasing in $x, x \in \mathbb{R}_+$ and nonincreasing in $y, y \in \mathbb{R}_+$. Now, by (i) of Lemma 2.2, we obtain

$$(2.10) \quad v(x, y) \leq e(x, y) \exp \left(\int_0^x \int_y^\infty d(s, t) p(s, t) c(s, t) dt ds \right).$$

Using (2.10) in (2.9) we get the required inequality in (2.2). □

Theorem 2.4. *Let $u(x, y), a(x, y), b(x, y), c(x, y), d(x, y)$ be nonnegative continuous functions defined for $x, y \in \mathbb{R}_+$ and let $g \in S$. Define a function $z(x, y)$ by*

$$z(x, y) = a(x, y) + c(x, y) \int_x^\infty \int_y^\infty d(s, t) u(s, t) dt ds$$

with $z(x, y)$ is nonincreasing in x and $z(x, y) \geq 1$ for $x, y \in \mathbb{R}_+$. If

$$u(x, y) \leq z(x, y) + \int_x^\beta b(s, y) g(u(s, y)) ds$$

for $\beta, x, y \in \mathbb{R}_+$ and $\beta \geq x$, then

$$u(x, y) \leq \bar{p}(x, y) \left[a(x, y) + c(x, y) \bar{e}(x, y) \times \exp \left(\int_x^\infty \int_y^\infty d(s, t) \bar{p}(s, t) c(s, t) dt ds \right) \right]$$



**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents




Go Back

Close

Quit

Page 9 of 17



**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 10 of 17

for $x, y \in \mathbb{R}_+$, where

$$(2.11) \quad \bar{p}(x, y) = G^{-1} \left(G(1) + \int_x^\beta b(s, y) ds \right),$$

$$\bar{e}(x, y) = \int_x^\infty \int_y^\infty d(s, t) \bar{p}(s, t) a(s, t) dt ds.$$

G is defined in (2.5), G^{-1} is the inverse function of G , and

$$G(1) + \int_x^\beta b(s, y) ds \in \text{Dom} (G^{-1}).$$

The details of the proof of Theorem 2.4 follows by an argument similar to that in the proofs of Theorem 2.3 with suitable changes. We omit the details.

Theorem 2.5. Let $u(x, y), a(x, y), b(x, y), c(x, y)$ be nonnegative continuous functions defined for $x, y \in \mathbb{R}_+$ and $F : \mathbb{R}_+^3 \rightarrow \mathbb{R}_+$ be a continuous function which satisfies the condition


$$(2.12) \quad 0 \leq F(x, y, u) - F(x, y, v) \leq K(x, y, v)(u - v)$$

for $u \geq v \geq 0$, where $K(x, y, v)$ is a nonnegative continuous function defined for $x, y, v \in \mathbb{R}_+$. And let $g \in S$. Define a function $z(x, y)$ by

$$z(x, y) = a(x, y) + c(x, y) \int_0^x \int_y^\infty F(s, t, u(s, t)) dt ds$$

with nondecreasing in x and $z(x, y) \geq 1$ for $x, y \in \mathbb{R}_+$. If

$$(2.13) \quad u(x, y) \leq z(x, y) + \int_\alpha^x b(s, y) g(u(s, y)) ds$$



On Certain New Integral Inequalities and their Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 11 of 17

for $\alpha, x, y \in \mathbb{R}_+$ and $\alpha \leq x$, then

$$(2.14) \quad u(x, y) \leq p(x, y) \left[a(x, y) + c(x, y)A(x, y) \right. \\ \left. \times \exp \left(\int_0^x \int_y^\infty K(s, t, p(s, t)a(s, t)) p(s, t)c(s, t) dt ds \right) \right]$$

for $x, y \in \mathbb{R}_+$, where $p(x, y)$ is defined by (2.3) and

$$(2.15) \quad A(x, y) = \int_0^x \int_y^\infty F(s, t, p(s, t)a(s, t)) dt ds,$$

$G(u) = \int_{u_0}^u (ds/g(s))$, $u \geq u_0 > 0$, G^{-1} is the inverse function of G , and

$$G(1) + \int_\alpha^x b(s, y) ds \in \text{Dom} (G^{-1}).$$

Proof. The proof of this theorem follows by argument similar to that given in the proof of Theorem 2.3. Let $z(x, y)$ is a nonnegative, continuous, nondecreasing and let $g \in S$, then, we observe that


$$u(x, y) \leq z(x, y)p(x, y),$$

where $p(x, y)$ is defined by (2.3). From the definition of $z(x, y)$ we have

$$(2.16) \quad u(x, y) \leq p(x, y) (a(x, y) + c(x, y)w(x, y)),$$

where $w(x, y)$ is defined by

$$w(x, y) = \int_0^x \int_y^\infty F(s, t, u(s, t)) dt ds.$$



On Certain New Integral
Inequalities and their
Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 12 of 17

From (2.12) and (2.16) we get

$$\begin{aligned}
 w(x, y) &\leq \int_0^x \int_y^\infty \left[F(s, t, p(s, t) (a(s, t) + c(s, t)w(s, t))) \right. \\
 &\quad \left. + F(s, t, p(s, t)a(s, t)) - F(s, t, p(s, t)a(s, t)) \right] dt ds \\
 &\leq A(x, y) + \int_0^x \int_y^\infty K((s, t, p(s, t)a(s, t)) p(s, t)c(s, t)w(s, t) dt ds,
 \end{aligned}$$

where $A(x, y)$ is defined by (2.15). Clearly, $A(x, y)$ is nonnegative, continuous, nondecreasing in x , $x \in \mathbb{R}_+$ and nonincreasing in y , $y \in \mathbb{R}_+$. Now, by (i) of Lemma 2.2, we obtain

$$\begin{aligned}
 (2.17) \quad w(x, y) &\leq A(x, y) \\
 &\quad \times \exp \left(\int_0^x \int_y^\infty K((s, t, p(s, t)a(s, t)) p(s, t)c(s, t) dt ds \right).
 \end{aligned}$$


Using (2.16) in (2.17) we get the required inequality in (2.14). \square

Theorem 2.6. *Let the assumptions of Theorem 2.5 be fulfilled. Define a function $z(x, y)$ by*

$$z(x, y) = a(x, y) + c(x, y) \int_x^\infty \int_y^\infty F(s, t, u(s, t)) dt ds,$$

with nonincreasing in x and $z(x, y) \geq 1$ for $x, y \in \mathbb{R}_+$. If

$$u(x, y) \leq z(x, y) + \int_x^\beta b(s, y)g(u(s, y)) ds$$



for $\beta, x, y \in \mathbb{R}_+$ and $\beta \geq x$, then

$$u(x, y) \leq \bar{p}(x, y) \left[a(x, y) + c(x, y) \bar{A}(x, y) \right. \\ \left. \times \exp \left(\int_x^\infty \int_y^\infty K(s, t, \bar{p}(s, t) a(s, t)) \bar{p}(s, t) c(s, t) dt ds \right) \right]$$

for $x, y \in \mathbb{R}_+$, where $\bar{p}(x, y)$ is defined by (2.11) and

$$\bar{A}(x, y) = \int_x^\infty \int_y^\infty F(s, t, \bar{p}(s, t) a(s, t)) dt ds.$$

G is defined in (2.5), G^{-1} is the inverse function of G , and

$$G(1) + \int_x^\beta b(s, y) ds \in \text{Dom}(G^{-1}).$$

The details of the proof of Theorem 2.6 follows by an argument similar to that in the proofs of Theorem 2.5 with suitable changes. We omit the details.

**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 13 of 17

3. Some Applications

In this section we present some immediate applications of Theorem 2.3 to study certain properties of solutions of the following terminal value problem for the hyperbolic partial differential equation

$$(3.1) \quad u_{xy}(x, y) = h(x, y, u(x, y)) + r(x, y),$$

$$(3.2) \quad u(x, \infty) = \sigma_\infty(x), u(0, y) = \tau(y), u(0, \infty) = k,$$

where $h : \mathbb{R}_+^2 \times \mathbb{R} \rightarrow \mathbb{R}, r : \mathbb{R}_+^2 \rightarrow \mathbb{R}, \sigma_\infty, \tau(y) : \mathbb{R}_+ \rightarrow \mathbb{R}$ are continuous functions and k is a real constant.

The following example deals with the estimate on the solution of the partial differential equation (3.1) with the conditions (3.2).


Example 3.1. Let $c(x, y)$ continuous, nonnegative, nondecreasing in x and non-increasing in y for $x, y \in \mathbb{R}_+$, and let

$$(3.3) \quad |h(x, y, u)| \leq c(x, y)d(x, y) |u|,$$

$$(3.4) \quad \left| \sigma_\infty(x) + \tau(y) - k - \int_0^x \int_y^\infty r(s, t) dt ds \right| \leq a(x, y) + \int_\alpha^x b(s, y)g(|u|) ds,$$

where $a(x, y), b(x, y), d(x, y), g$ are as defined in Theorem 2.3. If $u(x, y)$ is a solution of (3.1) with the conditions (3.2), then it can be written as (see [1, p. 80])

$$(3.5) \quad u(x, y) = \sigma_\infty(x) + \tau(y) - k - \int_0^x \int_y^\infty (h(s, t, u(s, t)) + r(s, t)) dt ds$$



On Certain New Integral Inequalities and their Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents




Go Back

Close

Quit

Page 14 of 17



for $x, y \in \mathbb{R}$. From (3.3), (3.4), (3.5) we get

$$(3.6) \quad |u(x, y)| \leq a(x, y) + \int_{\alpha}^x b(s, y)g(|u|) ds \\ + c(x, y) \int_0^x \int_y^{\infty} d(s, t)|u| dt ds.$$

Now, a suitable application of Theorem 2.3 to (3.6) yields the required estimate following

$$|u(x, y)| \leq p(x, y) \left[a(x, y) + c(x, y)e(x, y) \right. \\ \left. \times \exp \left(\int_0^x \int_y^{\infty} d(s, t)p(s, t)c(s, t) dt ds \right) \right]$$

for $x, y \in \mathbb{R}_+$, where $e(x, y), p(x, y)$ are define in Theorem 2.3.

Our next result deals with the uniqueness of the solution of the partial differential equation (3.1) with the conditions (3.2).

Example 3.2. Suppose that the function h in (3.1) satisfies the condition

$$(3.7) \quad |h(x, y, u) - h(x, y, v)| \leq c(x, y)d(x, y) |u - v|,$$

where $c(x, y), d(x, y)$ is as defined in Theorem 2.3 with $c(x, y)$ is nonincreasing in y . Let $u(x, y), v(x, y)$ be two solutions of equation (3.1) with the conditions (3.2). From (3.5), (3.7) we have

$$(3.8) \quad |u(x, y) - v(x, y)| \leq c(x, y) \int_0^x \int_y^{\infty} d(s, t)|u(s, t) - v(s, t)| dt ds.$$

Now a suitable application of Theorem 2.3 yields $u(x, y) = v(x, y)$, that is, there is at most one solution to the problem (3.1) with the conditions (3.2).

**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back


Close

Quit

Page 15 of 17

References

- [1] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, 1992.
- [2] P.R. BEESACK, Gronwall inequalities, *Carleton University Mathematical Lecture Notes*, No. 11, 1987.
- [3] S.S. DRAGOMIR AND N.M. IONESCU, On nonlinear integral inequalities in two independent variables, *Studia Univ. Babeş-Bolyai, Math.*, **34** (1989), 11–17.
- [4] A. MATE AND P. NEVAL, Sublinear perturbations of the differential equation $y^{(n)} = 0$ and of the analogous difference equation, *J. Differential Equations*, **52** (1984), 234–257.
- [5] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, Boston, London, 1991.
- [6] B.G. PACHPATTE, *Inequalities for Differential and Integral Equations*, Academic Press, New York, 1998.
- [7] B.G. PACHPATTE, On some new discrete inequalities and their applications, *Proc. Nat. Acad. Sci., India*, **46**(A) (1976), 255–262.
- [8] B.G. PACHPATTE, On certain new finite difference inequalities, *Indian J. Pure Appl. Math.*, **24** (1993), 373–384.
- [9] B.G. PACHPATTE, Some new finite difference inequalities, *Computer Math. Appl.*, **28** (1994), 227–241.



On Certain New Integral
Inequalities and their
Applications

S.S. Dragomir and Young-Ho Kim

Title Page

Contents




Go Back

Close

Quit

Page 16 of 17

- [10] B.G. PACHPATTE, On some new discrete inequalities useful in the theory of partial finite difference equations, *Ann. Differential Equations*, **12** (1996), 1–12.
- [11] B.G. PACHPATTE, On some fundamental integral inequalities and their discrete analogues, *J. Ineq. Pure Appl. Math.*, **2**(2) (2001), Article 15. [ONLINE: http://jipam.vu.edu.au/v2n2/046_00.html]



**On Certain New Integral
Inequalities and their
Applications**

S.S. Dragomir and Young-Ho Kim

Title Page

Contents



Go Back

Close

Quit

Page 17 of 17