

# Journal of Inequalities in Pure and Applied Mathematics

## AN INEQUALITY ABOUT $q$ -SERIES

MINGJIN WANG

Department of Mathematics  
East China Normal University  
Shanghai, 200062  
People's Republic of China

*EMail:* [wmj@jpu.edu.cn](mailto:wmj@jpu.edu.cn)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
037-06



---

volume 7, issue 4, article 136,  
2006.

*Received 10 February, 2006;  
accepted 16 March, 2006.*

*Communicated by: H. Bor*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

In this paper, we first generalize the traditional notation  $(a; q)_n$  to  $[g(x); q]_n$  and then obtain an inequality about  $q$ -series and some infinite products by means of the new conception. Because many  $q$ -series are not summable, our results are useful to study  $q$ -series and its application.

*2000 Mathematics Subject Classification:* Primary 60E15; Secondary 33D15.

*Key words:* Inequality,  $q$ -series.

## Contents

1	Introduction .....	3
2	An Inequality About $q$ -Series .....	4
	References	



---

### An Inequality About $q$ -Series

Mingjin Wang

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

# 1. Introduction

$q$ -series, which are also called basic hypergeometric series, play an very important role in many fields. Such as affine root systems, Lie algebras and groups, number theory, orthogonal polynomials, physics (such as representations of quantum groups and Baxter's work on the hard hexagon model). Most of the research work on  $q$ -series is to set up identity. But there are also great many  $q$ -series whose sums cannot be obtained easily. On these occasions, we must use other methods to study  $q$ -series. Using inequalities is one of the choices. In this paper, we obtain an inequality about  $q$ -series and some infinite products. Our results are useful for the study of  $q$ -series.



---

An Inequality About  $q$ -Series

Mingjin Wang

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

## 2. An Inequality About $q$ -Series

In this section we will introduce a new concept and obtain an inequality about  $q$ -series. First we give some lemmas.

**Lemma 2.1.** *If  $0 < q < 1$ ,  $0 < a < 1$ , then for any natural number  $n$ , we have*

$$1 < \frac{1 - aq^n}{1 - q^n} \leq \frac{1 - aq}{1 - q}.$$

*Proof.* Let  $g(x) = \frac{1-ax}{1-x}$ ,  $0 \leq x < 1$ , then  $g'(x) = \frac{1-a}{(1-x)^2} > 0$ . So  $g(x)$  is a strictly increasing function on  $[0, 1)$ . For any natural number  $n$ , we have

$$0 < q^n \leq q < 1.$$

So,

$$g(0) < g(q^n) \leq g(q).$$


That is

$$1 < \frac{1 - aq^n}{1 - q^n} \leq \frac{1 - aq}{1 - q}.$$

□

**Lemma 2.2.** *If  $0 < a < \frac{1-q}{1+q}$ ,  $0 < q < 1$ , then for any real number  $0 < x \leq 1$ , we have*

$$1 + ax \left( ax - \frac{2(1 - aqx)}{1 - q} \right) > 0.$$



---

An Inequality About  $q$ -Series

Mingjin Wang

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

*Proof.* Let

$$g(x) = 1 + ax \left( ax - \frac{2(1 - aqx)}{1 - q} \right) = 1 + \frac{a}{1 - q} [a(1 + q)x^2 - 2x].$$

Under the condition  $0 < a < \frac{1-q}{1+q}$ , we know  $0 < a(1 + q) < 1 - q$ , so

$$g'(x) = \frac{2a}{1 - q} [a(1 + q)x - 1] < \frac{2a}{1 - q} [(1 - q)x - 1].$$

Since  $0 < 1 - q < 1$ ,  $0 < x \leq 1$ , we know  $0 < (1 - q)x < 1$ . Therefore  $g'(x) < 0$  and  $g(x)$  is a strictly decreasing function on  $(0, 1]$ . We have

$$g(x) > g(1) = 1 + a \left( a - \frac{2(1 - aq)}{1 - q} \right) = \frac{1}{1 - q} [(1 + q)a^2 - 2a + (1 - q)].$$

Letting

$$(1 + q)a^2 - 2a + (1 - q) = 0,$$

we have

$$a_1 = \frac{1 - q}{1 + q}, \quad a_2 = 1.$$


So, when  $0 < a < \frac{1-q}{1+q}$ ,

$$(1 + q)a^2 - 2a + (1 - q) > 0,$$

that is,

$$1 + ax \left( ax - \frac{2(1 - aqx)}{1 - q} \right) > 0.$$

□



---

**An Inequality About  $q$ -Series**

Mingjin Wang

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

**Lemma 2.3.** If  $0 < q < \sqrt{2} - 1$ , we have

$$q < \frac{1 - q}{1 + q}.$$

*Proof.* Let

$$g(q) = q(1 + q) - (1 - q) = (q + 1 + \sqrt{2})(q + 1 - \sqrt{2}).$$

When  $0 < q < \sqrt{2} - 1$ ,  $g(q) < 0$ , so we have

$$q < \frac{1 - q}{1 + q}.$$

□

**Definition 2.1.** Suppose  $g(x)$  is a function on  $[0, 1]$ , we denote  $[g(x); q]_n$  by

$$[g(x); q]_n = (1 - g(q^0))(1 - g(q^1)) \cdots (1 - g(q^{n-1})).$$


We also use the notation  $[g(x); q]_\infty$  to express infinite product. That is

$$[g(x); q]_\infty = (1 - g(q^0))(1 - g(q^1)) \cdots (1 - g(q^n)) \cdots .$$

If  $g(x) = ax$ , then

$$[g(x); q]_n = (1 - a)(1 - aq) \cdots (1 - aq^{n-1}) = (a; q)_n.$$

So  $[g(x); q]_n$  is the expansion of  $(a; q)_n$ , where  $(a; q)_n$  is the  $q$ -shifted factorial. Please note that our notation  $[g(x); q]_n$  in this paper is different from traditional notation  $(a; q)_n$ .



Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

**Theorem 2.4.** Suppose  $0 < a < \frac{1-q}{1+q}$ ,  $0 < q < \sqrt{2} - 1$ ,  $0 < z < 1$ , then the following inequality holds

$$(2.1) \quad \frac{[g_2(x, a); q]_\infty}{(1-z)[g_1(x, q); q]_\infty} \leq \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n \leq \frac{[g_1(x, a); q]_\infty}{(1-z)[g_2(x, q); q]_\infty},$$

where

$$g_1(x, a) = -ax(ax - 2)z,$$

$$g_2(x, a) = -ax \left( ax - \frac{2(1 - aqx)}{1 - q} \right) z.$$

When  $a = q$ , the equality holds.


*Proof.* Let

$$f(a, z) = \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n,$$

since

$$1 - a = (1 - aq^n) - a(1 - q^n),$$

$$\begin{aligned} f(a, z) &= 1 + \sum_{n=1}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n \\ &= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_n^2} [(1 - aq^n) - a(1 - q^n)]^2 z^n \end{aligned}$$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

$$\begin{aligned}
&= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_n^2} [(1 - aq^n)^2 - 2a(1 - q^n)(1 - aq^n) + a^2(1 - q^n)^2] z^n \\
&= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_n^2}{(q; q)_n^2} z^n + a^2 \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n \\
&\quad - 2a \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_n^2} (1 - q^n)(1 - aq^n) z^n \\
&= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_n^2}{(q; q)_n^2} z^n + a^2 \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n - 2a \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} \frac{1 - aq^n}{1 - q^n} z^n.
\end{aligned}$$


From Lemma 2.1, we know  $\frac{1-aq^n}{1-q^n} > 1$ . So

$$\begin{aligned}
f(a, z) &= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_n^2}{(q; q)_n^2} z^n + a^2 \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n - 2a \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} \frac{1 - aq^n}{1 - q^n} z^n \\
&\leq f(aq, z) + a^2 z f(aq, z) - 2a \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n \\
&= f(aq, z) + a^2 z f(aq, z) - 2az f(aq, z) \\
&= (1 + a(a - 2)z) f(aq, z).
\end{aligned}$$

By iterating this functional inequality  $n - 1$  times we get that

$$f(a, z) \leq [g_1(x, a); q]_n f(aq^n, z), \quad n = 1, 2, \dots,$$

where  $g_1(x, a) = -ax(ax - 2)z$ . Which on letting  $n \rightarrow \infty$  and using  $q^n \rightarrow 0$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16



gives


$$(2.2) \quad f(a, z) \leq [g_1(x, a); q]_{\infty} f(0, z).$$

Again from Lemma 2.1, we know  $\frac{1-aq^n}{1-q^n} \leq \frac{1-aq}{1-q}$ . So

$$\begin{aligned} f(a, z) &= 1 + \sum_{n=1}^{\infty} \frac{(aq; q)_n^2}{(q; q)_n^2} z^n + a^2 \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n \\ &\quad - 2a \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} \frac{1-aq^n}{1-q^n} z^n \\ &\geq f(aq, z) + a^2 z f(aq, z) - 2a \frac{1-aq}{1-q} \sum_{n=1}^{\infty} \frac{(aq; q)_{n-1}^2}{(q; q)_{n-1}^2} z^n \\ &= \left( 1 + a^2 z - 2a z \frac{1-aq}{1-q} \right) f(aq, z) \\ &= \left( 1 + a \left( a - \frac{2(1-aq)}{1-q} \right) z \right) f(aq, z). \end{aligned}$$

Using Lemma 2.2, we know that, for any natural number  $n$

$$\begin{aligned} &1 + aq^n \left( aq^n - \frac{2(1-aq^{n+1})}{1-q} \right) z \\ &= z \left[ \frac{1}{z} + aq^n \left( aq^n - \frac{2(1-aq^{n+1})}{1-q} \right) \right] \\ &\geq z \left[ 1 + aq^n \left( aq^n - \frac{2(1-aq^{n+1})}{1-q} \right) \right] > 0. \end{aligned}$$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

Let  $g_2(x, a) = -ax \left( ax - \frac{2(1-ax)}{1-q} \right) z$ , and by iterating this functional inequality  $n - 1$  times we get that

$$f(a, z) \geq [g_2(x, a); q]_n f(aq^n, z), \quad n = 1, 2, \dots$$

Which on letting  $n \rightarrow \infty$  and using  $q^n \rightarrow 0$  gives

$$(2.3) \quad f(a, z) \geq [g_2(x, a); q]_\infty f(0, z).$$

Combined with (2.2) and (2.3) gives

$$(2.4) \quad [g_2(x, a); q]_\infty f(0, z) \leq f(a, z) \leq [g_1(x, a); q]_\infty f(0, z).$$

Using Lemma 2.3, when  $0 < q < \sqrt{2} - 1$ , we have  $q < \frac{1-q}{1+q}$ . So, let  $a = q$ , also combining (2.2) and (2.3) gives the following inequality


$$\frac{f(q, z)}{[g_1(x, q); q]_\infty} \leq f(0, z) \leq \frac{f(q, z)}{[g_2(x, q); q]_\infty}.$$

Because of  $f(q, z) = \frac{1}{1-z}$ , we have

$$(2.5) \quad \frac{1}{(1-z)[g_1(x, q); q]_\infty} \leq f(0, z) \leq \frac{1}{(1-z)[g_2(x, q); q]_\infty}.$$

(2.4) and (2.5) yield the following inequality

$$\frac{[g_2(x, a); q]_\infty}{(1-z)[g_1(x, q); q]_\infty} \leq f(a, z) \leq \frac{[g_1(x, a); q]_\infty}{(1-z)[g_2(x, q); q]_\infty}.$$



That is

$$\frac{[g_2(x, a); q]_\infty}{(1-z)[g_1(x, q); q]_\infty} \leq \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n \leq \frac{[g_1(x, a); q]_\infty}{(1-z)[g_2(x, q); q]_\infty}.$$

If  $a = q$ , we have

$$\frac{[g_2(x, a); q]_\infty}{(1-z)[g_1(x, q); q]_\infty} = \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n = \frac{[g_1(x, a); q]_\infty}{(1-z)[g_2(x, q); q]_\infty} = \frac{1}{1-z}.$$


So the equality holds. We complete our proof.  $\square$

**Corollary 2.5.** *Under the conditions of the theorem, we have*

$$(2.6) \quad \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_\infty}{(qz; q)_\infty} \right]^2 z^n \leq \frac{\left| \begin{array}{cc} [g_1(x, a); q]_\infty & [g_2(x, q); q]_\infty \\ (az; q)_\infty^2 & (qz; q)_\infty^2 \end{array} \right|}{[g_2(x, q); q]_\infty (z; q)_\infty (qz; q)_\infty}.$$

*Proof.* Since

$$\begin{aligned} & \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_\infty}{(qz; q)_\infty} \right]^2 z^n \\ &= \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n^2}{(q; q)_n^2} - 2 \frac{(a; q)_n (az; q)_\infty}{(q; q)_n (qz; q)_\infty} + \frac{(az; q)_\infty^2}{(qz; q)_\infty^2} \right] z^n \\ &= \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n - 2 \frac{(az; q)_\infty}{(qz; q)_\infty} \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} z^n + \frac{(az; q)_\infty^2}{(qz; q)_\infty^2} \sum_{n=0}^{\infty} z^n \end{aligned}$$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n - 2 \frac{(az; q)_{\infty} (az; q)_{\infty}}{(qz; q)_{\infty} (z; q)_{\infty}} + \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \frac{1}{1-z} \\
&= \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n - 2 \frac{1}{1-z} \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} + \frac{1}{1-z} \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \\
&= \sum_{n=0}^{\infty} \frac{(a; q)_n^2}{(q; q)_n^2} z^n - \frac{1}{1-z} \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \\
&\leq \frac{[g_1(x, a); q]_{\infty}}{(1-z)[g_2(x, q); q]_{\infty}} - \frac{1}{1-z} \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \\
&= \frac{\begin{vmatrix} [g_1(x, a); q]_{\infty} & [g_2(x, q); q]_{\infty} \\ (az; q)_{\infty}^2 & (qz; q)_{\infty}^2 \end{vmatrix}}{[g_2(x, q); q]_{\infty} (z; q)_{\infty} (qz; q)_{\infty}}
\end{aligned}$$

we gain the inequality we seek.  $\square$


**Theorem 2.6.** *Under the conditions of the theorem, the following inequality holds*

$$(2.7) \quad \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \leq \frac{[g_1(x, a); q]_{\infty}}{[g_2(x, q); q]_{\infty}}.$$

*Proof.* From the proof of (2.6), we have

$$\frac{[g_1(x, a); q]_{\infty}}{(1-z)[g_2(x, q); q]_{\infty}} - \frac{1}{1-z} \frac{(az; q)_{\infty}^2}{(qz; q)_{\infty}^2} \geq \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right]^2 z^n \geq 0$$

so the inequality (2.7) holds.  $\square$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

**Corollary 2.7.** Suppose  $0 < a < \frac{1-q}{1+q}$ ,  $0 < b < \frac{1-q}{1+q}$ ,  $0 < q < \sqrt{2}-1$ ,  $0 < z < 1$ , then the following inequality holds


$$(2.8) \quad \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n^2} z^n \leq \frac{(az, bz; q)_{\infty}}{(z; q)_{\infty}^2} + \frac{\left| \frac{[g_1(x, a); q]_{\infty}}{(az; q)_{\infty}^2} \frac{[g_2(x, q); q]_{\infty}}{(qz; q)_{\infty}^2} \right|^{\frac{1}{2}} \cdot \left| \frac{[g_1(x, b); q]_{\infty}}{(bz; q)_{\infty}^2} \frac{[g_2(x, q); q]_{\infty}}{(qz; q)_{\infty}^2} \right|^{\frac{1}{2}}}{[g_2(x, q); q]_{\infty} (z; q)_{\infty} (qz; q)_{\infty}}.$$

*Proof.* Noting that

$$\sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n^2} z^n > 0 \quad \text{and} \quad \frac{(az, bz; q)_{\infty}}{(z; q)_{\infty}^2} > 0,$$

we have

$$(2.9) \quad \begin{aligned} & \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n^2} z^n - \frac{(az, bz; q)_{\infty}}{(z; q)_{\infty}^2} \\ & \leq \left| \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n^2} z^n - \frac{(az, bz; q)_{\infty}}{(z; q)_{\infty}^2} \right| \\ & = \left| \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right] \left[ \frac{(b; q)_n}{(q; q)_n} - \frac{(bz; q)_{\infty}}{(qz; q)_{\infty}} \right] z^n \right| \\ & \leq \sum_{n=0}^{\infty} \left| \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right| \cdot \left| \frac{(b; q)_n}{(q; q)_n} - \frac{(bz; q)_{\infty}}{(qz; q)_{\infty}} \right| z^n \end{aligned}$$



An Inequality About  $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

$$= \sum_{n=0}^{\infty} \left| \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right| z^{\frac{n}{2}} \cdot \left| \frac{(b; q)_n}{(q; q)_n} - \frac{(bz; q)_{\infty}}{(qz; q)_{\infty}} \right| z^{\frac{n}{2}}.$$

Using the Cauchy inequality and (2.6), we have


$$(2.10) \quad \sum_{n=0}^{\infty} \left| \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right| z^{\frac{n}{2}} \cdot \left| \frac{(b; q)_n}{(q; q)_n} - \frac{(bz; q)_{\infty}}{(qz; q)_{\infty}} \right| z^{\frac{n}{2}}$$

$$\leq \left\{ \sum_{n=0}^{\infty} \left[ \frac{(a; q)_n}{(q; q)_n} - \frac{(az; q)_{\infty}}{(qz; q)_{\infty}} \right]^2 z^n \right\}^{\frac{1}{2}}$$

$$\cdot \left\{ \sum_{n=0}^{\infty} \left[ \frac{(b; q)_n}{(q; q)_n} - \frac{(bz; q)_{\infty}}{(qz; q)_{\infty}} \right]^2 z^n \right\}^{\frac{1}{2}}$$

$$\leq \left\{ \frac{\begin{vmatrix} [g_1(x, a); q]_{\infty} & [g_2(x, q); q]_{\infty} \\ (az; q)_{\infty}^2 & (qz; q)_{\infty}^2 \end{vmatrix}}{[g_2(x, q); q]_{\infty}(z; q)_{\infty}(qz; q)_{\infty}} \right\}^{\frac{1}{2}}$$

$$\cdot \left\{ \frac{\begin{vmatrix} [g_1(x, b); q]_{\infty} & [g_2(x, q); q]_{\infty} \\ (bz; q)_{\infty}^2 & (qz; q)_{\infty}^2 \end{vmatrix}}{[g_2(x, q); q]_{\infty}(z; q)_{\infty}(qz; q)_{\infty}} \right\}^{\frac{1}{2}}$$



## An Inequality About $q$ -Series

Mingjin Wang

Title Page

Contents



Go Back

Close

Quit


Page 14 of 16

$$= \frac{\left| \begin{matrix} [g_1(x, a); q]_\infty & [g_2(x, q); q]_\infty \\ (az; q)_\infty^2 & (qz; q)_\infty^2 \end{matrix} \right|^{\frac{1}{2}} \cdot \left| \begin{matrix} [g_1(x, b); q]_\infty & [g_2(x, q); q]_\infty \\ (bz; q)_\infty^2 & (qz; q)_\infty^2 \end{matrix} \right|^{\frac{1}{2}}}{[g_2(x, q); q]_\infty (z; q)_\infty (qz; q)_\infty}.$$

Combining (2.9) and (2.10) gives

$$\sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n^2} z^n \leq \frac{(az, bz; q)_\infty}{(z; q)_\infty^2} + \frac{\left| \begin{matrix} [g_1(x, a); q]_\infty & [g_2(x, q); q]_\infty \\ (az; q)_\infty^2 & (qz; q)_\infty^2 \end{matrix} \right|^{\frac{1}{2}} \cdot \left| \begin{matrix} [g_1(x, b); q]_\infty & [g_2(x, q); q]_\infty \\ (bz; q)_\infty^2 & (qz; q)_\infty^2 \end{matrix} \right|^{\frac{1}{2}}}{[g_2(x, q); q]_\infty (z; q)_\infty (qz; q)_\infty}.$$

This is the inequality we seek. □




---

### An Inequality About $q$ -Series

Mingjin Wang

---

Title Page

Contents



Go Back


Close

Quit

Page 15 of 16

## References

- [1] G.H. HARDY, J.E. LITTLEWOOD AND G. POLYA, *Inequalities, Edition 1 and 2*. Cambridge University Press, Cambridge, 1934, 1952.
- [2] G. GASPER, *Lecture Notes for an Introductory Minicourse on  $q$ -Series*, September 19, 1995 version.
- [3] G. GASPER AND M. RAHMAN, *Basic Hypergeometric Series, Encyclopedia of Mathematics and Its Applications, 35*. Cambridge University Press, Cambridge and New York, 1990.



---

An Inequality About  $q$ -Series

Mingjin Wang

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16