


A GENERALIZATION OF THE PRE-GRÜSS INEQUALITY AND APPLICATIONS TO SOME QUADRATURE FORMULAE

NENAD UJEVIĆ

Department of Mathematics
University of Split
Teslina 12/III
21000 Split, Croatia.
EMail: ujevic@pmfst.hr



volume 3, issue 1, article 13,
2002.

*Received 03 May, 2001;
accepted 26 October, 2001.*

Communicated by: P. Cerone

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract


A generalization of the pre-Grüss inequality is presented. It is applied to estimations of remainders of some quadrature formulas.

2000 Mathematics Subject Classification: 26D10, 41A55.

Key words: Pre-Grüss inequality, Generalization, Quadrature formulae.

Contents

| | | |
|---|--------------------|---|
| 1 | Introduction | 3 |
| 2 | Main Results | 6 |
| | References | |



A Generalization of the Pre-Grüss Inequality and Applications to some Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. Introduction

In recent years a number of authors have written about generalizations of Ostrowski's inequality. For example, this topic is considered in [1], [2], [5], [7], [9] and [12]. In this way some new types of inequalities are formed, such as inequalities of Ostrowski-Grüss type, inequalities of Ostrowski-Chebyshev type, etc. An important role in forming these inequalities is played by the pre-Grüss inequality. This paper develops a new approach to the topic obtaining better results than the approach using the pre-Grüss inequality. It presents new, improved versions of the mid-point and trapezoidal inequality. The mid-point inequality is considered in [1], [2], [3], [7] and [9], while the trapezoidal inequality is considered in [4], [5], [7] and [9].

In [11] we can find the pre-Grüss inequality:

$$(1.1) \quad T(f, g)^2 \leq T(f, f)T(g, g),$$

where $f, g \in L_2(a, b)$ and $T(f, g)$ is the Chebyshev functional:

$$(1.2) \quad T(f, g) = \frac{1}{b-a} \int_a^b f(t)g(t)dt - \frac{1}{(b-a)^2} \int_a^b f(t)dt \int_a^b g(t)dt.$$

If there exist constants $\gamma, \delta, \Gamma, \Delta \in \mathbb{R}$ such that

$$\delta \leq f(t) \leq \Delta \text{ and } \gamma \leq g(t) \leq \Gamma, t \in [a, b]$$

then, using (1.1), we get the Grüss inequality:

$$(1.3) \quad |T(f, g)| \leq \frac{(\Delta - \delta)(\Gamma - \gamma)}{4}.$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

Specially, we have


$$(1.4) \quad T(f, f) \leq \frac{(\Delta - \delta)^2}{4}.$$

Using the above inequalities we get the following inequalities:

$$(1.5) \quad \left| f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt \right| \\ \leq \frac{(b-a)^2}{2\sqrt{3}} \left[\frac{1}{b-a} \|f'\|_2^2 - \left(\frac{f(b)-f(a)}{b-a}\right)^2 \right]^{\frac{1}{2}} \\ \leq \frac{(b-a)^2}{4\sqrt{3}} (\Gamma - \gamma)$$

where $f : [a, b] \rightarrow \mathbb{R}$ is an absolutely continuous function whose derivative $f' \in L_2(a, b)$ and $\gamma \leq f'(t) \leq \Gamma, t \in [a, b]$. As usual, $\|\cdot\|_2$ is the norm in $L_2(a, b)$. Further,

$$(1.6) \quad \left| \frac{f(a)+f(b)}{2}(b-a) - \int_a^b f(t)dt \right| \\ \leq \frac{(b-a)^2}{2\sqrt{3}} \left[\frac{1}{b-a} \|f'\|_2^2 - \left(\frac{f(b)-f(a)}{b-a}\right)^2 \right]^{\frac{1}{2}} \\ \leq \frac{(b-a)^2}{4\sqrt{3}} (\Gamma - \gamma)$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit


Page 4 of 20

and

$$(1.7) \quad \left| \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt \right| \\ \leq \frac{(b-a)^2}{4\sqrt{3}} \left[\frac{1}{b-a} \|f'\|_2^2 - \left(\frac{f(b) - f(a)}{b-a} \right)^2 \right]^{\frac{1}{2}} \\ \leq \frac{(b-a)^2}{8\sqrt{3}} (\Gamma - \gamma)$$

where the function f satisfies the above conditions. The inequalities (1.5)-(1.7) are considered (and proved) in [2], [9] and [12].

In this paper we generalize (1.1). We use the generalization to improve the above inequalities.



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

2. Main Results

Lemma 2.1. Let $f, g, \Psi_i \in L_2(a, b)$, $i = 0, 1, 2, \dots, n$, where $\Psi_i^0 = \Psi_i(t) / \|\Psi_i\|_2$ are orthonormal functions. If $S_n(f, g)$ is defined by

$$S_n(f, g) = \int_a^b f(t)g(t)dt - \sum_{i=0}^n \int_a^b f(s)\Psi_i^0(s)ds \int_a^b g(s)\Psi_i^0(s)ds$$

then we have

$$|S_n(f, g)| \leq S_n(f, f)^{\frac{1}{2}} S_n(g, g)^{\frac{1}{2}}.$$


The proof follows by the known inequality holding in inner product spaces $(H, \langle \cdot, \cdot \rangle)$

$$\left| \langle x, y \rangle - \sum_{i=0}^n \langle x, l_i \rangle \langle l_i, y \rangle \right|^2 \leq \left(\|x\|^2 - \sum_{i=0}^n |\langle x, l_i \rangle|^2 \right) \left(\|y\|^2 - \sum_{i=0}^n |\langle l_i, y \rangle|^2 \right),$$

where $x, y \in H$ and $\{l_i\}_{i=0, n}$ is an orthonormal family in H , i.e., $(l_i, l_j) = \delta_{ij}$ for $i, j \in \{0, \dots, n\}$.

We here use only the case $n = 1$. We choose $\Psi_0^0(t) = 1/\sqrt{b-a}$, $\Psi_1(t) = \Psi(t)$ and denote $S_1(g, h) = S_\Psi(g, h)$ such that

$$(2.1) \quad S_\Psi(g, h) = \int_a^b g(t)h(t)dt - \frac{1}{b-a} \int_a^b g(t)dt \int_a^b h(t)dt - \int_a^b g(t)\Psi_0(t)dt \int_a^b h(t)\Psi_0(t)dt$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 6 of 20

where $g, h, \Psi \in L_2(a, b)$, $\Psi_0(t) = \Psi(t) / \|\Psi\|_2$ and

$$(2.2) \quad \int_a^b \Psi(t) dt = 0.$$

Lemma 2.2. *With the above notations we have*

$$(2.3) \quad |S_\Psi(g, h)| \leq S_\Psi(g, g)^{\frac{1}{2}} S_\Psi(h, h)^{\frac{1}{2}}.$$

It is obvious that

$$(2.4) \quad S_\Psi(g, h) = (b - a)T(g, h) - \int_a^b g(t)\Psi_0(t)dt \int_a^b h(t)\Psi_0(t)dt$$

so that $S_\Psi(g, h)$ is a generalization of the Chebyshev functional.

We also define the functions:


$$(2.5) \quad \Phi(t) = \begin{cases} t - \frac{2a+b}{3}, & t \in [a, \frac{a+b}{2}] \\ t - \frac{a+2b}{3}, & t \in (\frac{a+b}{2}, b] \end{cases}$$

and

$$(2.6) \quad \chi(t) = \begin{cases} t - \frac{5a+b}{6}, & t \in [a, \frac{a+b}{2}] \\ t - \frac{a+5b}{6}, & t \in (\frac{a+b}{2}, b]. \end{cases}$$

It is not difficult to verify that

$$(2.7) \quad \int_a^b \Phi(t)dt = \int_a^b \chi(t)dt = 0$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

and

$$(2.8) \quad \|\Phi\|_2^2 = \|\chi\|_2^2 = \frac{(b-a)^3}{36}.$$

We define

$$(2.9) \quad \Phi_0(t) = \frac{\Phi(t)}{\|\Phi\|_2}, \quad \chi_0(t) = \frac{\chi(t)}{\|\chi\|_2}.$$

Integrating by parts, we have


$$(2.10) \quad \begin{aligned} Q(f; a, b) &= \int_a^b \Phi_0(t) f'(t) dt \\ &= \frac{2}{\sqrt{b-a}} \left[f(a) + f\left(\frac{a+b}{2}\right) + f(b) - \frac{3}{b-a} \int_a^b f(t) dt \right] \end{aligned}$$

and

$$(2.11) \quad \begin{aligned} P(f; a, b) &= \int_a^b \chi_0(t) f'(t) dt \\ &= \frac{1}{\sqrt{b-a}} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) - \frac{6}{b-a} \int_a^b f(t) dt \right]. \end{aligned}$$

Remark 2.1. *It is obvious that*

$$(2.12) \quad S_{\Psi}(g, g) = (b-a)T(g, g) - \left(\int_a^b g(t) \Psi_0(t) dt \right)^2 \leq (b-a)T(g, g).$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 8 of 20

Theorem 2.3. (Mid-point inequality) Let $I \subset \mathbb{R}$ be a closed interval and $a, b \in \text{Int } I$, $a < b$. If $f : I \rightarrow \mathbb{R}$ is an absolutely continuous function whose derivative $f' \in L_2(a, b)$ then we have

$$(2.13) \quad \left| f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{2\sqrt{3}} C_1,$$

where

$$(2.14) \quad C_1 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - [Q(f; a, b)]^2 \right\}^{\frac{1}{2}}$$

and $Q(f; a, b)$ is defined by (2.10).

Proof. We define


$$(2.15) \quad p(t) = \begin{cases} t - a, & t \in [a, \frac{a+b}{2}] \\ t - b, & t \in (\frac{a+b}{2}, b]. \end{cases}$$

Then we have

$$(2.16) \quad \int_a^b p(t)dt = 0$$

and

$$(2.17) \quad \|p\|_2^2 = \int_a^b p(t)^2 dt = \frac{(b-a)^3}{12}.$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 9 of 20

We now calculate

$$(2.18) \quad \int_a^b p(t)\Phi(t)dt \\ = \int_a^{\frac{a+b}{2}} (t-a) \left(t - \frac{2a+b}{3}\right) dt + \int_{\frac{a+b}{2}}^b (t-b) \left(t - \frac{a+2b}{3}\right) dt = 0.$$

Integrating by parts, we have


$$(2.19) \quad \int_a^b p(t)f'(t)dt = \int_a^{\frac{a+b}{2}} (t-a)f'(t)dt + \int_{\frac{a+b}{2}}^b (t-b)f'(t)dt \\ = f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt.$$

Using (2.16), (2.18) and (2.19) we get

$$(2.20) \quad S_{\Phi}(p, f') = \int_a^b p(t)f'(t)dt - \frac{1}{b-a} \int_a^b p(t)dt \int_a^b f'(t)dt \\ - \int_a^b f'(t)\Phi_0(t)dt \int_a^b p(t)\Phi_0(t)dt \\ = f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt.$$

From (2.20) and (2.3) it follows that

$$(2.21) \quad \left| f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt \right| \leq S_{\Phi}(f', f')^{\frac{1}{2}} S_{\Phi}(p, p)^{\frac{1}{2}}.$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

From (2.16)-(2.18) we get

$$(2.22) \quad S_{\Phi}(p, p) = \|p\|_2^2 - \frac{1}{b-a} \left(\int_a^b p(t) dt \right)^2 - \left(\int_a^b p(t) \Phi_0(t) dt \right)^2 \\ = \frac{(b-a)^3}{12}.$$

We also have

$$(2.23) \quad C_1^2 = S_{\Phi}(f', f').$$

From (2.21)-(2.23) we easily find that (2.13) holds. \square

Remark 2.2. *It is not difficult to see that (2.13) is better than the first estimation in (1.5).*

Theorem 2.4. *(Trapezoidal inequality) Under the assumptions of Theorem 2.3 we have*

$$(2.24) \quad \left| \frac{f(a) + f(b)}{2} (b-a) - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{2\sqrt{3}} C_2,$$

where

$$(2.25) \quad C_2 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - [P(f; a, b)]^2 \right\}^{\frac{1}{2}}$$

and $P(f; a, b)$ is defined by (2.11).



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 11 of 20

Proof. Let $p(t)$ be defined by (2.15). We calculate


$$\begin{aligned}
 (2.26) \quad & \int_a^b p(t)\chi(t)dt \\
 &= \int_a^{\frac{a+b}{2}} (t-a) \left(t - \frac{5a+b}{6}\right) dt + \int_{\frac{a+b}{2}}^b (t-b) \left(t - \frac{a+5b}{6}\right) dt \\
 &= \frac{(b-a)^3}{24}.
 \end{aligned}$$

Integrating by parts, we have

$$\begin{aligned}
 (2.27) \quad & \int_a^b f'(t)\chi(t)dt \\
 &= \int_a^{\frac{a+b}{2}} \left(t - \frac{5a+b}{6}\right) f'(t)dt + \int_{\frac{a+b}{2}}^b \left(t - \frac{a+5b}{6}\right) f'(t)dt \\
 &= \frac{f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)}{6}(b-a) - \int_a^b f(t)dt.
 \end{aligned}$$

Using (2.16), (2.19), (2.26), (2.27) and (2.8) we get

$$\begin{aligned}
 (2.28) \quad S_\chi(f', p) &= \int_a^b p(t)f'(t)dt - \frac{1}{b-a} \int_a^b f'(t)dt \int_a^b p(t)dt \\
 &\quad - \int_a^b p(t)\chi_0(t)dt \int_a^b f'(t)\chi_0(t)dt
 \end{aligned}$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents




Go Back

Close

Quit

Page 12 of 20



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

$$\begin{aligned}
 &= f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt \\
 &\quad - \frac{3}{2} \left[\frac{f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)}{6} (b-a) - \int_a^b f(t)dt \right] \\
 &= -\frac{1}{2}(b-a) \frac{f(a) + f(b)}{2} + \frac{1}{2} \int_a^b f(t)dt.
 \end{aligned}$$

From (2.3) and (2.28) it follows that

$$(2.29) \quad \left| \frac{f(a) + f(b)}{2} (b-a) - \int_a^b f(t)dt \right| \leq 2S_\chi(f', f')^{\frac{1}{2}} S_\chi(p, p)^{\frac{1}{2}}.$$

We have

$$\begin{aligned}
 (2.30) \quad S_\chi(p, p) &= \|p\|_2^2 - \frac{1}{b-a} \left(\int_a^b p(t)dt \right)^2 - \left(\int_a^b p(t)\chi_0(t)dt \right)^2 \\
 &= \frac{(b-a)^3}{48}
 \end{aligned}$$

and

$$(2.31) \quad C_2^2 = S_\chi(f', f').$$

From (2.29)-(2.31) we easily get (2.24). □

Remark 2.3. We see that (2.24) is better than the first estimation in (1.6).

We now consider a simple quadrature rule of the form

$$(2.32) \quad \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt$$

$$= \frac{1}{2} \left[f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right] (b-a) - \int_a^b f(t)dt = R(f).$$


It is not difficult to see that (2.32) is a convex combination of the mid-point quadrature rule and the trapezoidal quadrature rule. In [5] it is shown that (2.32) has a better estimation of error than the well-known Simpson quadrature rule (when we estimate the error in terms of the first derivative f' of integrand f). We here have a similar case.

Theorem 2.5. *Under the assumptions of Theorem 2.3 we have*

$$(2.33) \quad \left| \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{4\sqrt{3}} C_3,$$

where

$$(2.34) \quad C_3 = \left[\|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - \frac{1}{b-a} \left(f(a) - 2f\left(\frac{a+b}{2}\right) + f(b) \right)^2 \right]^{\frac{1}{2}}.$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 14 of 20

Proof. We define

$$(2.35) \quad \eta(t) = \begin{cases} 1, & t \in [a, \frac{a+b}{2}] \\ -1, & t \in (\frac{a+b}{2}, b] \end{cases},$$

$$(2.36) \quad \eta_0(t) = \frac{\eta(t)}{\|\eta\|_2}.$$

We easily find that


$$(2.37) \quad \int_a^b \eta(t) dt = 0, \quad \|\eta\|_2^2 = b - a.$$

Let $p(t)$ be defined by (2.15). Then we have

$$(2.38) \quad \begin{aligned} \int_a^b p(t)\eta(t)dt &= \int_a^{\frac{a+b}{2}} (t-a)dt - \int_{\frac{a+b}{2}}^b (t-b)dt \\ &= \frac{(b-a)^2}{4}. \end{aligned}$$

We also have

$$(2.39) \quad \int_a^b f'(t)\eta(t)dt = -f(a) + 2f\left(\frac{a+b}{2}\right) - f(b).$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

From (2.37)-(2.39) we get


$$\begin{aligned}
 (2.40) \quad S_{\eta}(f', p) &= \int_a^b p(t)f'(t)dt - \frac{1}{b-a} \int_a^b f'(t)dt \int_a^b p(t)dt \\
 &\quad - \int_a^b p(t)\eta_0(t)dt \int_a^b f'(t)\eta_0(t)dt \\
 &= f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t)dt \\
 &\quad - \frac{b-a}{4} \left[-f(a) + 2f\left(\frac{a+b}{2}\right) - f(b)\right] \\
 &= \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt.
 \end{aligned}$$

From (2.3) and (2.40) it follows that

$$\begin{aligned}
 (2.41) \quad \left| \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt \right| \\
 \leq S_{\eta}(f', f')^{\frac{1}{2}} S_{\eta}(p, p)^{\frac{1}{2}}.
 \end{aligned}$$

We now calculate

$$\begin{aligned}
 (2.42) \quad S_{\eta}(p, p) &= \|p\|_2^2 - \frac{1}{b-a} \left(\int_a^b p(t)dt \right)^2 - \left(\int_a^b p(t)\eta_0(t)dt \right)^2 \\
 &= \frac{(b-a)^3}{48}.
 \end{aligned}$$



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

We also have

$$(2.43) \quad C_3^2 = S_\eta(f', f').$$

From (2.41)-(2.43) we easily get (2.33). \square

Remark 2.4. *It is not difficult to see that (2.33) is better than the first estimation in (1.7).*

Finally, in [12] we can find the next inequality

$$(2.44) \quad \left| \frac{f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)}{6}(b-a) - \int_a^b f(t)dt \right| \leq \frac{(b-a)^2}{12}(\Gamma - \gamma),$$


where $f : I \rightarrow \mathbb{R}$, ($I \subset \mathbb{R}$ is an open interval, $a < b$, $a, b \in I$) is a differentiable function, f' is integrable and there exist constants $\gamma, \Gamma \in \mathbb{R}$ such that $\gamma \leq f'(t) \leq \Gamma$, $t \in [a, b]$.

Inequality (2.44) is a variant of the Simpson's inequality. On the other hand, we have

$$(2.45) \quad \left| \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t)dt \right| \leq \frac{(b-a)^2}{8\sqrt{3}}(\Gamma - \gamma).$$

Inequality (2.45) follows from (2.33), since

$$(2.46) \quad S_\eta(f', f') \leq (b-a) \left(\frac{\Gamma - \gamma}{2} \right)^2$$



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20

and (2.46) follows from (2.4) and (1.4).


From (2.44) and (2.45) we see that the simple 3-point quadrature rule (2.32) has a better estimation of error than the well-known 3-point Simpson quadrature rule. Note that the estimations are expressed in terms of the first derivative f' of integrand.

Finally, the following remark is valid.

Remark 2.5. *The considered case $n = 1$ illustrates how to apply Lemma 2.1 to quadrature formulas. It is also shown that the derived results are better than some recently obtained results. We can use Lemma 2.1 to derive further improvements of the obtained results. However, in such a case we must require*

$$\int_a^b g(t)\Psi_i^0(t)dt = 0, i = 0, 1, 2, \dots, n.$$

Thus, the construction of such a finite sequence $\{\Psi_i^0\}_0^n$ can be complicated. However, if we really need better error bounds, without taking into account possible complications, then we can apply the procedure described in this section.



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 18 of 20

References

- [1] G.A. ANASTASSIOU, Ostrowski type inequalities, *Proc. Amer. Math. Soc.*, **123**(12) (1995), 3775–3781.
- [2] N.S. BARNETT, S.S. DRAGOMIR AND A. SOFO, Better bounds for an inequality of the Ostrowski type with applications, *RGMIA Research Report Collection*, **3**(1) (2000), Article 11.
- [3] P. CERONE AND S.S. DRAGOMIR, Midpoint-type rules from an inequalities point of view, *Handbook of Analytic-Computational Methods in Applied Mathematics*, Editor: G. Anastassiou, CRC Press, New York, (2000), 135–200.
- [4] P. CERONE AND S.S. DRAGOMIR, Trapezoidal-type rules from an inequalities point of view, *Handbook of Analytic-Computational Methods in Applied Mathematics*, Editor: G. Anastassiou, CRC Press, New York, (2000), 65–134.
- [5] S.S. DRAGOMIR, P. CERONE AND J. ROUMELIOTIS, A new generalization of Ostrowski integral inequality for mappings whose derivatives are bounded and applications in numerical integration and for special means, *Appl. Math. Lett.*, **13** (2000), 19–25.
- [6] S.S. DRAGOMIR AND T.C. PEACHY, New estimation of the remainder in the trapezoidal formula with applications, *Stud. Univ. Babeş-Bolyai Math.*, **XLV** (4), (2000), 31–42.
- [7] S.S. DRAGOMIR AND S. WANG, An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special



A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 19 of 20

means and some numerical quadrature rules, *Comput. Math. Appl.*, **33** (1997), 15–20.

- [8] A. GHIZZETTI AND A. OSSICINI, *Quadrature Formulae*, Birkhäuser Verlag, Basel/Stuttgart, 1970.
- [9] M. MATIĆ, J. PEČARIĆ AND N. UJEVIĆ, Improvement and further generalization of some inequalities of Ostrowski-Grüss type, *Comput. Math. Appl.*, **39** (2000), 161–179.
- [10] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Acad. Publ., Dordrecht/Boston/Lancaster/Tokyo, 1991.
- [11] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Acad. Publ., Dordrecht/Boston/Lancaster/Tokyo, 1993.
- [12] C.E.M. PEARCE, J. PEČARIĆ, N. UJEVIĆ AND S. VAROŠANEC, Generalizations of some inequalities of Ostrowski-Grüss type, *Math. Inequal. Appl.*, **3**(1) (2000), 25–34.



**A Generalization of the
Pre-Grüss Inequality and
Applications to some
Quadrature Formulae**

Nenad Ujević

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20