# Journal of Inequalities in Pure and Applied Mathematics

## NEW OSTROWSKI TYPE INEQUALITIES INVOLVING THE PRODUCT OF TWO FUNCTIONS

### B.G. PACHPATTE

57, Shri Niketen Coloney Near Abhinay Talkies Aurangabad-431001 Maharashtra, India

EMail: bgpachpatte@gmail.com



volume 7, issue 3, article 104, 2006.

Received 18 August, 2005; accepted 16 February, 2006.

Communicated by: G.V. Milovanović



©2000 Victoria University ISSN (electronic): 1443-5756 045-06

## Abstract

In this paper we establish new Ostrowski type inequalities involving product of two functions. The analysis used in the proofs is elementary and based on the use of the integral identity recently established by Dedić, Pečarić and Ujević.

#### 2000 Mathematics Subject Classification: 26D10, 26D15.

Key words: Ostrowski type inequalities, Product of two functions, Integral identity, Harmonic sequence.

## Contents

1	Introduction	3
2	Statement of Results	5
3	Proofs of Theorems 2.1 and 2.2	8
References		



New Ostrowski Type Inequalities Involving the Product of Two Functions

#### B.G. Pachpatte



## 1. Introduction

In 1938, Ostrowski [7, p. 468] proved the following inequality:

(1.1) 
$$\left| f(x) - \frac{1}{b-a} \int_{a}^{b} f(t) dt \right| \leq \left[ \frac{1}{4} + \left( \frac{x - \frac{a+b}{2}}{b-a} \right)^{2} \right] (b-a) M,$$

for all  $x \in [a, b]$ , where  $f : [a, b] \to \mathbb{R}$  is a differentiable function such that  $|f'(x)| \le M$  for all  $x \in [a, b]$ .

In 1992, Fink [4] and earlier in 1976, Milovanović and Pečarić [6] have obtained some interesting generalizations of (1.1) in the form

(1.2) 
$$\left| \frac{1}{n} \left( f(x) + \sum_{k=1}^{n-1} F_k(x) \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \le C(n, p, x) \left\| f^{(n)} \right\|_{\infty},$$

where

$$F_k(x) = \frac{n-k}{k!} \cdot \frac{f^{(k-1)}(a) (x-a)^k - f^{(k-1)}(b) (x-b)^k}{b-a},$$

as usual  $\frac{1}{p} + \frac{1}{p'} = 1$  with p' = 1 for  $p = \infty$ ,  $p' = \infty$  for p = 1 and


$$||f||_{p} = \left(\int_{a}^{b} |f(t)|^{p} dt\right)^{\frac{1}{p}}.$$

In fact, Milovanović and Pečarić [6] (see also [7, p. 469]) have proved that

$$C(n, \infty, x) = \frac{(x-a)^{n+1} + (b-x)^{n+1}}{n(n+1)!(b-a)},$$



New Ostrowski Type Inequalities Involving the Product of Two Functions



J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

while Fink [4] (see also [7, p. 473]) proved that the inequality (1.2) holds provided  $f^{(n-1)}$  is absolutely continuous on [a, b] and  $f^{(n)} \in L_p[a, b]$ , with

$$C(n, p, x) = \frac{\left[ (x-a)^{np'+1} + (b-x)^{np'+1} \right]^{\frac{1}{p'}}}{n! (b-a)} B((n-1)p'+1, p'+1)^{\frac{1}{p'}},$$

for 1 , B is the beta function, and

$$C(n, 1, x) = \frac{(n-1)^{n-1}}{n^n n! (b-a)} \max\left[ (x-a)^n, (b-x)^n \right].$$

Recently, Pachpatte [10] and Dedić, Pečarić and Ujević [3] (see also [2]) have given some generalizations of Milovanić-Pečarić [6] and Fink [4] inequalities. Motivated by the results in [10] and [3], in this paper we establish new Ostrowski type inequalities involving the product of two functions. The analysis used in the proofs is based on the integral identity proved in [3] and our results provide new estimates on these types of inequalities.





Page 4 of 13

J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

## 2. Statement of Results

Let  $(P_n)$  be a harmonic sequence of polynomials, that is,  $P'_n = P_{n-1}$ ,  $n \ge 1$ ,  $P_0 = 1$ . Furthermore, let  $I \subset \mathbb{R}$  be a segment and  $h : I \to \mathbb{R}$  be such that  $h^{(n-1)}$  is absolutely continuous for some  $n \ge 1$ . We use the notation

$$L[h(x)] = \frac{1}{n} \left[ h(x) + \sum_{k=1}^{n-1} (-1)^k P_k(x) h^{(k)}(x) + \sum_{k=1}^{n-1} \frac{(-1)^k (n-k)}{b-a} \left[ P_k(a) h^{(k-1)}(a) - P_k(b) h^{(k-1)}(b) \right] \right],$$

to simplify the details of presentation. For n = 1 the above sums are defined to be zero. In a recent paper [3], Dedić, Pečarić and Ujević proved the following identity (see also [2]):

(2.1) 
$$L[h(x)] - \frac{1}{b-a} \int_{a}^{b} h(t) dt = \frac{(-1)^{n+1}}{n(b-a)} \int_{a}^{b} P_{n-1}(t) e(t,x) h^{(n)}(t) dt,$$

where


(2.2) 
$$e(t,x) = \begin{cases} t-a & \text{if } t \in [a,x], \\ t-b & \text{if } t \in (x,b]. \end{cases}$$

For the harmonic sequence of polynomials  $P_k(t) = \frac{(t-x)^k}{k!}$ ,  $k \ge 0$  the identity (2.1) reduces to the main identity derived by Fink in [4] (see also [3, p. 177]).

Our main results are given in the following theorems.



New Ostrowski Type Inequalities Involving the Product of Two Functions



J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

**Theorem 2.1.** Let  $(P_n)$  be a harmonic sequence of polynomials and  $f, g : [a,b] \to \mathbb{R}$  be such that  $f^{(n-1)}, g^{(n-1)}$  are absolutely continuous for some  $n \ge 1$  and  $f^{(n)}, g^{(n)} \in L_p[a,b], 1 \le p \le \infty$ . Then the inequality

$$(2.3) \quad \left| g(x) L[f(x)] + f(x) L[g(x)] - \frac{1}{b-a} \left[ g(x) \int_{a}^{b} f(t) dt + f(x) \int_{a}^{b} g(t) dt \right] \right| \\ \leq D(n, p, x) \left[ |g(x)| \| f^{(n)} \|_{p} + |f(x)| \| g^{(n)} \|_{p} \right],$$

*holds for all*  $x \in [a, b]$ *, where* 

(2.4) 
$$D(n, p, x) = \frac{1}{n(b-a)} \|P_{n-1}e(\cdot, x)\|_{p'},$$

e(t, x) is given by (2.2) and p, p' are as explained in Section 1. **Theorem 2.2.** Let  $(P_n)$ ,  $f, g, f^{(n)}, g^{(n)}$  and p be as in Theorem 2.1. Then the inequality

(2.5) 
$$\begin{aligned} \left| L\left[f\left(x\right)\right] L\left[g\left(x\right)\right] \\ &- \frac{1}{b-a} \left[ L\left[g\left(x\right)\right] \int_{a}^{b} f\left(t\right) dt + L\left[f\left(x\right)\right] \int_{a}^{b} g\left(t\right) dt \right] \\ &+ \left(\frac{1}{b-a} \int_{a}^{b} f\left(t\right) dt\right) \left(\frac{1}{b-a} \int_{a}^{b} g\left(t\right) dt\right) \right| \\ &\leq \left\{ D\left(n,p,x\right) \right\}^{2} \left\| f^{(n)} \right\|_{p} \left\| g^{(n)} \right\|_{p}, \end{aligned}$$



New Ostrowski Type Inequalities Involving the Product of Two Functions



J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

holds for all  $x \in [a, b]$ , where D(n, p, x) and p' are as in Theorem 2.1.

**Remark 1.** If we take g(t) = 1 and hence  $g^{(n-1)}(t) = 0$  for  $n \ge 2$  in Theorem 2.1, then we get a variant of the Ostrowski type inequality given by Dedić, Pečarić and Ujević in [3, p. 180]. We note that the inequality established in Theorem 2.2 is similar to the inequality given by Pachpatte in [9, Theorem 2].



New Ostrowski Type Inequalities Involving the Product of Two Functions

#### B.G. Pachpatte



## 3. Proofs of Theorems 2.1 and 2.2

*Proof of Theorem 2.1.* From the hypotheses we have the following identities (see [3, p. 176]):


(3.1) 
$$L[f(x)] - \frac{1}{b-a} \int_{a}^{b} f(t) dt = \frac{(-1)^{n-1}}{n(b-a)} \int_{a}^{b} P_{n-1}(t) e(t,x) f^{(n)}(t) dt$$

and


(3.2) 
$$L[g(x)] - \frac{1}{b-a} \int_{a}^{b} g(t) dt = \frac{(-1)^{n-1}}{n(b-a)} \int_{a}^{b} P_{n-1}(t) e(t,x) g^{(n)}(t) dt.$$

Multiplying (3.1) and (3.2) by g(x) and f(x) respectively and adding the resulting identities we have

$$(3.3) \quad g(x) L[f(x)] + f(x) L[g(x)] \\ - \frac{1}{b-a} \left[ g(x) \int_{a}^{b} f(t) dt + f(x) \int_{a}^{b} g(t) dt \right] \\ = \frac{(-1)^{n-1}}{n(b-a)} \left[ g(x) \int_{a}^{b} P_{n-1}(t) e(t,x) f^{(n)}(t) dt \\ + f(x) \int_{a}^{b} P_{n-1}(t) e(t,x) g^{(n)}(t) dt \right]$$



$$\left| g(x) L[f(x)] + f(x) L[g(x)] - \frac{1}{b-a} \left[ g(x) \int_{a}^{b} f(t) dt + f(x) \int_{a}^{b} g(t) dt \right] \right|$$



Page 8 of 13

٠

J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

$$\leq \frac{1}{n(b-a)} \left[ |g(x)| \int_{a}^{b} |P_{n-1}(t) e(t,x) f^{(n)}(t)| dt \right. \\ \left. + |f(x)| \int_{a}^{b} |P_{n-1}(t) e(t,x) g^{(n)}(t)| dt \right] \\ \leq \frac{1}{n(b-a)} \left[ |g(x)| \left\{ \int_{a}^{b} |P_{n-1}(t) e(t,x)|^{p'} dt \right\}^{\frac{1}{p'}} \left\{ \int_{a}^{b} |f^{(n)}(t)|^{p} dt \right\}^{\frac{1}{p}} \\ \left. + |f(x)| \left\{ \int_{a}^{b} |P_{n-1}(t) e(t,x)|^{p'} dt \right\}^{\frac{1}{p'}} \left\{ \int_{a}^{b} |g^{(n)}(t)|^{p} dt \right\}^{\frac{1}{p}} \right] \\ = D(n,p,x) \left[ |g(x)| ||f^{(n)}||_{p} + |f(x)| ||g^{(n)}||_{p} \right].$$

The proof of Theorem 2.1 is complete.

*Proof of Theorem* 2.2. Multiplying the left sides and the right sides of (3.1) and (3.2) we get

$$(3.4) \quad L[f(x)] L[g(x)] \\ - \frac{1}{b-a} \left[ L[g(x)] \int_{a}^{b} f(t) dt + L[f(x)] \int_{a}^{b} g(t) dt \right] \\ + \left( \frac{1}{b-a} \int_{a}^{b} f(t) dt \right) \left( \frac{1}{b-a} \int_{a}^{b} g(t) dt \right) \\ = \frac{(-1)^{2n-2}}{n^{2} (b-a)^{2}} \left\{ \int_{a}^{b} P_{n-1}(t) e(t,x) f^{(n)}(t) dt \right\}$$



New Ostrowski Type Inequalities Involving the Product of Two Functions



J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

$$\times \left\{ \int_{a}^{b} P_{n-1}\left(t\right) e\left(t,x\right) g^{\left(n\right)}\left(t\right) dt \right\}.$$

From (3.4) and following the proof of Theorem 2.1 given above with suitable modifications, we get the required inequality in (2.4). The proof of Theorem 2.2 is complete.  $\Box$ 

**Remark 2.** Dividing both sides of (3.3) and (3.4) by (b - a) and integrating the resulting identities with respect to x over [a, b], then using the properties of modulus and Hölder's integral inequality, we get the following inequalities

$$(3.5) \quad \left| \frac{1}{b-a} \int_{a}^{b} \left[ g\left( x \right) L\left[ f\left( x \right) \right] + f\left( x \right) L\left[ g\left( x \right) \right] \right] dx \\ -2 \left( \frac{1}{b-a} \int_{a}^{b} f\left( t \right) dt \right) \left( \frac{1}{b-a} \int_{a}^{b} g\left( t \right) dt \right) \right| \\ \leq \frac{1}{b-a} \int_{a}^{b} D\left( n, p, x \right) \left[ \left| g\left( x \right) \right| \left\| f^{(n)} \right\|_{p} + \left| f\left( x \right) \right| \left\| g^{(n)} \right\|_{p} \right] dx,$$

and

$$(3.6) \quad \left| \frac{1}{b-a} \int_{a}^{b} L\left[f\left(x\right)\right] L\left[g\left(x\right)\right] dx \\ - \left[ \left(\frac{1}{b-a} \int_{a}^{b} L\left[f\left(x\right)\right] dx \right) \left(\frac{1}{b-a} \int_{a}^{b} g\left(x\right) dx \right) \\ + \left(\frac{1}{b-a} \int_{a}^{b} L\left[g\left(x\right)\right] dx \right) \left(\frac{1}{b-a} \int_{a}^{b} f\left(x\right) dx \right) \right]$$



New Ostrowski Type Inequalities Involving the Product of Two Functions



J. Ineq. Pure and Appl. Math. 7(3) Art. 104, 2006 http://jipam.vu.edu.au

$$+\left(\frac{1}{b-a}\int_{a}^{b}f\left(x\right)dx\right)\left(\frac{1}{b-a}\int_{a}^{b}g\left(x\right)dx\right)\Big|$$
$$\leq \frac{1}{b-a}\left\|f^{(n)}\right\|_{p}\left\|g^{(n)}\right\|_{p}\int_{a}^{b}\left\{D\left(n,p,x\right)\right\}^{2}dx.$$

We note that the inequalities obtained in (3.5) and (3.6) are respectively similar to the well known Grüss [5] and Čebyšev [1] inequalities (see also [8]) and we believe that these inequalities are new to the literature.



New Ostrowski Type Inequalities Involving the Product of Two Functions

#### B.G. Pachpatte



## References

- P.L. ČEBYŠEV, Sur les expressions approximatives des intégrales définies par les aures prises entre les mêmes limites, *Proc. Math. Soc. Charkov*, 2 (1882), 93–98.
- [2] Lj. DEDIĆ, M. MATIĆ AND J. PEČARIĆ, On some generalizations of Ostrowski inequality for Lipschitz functions and functions of bounded variation, *Math. Inequal. Appl.*, 3(1) (2000), 1–14.
- [3] Lj. DEDIĆ, J. PEČARIĆ AND N. UJEVIĆ, On generalizations of Ostrowski inequality and some related results, *Czechoslovak Math. J.*, 53(128) (2003), 173–189.
- [4] A.M. FINK, Bounds of the deviation of a function from its avarages, *Czechoslovak Math. J.*, **42**(117) (1992), 289–310.
- [5] G. GRÜSS, Über das maximum des absoluten Betraages von  $\frac{1}{b-a}\int_{a}^{b} f(x) g(x) dx \frac{1}{(b-a)^{2}}\int_{a}^{b} f(x) dx \int_{a}^{b} g(x) dx$ , Math. Z., **39** (1935), 215–226.
- [6] G.V. MILOVANOVIĆ AND J.E. PEČARIĆ, On generalizations of the inequality of A. Ostrowski and related applications, *Univ. Beograd. Publ. Elektrotehn. Fak., Ser. Mat. Fiz.*, No. 544-576 (1976), 155–158.
- [7] D.S. MITRINOVIĆ, J. PEČARIĆ AND A.M. FINK, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, 1991.



New Ostrowski Type Inequalities Involving the Product of Two Functions

B.G. Pachpatte



- [8] D.S. MITRINOVIĆ, J. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [9] B.G. PACHPATTE, New integral inequalities for differentiable functions, *Tamkang J. Math.*, **34**(3) (2003), 249–253.
- [10] B.G. PACHPATTE, On a new generalization of Ostrowski's inequality, J. Inequal. Pure and Appl. Math., 5(2) (2004), Art. 36. [ONLINE: http: //jipam.vu.edu.au/article.php?sid=378]



New Ostrowski Type Inequalities Involving the Product of Two Functions

#### B.G. Pachpatte

