



**MOMENTS INEQUALITIES OF A RANDOM VARIABLE DEFINED OVER A
FINITE INTERVAL**

PRANESH KUMAR

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE,
UNIVERSITY OF NORTHERN BRITISH COLUMBIA,
PRINCE GEORGE,
BC V2N 4Z9, CANADA
kumarp@unbc.ca

Received 01 October, 2001; accepted 03 April, 2002.

Communicated by C.E.M. Pearce

ABSTRACT. Some estimations and inequalities are given for the higher order central moments of a random variable taking values on a finite interval. An application is considered for estimating the moments of a truncated exponential distribution.

Key words and phrases: Random variable, Finite interval, Central moments, Hölder's inequality, Grüss inequality.

2000 *Mathematics Subject Classification.* 60E15, 26D15.

1. INTRODUCTION

Distribution functions and density functions provide complete descriptions of the distribution of probability for a given random variable. However they do not allow us to easily make comparisons between two different distributions. The set of moments that uniquely characterizes the distribution under reasonable conditions are useful in making comparisons. Knowing the probability function, we can determine the moments, if they exist. There are, however, applications wherein the exact forms of probability distributions are not known or are mathematically intractable so that the moments can not be calculated. As an example, an application in insurance in connection with the insurer's payout on a given contract or group of contracts follows a mixture or compound probability distribution that may not be known explicitly. It is this problem that motivates to find alternative estimations for the moments of a probability distribution. Based on the mathematical inequalities, we develop some estimations of the moments of a random variable taking its values on a finite interval.

Set X to denote a random variable whose probability function is $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}_+$ and its associated distribution function $F : [a, b] \rightarrow [0, 1]$.

ISSN (electronic): 1443-5756

© 2002 Victoria University. All rights reserved.

This research was supported by a grant from the Natural Sciences and Engineering Research Council of Canada. Thanks are due to the referee and Prof. Sever Dragomir for their valuable comments that helped in improving the paper.

069-01

Denote by M_r the r^{th} central moment of the random variable X defined as

$$(1.1) \quad M_r = \int_a^b (t - \mu)^r dF, \quad r = 0, 1, 2, \dots,$$

where μ is the mean of the random variable X . It may be noted that $M_0 = 1$, $M_1 = 0$ and $M_2 = \sigma^2$, the variance of the random variable X .

When reference is made to the r^{th} moment of a particular distribution, we assume that the appropriate integral (1.1) converges for that distribution.

2. RESULTS INVOLVING HIGHER MOMENTS

We first prove the following theorem for the higher central moments of the random variable X .

Theorem 2.1. *For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,*

$$(2.1) \quad \int_a^b (b-t)(t-a)^m dF \\ = \sum_{k=0}^m \binom{m}{k} (\mu-a)^k [(b-\mu)M_{m-k} - M_{m-k+1}], \quad m = 1, 2, 3, \dots$$

Proof. Expressing the left hand side of (2.1) as

$$\int_a^b (b-t)(t-a)^m dF = \int_a^b [(b-\mu) - (t-\mu)][(t-\mu) + (\mu-a)]^m dF,$$

and using the binomial expansion

$$[(t-\mu) + (\mu-a)]^m = \sum_{k=0}^m \binom{m}{k} (\mu-a)^k (t-\mu)^{m-k},$$

we get

$$\int_a^b (b-t)(t-a)^m dF \\ = \int_a^b [(b-\mu) - (t-\mu)] \left[\sum_{k=0}^m \binom{m}{k} (\mu-a)^k (t-\mu)^{m-k} \right] dF \\ = \sum_{k=0}^m \binom{m}{k} (b-\mu)(\mu-a)^k \cdot \int_a^b (t-\mu)^{m-k} dF \\ - \sum_{k=0}^m \binom{m}{k} (\mu-a)^k \cdot \int_a^b (t-\mu)^{m-k+1} dF,$$

and hence the theorem. □

In practice numerical moments of order higher than the fourth are rarely considered, therefore, we now derive the results for the first four central moments of the random variable X based on Theorem 2.1.

Corollary 2.2. *For $m = 1, k = 0, 1$ in (2.1), we have*

$$(2.2) \quad \int_a^b (b-t)(t-a) dF = (b-\mu)(\mu-a) - M_2.$$

This is a result in Theorem 1 by Barnett and Dragomir [1].

Corollary 2.3. For $m = 2, k = 0, 1, 2$ in (2.1),

$$(2.3) \quad \int_a^b (b-t)(t-a)^2 dF = (b-\mu)(\mu-a)^2 + [(b-\mu) - 2(\mu-a)]M_2 - M_3.$$

Corollary 2.4. For $m = 3, k = 0, 1, 2, 3$, we have from (2.1)

$$(2.4) \quad \int_a^b (b-t)(t-a)^3 dF = (b-\mu)(\mu-a)^3 + 3(\mu-a)[(b-\mu) - (\mu-a)]M_2 \\ + [(b-\mu) - 3(\mu-a)]M_3 - M_4.$$

3. SOME ESTIMATIONS FOR THE CENTRAL MOMENTS

We apply Hölder's inequality [4] and results of Barnett and Dragomir [1] to derive the bounds for the central moments of the random variable X .

Theorem 3.1. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$, we have

$$(3.1) \quad \int_a^b (b-t)^r (t-a)^s dF \leq \begin{cases} (b-a)^{r+s+1} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \cdot \|f\|_\infty, \\ (b-a)^{2+\frac{1}{q}} [B(rq+1, sq+1)] \cdot \|f\|_p, \end{cases}$$

for $p > 1, \frac{1}{p} + \frac{1}{q} = 1, r, s \geq 0$.

Proof. Let $t = a(1-u) + bu$. Then

$$\int_a^b (b-t)^r (t-a)^s dt = (b-a)^{r+s+1} \cdot \int_0^1 (1-u)^r u^s du.$$

Since $\int_0^1 u^s (1-u)^r du = \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)}$,

$$\int_a^b (b-t)^r (t-a)^s dt = (b-a)^{r+s+1} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)}.$$

Using the property of definite integral,

$$(3.2) \quad \int_a^b (b-t)^s (t-a)^r dF \geq 0, \quad \text{for } r, s \geq 0,$$

we get,

$$\int_a^b (b-t)^s (t-a)^r dF \\ \leq \|f\|_\infty \int_a^b (b-t)^s (t-a)^r dt, \\ = (b-a)^{r+s+1} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \cdot \|f\|_\infty \quad \text{for } r, s \geq 0,$$

the first inequality in (3.1).

Now applying the Hölder's integral inequality,

$$\begin{aligned} & \int_a^b (b-t)^s (t-a)^r dF \\ & \leq \left[\int_a^b f^p(t) dt \right]^{\frac{1}{p}} \left[\int_a^b (b-t)^{sq} (t-a)^{r q} dt \right]^{\frac{1}{q}} \\ & = (b-a)^{2+\frac{1}{q}} [B(rq+1, sq+1)] \cdot \|f\|_p, \end{aligned}$$

the second inequality in (3.1). □

Theorem 3.2. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,

$$\begin{aligned} (3.3) \quad & m(b-a)^{r+s+1} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \\ & \leq \int_a^b (b-t)^s (t-a)^r dF \\ & \leq M(b-a)^{r+s+1} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)}, r, s \geq 0. \end{aligned}$$

Proof. Noting that if $m \leq f \leq M$, a.e. on $[a, b]$, then

$$m(b-t)^s (t-a)^r \leq (b-t)^s (t-a)^r f \leq M(b-t)^s (t-a)^r,$$

a.e. on $[a, b]$ and by integrating over $[a, b]$, we prove the theorem. □

3.1. Bounds for the Second Central Moment M_2 (Variance). It is seen from (2.2) and (3.2) that the upper bound for M_2 , variance of the random variable X , is

$$(3.4) \quad M_2 \leq (b-\mu)(\mu-a).$$

Considering $x = (b-\mu)$ and $y = (\mu-a)$ in the elementary result

$$xy \leq \frac{(x+y)^2}{4}, \quad x, y \in \mathbb{R},$$

we have

$$(3.5) \quad M_2 \leq \frac{(b-a)^2}{4},$$

and thus,

$$(3.6) \quad 0 \leq M_2 \leq (b-\mu)(\mu-a) \leq \frac{(b-a)^2}{4}.$$

From (2.2) and (3.1), we get

$$(b-\mu)(\mu-a) - M_2 \leq \frac{(b-a)^3}{6} \|f\|_\infty,$$

$$(b-\mu)(\mu-a) - M_2 \leq \|f\|_p (b-a)^{2+\frac{1}{q}} [B(q+1, q+1)], \quad p > 1, \quad \frac{1}{p} + \frac{1}{q} = 1.$$

Other estimations for M_2 from (2.2) and (3.1) are

$$m \frac{(b-a)^3}{6} \leq (b-\mu)(\mu-a) - M_2 \leq M \frac{(b-a)^3}{6}, \quad m \leq f \leq M,$$

resulting in

$$(3.7) \quad M_2 \leq (b-\mu)(\mu-a) - m \frac{(b-a)^3}{6}, \quad m \leq f \leq M.$$

3.2. Bounds for the Third Central Moment M_3 . From (2.3) and (3.2), the upper bound for M_3

$$M_3 \leq (b - \mu)(\mu - a)^2 + [(b - \mu) - 2(\mu - a)]M_2.$$

Further we obtain from (2.3) and (3.4),

$$(3.8) \quad M_3 \leq (b - \mu)(\mu - a)(a + b - 2\mu),$$

from (2.3) and (3.5),

$$(3.9) \quad M_3 \leq \frac{1}{4}[(b - \mu)^3 + (b - \mu)(\mu - a)^2 - 2(\mu - a)^3],$$

and from (2.3) and (3.7),

$$(3.10) \quad M_3 \leq (b - \mu)(\mu - a)(a + b - 2\mu) - \frac{m(b - a)^3(b + \mu - 2a)}{6}.$$

3.3. Bounds for the Fourth Central Moment M_4 . The upper bounds for M_4 from (2.4) and (3.2)

$$M_4 \leq (b - \mu)(\mu - a)^3 + 3(\mu - a)[(b - \mu) - (\mu - a)]M_2 + [(b - \mu) - 3(\mu - a)]M_3.$$

Using (2.4), (3.4) and (3.8), we have

$$(3.11) \quad M_4 \leq (b - \mu)(\mu - a)[(b - a)^2 - 3(b - \mu)(\mu - a)],$$

from (2.4), (3.5) and (3.9),

$$(3.12) \quad M_4 \leq \frac{1}{4}[(b - \mu)^4 + 4(b - \mu)^2(\mu - a)^2 - 4(b - \mu)(\mu - a)^3 + 3(\mu - a)^4],$$

and from (2.4), (3.7) and (3.10),

$$(3.13) \quad M_4 \leq (b - \mu)(\mu - a)[(\mu - a)^2 + (a + b - 2\mu)(a + b - 4\mu) + 3(b - \mu)(a + b - 2\mu)] - \frac{m(b - a)^3(a + b - 2\mu)(b - 2a - \mu)}{6}.$$

4. RESULTS BASED ON THE GRÜSS TYPE INEQUALITY

We prove the following theorem based on the pre-Grüss inequality:

Theorem 4.1. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,

$$(4.1) \quad \left| \int_a^b (b - t)^r (t - a)^s f(t) dt - (b - a)^{r+s} \cdot \frac{\Gamma(r + 1)\Gamma(s + 1)}{\Gamma(r + s + 2)} \right| \\ \leq \frac{1}{2}(M - m)(b - a)^{r+s+1} \left[\frac{\Gamma(2r + 1)\Gamma(2s + 1)}{\Gamma(2r + 2s + 2)} - \left(\frac{\Gamma(r + 1)\Gamma(s + 1)}{\Gamma(r + s + 2)} \right)^2 \right]^{\frac{1}{2}},$$

where $m \leq f \leq M$ a.e. on $[a, b]$ and $r, s \geq 0$.

Proof. We apply the following pre-Grüss inequality [4]:

$$(4.2) \quad \left| \int_a^b h(t)g(t)dt - \frac{1}{b - a} \int_a^b h(t) dt \cdot \frac{1}{b - a} \int_a^b g(t)dt \right| \\ \leq \frac{1}{2}(\phi - \gamma) \cdot \left[\frac{1}{b - a} \int_a^b g^2(t)dt - \left(\frac{1}{b - a} \int_a^b g(t)dt \right)^2 \right]^{\frac{1}{2}},$$

provided the mappings $h, g : [a, b] \rightarrow \mathbb{R}$ are measurable, all integrals involved exist and are finite and $\gamma \leq h \leq \phi$ a.e. on $[a, b]$.

Let $h(t) = f(t)$, $g(t) = (b-t)^r(t-a)^s$ in (4.2). Then

$$(4.3) \quad \left| \int_a^b (b-t)^r(t-a)^s f(t) dt - \frac{1}{b-a} \int_a^b f(t) dt \cdot \frac{1}{b-a} \int_a^b (b-t)^r(t-a)^s dt \right| \\ \leq \frac{1}{2}(M-m) \cdot \left[\frac{1}{b-a} \int_a^b \{(b-t)^r(t-a)^s\}^2 dt - \left(\frac{1}{b-a} \int_a^b (b-t)^r(t-a)^s dt \right)^2 \right]^{\frac{1}{2}},$$

where $m \leq f \leq M$ a.e. on $[a, b]$.

On substituting from (3.2) into (4.3), we prove the theorem. \square

Corollary 4.2. For $r = s = 1$ in (4.2),

$$\left| \int_a^b (b-t)(t-a)f(t) dt - \frac{(b-a)^2}{6} \right| \leq \frac{(M-m)(b-a)^3}{12\sqrt{5}},$$

a result (2.7) in Theorem 1 by Barnett and Dragomir [1].

We have the following lemma based on the pre-Grüss inequality:

Lemma 4.3. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,

$$(4.4) \quad \left| \int_a^b (b-t)^r(t-a)^s f(t) dt - (b-a)^{r+s} \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \right| \\ \leq \frac{1}{2}(M-m) \left[(b-a) \int_a^b f^2(t) dt - 1 \right]^{\frac{1}{2}},$$

where $m \leq f \leq M$ a.e. on $[a, b]$ and $r, s \geq 0$.

Proof. We choose $h(t) = (b-t)^r(t-a)^s$, $g(t) = f(t)$ in the pre-Grüss inequality (4.2) to prove this lemma. \square

We now prove the following theorems based on Lemma 4.3:

Theorem 4.4. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,

$$(4.5) \quad \left| \int_a^b (b-t)^r(t-a)^s f(t) dt - (b-a)^{r+s} \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \right| \leq \frac{1}{4}(b-a)(M-m)^2,$$

where $m \leq f \leq M$ a.e. on $[a, b]$ and $r, s \geq 0$.

Proof. Barnett and Dragomir [3] established the following identity:

$$(4.6) \quad \frac{1}{b-a} \int_a^b f(t)g(t) dt = p + \left(\frac{1}{b-a} \right)^2 \cdot \int_a^b f(t) dt \cdot \int_a^b g(t) dt,$$

where

$$|p| \leq \frac{1}{4}(\Gamma - \gamma)(\Phi - \phi), \quad \text{and} \quad \Gamma < f < \gamma, \Phi < g < \phi.$$

By taking $g = f$ in (4.6), we get

$$(4.7) \quad \frac{1}{b-a} \int_a^b f^2(t) dt = p + \left(\frac{1}{b-a} \right)^2, \quad \text{where} \quad |p| \leq \frac{1}{4}(M-m), \quad M < f < m.$$

Thus, (4.4) and (4.7) prove the theorem. \square

Another inequality based on a result from Barnett and Dragomir [3] follows:

Theorem 4.5. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$,

$$(4.8) \quad \left| \int_a^b (b-t)^r (t-a)^s f(t) dt - (b-a)^{r+s} \cdot \frac{\Gamma(r+1)\Gamma(s+1)}{\Gamma(r+s+2)} \right| \leq \frac{1}{4} M(M-m)(b-a),$$

where $m \leq f \leq M$ a.e. on $[a, b]$ and $r, s \geq 0$.

Proof. Barnett and Dragomir [3] have established the following inequality:

$$(4.9) \quad \left| \frac{1}{b-a} \int_a^b f^n(t) dt - \left(\frac{1}{b-a} \right)^n \right| \leq \frac{\Gamma^2}{4(b-a)^{n-2}} \left[\frac{\Gamma^{n-1} \cdot (b-a)^{n-1} - 1}{\Gamma \cdot (b-a) - 1} \right],$$

where $\gamma < f < \Gamma$.

From (4.9), we get

$$\left[\left| \frac{1}{b-a} \int_a^b f^2(t) dt - \left(\frac{1}{b-a} \right)^2 \right| \right]^{\frac{1}{2}} \leq \frac{M}{2}, m \leq f \leq M.$$

and substituting in (4.4) proves the theorem. □

5. RESULTS BASED ON THE HÖLDER'S INTEGRAL INEQUALITY

We consider the Hölder's integral inequality [4] and for $t \in [a, b]$, $\frac{1}{p} + \frac{1}{q} = 1$, $p > 1$,

$$(5.1) \quad \begin{aligned} & \left| \int_a^t (t-u)^n f^{(n+1)}(u) du \right| \\ & \leq \left(\int_a^t |f^{(n+1)}(u)| du \right)^{\frac{1}{p}} \cdot \left(\int_a^t (t-u)^{nq} du \right)^{\frac{1}{q}} \\ & \leq \|f^{(n+1)}\|_p \cdot \left[\frac{(t-a)^{nq+1}}{nq+1} \right]^{\frac{1}{q}}. \end{aligned}$$

On applying (5.1), we have the theorem:

Theorem 5.1. For the random variable X with distribution function $F : [a, b] \rightarrow [0, 1]$, suppose that the density function $f : [a, b]$ is n -times differentiable and $f^{(n)}$ ($n \geq 0$) is absolutely continuous on $[a, b]$. Then,

$$(5.2) \quad \left| \int_a^b (t-a)^r (b-t)^s f(t) dt - \sum_{k=0}^n (b-a)^{r+s+k+1} \cdot \frac{\Gamma(s+1)\Gamma(r+k+1)}{\Gamma(r+s+k+2)} \right|$$

$$\leq \frac{1}{n!} \cdot \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{n+1} \cdot (b-a)^{r+s+n+2} \cdot \frac{\Gamma(r+n+2)\Gamma(s+1)}{\Gamma(r+s+n+3)}, & \text{if } f^{(n+1)} \in L_\infty[a, b], \\ \frac{\|f^{(n+1)}\|_p}{(nq+1)^{1/q}} \cdot (b-a)^{r+s+n+\frac{1}{q}+1} \cdot \frac{\Gamma(r+n+\frac{1}{q}+1)\Gamma(s+1)}{\Gamma(r+s+n+\frac{1}{q}+2)}, & \text{if } f^{(n+1)} \in L_p[a, b], p > 1, \\ \|f^{(n+1)}\|_1 \cdot (b-a)^{r+s+n+1} \cdot \frac{\Gamma(r+n+1)\Gamma(s+1)}{\Gamma(r+s+n+2)}, & \text{if } f^{(n+1)} \in L_1[a, b], \end{cases}$$

where $\|\cdot\|_p$ ($1 \leq p \leq \infty$) are the Lebesgue norms on $[a, b]$, i.e.,

$$\|g\|_\infty := \text{ess sup}_{t \in [a,b]} |g(t)|, \text{ and } \|g\|_p := \left(\int_a^b |g(t)|^p dt \right)^{\frac{1}{p}}, \quad (p \geq 1).$$

Proof. Using the Taylor's expansion of f about a :

$$f(t) = \sum_{k=0}^n \frac{(t-a)^k}{k!} f^{(k)}(a) + \frac{1}{n!} \int_a^t (t-u)^n f^{(n+1)}(u) du, \quad t \in [a, b],$$

we have

$$(5.3) \quad \int_a^b (t-a)^r (b-t)^s f(t) dt = \sum_{k=0}^n \left[\int_a^b (t-a)^{r+k} (b-t)^s dt \cdot \frac{f^{(k)}(a)}{k!} \right] \\ + \left[\frac{1}{n!} \int_a^b (t-a)^r (b-t)^s \left(\int_a^t (t-u)^n f^{(n+1)}(u) du \right) dt \right].$$

Applying the transformation $t = (1-x)a + xb$, we have

$$(5.4) \quad \int_a^b (t-a)^{r+k} (b-t)^s dt = (b-a)^{r+s+k+1} \cdot \frac{\Gamma(s+1)\Gamma(r+k+1)}{\Gamma(r+s+k+2)}.$$

For $t \in [a, b]$, it may be seen that

$$(5.5) \quad \left| \int_a^t (t-u)^n f^{(n+1)}(u) du \right| \leq \int_a^t |(t-u)^n| |f^{(n+1)}(u)| du \\ \leq \sup_{u \in [a, b]} |f^{(n+1)}(u)| \cdot \int_a^t (t-u)^n du \\ \leq \|f^{(n+1)}\|_{\infty} \cdot \frac{(t-a)^{n+1}}{n+1}.$$

Further, for $t \in [a, b]$,

$$(5.6) \quad \left| \int_a^t (t-u)^n f^{(n+1)}(u) du \right| \leq \int_a^t (t-u)^n |f^{(n+1)}(u)| du \\ \leq (t-a)^n \int_a^t |f^{(n+1)}(u)| du \\ \leq \|f^{(n+1)}\| \cdot (t-a)^n.$$

Let

$$(5.7) \quad M(a, b) := \frac{1}{n!} \int_a^b (t-a)^r (b-t)^s \left(\int_a^t (t-u)^n f^{(n+1)}(u) du \right) dt.$$

Then (5.1) and (5.5) to (5.7) result in

$$(5.8) \quad M(a, b) \leq \frac{1}{n!} \cdot \begin{cases} \frac{\|f^{(n+1)}\|_{\infty}}{n+1} \cdot \int_a^b (t-a)^{r+n+1} (b-t)^s dt, & \text{if } f^{(n+1)} \in L_{\infty}[a, b], \\ \frac{\|f^{(n+1)}\|_p}{(nq+1)^{1/q}} \cdot \int_a^b (t-a)^{r+n+\frac{1}{q}} (b-t)^s dt, & \text{if } f^{(n+1)} \in L_p[a, b], \quad p > 1, \\ \|f^{(n+1)}\|_1 \cdot \int_a^b (t-a)^{r+n} (b-t)^s dt, & \text{if } f^{(n+1)} \in L_1[a, b]. \end{cases}$$

Using (5.3), (5.4) and (5.8), we prove the theorem. \square

Corollary 5.2. *Considering $r = s = 1$, the inequality (5.8) leads to*

$$M(a, b) \leq \frac{1}{n!} \begin{cases} \frac{\|f^{(n+1)}\|_{\infty}}{(n+1)} \cdot \frac{(b-a)^{n+4}}{(n+3)(n+4)}, & \text{if } f^{(n+1)} \in L_{\infty}[a, b], \\ \frac{\|f^{(n+1)}\|_p}{(nq+1)^{\frac{1}{q}}} \cdot \frac{(b-a)^{n+\frac{1}{q}+3}}{\left(n+\frac{1}{q}+2\right)\left(n+\frac{1}{q}+3\right)}, & \text{if } f^{(n+1)} \in L_p[a, b], p > 1, \\ \|f^{(n+1)}\|_1 \cdot \frac{(b-a)^{n+3}}{(n+2)(n+3)}, & \text{if } f^{(n+1)} \in L_1[a, b], \end{cases}$$

which is Theorem 3 of Barnett and Dragomir [1].

6. APPLICATION TO THE TRUNCATED EXPONENTIAL DISTRIBUTION

The truncated exponential distribution arises frequently in applications particularly in insurance contracts with caps and deductible and in the field of life-testing. A random variable X with distribution function

$$F(x) = \begin{cases} 1 - e^{-\lambda x} & \text{for } 0 \leq x < c, \\ 1 & \text{for } x \geq c, \end{cases}$$

is a truncated exponential distribution with parameters λ and c .

The density function for X :

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{for } 0 \leq x < c \\ 0 & \text{for } x \geq c \end{cases} + e^{-\lambda c} \cdot \delta_c(x),$$

where δ_c is the delta function at $x = c$. This distribution is therefore mixed with a continuous distribution $f(x) = \lambda e^{-\lambda x}$ on the interval $0 \leq x < c$ and a point mass of size $e^{-\lambda c}$ at $x = c$.

The moment generating function for the random variable X :

$$\begin{aligned} M_X(t) &= \int_0^c e^{tx} \cdot \lambda e^{-\lambda x} dx + e^{tc} \cdot e^{-\lambda c} \\ &= \begin{cases} \frac{\lambda - te^{-c(\lambda-t)}}{\lambda - t}, & \text{for } t \neq \lambda, \\ \lambda c + 1, & \text{for } t = \lambda. \end{cases} \end{aligned}$$

For further calculations in what follows, we assume $t \neq \lambda$. From the moment generating function $M_X(t)$, we have:

$$\begin{aligned} E(X) &= \frac{1 - e^{-\lambda c}}{\lambda}, \\ E(X^2) &= \frac{2[1 - (1 + \lambda c)e^{-\lambda c}]}{\lambda^2}, \\ E(X^3) &= \frac{3[2 - (2 + 2\lambda c + \lambda^2 c^2)e^{-\lambda c}]}{\lambda^3}, \\ E(X^4) &= \frac{4[6 - (6 + 6\lambda c + 3\lambda^2 c^2 + \lambda^3 c^3)e^{-\lambda c}]}{\lambda^4}. \end{aligned}$$

The higher order central moments are:

$$M_k = \sum_{i=0}^k \binom{k}{i} E(X^i) \cdot \mu^{k-i}, \text{ for } k = 2, 3, 4, \dots,$$

in particular,

$$\begin{aligned} M_2 &= \frac{1 - 2\lambda c e^{-\lambda c} - e^{-2\lambda c}}{\lambda^2}, \\ M_3 &= \frac{16 - 3e^{-\lambda c}(10 + 4\lambda c + \lambda^2 c^2) + 6e^{-2\lambda c}(3 + \lambda c) - 4e^{-3\lambda c}}{\lambda^3}, \\ M_4 &= \frac{65 - 4e^{-\lambda c}(32 + 15\lambda c + 6\lambda^2 c^2 + \lambda^3 c^3)}{\lambda^4} \\ &\quad + \frac{3e^{-2\lambda c}(30 + 16\lambda c + 4\lambda^2 c^2) - 4e^{-3\lambda c}(8 + 3\lambda c) + 5e^{-4\lambda c}}{\lambda^4}. \end{aligned}$$

Using the moment-estimation inequality (3.6), the upper bound for M_2 , in terms of the parameters λ and c of the distribution:

$$\hat{M}_2 \leq \frac{(1 - e^{-\lambda c})(\lambda c - 1 + e^{-\lambda c})}{\lambda^2}.$$

The upper bounds for M_3 using (3.8)

$$\hat{M}_3 \leq \frac{(2 - 3\lambda c + \lambda^2 c^2) - e^{-\lambda c}(6 - 6\lambda c + \lambda^2 c^2) + 3e^{-2\lambda c}(2 - \lambda c) - 2e^{-3\lambda c}}{\lambda^3},$$

and using (3.9)

$$\hat{M}_3 \leq \frac{(-3 + 4\lambda c - 3\lambda^2 c^2 + \lambda^3 c^3) + e^{-\lambda c}(9 - 8\lambda c + 3\lambda^2 c^2) - e^{-2\lambda c}(9 - 4\lambda c) + 3e^{-3\lambda c}}{4\lambda^3}.$$

The upper bounds for M_4 using (3.11)

$$\begin{aligned} \hat{M}_4 &\leq \frac{(-3 + 6\lambda c - 4\lambda^2 c^2 + \lambda^3 c^3) + e^{-\lambda c}(12 - 18\lambda c + 8\lambda^2 c^2 - \lambda^3 c^3)}{\lambda^4} \\ &\quad - \frac{2e^{-2\lambda c}(9 + 9\lambda c + 2\lambda^2 c^2) - 6e^{-3\lambda c}(2 - \lambda c) + 3e^{-4\lambda c}}{\lambda^4}, \end{aligned}$$

and from (3.12),

$$\begin{aligned} \hat{M}_4 &\leq \frac{(12 - 16\lambda c + 103\lambda^2 c^2 - 4\lambda^3 c^3 + \lambda^4 c^4) - 4e^{-\lambda c}(12 - 12\lambda c + 5\lambda^2 c^2 - \lambda^3 c^3)}{4\lambda^4} \\ &\quad + \frac{2e^{-2\lambda c}(36 + 24\lambda c + 5\lambda^2 c^2) - 16e^{-3\lambda c}(3 - \lambda c) + 12e^{-4\lambda c}}{4\lambda^4}. \end{aligned}$$

REFERENCES

- [1] N.S.BARNETT AND S.S.DRAGOMIR, Some elementary inequalities for the expectation and variance of a random variable whose pdf is defined on a finite interval, *RGMA Res. Rep. Coll.*, **2**(7) (1999). [ONLINE] <http://rgmia.vu.edu.au/v1n2.html>
- [2] N.S.BARNETT AND S.S.DRAGOMIR, Some further inequalities for univariate moments and some new ones for the covariance, *RGMA Res. Rep. Coll.*, **3**(4) (2000). [ONLINE] <http://rgmia.vu.edu.au/v3n4.html>
- [3] N.S. BARNETT, P. CERONE, S.S. DRAGOMIR AND J. ROUMELIOTIS, Some inequalities for the dispersion of a random variable whose pdf is defined on a finite interval, *J. Ineq. Pure & Appl. Math.*, **2**(1) (2001), 1–18. [ONLINE] <http://jipam.vu.edu.au/v2n1.html>

- [4] J.E. PEČARIĆ, F. PROSCHAN AND Y.L. TONG, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, 1992.