

Journal of Inequalities in Pure and Applied Mathematics

A q -ANALOGUE OF AN INEQUALITY DUE TO KONRAD KNOPP

MEHMET ZEKI SARIKAYA AND HÜSEYİN YILDIRIM

Department of Mathematics
Faculty of Science and Arts
Kocatepe University
Afyon-TURKEY

EMail: sarikaya@aku.edu.tr

EMail: hyildir@aku.edu.tr

©2000 Victoria University
ISSN (electronic): 1443-5756
085-06



volume 7, issue 5, article 178,
2006.

*Received 25 January, 2006;
accepted 22 March, 2006.*

Communicated by: L. Pick

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this note, we obtain some new generalizations of the Hardy's integral inequality by using a fairly elementary analysis. These inequalities generalize some known results and simplify the proofs of some existing results.

2000 Mathematics Subject Classification: 26D10 and 26D15.

Key words: Hardy integral inequalities and Hölder's inequality.

Contents

1	Introduction	3
2	Main Results	5
	References	



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

1. Introduction

In [4], Hardy proved the following inequality. If $p > 1$, $f \geq 0$ and

$$F(x) = \int_0^x f(t)dt,$$

then

$$(1.1) \quad \int_0^\infty \left(\frac{F}{x}\right)^p dx < q^p \int_0^\infty f^p(x)dx$$

unless $f \equiv 0$. The constant $q = p(p-1)^{-1}$ is the best possible. This inequality plays an important role in analysis and its applications. It is obvious that, for parameters a and b such that $0 < a < b < \infty$, the following inequality is also valid

$$(1.2) \quad \int_a^b \left(\frac{F}{x}\right)^p dx < q^p \int_a^b f^p(x)dx,$$

where $0 < \int_0^\infty f^p(x)dx < \infty$. The classical Hardy inequality asserts that if $p > 1$ and f is a nonnegative measurable function on (a, b) , then (1.2) is true unless $f \equiv 0$ a.e. in (a, b) , where the constant here is best possible. This inequality remains true provided that $0 < a < b < \infty$.

In particular, Hardy [3] in 1928 gave a generalized form of the inequality (1.1) when he showed that for any $m \neq 1$, $p > 1$ and any integrable function $f(x) \geq 0$ on $(0, \infty)$ for which

$$F(x) = \begin{cases} \int_0^x f(t)dt & \text{for } m > 1, \\ \int_x^\infty f(t)dt & \text{for } m < 1, \end{cases}$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 3 of 12


then

$$(1.3) \quad \int_0^{\infty} x^{-m} F^p(x) dx < \left(\frac{p}{|m-1|} \right)^p \int_0^{\infty} x^{-m} [xf(x)]^p dx$$

unless $f \equiv 0$, where the constant is also best possible.

Because of their fundamental importance in the discipline over the years much effort and time has been devoted to the improvement and generalization of Hardy's inequalities (1.1), (1.2) and (1.3). These include, among others, the works in [1] – [9].

The objective of this paper is to obtain further generalizations of the classical Hardy integral inequality which will be useful in applications by using some elementary methods of analysis. Throughout this paper, the left-hand sides of the inequalities exist if the right-hand sides exist.



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

2. Main Results

The following theorems are the main results of the present paper.

Theorem 2.1. *Let $p > 1$, $m > 1$ be constants. Let $f(x)$ be a nonnegative and integrable function on $(0, \infty)$ and let $z(x)$ be differentiable in $(0, \infty)$ with $z'(x) > 0$ and $z(0_+) > 0$. Let $w(x)$ and $r(x)$ be positive and absolutely continuous functions on $(0, \infty)$. Let*

$$1 - \frac{1}{m-1} \frac{z(x)}{z'(x)} \frac{w'(x)}{w(x)} + \frac{p}{m-1} \frac{z(x)}{z'(x)} \frac{r'(x)}{r(x)} \geq \frac{1}{\lambda} > 0.$$

Let $a \in (0, \infty)$ be fixed and set

$$F(x) := \frac{1}{r(x)} \int_a^x \frac{r(t)z'(t)f(t)}{z(t)} dt, \quad x \in (0, \infty).$$

Then the inequality

$$(2.1) \quad \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_a^b w(x) \frac{z'(x)}{z^m(x)} f^p(x) dx$$

holds for all $b \geq a$.

Proof. Integrating the left-hand side of inequality (2.1) by parts gives

$$\begin{aligned} & \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \\ &= w(b) \frac{[z(b)]^{-m+1}}{-m+1} F^p(b) + \frac{1}{m-1} \int_a^b z^{-m+1} w' F^p dx \end{aligned}$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 5 of 12

$$-\frac{p}{m-1} \int_a^b z^{-m+1} w \frac{r'(x)}{r(x)} F^p dx + \frac{p}{m-1} \int_a^b w \frac{z'(x)}{z^m(x)} F^{p-1} f dx.$$

Since $m > 1$ and $F(b) \geq 0$, we have


$$\begin{aligned} & \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \left[1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} + \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \right] dx \\ &= w(b) \frac{[z(b)]^{-m+1}}{-m+1} F^p(b) + \frac{p}{m-1} \int_a^b w \frac{z'(x)}{z^m(x)} F^{p-1} f dx \\ &\leq \frac{p}{m-1} \int_a^b w \frac{z'(x)}{z^m(x)} F^{p-1} f dx. \end{aligned}$$

Here, using the assumption on λ , we have

$$\begin{aligned} & \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \frac{1}{\lambda} dx \\ &\leq \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \left[1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} + \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \right] dx \\ &\leq \frac{p}{m-1} \int_a^b w \frac{z'(x)}{z^m(x)} F^{p-1} f dx. \end{aligned}$$

By Hölder's inequality,

$$\int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \frac{\lambda p}{m-1} \left(\int_a^b w \frac{z'(x)}{z^m(x)} F^p dx \right)^{\frac{1}{q}} \left(\int_a^b w \frac{z'(x)}{z^m(x)} f^p dx \right)^{\frac{1}{p}}$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

and thus on simplification, we have

$$\int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_a^b w \frac{z'(x)}{z^m(x)} f^p dx.$$

This proves the theorem. □

Theorem 2.2. Let p, m, f, z, z', w and r be as in Theorem 2.1. Let

$$1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} - \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \geq \frac{1}{\lambda} > 0.$$

Let $a \in (0, \infty)$ be fixed and set

$$F(x) := r(x) \int_a^x \frac{z'(t) f(t)}{z(t) r(t)} dt, \quad x \in (0, \infty).$$

Then the inequality


$$(2.2) \quad \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_a^b w(x) \frac{z'(x)}{z^m(x)} f^p(x) dx$$

holds for all $b \geq a$.

Proof. This is similar to the proof of the Theorem 2.1. □

Theorem 2.3. Let p, m, f, z, z', w and r be as in Theorem 2.1. Let

$$1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} + \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \geq \frac{1}{\lambda} > 0.$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12

Let $a \in (0, \infty)$ be fixed and set

$$F(x) := \frac{1}{r(x)} \int_{\frac{x}{2}}^x \frac{r(t)z'(t)f(t)}{z(t)} dt, \quad x \in (0, \infty).$$

Then the inequality

$$(2.3) \quad \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_0^b w(x) \frac{[z(x)]^{p-m}}{[z'(x)]^{\frac{p}{q}}} |g(x)|^p dx$$

holds where


$$g(x) = \frac{1}{r(x)} \left(\frac{r(x)z'(x)f(x)}{z(x)} - \frac{1}{2} \frac{r(\frac{x}{2})z'(\frac{x}{2})f(\frac{x}{2})}{z(\frac{x}{2})} \right).$$

Proof. Upon integrating by parts we have

$$\begin{aligned} & \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \\ &= w(b) \frac{[z(b)]^{-m+1}}{-m+1} F^p(b) + \frac{1}{m-1} \int_0^b z^{-m+1} w' F^p dx \\ & - \frac{p}{m-1} \int_0^b z^{-m+1} w \frac{r'(x)}{r(x)} F^p dx + \frac{p}{m-1} \int_0^b z^{-m+1} w F^{p-1} |g(x)| dx, \end{aligned}$$

where

$$g(x) = \frac{1}{r(x)} \left(\frac{r(x)z'(x)f(x)}{z(x)} - \frac{1}{2} \frac{r(\frac{x}{2})z'(\frac{x}{2})f(\frac{x}{2})}{z(\frac{x}{2})} \right).$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12

Since $m > 1$ and $F(b) \geq 0$ we have


$$\begin{aligned} & \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \left[1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} + \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \right] dx \\ &= w(b) \frac{[z(b)]^{-m+1}}{-m+1} F^p(b) + \frac{p}{m-1} \int_0^b w \frac{z'(x)}{z^m(x)} F^{p-1} f dx \\ &\leq \frac{p}{m-1} \int_0^b z^{-m+1} w F^{p-1} |g(x)| dx. \end{aligned}$$

By the assumption on λ , we have

$$\begin{aligned} & \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \frac{1}{\lambda} dx \\ &\leq \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) \left[1 - \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} + \frac{p}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \right] dx \\ &\leq \frac{p}{m-1} \int_0^b \frac{z(x)}{z'(x)} w \frac{z'(x)}{z^m(x)} F^{p-1} |g(x)| dx. \end{aligned}$$

By Hölder's inequality

$$\begin{aligned} & \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \\ &\leq \frac{\lambda p}{m-1} \left(\int_0^b w \frac{z'(x)}{z^m(x)} F^p dx \right)^{\frac{1}{q}} \left(\int_0^b w \frac{z^{p-m}(x)}{[z'(x)]^{\frac{p}{q}}} |g(x)|^p dx \right)^{\frac{1}{p}} \end{aligned}$$



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 9 of 12

and thus on simplification, we have

$$\int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_0^b w \frac{z^{p-m}(x)}{[z'(x)]^{\frac{p}{q}}} |g(x)|^p dx.$$

This proves the theorem. □

Theorem 2.4. Let p, m, f, z, z', w and r be as in Theorem 2.1. Let

$$1 - \frac{1}{m-1} \frac{z(x)}{z'(x)} \frac{w'(x)}{w(x)} + \frac{p}{m-1} \frac{z(x)}{z'(x)} \frac{r'(x)}{r(x)} \geq \frac{1}{\lambda} > 0.$$

Let $a \in (0, \infty)$ be fixed and set

$$F(x) := r(x) \int_{\frac{x}{2}}^x \frac{z'(t)f(t)}{z(t)r(t)} dt, \quad x \in (0, \infty).$$


Then the inequality

$$(2.4) \quad \int_0^b w(x) \frac{z'(x)}{z^m(x)} F^p(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_0^b w(x) \frac{[z(x)]^{p-m}}{[z'(x)]^{\frac{p}{q}}} |g(x)|^p dx$$

holds where

$$g(x) = r(x) \left(\frac{r(x)z'(x)f(x)}{z(x)} - \frac{1}{2} \frac{r(\frac{x}{2})z'(\frac{x}{2})f(\frac{x}{2})}{z(\frac{x}{2})} \right).$$

Proof. This is similar to the proof of the Theorem 2.3. □



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 10 of 12

Theorem 2.5. Let $p > q > 0$, $\frac{1}{p} + \frac{1}{q} = 1$, and $m < 1$ be real numbers. Let $z(x)$ be differentiable in $(0, \infty)$ with $z'(x) > 0$ and $z(0_+) > 0$, let $w(x)$ and $r(x)$ be positive and absolutely continuous functions on $(0, \infty)$, n and let $f : [0, \infty) \rightarrow [0, \infty)$ be integrable so that

$$1 + \frac{1}{m-1} \frac{z(x) w'(x)}{z'(x) w(x)} - \frac{p}{q} \frac{1}{m-1} \frac{z(x) r'(x)}{z'(x) r(x)} \geq \frac{1}{\lambda} > 0$$

a.e. for some $\lambda > 0$. Let $b \in (0, \infty)$ be fixed and set

$$F(x) := \frac{1}{r(x)} \int_x^b \frac{r(t) z'(t) f(t)}{z(t)} dt, \quad x \in (0, \infty).$$

Then the following inequality

$$(2.5) \quad \int_a^b w(x) \frac{z'(x)}{z^m(x)} F^{\frac{p}{q}}(x) dx \leq \left(\frac{\lambda p}{m-1} \right)^p \int_a^b w(x) \frac{z'(x)}{z^m(x)} f^{\frac{p}{q}}(x) dx$$

holds for all $0 \leq a \leq b$.

Proof. This is similar to the proof of the Theorem 2.1. □



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 11 of 12

References

- [1] W.S. CHEUNG, Z. HANJS AND J. PEČARIĆ, Some Hardy-type inequalities, *J. Math. Anal. Appl.*, **250**(2) (2000), 621–634.
- [2] G.H. HARDY, J.E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge University Press, (1952).
- [3] G.H. HARDY, Notes on some points in the integral calculus, *Messenger Math.*, **57** (1928), 12–16.
- [4] G.H. HARDY, Notes on a theorem of Hilbert, *Math. Z.*, **6** (1920), 314–317.
- [5] B.G. PACHPATTE, On some variants of Hardy’s inequality, *J. Math. Anal. Appl.*, **124** (1987), 495–501.
- [6] B.G. PACHPATTE, On some generalizations of Hardy’s integral inequality, *J. Math. Anal. Appl.*, **234**(1) (1999), 15–30.
- [7] B.G. PACHPATTE, On a new class of Hardy type inequalities, *Proc. R. Soc. Edin.*, **105A** (1987), 265–274.
- [8] B. YANG, Z. ZENG AND L. DEBNATH, On New Generalizations of Hardy’s Integral Inequality, *J. Math. Anal. Appl.*, **217**(1) (1998), 321–327.
- [9] J.E. PEČARIĆ AND E.R. LOVE, Still more generalizations of Hardy’s inequality, *J. Austral. Math. Soc. Ser. A*, **58** (1995), 1–11.



Some Hardy Type Integral Inequalities

Mehmet Zeki Sarikaya and
Hüseyin Yildirim

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12