

A GENERALIZATION OF CONSTANTIN'S INTEGRAL INEQUALITY AND ITS DISCRETE ANALOGUE

EN-HAO YANG AND MAN-CHUN TAN

Department of Mathematics
Jinan University
Guangzhou 510632, People's Republic of China
EMail: tanmc@jnu.edu.cn

Received: 21 March, 2007

Accepted: 08 May, 2007

Communicated by: S.S. Dragomir

2000 AMS Sub. Class.: 26D10, 26D15, 39A12, 45D05.

Key words: Nonlinear integral inequality, Discrete analogue, Bound on solutions.

Abstract: A generalization of Constantin's integral inequality and its discrete analogue are established. A discrete analogue of Okrasinsky's model for the infiltration phenomena of a fluid is also discussed to convey the usefulness of the discrete inequality obtained.

Acknowledgements: The work that is described in this paper was jointly supported by grants from the National Natural Science Foundation of China (No. 50578064), the Natural Science Foundation of Guangdong Province, China (No.06025219), and the Science Foundation of Key Discipline of the State Council Office of Overseas Chinese Affairs of China.





Integral Inequality and
Its Discrete Analogue

En-hao Yang and Man-chun Tan

vol. 8, iss. 2, art. 57, 2007

[Title Page](#)

[Contents](#)



Page 1 of 20

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Nonlinear Integral Inequality	5
3	Discrete Analogue	11
4	Discrete Model of Infiltration	17



**Integral Inequality and
Its Discrete Analogue**

En-hao Yang and Man-chun Tan

vol. 8, iss. 2, art. 57, 2007

Title Page

Contents



Page 2 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 20

Go Back

Full Screen

Close

1. Introduction

L. Ou-Iang [9] studied the boundedness of solutions for some nonautonomous second order linear differential equations by means of a nonlinear integral inequality. This integral inequality had been frequently used by authors to obtain global existence, uniqueness and stability properties of various nonlinear differential equations. A number of generalizations and discrete analogues of this inequality and their new applications have appeared in the literature. See, for example, B.G. Pachpatte ([10] – [12]) and the present author [13][14] and the references given therein.

In 1996, A. Constantin [2] established the following interesting alternative result for a generalized Ou-Iang type integral inequality given by B. G. Pachpatte [12]:

Theorem A. *Let $T > 0, k \geq 0$, and $u, f, g \in C([0, T], \mathbb{R}_+)$, $\mathbb{R}_+ = [0, \infty)$. Further, let $w \in C(\mathbb{R}_+, \mathbb{R}_+)$ be nondecreasing, $w(r) > 0$ for $r > 0$ and $\int_{r_0}^{\infty} \frac{ds}{w(s)} = \infty$ hold for some number $r_0 > 0$. Then the integral inequality*

$$u^2(t) \leq k^2 + 2 \int_0^t \left\{ f(s)u(s) \left[u(s) + \int_0^s g(\xi)w(u(\xi)) d\xi \right] \right\} ds, \quad t \in [0, T]$$

implies

$$u(t) \leq k + \int_0^t f(s)G^{-1} \left\{ G(k) + \int_0^s [f(\xi) + g(\xi)] d\xi \right\} ds, \quad t \in [0, T],$$


where G^{-1} denotes the inverse function of G and

$$G(r) := \int_{r_0}^r \frac{ds}{w(s) + s}, \quad r \geq r_0, \quad 1 > r_0 > 0.$$

Applying the above result and a topological transversality theorem, A. Granas [4] proved a nonlocal existence theorem for a certain class of initial value problems of

nonlinear integrodifferential equations. We refer to D. O'Regan and M. Meehan [6] for more existence results obtained by means of topological transversality theorems.

The purpose of the present paper is to obtain a new generalization of Constantin's inequality and its discrete analogue. The integral inequality obtained can be used to study some more general initial value problems by following the same argument as that applied in Constantin [2]. A discrete analogue of W.Okrasinsky's mathematical model for the infiltration phenomena of a fluid (see [7] and [8]) is discussed to convey the usefulness of the discrete inequality given in the paper.



**Integral Inequality and
Its Discrete Analogue**

En-hao Yang and Man-chun Tan

vol. 8, iss. 2, art. 57, 2007

Title Page

Contents



Page 4 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 5 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

2. Nonlinear Integral Inequality

Theorem 2.1. Let $u, c \in C(\mathbb{R}_+, \mathbb{R}_+)$ with c nondecreasing, and $\varphi \in C^1(\mathbb{R}_+, \mathbb{R}_+)$ with φ' nonnegative and nondecreasing. Let $f(t, \xi), g(t, \xi), h(t, \xi) \in C(\mathbb{R}_+ \times \mathbb{R}_+, \mathbb{R}_+)$ be nondecreasing in t for every ξ fixed. Further, let $w \in C(\mathbb{R}_+, \mathbb{R}_+)$ be nondecreasing, $w(r) > 0$ for $r > 0$ and $\int_{r_0}^{\infty} \frac{ds}{w(s)} = \infty$ hold for some number $r_0 > 0$. Then the integral inequality

$$(2.1) \quad \varphi[u(t)] \leq c(t) + \int_0^t \left\{ f(t, s)\varphi'[u(s)] \right. \\ \left. \times \left[u(s) + \int_0^s g(s, \xi)w(u(\xi))d\xi \right] + h(t, s)\varphi'[u(s)] \right\} ds, \quad t \in [0, T],$$

implies

$$(2.2) \quad u(t) \leq K(t) + \int_0^t f(t, s) \\ \times G^{-1} \left\{ G(K(t)) + \int_0^s [f(t, \xi) + g(t, \xi)] d\xi \right\} ds, \quad t \in [0, T],$$

herein

$$(2.3) \quad K(t) = \varphi^{-1}[c(t)] + \int_0^t h(t, s)ds,$$

G^{-1}, φ^{-1} denote the inverse function of G, φ , respectively, and

$$(2.4) \quad G(r) := \int_{r_0}^r \frac{ds}{w(s) + s}, \quad r \geq r_0, \quad 1 > r_0 > 0.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 6 of 20

Go Back

Full Screen

Close

Note that, by Constatin [1] the above function G is positive, strictly increasing and satisfies the condition $G(r) \rightarrow \infty$ as $r \rightarrow \infty$.

Proof. Letting $t = 0$ in (2.1), we observe that inequality (2.2) holds trivially for $t = 0$. Fixing an arbitrary number $t_0 \in (0, T)$, we define on $[0, t_0]$ a positive function $z(t)$ by

$$(2.5) \quad z(t) = c(t_0) + \varepsilon + \int_0^t \left\{ f(t_0, s) \varphi' [u(s)] \right. \\ \left. \times \left[u(s) + \int_0^s g(t_0, \xi) w(u(\xi)) d\xi \right] + h(t_0, s) \varphi' [u(s)] \right\} ds,$$

where $\varepsilon > 0$ is an arbitrary small constant. By inequality (2.1) we have

$$(2.6) \quad u(t) \leq \varphi^{-1} [z(t)], \quad t \in [0, t_0].$$

From (2.5) we derive by differentiation

$$z'(t) = f(t_0, t) \varphi' [u(t)] \left[u(t) + \int_0^t g(t_0, \xi) w [u(\xi)] d\xi \right] + h(t_0, t) \varphi' [u(t)] \\ \leq \varphi' [\varphi^{-1} [z(t)]] \left\{ f(t_0, t) \left[\varphi^{-1} [z(t)] + \int_0^t g(t_0, \xi) w (\varphi^{-1} [z(\xi)]) d\xi \right] + h(t_0, t) \right\},$$

for $t \in [0, t_0]$, since φ' is nonnegative and nondecreasing. Hence we obtain

$$\frac{d}{dt} \varphi^{-1} [z(t)] = \frac{z'(t)}{\varphi' [\varphi^{-1} [z(t)]]} \\ \leq f(t_0, t) \left[\varphi^{-1} [z(t)] + \int_0^t g(t_0, \xi) w (\varphi^{-1} [z(\xi)]) d\xi \right] + h(t_0, t), \quad t \in [0, t_0],$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 7 of 20

Go Back

Full Screen

Close

Integrating both sides of the last relation from 0 to t , we get

$$\begin{aligned} \varphi^{-1}[z(t)] &\leq \varphi^{-1}[z(0)] + \int_0^{t_0} h(t_0, s) ds \\ &+ \int_0^t f(t_0, s) \left[\varphi^{-1}[z(s)] + \int_0^s g(t_0, \xi) w(\varphi^{-1}[z(\xi)]) d\xi \right] ds, \quad t \in [0, t_0]. \end{aligned}$$

Define a function $v(t)$, $0 \leq t \leq t_0$, by the right member of the last relation, we have

$$(2.7) \quad \varphi^{-1}[z(t)] \leq v(t), \quad t \in [0, t_0],$$

where

$$(2.8) \quad v(0) = \varphi^{-1}[z(0)] + \int_0^{t_0} h(t_0, s) ds.$$

By differentiation we derive

$$(2.9) \quad \begin{aligned} v'(t) &= f(t_0, t) \left[\varphi^{-1}[z(t)] + \int_0^t g(t_0, \xi) w(\varphi^{-1}[z(\xi)]) d\xi \right] \\ &\leq f(t_0, t) \left[v(t) + \int_0^t g(t_0, \xi) w(v(\xi)) d\xi \right] = f(t_0, t) \Omega(t), \quad t \in [0, t_0]. \end{aligned}$$

where

$$\Omega(t) = \left[v(t) + \int_0^t g(t_0, \xi) w(v(\xi)) d\xi \right].$$

Hence we have $v(t) \leq \Omega(t)$,

$$(2.10) \quad \Omega(0) = v(0) = \varphi^{-1}[z(0)] + \int_0^{t_0} h(t_0, s) ds,$$

and

$$\begin{aligned}\Omega'(t) &= v'(t) + g(t_0, t)w(v(t)) \\ &\leq f(t_0, t)\Omega(t) + g(t_0, t)w(\Omega(t)), \quad t \in [0, t_0].\end{aligned}$$

Because $\Omega(t)$, and hence $w(\Omega(t))$, is positive on $[0, t_0]$, the last inequality can be rewritten as

$$(2.11) \quad \frac{\Omega'(t)}{\Omega(t) + w(\Omega(t))} \leq f(t_0, t) + g(t_0, t), \quad t \in [0, t_0].$$

Integrating both sides of the last relation from 0 to t and in view of the definition of G , we obtain


$$G[\Omega(t)] - G[\Omega(0)] \leq \int_0^t [f(t_0, s) + g(t_0, s)] ds, \quad t \in [0, t_0].$$

By (2.10) and the fact that $G(r) \rightarrow \infty$ as $r \rightarrow \infty$, the last relation yields

$$\Omega(t) \leq G^{-1} \left\{ G \left[\varphi^{-1}[z(0)] + \int_0^{t_0} h(t_0, s) ds \right] + \int_0^t [f(t_0, s) + g(t_0, s)] ds \right\}, \quad t \in [0, t_0].$$

Substituting the last relation into (2.9), then integrating from 0 to t , we derive for $t \in [0, t_0]$ that

$$\begin{aligned}u(t) &\leq \varphi^{-1}[c(t_0) + \varepsilon] + \int_0^{t_0} h(t_0, s) ds \\ &\quad + \int_0^t f(t_0, s) G^{-1} \left\{ G \left[\varphi^{-1}[c(t_0) + \varepsilon] + \int_0^{t_0} h(t_0, s) ds \right] \right. \\ &\quad \left. + \int_0^s [f(t_0, \xi) + g(t_0, \xi)] d\xi \right\} ds,\end{aligned}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 8 of 20

Go Back

Full Screen

Close



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 9 of 20

Go Back

Full Screen

Close

where we used the relation $u(t) \leq \varphi^{-1}[z(t)] \leq v(t) \leq \Omega(t)$.

Taking $t = t_0$ and letting $\varepsilon \rightarrow 0$, from the last relation we have

$$u(t_0) \leq K(t_0) + \int_0^{t_0} f(t_0, s) G^{-1} \left\{ G[K(t_0)] + \int_0^s [f(t_0, \xi) + g(t_0, \xi)] d\xi \right\} ds,$$

where $K(t)$ is defined by (2.3). This means that the desired inequality (2.2) is valid when $t = t_0$. Since the choice of t_0 from $(0, T]$ is arbitrary, the proof of Theorem 2.1 is complete. \square

If $w(r) = r$ holds in Theorem 2.1, the inequality (2.11) can be replaced by the following sharper relation

$$\frac{\Omega'(t)}{\Omega(t)} \leq f(t_0, t) + g(t_0, t), \quad t \in [0, t_0],$$

and functions G, G^{-1} can be replaced by $H(r) = \ln(r/r_0)$, $H^{-1}(\eta) = r_0 e^\eta$, respectively. Hence we derive the following:

Corollary 2.2. *Under the conditions of Theorem 2.1, the integral inequality*

$$(2.12) \quad \varphi[u(t)] \leq c(t) + \int_0^t \left\{ f(t, s) \varphi'[u(s)] \right. \\ \left. \times \left[u(s) + \int_0^s g(s, \xi) u(\xi) d\xi \right] + h(t, s) \varphi'[u(s)] \right\} ds, \quad t \in [0, T],$$

implies

$$(2.13) \quad u(t) \leq \left(\varphi^{-1}[c(t)] + \int_0^t h(t, s) ds \right) \\ \times \left(1 + \int_0^t f(t, s) \left\{ \exp \int_0^s [f(t, \xi) + g(t, \xi)] d\xi \right\} ds \right), \quad t \in [0, T].$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

If $\varphi(\eta) = \eta^p$, $p > 1$, $c(t) = k^p \geq 0$ and $f(t, s), g(t, s), h(t, s)$ do not depend on the variable t , by Theorem 2.1 we have the following:


Corollary 2.3. Let $p > 1, k \geq 0$ be constants and $u, f, g \in C([0, T], \mathbb{R}_+)$. Then the integral inequality

$$u^p(t) \leq k^p + p \int_0^t \left\{ f(s)u^{p-1}(s) \left[u(s) + \int_0^s g(\xi)w(u(\xi)) d\xi \right] \right\} ds, \quad t \in [0, T]$$

implies

$$u(t) \leq k + \int_0^t f(s)G^{-1} \left\{ G(k) + \int_0^s [f(\xi) + g(\xi)] d\xi \right\} ds, \quad t \in [0, T].$$

Remark 1. Clearly, Constantin's Theorem A is the special case $p = 2$ of the last result.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 11 of 20

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

3. Discrete Analogue

In this section we will establish a discrete analogue of Theorem 2.1. Denote by \mathbb{N} the set of nonnegative integers and let $\mathbb{N}_0 = \{n \in \mathbb{N}: n \leq M\}$ for some natural number M . For simplicity, we denote by $K(P, Q)$ the class of functions defined on set P with range in set Q . For a function $u \in K(\mathbb{N}, \mathbb{R})$, $\mathbb{R} = (-\infty, \infty)$, we define the forward difference operator Δ by $\Delta u(n) = u(n+1) - u(n)$.

As usual, we suppose that the empty sum and empty product are zero and one, respectively. For instance,

$$\sum_{s=0}^{-1} p(s) = 0 \quad \text{and} \quad \prod_{s=0}^{-1} p(s) = 1$$

hold for any function $p(n)$, $n \in \mathbb{N}$.

Theorem 3.1. *Let the functions w, φ be as defined in Theorem 2.1 and $u, c \in K(\mathbb{N}, \mathbb{R}_+)$ with $c(n)$ nondecreasing. Further, let $f(n, s), g(n, s), h(n, s) \in K(\mathbb{N} \times \mathbb{N}, \mathbb{R}_+)$ be nondecreasing with respect to n for every s fixed. Then the discrete inequality*

$$(3.1) \quad \varphi[u(n)] \leq c(n) + \sum_{s=0}^{n-1} \left\{ f(n, s) \varphi'[u(s)] \right. \\ \left. \times \left[u(s) + \sum_{\xi=0}^{s-1} g(s, \xi) w(u(\xi)) \right] + h(n, s) \varphi'[u(s)] \right\}, \quad n \in \mathbb{N}_0,$$

implies

$$(3.2) \quad u(n) \leq L(n) + \sum_{s=0}^{n-1} f(n, s) G^{-1} \left\{ G[L(n)] + \sum_{\xi=0}^{s-1} [f(n, \xi) + g(n, \xi)] \right\}, \quad n \in \mathbb{N}_0,$$



Title Page

Contents



Page 12 of 20

Go Back

Full Screen

Close

where G, G^{-1} are as defined in Theorem 2.1 and

$$(3.3) \quad L(n) := \varphi^{-1}[c(n)] + \sum_{s=0}^{n-1} h(n, s).$$

Proof. Fixing an arbitrary positive integer $m \in (0, M)$, we define on the set $J := \{0, 1, \dots, m\}$ a positive function $z(n) \in K(J, (0, \infty))$ by

$$z(n) = c(m) + \varepsilon + \sum_{s=0}^{n-1} \left\{ f(m, s)\varphi'[u(s)] \right. \\ \left. \times \left[u(s) + \sum_{\xi=0}^{s-1} g(m, \xi)w(u(\xi)) \right] + h(m, s)\varphi'[u(s)] \right\},$$

where ε is an arbitrary positive constant, then $z(0) = c(m) + \varepsilon > 0$ and by (3.1) we have

$$(3.4) \quad u(n) \leq \varphi^{-1}[z(n)], \quad n \in J.$$

Using the last relation, we derive

$$\Delta z(n) = f(m, n)\varphi'[u(n)] \left[u(n) + \sum_{s=0}^{n-1} g(m, s)w(u(s)) \right] + h(m, n)\varphi'[u(n)] \\ \leq \varphi' [\varphi^{-1}[z(n)]] \\ \times \left\{ f(m, n) \left[\varphi^{-1}[z(n)] + \sum_{s=0}^{n-1} g(m, s)w(\varphi^{-1}[z(s)]) \right] + h(m, n) \right\}, \quad n \in J.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 20

Go Back

Full Screen

Close

By the mean value theorem and the last relation, we obtain

$$\begin{aligned} \Delta\varphi^{-1}[z(n)] &\leq \frac{\Delta z(n)}{\varphi'[\varphi^{-1}[z(n)]]} \\ &\leq f(m, n) \left[\varphi^{-1}[z(n)] + \sum_{s=0}^{n-1} g(m, s)w(\varphi^{-1}[z(s)]) \right] + h(m, n), \quad n \in J, \end{aligned}$$

since φ^{-1} and $z(n)$ are nondecreasing. Substituting $n = \xi$ in the last relation and then summing over $\xi = 0, 1, 2, \dots, n - 1$, we obtain


$$\begin{aligned} \varphi^{-1}[z(n)] &\leq \varphi^{-1}[z(0)] + \sum_{\xi=0}^{m-1} h(m, \xi) \\ &\quad + \sum_{\xi=0}^{n-1} f(m, \xi) \left[\varphi^{-1}[z(\xi)] + \sum_{s=0}^{\xi-1} g(m, s)w(\varphi^{-1}[z(s)]) \right], \end{aligned}$$

where $n \in J$, since $h(n, s)$ is nonnegative and $m \geq n$ holds. Now, defining by $v(n)$ the right member of the last relation, we have

$$v(0) = \varphi^{-1}[z(0)] + \sum_{\xi=0}^{m-1} h(m, \xi) = \varphi^{-1}[c(m) + \varepsilon] + \sum_{\xi=0}^{m-1} h(m, \xi)$$

and

$$(3.5) \quad \varphi^{-1}[z(n)] \leq v(n), \quad n \in J.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 14 of 20

Go Back

Full Screen

Close

By (3.5) we easily derive

$$\begin{aligned}\Delta v(n) &= f(m, n) \left[\varphi^{-1}[z(n)] + \sum_{s=0}^{n-1} g(m, s)w(\varphi^{-1}[z(s)]) \right] \\ &\leq f(m, n) \left[v(n) + \sum_{s=0}^{n-1} g(m, s)w(v(s)) \right], \quad n \in J,\end{aligned}$$

or

$$(3.6) \quad \Delta v(n) \leq f(m, n)y(n), \quad n \in J,$$

where

$$y(n) := v(n) + \sum_{s=0}^{n-1} g(m, s)w(v(s)), \quad n \in J.$$

Clearly, $y(0) = v(0)$ holds and by (3.6) we have

$$\Delta y(n) \leq \Delta v(n) + g(m, n)w(v(n)) \leq [f(m, n) + g(m, n)] [y(n) + w(y(n))],$$

i.e.,

$$\frac{\Delta y(n)}{y(n) + w(y(n))} \leq f(m, n) + g(m, n), \quad n \in J.$$

Because $y(n), w(r)$ are positive and nondecreasing, we have

$$\int_{y(0)}^{y(n)} \frac{ds}{s + w(s)} \leq \sum_{s=0}^{n-1} \frac{\Delta y(s)}{y(s) + w(y(s))} \leq \sum_{s=0}^{n-1} [f(m, s) + g(m, s)],$$

or

$$G[y(n)] - G[y(0)] \leq \sum_{s=0}^{n-1} [f(m, s) + g(m, s)], \quad n \in J.$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 15 of 20

[Go Back](#)

[Full Screen](#)

[Close](#)

Since $G(r) \rightarrow \infty$ as $r \rightarrow \infty$, the last relation yields

$$y(n) \leq G^{-1} \left\{ G \left[\varphi^{-1}[c(m) + \varepsilon] + \sum_{\xi=0}^{m-1} h(m, \xi) \right] + \sum_{s=0}^{n-1} [f(m, s) + g(m, s)] \right\}, \quad n \in J.$$

Substituting this relation into (3.6), setting $n = s$ and then summing over $s = 0, 1, \dots, n-1$, we have

$$v(n) \leq v(0) + \sum_{s=0}^{n-1} f(m, s) \\ \times G^{-1} \left\{ G \left[\varphi^{-1}[c(m) + \varepsilon] + \sum_{\xi=0}^{m-1} h(m, \xi) \right] + \sum_{\xi=0}^{s-1} [f(m, \xi) + g(m, \xi)] \right\}, \quad n \in J.$$

Because $u(n) \leq \varphi^{-1}[z(n)] \leq v(n)$, $n \in J$, by letting $n = m$ and $\varepsilon \rightarrow 0$ in the last relation, we obtain

$$u(m) \leq L(m) + \sum_{s=0}^{m-1} f(m, s) G^{-1} \left\{ G[L(m)] + \sum_{\xi=0}^{s-1} [f(m, \xi) + g(m, \xi)] \right\}.$$

This means that the desired inequality (3.2) is valid when $n = m$. Since $m \in (0, M)$ is chosen arbitrarily and by (3.1), inequality (3.2) holds also for $n = 0$. Thus the proof of Theorem 3.1 is complete. \square

The following result is a special case of Theorem 3.1 when $\varphi(\eta) = \eta^p$, $w(r) = r$:



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 16 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Corollary 3.2. Under the conditions of Theorem 3.1, the discrete inequality

$$(3.7) \quad u^p(n) \leq c^p(n) + p \sum_{s=0}^{n-1} \left\{ f(n, s)u(s) \left[u(s) + \sum_{\xi=0}^{s-1} g(s, \xi)u(\xi) \right] + h(n, s)u(s) \right\}, \quad n \in \mathbb{N}_0,$$

where $p > 1$ is a real number, implies that

$$(3.8) \quad u(n) \leq \left[c(n) + \sum_{s=0}^{n-1} h(n, s) \right] \times \left\{ 1 + \sum_{s=0}^{n-1} f(n, s) \exp \sum_{\xi=0}^{s-1} [f(n, \xi) + g(n, \xi)] \right\}, \quad n \in \mathbb{N}_0.$$

Note that, the particular case of Theorem 3.1 when $\varphi(\eta) = \eta^2$, $c(n) \equiv k^2 > 0$ and the functions $f(n, s)$, $g(n, s)$, $h(n, s)$ are independent of the variable n , yields a discrete analogue of the Constantin integral inequality.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 17 of 20

Go Back

Full Screen

Close

4. Discrete Model of Infiltration

The mathematical model of the infiltration phenomena of a fluid due to Okrasinsky [7] was studied in [2] (see, also [8]):

$$(4.1) \quad u^2(t) = L + \int_0^t P(t-s)u(s)ds, \quad t \in \mathbb{R}_+,$$


where $L > 0$ is a constant, $P \in C(\mathbb{R}_+, \mathbb{R}_+)$ and u denotes the height of the percolating fluid above the horizontal impervious base, multiplied by a positive number. This model describes the infiltration phenomena of a fluid from a cylindrical reservoir into an isotropic homogeneous porous medium. Under the condition “ P is differentiable and nondecreasing”, Constantin obtained the existence and uniqueness of a solution $u \in C^1(\mathbb{R}_+, (0, \infty))$ of equation (4.1). Some known results for equation (4.1) are also given in Constantin [3] and Lipovan [5].

We note here that, although the conclusions given therein are correct, the derivation of them has a small defect. Actually, since function P depends on both variables t, s , the integral inequality given in the lemma of [2] is not applicable. However, using our Theorem 2.1 and by following the same argument as used in [2] these conclusions can be reproved very easily.

Now we consider the discrete analogue of equation (4.1) without a differentiability requirement on the function P :

$$(4.2) \quad u^2(n) = L + \sum_{s=0}^{n-1} P(n,s)u(s), \quad n \in \mathbb{N},$$

where $L > 0$ is a constant, $u, P \in K(\mathbb{N}, \mathbb{R}_+)$ with P nondecreasing. The unique positive solution to equation (4.2) can be obtained by successive substitution. For



instance, by letting $n = 0, 1, 2$ successively in (4.2), we obtain

$$u(0) = \sqrt{L}, \quad u(1) = \sqrt{L + P(1)u(0)}, \quad u(2) = \sqrt{L + P(1)u(1) + P(2)u(0)}.$$

An application of Corollary 2.3 with $f(n, s) = g(n, s) \equiv 0, h(n, s) = P(n - s)$ to (4.1) yields an upper bound on $u(n)$ of the form

$$u(n) \leq \sqrt{L} + \sum_{s=0}^{n-1} P(n - s), \quad n \in \mathbb{N}.$$

**Integral Inequality and
Its Discrete Analogue**

En-hao Yang and Man-chun Tan

vol. 8, iss. 2, art. 57, 2007

Title Page

Contents



Page 18 of 20

Go Back

Full Screen


Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] A. CONSTANTIN, Solutions globales d'équations différentielles perturbées, *C. R. Acad. Sci. Paris*, **320** (1995), 1319–1322.
- [2] A. CONSTANTIN, Topological transversality: Application to an integro-differential equation, *J. Math. Anal. Appl.*, **197**(1) (1996), 855–863.
- [3] A. CONSTANTIN, Monotone iterative technique for a nonlinear integral equation, *J. Math. Anal. Appl.*, **205** (1997), 280–283.
- [4] A. GRANAS, Sur la méthode de continuité de Poincaré, *C. R. Acad. Sci. Paris*, **282** (1976), 983–985.
- [5] O. LIPOVAN, On a nonlinear integral equation, *Bul. Stiint. Univ. Politeh. Timis. Ser. Mat. Fiz.*, **44** (1999), 40–45.
- [6] D. O'REGAN AND M. MEEHAN, *Existence Theory for Nonlinear Integral and Integrodifferential Equations*, Dordrecht/Boston/London, Kluwer Acad. Publ., 1998.
- [7] W. OKRASINSKY, On the existence and uniqueness of non-negative solutions of a certain non-linear convolution equation, *Ann. Polon. Math.*, **38** (1979), 61–72.
- [8] W. OKRASINSKY, On a non-linear convolution equation occurring in the theory of water percolation, *Ann. Polon. Math.*, **37** (1980), 223–229.
- [9] L. OU-ANG, The boundedness of solutions of linear differential equations $y'' + A(t)y = 0$, *Shuxue Jinzhan*, **3** (1957), 409–418.
- [10] B.G. PACHPATTE, On a certain inequality arising in the theory of differential equations, *J. Math. Anal. Appl.*, **182**(1) (1994), 143–157.



Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 19 of 20

Go Back

Full Screen

Close



- [11] B.G. PACHPATTE, A note on certain integral inequalities with delay, *Period. Math. Hungar.*, **31** (1995), 229–234.
- [12] B.G. PACHPATTE, Some new inequalities related to certain inequalities in the theory of differential equations, *J. Math. Anal. Appl.*, **189**(1) (1995), 128–144.
- [13] E.H. YANG, Generalizations of Pachpatte’s integral and discrete inequalities, *Ann. Diff. Eqs.*, **13** (1997), 180–188.
- [14] E.H. YANG, On some nonlinear integral and discrete inequalities related to Ou-Iang’s inequality, *Acta Math. Sinica (New Series)*, **14**(3) (1998), 353–360.

Title Page

Contents



Page 20 of 20

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756