

ON AN INEQUALITY OF V. CSISZÁR AND T.F. MÓRI FOR CONCAVE FUNCTIONS OF TWO VARIABLES



BOŽIDAR IVANKOVIĆ

Faculty of Transport and Traffic Engineering
University of Zagreb, Vukelićeva 4
10000 Zagreb, Croatia
EMail: ivankovb@fpz.hr

JOSIP E. PEČARIĆ

Faculty of Textile Technology
University of Zagreb, Pierottijeva 6
10000 Zagreb, Croatia
EMail: pecaric@mahazu.hazu.hr

SAICHI IZUMINO

Faculty of Education
Toyama University
Gofuku, Toyama 930-8555, JAPAN
EMail: s-izumino@h5.dion.ne.jp

MASARU TOMINAGA

Toyama National College of Technology
13, Hongo-machi, Toyama-shi
939-8630, Japan
EMail: mtommy@toyama-nct.ac.jp

Received: 27 September, 2006

Accepted: 21 April, 2007

Communicated by: [I. Pinelis](#)

2000 AMS Sub. Class.: 26D15.

Key words: Diaz-Metcalf inequality, Hölder's inequality, Hadamard's inequality, Petrović's inequality, Giaccardi's inequality.

Abstract: V. Csiszár and T.F. Móri gave an extension of Diaz-Metcalf's inequality for concave functions. In this paper, we show its restatement. As its applications we first give a reverse inequality of Hölder's inequality. Next we consider two variable versions of Hadamard, Petrović and Giaccardi inequalities.

Inequality of V. Csiszár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Reverse Hölder's Inequality	5
3	Hadamard's Inequality	9
4	Petrović's Inequality	11
5	Giaccardi's Inequality	14



Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 2 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)



Page 3 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

1. Introduction

In this paper, let (X, Y) be a random vector with $P[(X, Y) \in D] = 1$ where $D := [a, A] \times [b, B]$ ($0 \leq a < A$ and $0 \leq b < B$). Let $E[X]$ be the expectation of a random variable X with respect to P . For a function $\phi : D \rightarrow \mathbb{R}$, we put

$$\Delta\phi = \Delta\phi(a, b, A, B) := \phi(a, b) - \phi(a, B) - \phi(A, b) + \phi(A, B).$$

In [1], V. Csizsár and T.F. Móri showed the following theorem as an extension of Diaz-Metcalf's inequality [2].

Theorem A. *Let $\phi : D \rightarrow \mathbb{R}$ be a concave function.*

We use the following notations:

$$\begin{aligned} \lambda_1 = \lambda_4 &:= \frac{\phi(A, b) - \phi(a, b)}{A - a}, & \mu_1 = \mu_3 &:= \frac{\phi(a, B) - \phi(a, b)}{B - b}, \\ \lambda_2 = \lambda_3 &:= \frac{\phi(A, B) - \phi(a, B)}{A - a}, & \mu_2 = \mu_4 &:= \frac{\phi(A, B) - \phi(A, b)}{B - b}, & \text{and} \\ \nu_1 &:= \frac{AB - ab}{(A - a)(B - b)}\phi(a, b) - \frac{b}{B - b}\phi(a, B) - \frac{a}{A - a}\phi(A, b), \\ \nu_2 &:= \frac{A}{A - a}\phi(a, B) + \frac{B}{B - b}\phi(A, b) - \frac{AB - ab}{(A - a)(B - b)}\phi(A, B), \\ \nu_3 &:= \frac{B}{B - b}\phi(a, b) - \frac{a}{A - a}\phi(A, B) + \frac{aB - Ab}{(A - a)(B - b)}\phi(a, B), \\ \nu_4 &:= \frac{A}{A - a}\phi(a, b) - \frac{b}{B - b}\phi(A, B) - \frac{aB - Ab}{(A - a)(B - b)}\phi(A, b). \end{aligned}$$

a) *Suppose that $\Delta\phi \geq 0$.*



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 18

Go Back

Full Screen

Close

$a - (i)$ If $(B - b)E[X] + (A - a)E[Y] \leq AB - ab$, then

$$\lambda_1 E[X] + \mu_1 E[Y] + \nu_1 \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])).$$

$a - (ii)$ If $(B - b)E[X] + (A - a)E[Y] \geq AB - ab$, then

$$\lambda_2 E[X] + \mu_2 E[Y] + \nu_2 \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])).$$

$b)$ Suppose that $\Delta\phi \leq 0$

$b - (iii)$ If $(B - b)E[X] + (A - a)E[Y] \leq aB - Ab$, then

$$\lambda_3 E[X] + \mu_3 E[Y] + \nu_3 \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])).$$

$b - (iv)$ If $(B - b)E[X] + (A - a)E[Y] \geq aB - Ab$, then

$$\lambda_4 E[X] + \mu_4 E[Y] + \nu_4 \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])).$$

Let us note that Theorem A can be given in the following form:

Theorem 1.1. Suppose that $\phi : D \rightarrow \mathbb{R}$ is a concave function.

$a)$ If $\Delta\phi \geq 0$, then

$$(1.1) \quad \max_{k=1,2} \{\lambda_k E[X] + \mu_k E[Y] + \nu_k\} \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])),$$

where λ_k, μ_k and ν_k ($k = 1, 2$) are defined in Theorem A.

$b)$ If $\Delta\phi \leq 0$, then

$$(1.2) \quad \max_{k=3,4} \{\lambda_k E[X] + \mu_k E[Y] + \nu_k\} \leq E[\phi(X, Y)] (\leq \phi(E[X], E[Y])),$$

where λ_k, μ_k and ν_k ($k = 3, 4$) are defined in Theorem A.

Remark 1. The inequality $E[\phi(X, Y)] \leq \phi(E[X], E[Y])$ is Jensen's inequality. So the inequalities in Theorem A represent reverse inequalities of it.

In this note, we shall give some applications of these results.



2. Reverse Hölder's Inequality

Let $p, q > 1$ be real numbers with $\frac{1}{p} + \frac{1}{q} = 1$. Then $\phi(x, y) := x^{\frac{1}{p}}y^{\frac{1}{q}}$ is a concave function on $(0, \infty) \times (0, \infty)$. For $0 < a < A$ and $0 < b < B$, $\Delta\phi$ is represented as follows:

$$\Delta\phi = a^{\frac{1}{p}}b^{\frac{1}{q}} - a^{\frac{1}{p}}B^{\frac{1}{q}} - A^{\frac{1}{p}}b^{\frac{1}{q}} + A^{\frac{1}{p}}B^{\frac{1}{q}} = \left(A^{\frac{1}{p}} - a^{\frac{1}{p}}\right) \left(B^{\frac{1}{q}} - b^{\frac{1}{q}}\right) (> 0).$$

Moreover, putting $A = B = 1$, and replacing X, Y, a and b by X^p, Y^q, α^p and β^q , respectively, in Theorem A, we have the following result:

Theorem 2.1. *Let $p, q > 1$ be real numbers with $\frac{1}{p} + \frac{1}{q} = 1$. Let $0 < \alpha \leq X \leq 1$ and $0 < \beta \leq Y \leq 1$.*

(i) *If $(1 - \beta^q)E[X^p] + (1 - \alpha^p)E[Y^q] \leq 1 - \alpha^p\beta^q$, then*

$$(2.1) \quad \frac{\beta(1 - \alpha)}{1 - \alpha^p}E[X^p] + \frac{\alpha(1 - \beta)}{1 - \beta^q}E[Y^q] + \frac{\alpha\beta(1 - \alpha^{p-1} - \beta^{q-1} + \alpha^{p-1}\beta^q + \alpha^p\beta^{q-1} - \alpha^p\beta^q)}{(1 - \alpha^p)(1 - \beta^q)} \leq E[XY].$$

(ii) *If $(1 - \beta^q)E[X^p] + (1 - \alpha^p)E[Y^q] \geq 1 - \alpha^p\beta^q$, then*

$$(2.2) \quad \frac{1 - \alpha}{1 - \alpha^p}E[X^p] + \frac{1 - \beta}{1 - \beta^q}E[Y^q] - \frac{1 - \alpha - \beta + \alpha\beta^q + \alpha^p\beta - \alpha^p\beta^q}{(1 - \alpha^p)(1 - \beta^q)} \leq E[XY].$$

By Theorem 2.1 we have the following inequality related to Hölder's inequality:

Title Page

Contents



Page 5 of 18

Go Back

Full Screen

Close



Theorem 2.2. Let $p, q > 1$ be real numbers with $\frac{1}{p} + \frac{1}{q} = 1$. If $0 < \alpha \leq X \leq 1$ and $0 < \beta \leq Y \leq 1$, then

$$(2.3) \quad \begin{aligned} p^{\frac{1}{p}} q^{\frac{1}{q}} (\beta - \alpha\beta^q)^{\frac{1}{p}} (\alpha - \alpha^p\beta)^{\frac{1}{q}} E[X^p]^{\frac{1}{p}} E[Y^q]^{\frac{1}{q}} \\ \leq (\beta - \alpha\beta^q) E[X^p] + (\alpha - \alpha^p\beta) E[Y^q] \\ \leq (1 - \alpha^p\beta^q) E[XY]. \end{aligned}$$

Proof. We have by Young's inequality

$$\begin{aligned} (\beta - \alpha\beta^q) E[X^p] + (\alpha - \alpha^p\beta) E[Y^q] \\ = \frac{1}{p} \cdot p(\beta - \alpha\beta^q) E[X^p] + \frac{1}{q} \cdot q(\alpha - \alpha^p\beta) E[Y^q] \\ \geq \{p(\beta - \alpha\beta^q) E[X^p]\}^{\frac{1}{p}} \{q(\alpha - \alpha^p\beta) E[Y^q]\}^{\frac{1}{q}} \\ = p^{\frac{1}{p}} q^{\frac{1}{q}} (\beta - \alpha\beta^q)^{\frac{1}{p}} (\alpha - \alpha^p\beta)^{\frac{1}{q}} E[X^p]^{\frac{1}{p}} E[Y^q]^{\frac{1}{q}}. \end{aligned}$$

Hence the first inequality holds. Next, we see that

$$(2.4) \quad -\gamma_1 := \frac{\alpha\beta(1 - \alpha^{p-1} - \beta^{q-1} + \alpha^{p-1}\beta^q + \alpha^p\beta^{q-1} - \alpha^p\beta^q)}{(1 - \alpha^p)(1 - \beta^q)} \geq 0$$

and

$$(2.5) \quad \gamma_2 := \frac{1 - \alpha - \beta + \alpha\beta^q + \alpha^p\beta - \alpha^p\beta^q}{(1 - \alpha^p)(1 - \beta^q)} \geq 0.$$

Indeed, we have $(1 - \alpha^p)(1 - \beta^q) > 0$ and moreover by Young's inequality

$$\begin{aligned} 1 - \alpha^{p-1} - \beta^{q-1} + \alpha^{p-1}\beta^q + \alpha^p\beta^{q-1} - \alpha^p\beta^q \\ = 1 - \alpha^p\beta^q - \alpha^{p-1}(1 - \beta^q) - \beta^{q-1}(1 - \alpha^p) \\ \geq 1 - \alpha^p\beta^q - \left(\frac{1}{p} + \frac{1}{q}\alpha^p\right)(1 - \beta^q) - \left(\frac{1}{q} + \frac{1}{p}\beta^q\right)(1 - \alpha^p) = 0 \end{aligned}$$

Inequality of V. Császár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 6 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 7 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

and

$$\begin{aligned} 1 - \alpha - \beta + \alpha\beta^q + \alpha^p\beta - \alpha^p\beta^q \\ &= 1 - \alpha^p\beta^q - \alpha(1 - \beta^q) - \beta(1 - \alpha^p) \\ &\geq 1 - \alpha^p\beta^q - \left(\frac{1}{q} + \frac{1}{p}\alpha^p\right)(1 - \beta^q) - \left(\frac{1}{p} + \frac{1}{q}\beta^q\right)(1 - \alpha^p) = 0. \end{aligned}$$

Multiplying both sides of (2.1) by γ_2 and those of (2.2) by $-\gamma_1$, respectively, and taking the sum of the two inequalities, we have

$$\left| \begin{array}{cc} \frac{\beta(1-\alpha)}{1-\alpha^p} & \gamma_1 \\ \frac{1-\alpha}{1-\alpha^p} & \gamma_2 \end{array} \right| E[X^p] + \left| \begin{array}{cc} \frac{\alpha(1-\beta)}{1-\beta^q} & \gamma_1 \\ \frac{1-\beta}{1-\beta^q} & \gamma_2 \end{array} \right| E[Y^q] \leq (\gamma_2 - \gamma_1)E[XY].$$

Here we note that from (2.4) and (2.5),

$$\left| \begin{array}{cc} \frac{\beta(1-\alpha)}{1-\alpha^p} & \gamma_1 \\ \frac{1-\alpha}{1-\alpha^p} & \gamma_2 \end{array} \right| = \frac{\beta(1-\alpha)(1-\beta)(1-\alpha\beta^{q-1})}{(1-\alpha^p)(1-\beta^q)},$$

$$\left| \begin{array}{cc} \frac{\alpha(1-\beta)}{1-\beta^q} & \gamma_1 \\ \frac{1-\beta}{1-\beta^q} & \gamma_2 \end{array} \right| = \frac{\alpha(1-\alpha)(1-\beta)(1-\alpha^{p-1}\beta)}{(1-\alpha^p)(1-\beta^q)}$$

and

$$\gamma_2 - \gamma_1 = \frac{(1-\alpha)(1-\beta)(1-\alpha^p\beta^q)}{(1-\alpha^p)(1-\beta^q)}.$$

Hence we have

$$\frac{\beta(1-\alpha)(1-\beta)(1-\alpha\beta^{q-1})}{(1-\alpha^p)(1-\beta^q)} E[X^p] + \frac{\alpha(1-\alpha)(1-\beta)(1-\alpha^{p-1}\beta)}{(1-\alpha^p)(1-\beta^q)} E[Y^q]$$

$$\leq \frac{(1-\alpha)(1-\beta)(1-\alpha^p\beta^q)}{(1-\alpha^p)(1-\beta^q)} E[XY]$$

and so the second inequality of (2.3) holds. \square

The second inequality is given in [5, p.124]. In (2.3), the first and the third terms yield the following Gheorghiu inequality [4, p.184], [5, p.124]:


Theorem B. *Let $p, q > 1$ be real numbers with $\frac{1}{p} + \frac{1}{q} = 1$. If $0 < \alpha \leq X \leq 1$ and $0 < \beta \leq Y \leq 1$, then*

$$(2.6) \quad E[X^p]^{\frac{1}{p}} E[Y^q]^{\frac{1}{q}} \leq \frac{1 - \alpha^p \beta^q}{p^{\frac{1}{p}} q^{\frac{1}{q}} (\beta - \alpha \beta^q)^{\frac{1}{p}} (\alpha - \alpha^p \beta)^{\frac{1}{q}}} E[XY].$$

We see that (2.3) is a kind of a refinement of (2.6). Theorem B gives us the next estimation.

Corollary 2.3. *Let $X = \{a_i\}$ and $Y = \{b_j\}$ be independent discrete random variables with distributions $P(X = a_i) = w_i$ and $P(Y = b_j) = z_j$. Suppose $0 < \alpha \leq X \leq 1$ and $0 < \beta \leq Y \leq 1$. $E[X^p]$, $E[Y^q]$ and $E[XY]$ are given by $\sum_{i=1}^n w_i a_i^p$, $\sum_{j=1}^n z_j b_j^q$ and $\sum_{i=1}^n \sum_{j=1}^n w_i z_j a_i b_j$, respectively. Then we have inequalities*

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n w_i z_j a_i b_j &\leq \left(\sum_{i=1}^n w_i a_i^p \right)^{\frac{1}{p}} \left(\sum_{j=1}^n z_j b_j^q \right)^{\frac{1}{q}} \\ &\leq \frac{1 - \alpha^p \beta^q}{p^{\frac{1}{p}} q^{\frac{1}{q}} (\beta - \alpha \beta^q)^{\frac{1}{p}} (\alpha - \alpha^p \beta)^{\frac{1}{q}}} \sum_{i=1}^n \sum_{j=1}^n w_i z_j a_i b_j. \end{aligned}$$



Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 8 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 9 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

3. Hadamard's Inequality

The following well-known inequality is due to Hadamard [5, p.11]: For a concave function $f : [a, b] \rightarrow \mathbb{R}$,

$$(3.1) \quad \frac{f(a) + f(b)}{2} \leq \frac{1}{b-a} \int_a^b f(t) dt \leq f\left(\frac{a+b}{2}\right).$$

Moreover, the following is an extension of the weighted version of Hadamard's inequality by Fejér ([3], [6, p.138]): Let g be a positive integrable function on $[a, b]$ with $g(a+t) = g(b-t)$ for $0 \leq t \leq \frac{1}{2}(a-b)$. Then

$$(3.2) \quad \frac{f(a) + f(b)}{2} \int_a^b g(t) dt \leq \int_a^b f(t)g(t) dt \leq f\left(\frac{a+b}{2}\right) \int_a^b g(t) dt.$$

Here we give an analogous result for a function of two variables.

Theorem 3.1. *Let X and Y be independent random variables such that*

$$(3.3) \quad E[X] = \frac{a+A}{2} \quad \text{and} \quad E[Y] = \frac{b+B}{2}$$

for $0 < a \leq X \leq A$ and $0 < b \leq Y \leq B$. If $\phi : D \rightarrow \mathbb{R}$ is a concave function, then

$$(3.4) \quad \min \left\{ \frac{\phi(A, b) + \phi(a, B)}{2}, \frac{\phi(a, b) + \phi(A, B)}{2} \right\} \leq E[\phi(X, Y)] \\ \leq \phi\left(\frac{a+A}{2}, \frac{b+B}{2}\right).$$

Proof. We only have to prove the case $\Delta\phi \geq 0$. Then with same notations as in Theorem A we have

$$\lambda_1 E[X] + \mu_1 E[Y] + \nu_1 = \lambda_2 E[X] + \mu_2 E[Y] + \nu_2 = \frac{\phi(A, b) + \phi(a, B)}{2}$$



by (3.3). Since $\Delta\phi \geq 0$, it is the same as the first expression in (3.4). Similarly calculation for $\Delta\phi \leq 0$ proves that the desired inequality (3.4) also holds. \square

We can obtain the following result as an extension of Hadamard's inequality (3.1) from Theorem 3.1 by letting X and Y be independent, uniformly distributed random variables on the intervals $[a, A]$ and $[b, B]$, respectively:

Corollary 3.2. *Let $0 < a < A$ and $0 < b < B$. If ϕ is a concave function, then*

$$\min \left\{ \frac{\phi(A, b) + \phi(a, B)}{2}, \frac{\phi(a, b) + \phi(A, B)}{2} \right\} \leq \frac{1}{(A-a)(B-b)} \int_a^A \int_b^B \phi(t, s) ds dt \\ \leq \phi \left(\frac{a+A}{2}, \frac{b+B}{2} \right).$$

By Theorem 3.1, we have the following analogue of (3.2) for a function of two variables:

Corollary 3.3. *Let $w : D \rightarrow \mathbb{R}$ be a nonnegative integrable function such that $w(s, t) = u(s)v(t)$ where $u : [a, A] \rightarrow \mathbb{R}$ is an integrable function with $u(s) = u(a + A - s)$, $\int_a^A u(s) ds = 1$ and $v : [b, B] \rightarrow \mathbb{R}$ is an integrable function such that $\int_b^B v(t) dt = 1$, $v(t) = v(b + B - t)$. If ϕ is a concave function, then*

$$\min \left\{ \frac{\phi(A, b) + \phi(a, B)}{2}, \frac{\phi(a, b) + \phi(A, B)}{2} \right\} \leq \int_a^A \int_b^B w(s, t) \phi(s, t) ds dt \\ \leq \phi \left(\frac{a+A}{2}, \frac{b+B}{2} \right).$$

Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.



4. Petrović's Inequality

The following is called Petrović's inequality for a concave function $f : [0, c] \rightarrow \mathbb{R}$:

$$f\left(\sum_{i=1}^n p_i x_i\right) \leq \sum_{i=1}^n p_i f(x_i) + (1 - P_n) f(0),$$

where $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{p} = (p_1, \dots, p_n)$ are n -tuples of nonnegative real numbers such that $\sum_{i=1}^n p_i x_i \geq x_k$ for $k = 1, \dots, n$, $\sum_{i=1}^n p_i x_i \in [0, c]$ and $P_n := \sum_{i=1}^n p_i$ (see [5, p.11] and [6]).

We give an analogous result for a function of two variables.

Theorem 4.1. Let $\mathbf{p} = (p_1, \dots, p_n)$ and $\mathbf{q} = (q_1, \dots, q_n)$ be n -tuples of nonnegative real numbers and put $P_n := \sum_{i=1}^n p_i (> 0)$ and $Q_n := \sum_{j=1}^n q_j (> 0)$. Suppose that $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ are n -tuples of nonnegative real numbers with $0 \leq x_k \leq \sum_{i=1}^n p_i x_i \leq c$ and $0 \leq y_k \leq \sum_{j=1}^n q_j y_j \leq d$ for $k = 1, 2, \dots, n$. Let $\phi : [0, c] \times [0, d] \rightarrow \mathbb{R}$ be a concave function.

a) Suppose

$$\phi(0, 0) + \phi\left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j\right) \geq \phi\left(\sum_{i=1}^n p_i x_i, 0\right) + \phi\left(0, \sum_{j=1}^n q_j y_j\right).$$

a – (i) If $\frac{1}{P_n} + \frac{1}{Q_n} \leq 1$, then

$$(4.1) \quad \frac{1}{P_n} \phi\left(\sum_{i=1}^n p_i x_i, 0\right) + \frac{1}{Q_n} \phi\left(0, \sum_{j=1}^n q_j y_j\right) + \left(1 - \frac{1}{P_n} - \frac{1}{Q_n}\right) \phi(0, 0) \\ \leq \frac{1}{P_n Q_n} \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j).$$

Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 11 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 12 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

a – (ii) If $\frac{1}{P_n} + \frac{1}{Q_n} \geq 1$, then

$$\begin{aligned} & \left(\frac{1}{P_n} + \frac{1}{Q_n} - 1 \right) \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) \\ & + \left(1 - \frac{1}{Q_n} \right) \phi \left(\sum_{i=1}^n p_i x_i, 0 \right) + \left(1 - \frac{1}{P_n} \right) \phi \left(0, \sum_{j=1}^n q_j y_j \right) \\ & \leq \frac{1}{P_n Q_n} \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j). \end{aligned}$$

b) Suppose

$$\phi(0, 0) + \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) \leq \phi \left(\sum_{i=1}^n p_i x_i, 0 \right) + \phi \left(0, \sum_{j=1}^n q_j y_j \right).$$

b – (iii) If $P_n \geq Q_n$, then

$$\begin{aligned} & \frac{1}{P_n} \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) \\ & + \left(\frac{1}{Q_n} - \frac{1}{P_n} \right) \phi \left(0, \sum_{j=1}^n q_j y_j \right) + \left(1 - \frac{1}{Q_n} \right) \phi(0, 0) \\ & \leq \frac{1}{P_n Q_n} \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j). \end{aligned}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 18

Go Back

Full Screen

Close

$b - (iv)$ If $Q_n \geq P_n$, then

$$\begin{aligned} & \frac{1}{Q_n} \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) \\ & - \left(\frac{1}{Q_n} - \frac{1}{P_n} \right) \phi \left(\sum_{i=1}^n p_i x_i, 0 \right) + \left(1 - \frac{1}{P_n} \right) \phi(0, 0) \\ & \leq \frac{1}{P_n Q_n} \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j). \end{aligned}$$

Proof. We put $a = b = 0$, $A = \sum_{i=1}^n p_i x_i$ and $B = \sum_{j=1}^n q_j y_j$ in Theorem A. Let $X = \{a_i\}$ and $Y = \{b_j\}$ be independent discrete random variables with distributions $P(X = x_i) = \frac{p_i}{P_n}$ and $P(Y = y_j) = \frac{q_j}{Q_n}$, $1 \leq i \leq n$, respectively. So we have the desired inequalities. \square

Specially, if $p_i = q_j = 1$ ($i, j = 1, \dots, n$) in Theorem 4.1, then we have the following:

Corollary 4.2. Suppose that $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ are n -tuples of nonnegative real numbers for $n \geq 2$ with $\sum_{i=1}^n x_i \in [0, c]$ and $\sum_{i=1}^n y_i \in [0, d]$. If $\phi : [0, c] \times [0, d] \rightarrow \mathbb{R}$ is a concave function, then

$$(4.2) \quad \phi \left(\sum_{i=1}^n x_i, 0 \right) + \phi \left(0, \sum_{j=1}^n y_j \right) + (n-2) \phi(0, 0) \leq \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n \phi(x_i, y_j).$$

5. Giaccardi's Inequality

In 1955, Giaccardi (cf. [5, p.11]) proved the following inequality for a convex function $f : [a, A] \rightarrow \mathbb{R}$,

$$\sum_{i=1}^n p_i f(x_i) \leq C \cdot f\left(\sum_{i=1}^n p_i x_i\right) + D \cdot (P_n - 1) \cdot f(x_0),$$

where

$$C = \frac{\sum_{i=1}^n p_i (x_i - x_0)}{\sum_{i=1}^n p_i x_i - x_0} \quad \text{and} \quad D = \frac{\sum_{i=1}^n p_i x_i}{\sum_{i=1}^n p_i x_i - x_0}$$

for a nonnegative n -tuple $\mathbf{p} = (p_1, \dots, p_n)$ with $P_n := \sum_{i=1}^n p_i$ and a real $(n+1)$ -tuple $\mathbf{x} = (x_0, x_1, \dots, x_n)$ such that for $k = 0, 1, \dots, n$

$$a \leq x_i \leq A, \quad (x_k - x_0) \left(\sum_{i=1}^n p_i x_i - x_0 \right) \geq 0,$$

$$a < \sum_{i=1}^n p_i x_i < A \quad \text{and} \quad \sum_{i=1}^n p_i x_i \neq x_0.$$

In this section, we discuss a generalization of Giaccardi's inequality to a function of two variables under similar conditions. Let $\mathbf{x} = (x_0, x_1, \dots, x_n)$ and $\mathbf{y} = (y_0, y_1, \dots, y_n)$ be nonnegative $(n+1)$ -tuples, and $\mathbf{p} = (p_1, p_2, \dots, p_n)$ and $\mathbf{q} = (q_1, q_2, \dots, q_n)$ be nonnegative n -tuples with

$$(5.1) \quad x_0 \leq x_k \leq \sum_{i=1}^n p_i x_i \quad \text{and} \quad y_0 \leq y_k \leq \sum_{j=1}^n q_j y_j \quad \text{for } k = 1, \dots, n.$$



Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 14 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 15 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

We use the following notations:

$$P_n := \sum_{i=1}^n p_i (\geq 0),$$

$$Q_n := \sum_{j=1}^n q_j (\geq 0),$$

$$K(X) := \frac{\sum_{i=1}^n p_i x_i - P_n x_0}{\sum_{i=1}^n p_i x_i - x_0},$$

$$K(Y) := \frac{\sum_{j=1}^n q_j y_j - Q_n y_0}{\sum_{j=1}^n q_j y_j - y_0},$$

$$L(X) := \frac{(P_n - 1) \sum_{i=1}^n p_i x_i}{\sum_{i=1}^n p_i x_i - x_0},$$

$$L(Y) := \frac{(Q_n - 1) \sum_{j=1}^n q_j y_j}{\sum_{j=1}^n q_j y_j - y_0},$$

$$M(X, Y) := \left\{ (P_n Q_n - P_n - Q_n) \sum_{i=1}^n p_i x_i \sum_{j=1}^n q_j y_j \right. \\ \left. + Q_n y_0 \sum_{i=1}^n p_i x_i + P_n x_0 \sum_{j=1}^n q_j y_j - P_n Q_n x_0 y_0 \right\} \\ \times \frac{1}{\left(\sum_{i=1}^n p_i x_i - x_0 \right) \left(\sum_{j=1}^n q_j y_j - y_0 \right)}$$

and

$$N(X, Y) \\ := \frac{(P_n - Q_n) \sum_{i=1}^n p_i x_i \sum_{j=1}^n q_j y_j - (P_n - 1) Q_n \sum_{i=1}^n p_i x_i y_0 + P_n (Q_n - 1) x_0 \sum_{j=1}^n q_j y_j}{\left(\sum_{i=1}^n p_i x_i - x_0 \right) \left(\sum_{j=1}^n q_j y_j - y_0 \right)}.$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 16 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

Then we have the following theorem:

Theorem 5.1. Let $\phi : [x_0, \sum_{i=1}^n p_i x_i] \times [y_0, \sum_{j=1}^n q_j y_j] \rightarrow \mathbb{R}$ be a concave function.

a) If

$$\phi(x_0, y_0) + \phi\left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j\right) \geq \phi\left(x_0, \sum_{j=1}^n q_j y_j\right) + \phi\left(\sum_{i=1}^n p_i x_i, y_0\right),$$

then

$$\begin{aligned} \max \left\{ Q_n K(X) \phi\left(\sum_{i=1}^n p_i x_i, y_0\right) + P_n K(Y) \phi\left(x_0, \sum_{j=1}^n q_j y_j\right) + M(X, Y) \phi(x_0, y_0), \right. \\ P_n L(Y) \phi\left(\sum_{i=1}^n p_i x_i, y_0\right) + Q_n L(X) \phi\left(x_0, \sum_{j=1}^n q_j y_j\right) \\ \left. - M(X, Y) \phi\left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j\right) \right\} \leq \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j). \end{aligned}$$

b) If

$$\phi(x_0, y_0) + \phi\left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j\right) \leq \phi\left(x_0, \sum_{j=1}^n q_j y_j\right) + \phi\left(\sum_{i=1}^n p_i x_i, y_0\right),$$

then

$$\begin{aligned} & \max \left\{ Q_n K(X) \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) \right. \\ & \quad \left. + P_n L(Y) \phi(x_0, y_0) + N(X, Y) \phi \left(x_0, \sum_{j=1}^n q_j y_j \right), \right. \\ & P_n K(Y) \phi \left(\sum_{i=1}^n p_i x_i, \sum_{j=1}^n q_j y_j \right) + Q_n L(X) \phi(x_0, y_0) \\ & \quad \left. - N(X, Y) \phi \left(\sum_{i=1}^n p_i x_i, y_0 \right) \right\} \leq \sum_{i=1}^n \sum_{j=1}^n p_i q_j \phi(x_i, y_j). \end{aligned}$$

Proof. Let X and Y be as they were in the proof of Theorem 4.1, and put $a = x_0$, $A = \sum_{i=1}^n p_i x_i$, $b = y_0$ and $B = \sum_{j=1}^n q_j y_j$, and use Theorem A. Then we have the desired inequalities of this theorem. \square



Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 17 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] V. CSISZÁR AND T.F. MÓRI, The convexity method of proving moment-type inequalities, *Statist. Probab. Lett.*, **66** (2004), 303–313.
- [2] J.B. DIAZ AND F.T. METCALF, Stronger forms of a class of inequalities of G. Pólya-G. Szegő, and L. V. Kantorovich, *Bull. Amer. Math. Soc.*, **69** (1963), 415–418.
- [3] L. FEJÉR, Über die Fourierreihen, II., *Math. Naturwiss, Ant. Ungar. Acad. Wiss*, **24** (1906), 369–390 (in Hungarian).
- [4] S. IZUMINO AND M. TOMINAGA, Estimations in Hölder’s type inequalities, *Math. Inequal. Appl.*, **4** (2001), 163–187.
- [5] D. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer. Acad. Pub., Boston, London 1993.
- [6] J.E. PEČARIĆ, F. PROSCHAN AND Y.L. TONG, *Convex Functions, Partial Orderings, and Statistical Applications*, Mathematics in Science and Engineering, Georgia Institute of Technology, 1992.



Inequality of V. Csizsár
and T.F. Móri

Božidar Ivanković, Saichi Izumino,
Josip E. Pečarić and Masaru Tominaga

vol. 8, iss. 3, art. 88, 2007

Title Page

Contents



Page 18 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756