

# Journal of Inequalities in Pure and Applied Mathematics

## A NOTE ON THE BEZIER VARIANT OF CERTAIN BERNSTEIN DURMEYER OPERATORS

M.K. GUPTA

Department of Mathematics  
Ch. Charan Singh University  
Meerut-250004, India.

*EMail:* [mkgupta2002@hotmail.com](mailto:mkgupta2002@hotmail.com)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
140-04



---

volume 6, issue 3, article 74,  
2005.

*Received 27 July, 2004;  
accepted 13 May, 2005.*

*Communicated by: R.N. Mohapatra*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

In the present note, we introduce a Bezier variant of a new type of Bernstein Durrmeyer operator, which was introduced by Gupta [3]. We estimate the rate of convergence by using the decomposition technique of functions of bounded variation and applying the optimum bound. It is observed that the analysis for our Bezier variant of new Bernstein Durrmeyer operators is different from the usual Bernstein Durrmeyer operators studied by Zeng and Chen [9].


*2000 Mathematics Subject Classification:* 41A30, 41A36.

*Key words:* Lebesgue integrable functions; Bernstein polynomials; Bezier variant; Functions of bounded variation.

The author is thankful to the referee for his suggestions leading to substantial improvements in the paper.

## Contents

1	Introduction .....	3
2	Auxiliary Results .....	7
3	Main Result .....	9
	References	



### A Note on the Bezier Variant of Certain Bernstein Durrmeyer Operators

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 2 of 17

# 1. Introduction

Durrmeyer [1] introduced the integral modification of Bernstein polynomials to approximate Lebesgue integrable functions on the interval  $[0, 1]$ . The operators introduced by Durrmeyer are defined by

$$(1.1) \quad D_n(f, x) = (n + 1) \sum_{k=0}^n p_{n,k}(x) \int_0^1 p_{n,k}(t) f(t) dt, \quad x \in [0, 1],$$

where  $p_{n,k}(x) = \binom{n}{k} x^k (1 - x)^{n-k}$ .

Gupta [3] introduced a different Durrmeyer type modification of the Bernstein polynomials and estimated the rate of convergence for functions of bounded variation. The operators introduced in [3] are defined by

$$(1.2) \quad B_n(f, x) = n \sum_{k=0}^n p_{n,k}(x) \int_0^1 b_{n,k}(t) f(t) dt, \quad x \in [0, 1],$$

where

$$p_{n,k}(x) = (-1)^k \frac{x^k}{k!} \phi_n^{(k)}(x), \quad b_{n,k}(t) = (-1)^{k+1} \frac{t^k}{k!} \phi_n^{(k+1)}(t)$$

and

$$\phi_n(x) = (1 - x)^n.$$

It is easily verified that the values of  $p_{n,k}(x)$  used in (1.1) and (1.2) are same. Also it is easily verified that

$$\sum_{k=0}^n p_{n,k}(x) = 1, \quad \int_0^1 b_{n,k}(t) dt = 1 \quad \text{and} \quad b_{n,n}(t) = 0.$$



A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 3 of 17

By considering the integral modification of Bernstein polynomials in the form (1.2) some approximation properties become simpler in the analysis. So it is significant to study further on the different integral modification of Bernstein polynomials introduced by Gupta [3]. For  $\alpha \geq 1$ , we now define the Bezier variant of the operators (1.2), to approximate Lebesgue integrable functions on the interval  $[0, 1]$  as

$$(1.3) \quad B_{n,\alpha}(f, x) = \sum_{k=0}^n Q_{n,k}^{(\alpha)}(x) \int_0^1 b_{n,k}(t) f(t) dt, \quad x \in [0, 1],$$

where

$$Q_{n,k}^{(\alpha)}(x) = J_{n,k}^{\alpha}(x) - J_{n,k+1}^{\alpha}(x)$$


and

$$J_{n,k}(x) = \sum_{j=k}^n p_{n,j}(x),$$

when  $k \leq n$  and 0 otherwise.

Some important properties of  $J_{n,k}(x)$  are as follows:

- (i)  $J_{n,k}(x) - J_{n,k+1}(x) = p_{n,k}(x), k = 0, 1, 2, 3, \dots;$
- (ii)  $J'_{n,k}(x) = np_{n-1,k-1}(x), k = 1, 2, 3, \dots;$
- (iii)  $J_{n,k}(x) = n \int_0^x p_{n-1,k-1}(u) du, k = 1, 2, 3, \dots;$
- (iv)  $J_{n,0}(x) > J_{n,1}(x) > J_{n,2}(x) > \dots > J_{n,n}(x) > 0, 0 < x < 1.$




---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 17

For every natural number  $k$ ,  $J_{n,k}(x)$  increases strictly from 0 to 1 on  $[0, 1]$ .  
 Alternatively we may rewrite the operators (1.3) as


$$(1.4) \quad B_{n,\alpha}(f, x) = \int_0^1 K_{n,\alpha}(x, t)f(t)dt, \quad 0 \leq x \leq 1,$$

where

$$K_{n,\alpha}(x, t) = \sum_{k=0}^n Q_{n,k}^{(\alpha)}(x)b_{n,k}(t).$$

It is easily verified that  $B_{n,\alpha}(f, x)$  are linear positive operators,  $B_{n,\alpha}(1, x) = 1$  and for  $\alpha = 1$ , the operators  $B_{n,1}(f, x) \equiv B_n(f, x)$ , i.e. the operators (1.3) reduce to the operators (1.2). For further properties of  $Q_{n,k}^{(\alpha)}(x)$ , we refer the readers to [3].

Guo [2] studied the rate of convergence for bounded variation functions for Bernstein Durrmeyer operators. Zeng and Chen [9] were the first to estimate the rate of convergence for the Bezier variant of Bernstein Durrmeyer operators. Several other Bezier variants of some summation-integral type operators were studied in [4], [6] and [8] etc. It is well-known that Bezier basis functions play an important role in computer aided design. Moreover the recent work on different Bernstein Bezier type operators motivated us to study further in this direction. The advantage of the operators  $B_{n,\alpha}(f, x)$  over the Bernstein Durrmeyer operators considered in [9] is that one does not require the results of the type Lemma 3 and Lemma 4 of [9]. Consequently some approximation formulae become simpler. Further for  $\alpha = 1$ , these operators provide improved estimates over the main results of [2] and [3]. In the present paper, we estimate




---

**A Note on the Bezier Variant of  
 Certain Bernstein Durrmeyer  
 Operators**

M.K. Gupta

---

Title Page

Contents




Go Back

Close

Quit

Page 5 of 17

the rate of point wise convergence of the operators  $B_{n,\alpha}(f, x)$  at those points  $x \in (0, 1)$  at which one sided limits  $f(x-)$  and  $f(x+)$  exist.



---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 17

## 2. Auxiliary Results

In this section we give certain results, which are necessary to prove the main result.

**Lemma 2.1 ([3]).** *If  $n$  is sufficiently large, then*

$$\frac{x(1-x)}{n} \leq B_n((t-x)^2, x) \leq \frac{2x(1-x)}{n}.$$

**Lemma 2.2 ([4]).** *For every  $0 \leq k \leq n$ ,  $x \in (0, 1)$  and for all  $n \in \mathbb{N}$ , we have*

$$p_{n,k}(x) \leq \frac{1}{\sqrt{2enx(1-x)}}.$$

**Lemma 2.3.** *For all  $x \in (0, 1)$ , there holds*

$$Q_{n,k}^{(\alpha)}(x) \leq \alpha \cdot p_{n,k}(x) \leq \frac{\alpha}{\sqrt{2enx(1-x)}}.$$


*Proof.* Using the well known inequality  $|a^\alpha - b^\alpha| \leq \alpha |a - b|$ , ( $0 \leq a, b \leq 1$ ,  $\alpha \geq 1$ ) and by Lemma 2.2, we obtain

$$Q_{n,k}^{(\alpha)}(x) \leq \alpha p_{n,k}(x) \leq \frac{\alpha}{\sqrt{2enx(1-x)}}.$$

□

**Lemma 2.4.** *Let  $x \in (0, 1)$  and  $K_{n,\alpha}(x, t)$  be the kernel defined by (1.4). Then for  $n$  sufficiently large, we have*

$$(2.1) \quad \lambda_{n,\alpha}(x, y) := \int_0^y K_{n,\alpha}(x, t) dt \leq \frac{2\alpha \cdot x(1-x)}{n(x-y)^2}, \quad 0 \leq y < x,$$



A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 7 of 17


and

$$(2.2) \quad 1 - \lambda_{n,\alpha}(x, z) := \int_z^1 K_{n,\alpha}(x, t) dt \leq \frac{2\alpha \cdot x(1-x)}{n(z-x)^2}, \quad x < z < 1.$$

*Proof.* We first prove (2.1), as follows

$$\begin{aligned} \int_0^y K_{n,\alpha}(x, t) dt &\leq \int_0^y K_{n,\alpha}(x, t) \frac{(x-t)^2}{(x-y)^2} dt \\ &\leq \frac{1}{(x-y)^2} B_{n,\alpha}((t-x)^2, x) \\ &\leq \frac{\alpha \cdot B_{n,1}((t-x)^2, x)}{(x-y)^2} \leq \frac{2\alpha \cdot x(1-x)}{n(x-y)^2}, \end{aligned}$$

by Lemma 2.1. The proof of (2.2) is similar. □



---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 17



### 3. Main Result

In this section we prove the following main theorem.

**Theorem 3.1.** *Let  $f$  be a function of bounded variation on the interval  $[0, 1]$  and suppose  $\alpha \geq 1$ . Then for every  $x \in (0, 1)$  and  $n$  sufficiently large, we have*

$$\left| B_{n,\alpha}(f, x) - \left[ \frac{1}{\alpha+1} f(x+) + \frac{\alpha}{\alpha+1} f(x-) \right] \right| \leq \frac{\alpha}{\sqrt{2enx(1-x)}} |f(x+) - f(x-)| + \frac{5\alpha}{nx(1-x)} \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^{x+(1-x)/\sqrt{k}} (g_x),$$


where

$$g_x(t) = \begin{cases} f(t) - f(x-), & \text{for } 0 \leq t < x \\ 0, & \text{for } t = x \\ f(t) - f(x+), & \text{for } x < t \leq 1 \end{cases}$$

and  $\bigvee_a^b(g_x)$  is the total variation of  $g_x$  on  $[a, b]$ .

*Proof.* Clearly

$$(3.1) \quad \left| B_{n,\alpha}(f, x) - \left\{ \frac{1}{\alpha+1} f(x+) + \frac{\alpha}{\alpha+1} f(x-) \right\} \right| \leq |B_{n,\alpha}(g_x, x)| + \frac{1}{2} \left| B_{n,\alpha}(\text{sgn}(t-x), x) + \frac{\alpha-1}{\alpha+1} \right| |f(x+) - f(x-)|.$$



A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 9 of 17

First, we have


$$\begin{aligned}
 B_{n,\alpha}(\operatorname{sgn}(t-x), x) &= \int_x^1 K_{n,\alpha}(x, t) dt - \int_0^x K_{n,\alpha}(x, t) dt \\
 &= \int_0^1 K_{n,\alpha}(x, t) dt - 2 \int_0^x K_{n,\alpha}(x, t) dt \\
 &= 1 - 2 \int_0^x K_{n,\alpha}(x, t) dt = -1 + 2 \int_x^1 K_{n,\alpha}(x, t) dt.
 \end{aligned}$$

Using Lemma 2.2, Lemma 2.3 and the fact that

$$\sum_{j=0}^k p_{n,j}(x) = \int_x^1 b_{n,k}(t) dt,$$

we have

$$\begin{aligned}
 B_{n,\alpha}(\operatorname{sgn}(t-x), x) &= -1 + 2 \sum_{k=0}^n Q_{n,k}^{(\alpha)}(x) \int_x^1 b_{n,k}(t) dt \\
 &= -1 + 2 \sum_{k=0}^n Q_{n,k}^{(\alpha)}(x) \sum_{j=0}^k p_{n,j}(x) \\
 &= -1 + 2 \sum_{j=0}^n p_{n,j}(x) \sum_{k=j}^n Q_{n,k}^{(\alpha)}(x) \\
 &= -1 + 2 \sum_{j=0}^n p_{n,j}(x) J_{n,j}^{\alpha}(x).
 \end{aligned}$$



A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 10 of 17

Since

$$\sum_{j=0}^n Q_{n,j}^{(\alpha+1)}(x) = 1,$$

therefore we have

$$B_{n,\alpha}(\operatorname{sgn}(t-x), x) + \frac{\alpha-1}{\alpha+1} = 2 \sum_{j=0}^n p_{n,j}(x) J_{n,j}^\alpha(x) - \frac{2}{\alpha+1} \sum_{j=0}^n Q_{n,j}^{(\alpha+1)}(x).$$

By the mean value theorem, it follows


$$Q_{n,j}^{(\alpha+1)}(x) = J_{n,j}^{\alpha+1}(x) - J_{n,j+1}^{\alpha+1}(x) = (\alpha+1)p_{n,j}(x)\gamma_{n,j}^\alpha(x),$$

where

$$J_{n,j+1}^\alpha(x) < \gamma_{n,j}^\alpha(x) < J_{n,j}^\alpha(x).$$

Hence

$$\begin{aligned} \left| B_{n,\alpha}(\operatorname{sgn}(t-x), x) + \frac{\alpha-1}{\alpha+1} \right| &\leq 2 \sum_{j=0}^n p_{n,j}(x) (J_{n,j}^\alpha(x) - \gamma_{n,j}^\alpha(x)) \\ &\leq 2 \sum_{j=0}^n p_{n,j}(x) (J_{n,j}^\alpha(x) - J_{n,j+1}^\alpha(x)) \\ &\leq 2\alpha \sum_{j=0}^n p_{n,j}^2(x), \end{aligned}$$



**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 11 of 17

where we have used the inequality  $b^\alpha - a^\alpha < \alpha(b - a)$ ,  $0 \leq a, b \leq 1$  and  $\alpha \geq 1$ . Applying Lemma 2.2, we get

$$(3.2) \quad \left| B_{n,\alpha}(\operatorname{sgn}(t - x), x) + \frac{\alpha - 1}{\alpha + 1} \right| = \frac{2\alpha}{\sqrt{2enx(1 - x)}}, \quad x \in (0, 1).$$

Next we estimate  $B_{n,\alpha}(g_x, x)$ . By a Lebesgue-Stieltjes integral representation, we have


$$(3.3) \quad \begin{aligned} B_{n,\alpha}(g_x, x) &= \int_0^1 K_{n,\alpha}(x, t) g_x(t) dt \\ &= \left( \int_{I_1} + \int_{I_2} + \int_{I_3} \right) K_{n,\alpha}(x, t) g_x(t) dt \\ &= E_1 + E_2 + E_3, \quad \text{say,} \end{aligned}$$

where  $I_1 = [0, x - x/\sqrt{n}]$ ,  $I_2 = [x - x/\sqrt{n}, x + (1 - x)/\sqrt{n}]$  and  $I_3 = [x + (1 - x)/\sqrt{n}, 1]$ . We first estimate  $E_1$ . Writing  $y = x - x/\sqrt{n}$  and using Lebesgue-Stieltjes integration by parts, we have

$$E_1 = \int_0^y g_x(t) d_t(\lambda_{n,\alpha}(x, t)) = g_x(y+) \lambda_{n,\alpha}(x, y) - \int_0^y \lambda_{n,\alpha}(x, t) d_t(g_x(t)).$$

Since  $|g_x(y+)| \leq \bigvee_{y+}^x(g_x)$ , it follows that

$$|E_1| \leq \bigvee_{y+}^x(g_x) \lambda_{n,\alpha}(x, y) + \int_0^y \lambda_{n,\alpha}(x, t) d_t \left( -\bigvee_t^x(g_x) \right).$$




---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 17

By using (2.1) of Lemma 2.4, we get

$$|E_1| \leq \bigvee_{y+}^x (g_x) \frac{2\alpha \cdot x(1-x)}{n(x-y)^2} + \frac{2\alpha \cdot x(1-x)}{n} \int_0^y \frac{1}{(x-t)^2} dt \left( -\bigvee_t^x (g_x) \right).$$

Integrating by parts the last term we have after simple computation

$$|E_1| \leq \frac{2\alpha \cdot x(1-x)}{n} \left[ \frac{\bigvee_0^x (g_x)}{x^2} + 2 \int_0^y \frac{\bigvee_t^x (g_x)}{(x-t)^3} dt \right].$$

Now replacing the variable  $y$  in the last integral by  $x - x/\sqrt{t}$ , we obtain


$$(3.4) \quad |E_1| \leq \frac{2\alpha(1-x)}{nx} \left[ \bigvee_0^x (g_x) + \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^x (g_x) \right] \\ \leq \frac{4\alpha}{nx(1-x)} \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^x (g_x).$$

Using a similar method and (2.2) of Lemma 2.4, we get

$$(3.5) \quad |E_3| \leq \frac{4\alpha}{nx(1-x)} \sum_{k=1}^n \bigvee_x^{x+(1-x)/\sqrt{k}} (g_x).$$

Finally we estimate  $E_2$ . For  $t \in [x - x/\sqrt{n}, x + (1-x)/\sqrt{n}]$ , we have

$$|g_x(t)| = |g_x(t) - g_x(x)| \leq \bigvee_{x-x/\sqrt{n}}^{x+(1-x)/\sqrt{n}} (g_x),$$



**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 13 of 17

and therefore

$$|E_2| \leq \bigvee_{x-x/\sqrt{k}}^{x+(1-x)/\sqrt{k}} (g_x) \int_{x-x/\sqrt{k}}^{x+(1-x)/\sqrt{k}} d_t(\lambda_{n,\alpha}(x, t))$$

Since  $\int_a^b d_t \lambda_n(x, t) \leq 1$ , for all  $(a, b) \subseteq [0, 1]$ , therefore

$$(3.6) \quad |E_2| \leq \bigvee_{x-x/\sqrt{n}}^{x+(1-x)/\sqrt{n}} (g_x).$$


Collecting the estimates (3.3) – (3.6), we have

$$(3.7) \quad |B_{n,\alpha}(g_x, x)| \leq \frac{5\alpha}{nx(1-x)} \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^{x+(1-x)/\sqrt{k}} (g_x).$$

Combining the estimates of (3.1), (3.2) and (3.7), our theorem follows.  $\square$

For  $\alpha = 1$ , we obtain the following corollary, which is an improved estimate over the main results of [2] and [3].

**Corollary 3.2.** *Let  $f$  be a function of bounded variation on the interval  $[0, 1]$ .*




---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 14 of 17

Then for every  $x \in (0, 1)$  and  $n$  sufficiently large, we have

$$\left| B_n(f, x) - \frac{1}{2}[f(x+) + f(x-)] \right| \leq \frac{1}{\sqrt{2enx(1-x)}} |f(x+) - f(x-)| + \frac{5}{nx(1-x)} \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^{x+(1-x)/\sqrt{k}} (g_x).$$




---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back


Close

Quit

Page 15 of 17

## References

- [1] J.L. DURRMEYER, Une formule d'inversion de la transformee de Laplace: Application a la Theorie des Moments, These de 3e cycle, Faculte des Sciences de l' Universite de Paris 1967.
- [2] S. GUO, On the rate of convergence of Durrmeyer operator for function of bounded variation, *J. Approx. Theory*, **51** (1987), 183–192.
- [3] V. GUPTA, A note on the rate of convergence of Durrmeyer type operators for functions of bounded variation, *Soochow J. Math.*, **23**(1) (1997), 115–118.
- [4] V. GUPTA, Rate of convergence on Baskakov Beta Bezier operators for functions of bounded variation, *Int. J. Math. and Math. Sci.*, **32**(8) (2002), 471–479.
- [5] V. GUPTA, Rate of approximation by new sequence of linear positive operators, *Comput. Math. Appl.*, **45**(12) (2003), 1895–1904.
- [6] V. GUPTA, Degree of approximation to function of bounded variation by Bezier variant of MKZ operators, *J. Math. Anal. Appl.*, **289**(1) (2004), 292–300.
- [7] X.M. ZENG, Bounds for Bernstein basis functions and Meyer- Konig-Zeller basis functions, *J. Math. Anal. Appl.*, **219** (1998), 364–376.
- [8] X.M. ZENG, On the rate of convergence of the generalized Szasz type operators for functions of bounded variation, *J. Math. Anal. Appl.*, **226** (1998), 309–325.



---

### A Note on the Bezier Variant of Certain Bernstein Durrmeyer Operators

M.K. Gupta

---

Title Page

Contents



Go Back


Close

Quit

Page 16 of 17



- [9] X.M. ZENG AND W. CHEN, On the rate of convergence of the generalized Durrmeyer type operators for functions of bounded variation, *J. Approx. Theory*, **102** (2000), 1–12.



---

**A Note on the Bezier Variant of  
Certain Bernstein Durrmeyer  
Operators**

M.K. Gupta

---

Title Page

Contents



Go Back

Close

Quit

Page 17 of 17