


## NEWER APPLICATIONS OF GENERALIZED MONOTONE SEQUENCES

L. LEINDLER

Bolyai Institute, University of Szeged,  
Aradi vértanúk tere 1  
H-6720 Szeged, Hungary

*EMail:* [leindler@math.u-szeged.hu](mailto:leindler@math.u-szeged.hu)



---

volume 7, issue 5, article 167,  
2006.

*Received 03 July, 2006;  
accepted 06 December, 2006.*

*Communicated by: H. Bor*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

A particular result of Telyakovskii is extended to the newly defined class of numerical sequences and a specific problem is also highlighted. A further analogous result is also proved.


*2000 Mathematics Subject Classification:* 42A20, 40A05, 26D15.

*Key words:* Sine coefficients, Special sequences, Integrability.

The author was partially supported by the Hungarian National Foundation for Scientific Research under Grant # T042462 and TS44782.

## Contents

1	Introduction .....	3
2	Results .....	7
3	Notions and Notations .....	9
4	Proofs .....	11
	References	



---

### Newer Applications of Generalized Monotone Sequences

L. Leindler

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

# 1. Introduction

Recently several papers, see [4], [5] and [6], have dealt with the issue of uniform convergence and boundedness of monotone decreasing sequences. Further results and extensions have also been reported by the author in [6].

In this paper we shall give two further results on boundedness for wider classes of monotone sequences. First we present some theorems which will be useful in the following sections of this paper. In Section 2 we state the main results, in Section 3, we provide definitions and notations and in Section 4 we give detailed proofs of the main theorem and corollary.

In [7] S.A. Telyakovskii proved the following useful theorem.

**Theorem 1.1.** *If a sequence  $\{n_m\}$  of natural numbers ( $n_1 = 1 < n_2 < n_3 < \dots$ ) is such that*


$$(1.1) \quad \sum_{j=m}^{\infty} \frac{1}{n_j} \leq \frac{A}{n_m}$$

for all  $m = 1, 2, \dots$ , where  $A > 1$ , then the estimate

$$(1.2) \quad \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} \frac{\sin kx}{k} \right| \leq KA$$

holds for all  $x$ , where  $K$  is an absolute positive constant.

In [4], the author showed that the sequence  $\{k^{-1}\}$  in (1.2) can be replaced by any sequence  $\mathbf{c} := \{c_k\}$  which belongs to the class  $R_0^+ BVS$ .



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 3 of 12

Recently in [5] and [6] we verified as well that the sequence  $\{k^{-1}\}$  can be replaced by sequences which belong to either of the classes  $\gamma RBVS$  and  $\gamma GBVS$ .

More precisely we proved:

**Theorem 1.2 (see [6]).** *Let  $\gamma := \{\gamma_n\}$  be a sequence of nonnegative numbers satisfying the condition  $\gamma_n = O(n^{-1})$ ; furthermore let  $\alpha := \{\alpha_n\}$  be a similar sequence with the condition  $\alpha_n = o(n^{-1})$ . If  $c := \{c_n\} \in \gamma GBVS$ , or belongs to  $\alpha GBVS$ , furthermore, if the sequence  $\{n_m\}$  satisfies (1.1), then the estimates*

$$(1.3) \quad \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} c_k \sin kx \right| \leq K(\mathbf{c}, \{n_m\}),$$


or

$$(1.4) \quad \sum_{j=m}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} c_k \sin kx \right| = o(1), \quad m \rightarrow \infty,$$

hold uniformly in  $x$ , respectively.

We note that, in general, (1.3) does not imply (1.4) see the Remark in [5]. We also note that, every quasi geometrically increasing sequence  $\{n_m\}$  satisfies the inequality (1.1) (see [3, Lemma 1]).

A consequence of Theorem 1.1 shows that not only series (1.2) but also the Fourier series of any function of bounded variation possesses the property analogous to (1.2) (see [7, Theorem 2]).



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

Utilizing these results Telyakovskii [7] proved another theorem, which is an interesting variation of a theorem by W.H. Young [8].

This theorem reads as follows.

**Theorem 1.3.** *If the function  $f \in L(0, 2\pi)$  and the function  $g$  is of bounded variation on  $[0, 2\pi]$ , then the estimate*

$$(1.5) \quad \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} (a_k \alpha_k + b_k \beta_k) \right| \leq KA \|f\|_{LV}(g)$$

is valid for any sequence  $\{n_m\}$  with (1.1), where  $a_k, b_k$  and  $\alpha_k, \beta_k$  are the Fourier coefficients of  $f$  and  $g$ , respectively.

One can see that if we consider the function of bounded variation


$$g(x) := \frac{\pi - x}{2} = \sum_{k=1}^{\infty} \frac{\sin kx}{k}, \quad 0 < x < 2\pi,$$

then (1.5) reduces to

$$(1.6) \quad \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} \frac{b_k}{k} \right| \leq KA \|f\|_L,$$

which strengthens the well-known result by H. Lebesgue [2, p. 102] that the series

$$\sum_{k=1}^{\infty} \frac{b_k}{k}$$



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back

Close


Quit

Page 5 of 12

converges for the functions  $f \in L(0, 2\pi)$ .

These observations are made in [7] as well.

We have recalled (1.6) because one of our aims is to show that the sequence  $\{k^{-1}\}$  appearing in (1.6) can be replaced, as was the case in (1.2), by any sequence  $\{\beta_k\} \in \gamma GBVS$ , if  $\gamma_n = O(n^{-1})$ .



---

**Newer Applications of  
Generalized Monotone  
Sequences**

L. Leindler

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

## 2. Results

We prove the following assertions.

**Theorem 2.1.** *If the function  $f \in L(0, 2\pi)$  with  $\{b_k\}$  Fourier sine coefficients, the sequence  $\{n_m\}$  is quasi geometrically increasing, and the sequence  $\{\beta_k\}$  belongs to  $\gamma$ GBVS or  $\alpha$ GBVS, where  $\alpha$  and  $\gamma$  have the same definition as in Theorem 1.2, then*

$$(2.1) \quad \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k \right| \leq K(\{n_m\}, \{\beta_k\}) \|f\|_L,$$

or


$$(2.2) \quad \sum_{j=m}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k \right| = o(1), \quad m \rightarrow \infty,$$

hold, respectively.

**Remark 1.** *It is clear that if a sine series with coefficients  $\{\beta_n\} \in \gamma$ GBVS and  $\gamma_n = O(n^{-1})$ , that is, if the function*

$$g(x) := \sum_{k=1}^{\infty} \beta_k \sin kx$$

had a bounded variation, then (2.1) would be a special case of (1.5).



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12


The author is unaware of such a result, or its converse. It is an interesting open question.

Utilizing our result (2.1) and the method of Telyakovskii used in [7] we can also obtain estimates for  $E_n(f)_L$  and  $\omega_\nu(f, \delta)_L$ .

**Corollary 2.2.** *If  $f(x)$ ,  $\gamma$ ,  $\{b_k\}$ ,  $\{\beta_k\}$  and  $\{n_m\}$  are as in Theorem 2.1, then for any  $n$  with  $n_i \leq n < n_{i+1}$  the following estimates*

$$\begin{aligned}
 (2.3) \quad \omega_\nu \left( f, \frac{1}{n} \right)_L &\geq K(\nu) E_n(f)_L \\
 &\geq K(\nu, \{n_m\}, \{\beta_k\}) \left( \left| \sum_{k=n+1}^{n_{i+1}-1} b_k \beta_k \right| + \sum_{j=i+1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k \right| \right)
 \end{aligned}$$

*hold.*




---

**Newer Applications of  
Generalized Monotone  
Sequences**

L. Leindler

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12



### 3. Notions and Notations

A *positive* null-sequence  $\mathbf{c} := \{c_n\}$  ( $c_n \rightarrow 0$ ) belongs to the family of *sequences of rest bounded variation*, and briefly we write  $\mathbf{c} \in R_0^+ BVS$ , if

$$\sum_{n=m}^{\infty} |\Delta c_n| \leq K c_m, \quad (\Delta c_n = c_n - c_{n+1}),$$

holds for all  $m \in \mathbb{N}$ , where  $K = K(\mathbf{c})$  is a constant depending only on  $\mathbf{c}$ .

In this paper we shall use  $K$  to designate either an absolute constant or a constant depending on the indicated parameters, not necessarily the same at each occurrence.

Let  $\gamma := \{\gamma_n\}$  be a given *positive* sequence. A null-sequence  $\mathbf{c}$  of *real numbers* satisfying the inequality

$$\sum_{n=m}^{\infty} |\Delta c_n| \leq K \gamma_m$$

is said to be a *sequence of  $\gamma$  rest bounded variation*, represented by  $\mathbf{c} \in \gamma RBVS$ .


If  $\gamma$  is a given sequence of *nonnegative* numbers, the terms  $c_n$  are real and the inequality

$$\sum_{n=m}^{2m} |\Delta c_n| \leq K \gamma_m, \quad m = 1, 2, \dots$$

holds, then we write  $\mathbf{c} \in \gamma GBVS$ .

The class  $\gamma GBVS$  of sequences is wider than any one of the classes  $\gamma RBVS$  and  $GBVS$ . The class  $GBVS$  was defined in [1] by Le and Zhou with  $\gamma_m :=$

$\max_{m \leq n < m+N} |c_n|$ , where  $N$  is a natural number.



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back

Close

Quit


Page 9 of 12

A sequence  $\beta := \{\beta_n\}$  of positive numbers is called quasi geometrically increasing (decreasing) if there exist natural numbers  $\mu$  and  $K = K(\beta) \geq 1$  such that for all natural numbers  $n$ ,

$$\beta_{n+\mu} \geq 2\beta_n \text{ and } \beta_n \leq K \beta_{n+1} \quad \left( \beta_{n+\mu} \leq \frac{1}{2}\beta_n \text{ and } \beta_{n+1} \leq K \beta_n \right).$$

Let  $E_n(f)_L$  denote the best approximation of the function  $f$  in the metric  $L$  by trigonometric polynomials of order  $n$ ; and  $t_n(f, x)$  be a polynomial of best approximation of  $f(x)$  in the metric  $L$  by trigonometric polynomials of order  $n$ .

Finally denote by  $\omega_\nu(f, \delta)_L$  the integral modulus of continuity of order  $\nu$  of  $f \in L$ .




---

**Newer Applications of  
Generalized Monotone  
Sequences**

L. Leindler

---

Title Page

Contents



Go Back

Close

Quit

Page 10 of 12

## 4. Proofs

In this section we detail proofs of Theorem 2.1 and Corollary 2.2.

*Proof of Theorem 2.1.* It is clear that

$$\sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k = \frac{1}{\pi} \int_0^{2\pi} \sum_{k=n_j}^{n_{j+1}-1} \beta_k \sin kx \, dx.$$

Thus

$$\left| \sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k \right| \leq \frac{1}{\pi} \int_0^{2\pi} |f(x)| \left| \sum_{k=n_j}^{n_{j+1}-1} \beta \sin kx \right| dx.$$

Let us sum up these inequalities and apply the estimate (1.3) with  $\beta_k$  in place of  $c_k$ , we get that

$$\begin{aligned} \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} b_k \beta_k \right| &\leq \frac{1}{\pi} \int_0^{2\pi} |f(x)| \sum_{j=1}^{\infty} \left| \sum_{k=n_j}^{n_{j+1}-1} \beta_k \sin kx \right| dx \\ &\leq K(\{\beta_k\}, \{n_m\}) \|f\|_L, \end{aligned}$$


which proves (2.1).

If we sum only from  $m$  to infinity and use the assertion (1.4) instead of (1.3), we clearly obtain (2.2).

Herewith Theorem 2.1 is proved.  $\square$

*Proof of Corollary 2.2.* It is easy to see that Jackson's theorem and the estimate (2.1) with  $f(x) - t_n(f, x)$  in place of  $f(x)$  yield (2.3) immediately.  $\square$

An itemized reasoning can be found in [7].



Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

Title Page

Contents



Go Back


Close

Quit

Page 11 of 12

## References

- [1] R.J. LE AND S.P. ZHOU, A new condition for uniform convergence of certain trigonometric series, *Acta Math. Hungar.*, **10**(1-2) (2005), 161–169.
- [2] H. LEBESGUE, *Leçons sur les Séries Trigonometriques*, Paris: Gauthier-Villars, 1906.
- [3] L. LEINDLER, On the utility of power-monotone sequences, *Publ. Math. Debrecen*, **55**(1-2) (1999), 169–176.
- [4] L. LEINDLER, On the uniform convergence and boundedness of a certain class of sine series, *Analysis Math.*, **27** (2001), 279–285.
- [5] L. LEINDLER, A note on the uniform convergence and boundedness of a new class of sine series, *Analysis Math.*, **31** (2005), 269–275.
- [6] L. LEINDLER, A new extension of monotone sequences and its applications, *J. Inequal. Pure and Appl. Math.*, **7**(1) (2006), Art. 39.
- [7] S.A. TELYAKOVSKIĬ, On partial sums of Fourier series of functions of bounded variation, *Proc. Steklov Inst. Math.*, **219** (1997), 372–381.
- [8] W.H. YOUNG, On the integration of Fourier series, *Proc. London Math. Soc.*, **9** (1911), 449–462.



---

Newer Applications of  
Generalized Monotone  
Sequences

L. Leindler

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12