

# Journal of Inequalities in Pure and Applied Mathematics

## CERTAIN INEQUALITIES FOR CONVEX FUNCTIONS

P.G. POPESCU AND J.L. DI AZ-BARRERO


Automatics and Computer Science Faculty  
Politehnica University, București, Romania.

*EMail:* [pgpopescu@yahoo.com](mailto:pgpopescu@yahoo.com)

Applied Mathematics III  
Universitat Politècnica de Catalunya  
Jordi Girona 1-3, C2, 08034 Barcelona, Spain

*EMail:* [jose.luis.diaz@upc.edu](mailto:jose.luis.diaz@upc.edu)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
206-05



---

volume 7, issue 2, article 41,  
2006.

*Received 12 July, 2005;*  
*accepted 09 December, 2005.*

*Communicated by: S.S. Dragomir*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract


Classical inequalities like Jensen and its reverse are used to obtain some elementary numerical inequalities for convex functions. Furthermore, imposing restrictions on the data points several new constrained inequalities are given.

*2000 Mathematics Subject Classification:* 26D15.

*Key words:* Convex functions, Numerical Inequalities, Inequalities with constraints.

## Contents

1	Introduction .....	3
2	Unconstrained Inequalities .....	4
3	Constrained Inequalities .....	8
	References	



---

### Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

# 1. Introduction

It is well known ([1], [2]) that a continuous function,  $f$ , convex in a real interval  $I \subseteq \mathbb{R}$  has the property


$$(1.1) \quad f\left(\frac{1}{P_n} \sum_{k=1}^n p_k a_k\right) \leq \frac{1}{P_n} \sum_{k=1}^n p_k f(a_k),$$

where  $a_k \in I$ ,  $1 \leq k \leq n$  are given data points and  $p_1, p_2, \dots, p_n$  is a set of nonnegative real numbers constrained by  $\sum_{k=1}^n p_k = P_n$ . If  $f$  is concave the preceding inequality is reversed.

A broad consideration of inequalities for convex functions can be found, among others, in ([3], [4]). Furthermore, in [5] a reverse of Jensen's inequality is presented. It states that if  $p_1, p_2, \dots, p_n$  are real numbers such that  $p_1 > 0, p_k \leq 0$  for  $2 \leq k \leq n$  and  $P_n > 0$ , then

$$(1.2) \quad f\left(\frac{1}{P_n} \sum_{k=1}^n p_k a_k\right) \geq \frac{1}{P_n} \sum_{k=1}^n p_k f(a_k)$$

holds, where  $f : I \rightarrow \mathbb{R}$  is a convex function in  $I$  and  $a_k \in I$ ,  $1 \leq k \leq n$  are such that  $\frac{1}{P_n} \sum_{k=1}^n p_k x_k \in I$ . If  $f$  is concave (1.2) is reversed. Our aim in this paper is to use the preceding results to get new inequalities for convex functions. In addition, when the  $x_k$ 's are constrained some inequalities are obtained.



Title Page

Contents



Go Back

Close

Quit

Page 3 of 12

## 2. Unconstrained Inequalities

In the sequel, applying the preceding results and some numerical identities, some elementary inequalities are obtained. We begin with:

**Theorem 2.1.** *Let  $a_0, a_1, \dots, a_n$  be nonnegative real numbers. Then, the following inequality*

$$\exp \left[ \sum_{k=0}^n \binom{n}{k} \frac{a_k}{2^n} \right] \leq \frac{1}{8^n} \left[ \sum_{k=0}^n \binom{n}{k} \frac{e^{a_k}}{(1+a_k)^2} \right] \left[ \sum_{k=0}^n \binom{n}{k} (1+a_k) \right]^2$$

holds.


*Proof.* Since  $f(t) = \frac{e^t}{(1+t)^2}$  is convex in  $[0, +\infty)$ , then setting  $p_k = \binom{n}{k} / 2^n$ ,  $0 \leq k \leq n$ , into (1.1) and taking into account the well known identity  $\sum_{k=0}^n \binom{n}{k} = 2^n$ , we have

$$\exp \left( \sum_{k=0}^n \binom{n}{k} \frac{a_k}{2^n} \right) \left[ 1 + \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} a_k \right]^{-2} \leq \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \frac{e^{a_k}}{(1+a_k)^2}.$$

After rearranging terms, the inequality claimed immediately follows and the proof is complete.  $\square$

**Theorem 2.2.** *Let  $p_1, p_2, \dots, p_n$  be a set of nonnegative real numbers constrained by  $\sum_{k=1}^j p_k = P_j$ . If  $a_1, a_2, \dots, a_n$  are positive real numbers, then*

$$\left[ \prod_{k=1}^n \left( a_k + \sqrt{1+a_k^2} \right)^{p_k} \right]^{\frac{1}{P_n}} \leq \frac{1}{P_n} \sum_{k=1}^n p_k a_k + \sqrt{1 + \left( \frac{1}{P_n} \sum_{k=1}^n p_k a_k \right)^2}$$



Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

holds.

*Proof.* Let  $f : (0, +\infty) \rightarrow \mathbb{R}$  be the function defined by  $f(t) = \ln(t + \sqrt{1 + t^2})$ . Then, we have  $f'(t) = \frac{1}{\sqrt{1+t^2}} > 0$  and  $f''(t) = -\frac{t}{\sqrt{(1+t^2)^3}} \leq 0$ . Therefore,  $f$  is concave and applying (1.1) yields

$$\begin{aligned} & \ln \left[ \frac{1}{P_n} \sum_{k=1}^n p_k a_k + \sqrt{1 + \left( \frac{1}{P_n} \sum_{k=1}^n p_k a_k \right)^2} \right] \\ & \geq \frac{1}{P_n} \sum_{k=1}^n p_k \ln \left( a_k + \sqrt{1 + a_k^2} \right) = \ln \left[ \prod_{k=1}^n \left( a_k + \sqrt{1 + a_k^2} \right)^{p_k} \right]^{\frac{1}{P_n}}. \end{aligned}$$

Taking into account that  $f(t) = \log(t)$  is injective, the statement immediately follows and this completes the proof.  $\square$


Setting  $p_k = \frac{1}{n}$ ,  $1 \leq k \leq n$  into the preceding result we get

**Corollary 2.3.** *Let  $a_1, a_2, \dots, a_n$  be a set of positive real numbers. Then*

$$\prod_{k=1}^n \left( a_k + \sqrt{1 + a_k^2} \right)^{1/n} \leq \frac{1}{n} \left( \sum_{k=1}^n a_k + \sqrt{n^2 + \left( \sum_{k=1}^n a_k \right)^2} \right)$$

holds.

Let  $T_n$  be the  $n^{\text{th}}$  triangular number defined by  $T_n = \frac{n(n+1)}{2}$ . Then, setting  $a_k = T_k$ ,  $1 \leq k \leq n$  into the preceding result, we get




---

**Certain Inequalities for Convex Functions**

P.G. Popescu and  
J.L. Díaz-Barrero

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 12

**Corollary 2.4.** For all  $n \geq 1$ ,

$$\prod_{k=1}^n \left( T_k + \sqrt{1 + T_k^2} \right)^{\frac{1}{n}} \leq \frac{1}{3} \left( T_{n+1} + \sqrt{9 + T_{n+1}^2} \right)$$

holds.

An interesting result involving Fibonacci numbers that can be proved using convex functions is the following

**Theorem 2.5.** Let  $n$  be a positive integer and  $\ell$  be a whole number. Then,

$$(F_1^\ell + F_2^\ell + \dots + F_n^\ell) \left[ \frac{1}{F_1^{\ell-4}} + \frac{1}{F_2^{\ell-4}} + \dots + \frac{1}{F_n^{\ell-4}} \right] \geq F_n^2 F_{n+1}^2$$

holds, where  $F_n$  is the  $n^{\text{th}}$  Fibonacci number defined by  $F_0 = 0$ ,  $F_1 = 1$  and for all  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ .


*Proof.* Taking into account that  $F_1^2 + F_2^2 + \dots + F_n^2 = F_n F_{n+1}$ , as is well known, and the fact that the function  $f : (0, \infty) \rightarrow \mathbb{R}$ , defined by  $f(t) = 1/t$  is convex, we get after setting  $p_i = \frac{F_i^2}{F_n F_{n+1}}$ ,  $1 \leq i \leq n$  and  $a_i = F_n F_i^{\ell-2}$ ,  $1 \leq i \leq n$  :

$$\frac{1}{\frac{F_1^\ell}{F_{n+1}} + \frac{F_2^\ell}{F_{n+1}} + \dots + \frac{F_n^\ell}{F_{n+1}}} \leq \frac{1}{F_n^2 F_{n+1}} \left[ \frac{1}{F_1^{\ell-4}} + \frac{1}{F_2^{\ell-4}} + \dots + \frac{1}{F_n^{\ell-4}} \right].$$

From the preceding expression immediately follows

$$(F_1^\ell + F_2^\ell + \dots + F_n^\ell) \left[ \frac{1}{F_1^{\ell-4}} + \frac{1}{F_2^{\ell-4}} + \dots + \frac{1}{F_n^{\ell-4}} \right] \geq F_n^2 F_{n+1}^2,$$

and this completes the proof. □



### Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

Finally, using the reverse Jensen's inequality, we state and prove:

**Theorem 2.6.** Let  $a_0, a_1, \dots, a_n$  be positive real numbers such that  $a_0 \geq a_1 \geq \dots \geq a_n$  and let  $p_0 = n(n+1)$  and  $p_k = -k, k = 1, 2, \dots, n$ . Then

$$(2.1) \quad \left( \sum_{k=0}^n p_k a_k \right) \left( \sum_{k=0}^n \frac{p_k}{a_k} \right) \leq \binom{n+1}{2}^2.$$


*Proof.* Setting  $f(t) = \frac{1}{t}$ , that is convex in  $(0, +\infty)$ , and taking into account that  $\sum_{k=1}^n k = \frac{n(n+1)}{2}$  from (1.2) we have

$$f \left( \frac{2}{n(n+1)} \sum_{k=0}^n p_k a_k \right) \geq \frac{2}{n(n+1)} \sum_{k=0}^n p_k f(a_k)$$

or

$$\left( \frac{2}{n(n+1)} \sum_{k=0}^n p_k a_k \right)^{-1} \geq \frac{2}{n(n+1)} \sum_{k=0}^n \frac{p_k}{a_k}$$

from which, after rearranging terms, (2.1) immediately follows and the proof is complete.  $\square$




---

Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12

### 3. Constrained Inequalities

In the sequel, imposing restrictions on  $x_1, x_2, \dots, x_n$ , some inequalities with constraints are given. We begin with the following.

**Theorem 3.1.** Let  $p_1, p_2, \dots, p_n \in [0, 1)$  be a set of real numbers constrained by  $\sum_{k=1}^j p_k = P_j$ . If  $x_1, x_2, \dots, x_n$  are positive real numbers such that  $\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n} = 1$ , then

$$\left( \sum_{k=1}^n p_k x_k \right) \left( \sum_{k=1}^n \frac{1}{x_k^{p_k}} \right) \geq P_n^2$$

holds.

*Proof.* Taking into account the weighted AM-HM inequality, we have

$$\frac{1}{P_n} \sum_{k=1}^n p_k x_k \geq \frac{P_n}{\sum_{k=1}^n \left( \frac{p_k}{x_k} \right)}.$$


Since  $0 \leq p_k < 1$  for  $1 \leq k \leq n$ , then  $\frac{p_k}{x_k} \leq \frac{1}{x_k^{p_k}}$ . From which, we get

$$\frac{P_n}{\sum_{k=1}^n \left( \frac{p_k}{x_k} \right)} \geq \frac{P_n}{\sum_{k=1}^n \frac{1}{x_k^{p_k}}}.$$

Then,

$$\frac{1}{P_n} \sum_{k=1}^n p_k x_k \geq \frac{P_n}{\sum_{k=1}^n \frac{1}{x_k^{p_k}}}$$

and the statement immediately follows.  $\square$



Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12



**Corollary 3.2.** If  $x_1, x_2, \dots, x_n$  are positive real numbers such that  $\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n} = 1$ , then

$$\frac{1}{n} \leq \sum_{k=1}^n \frac{1}{x_k^{1/x_k}}.$$

*Proof.* Setting  $p_k = 1/x_k$ ,  $1 \leq k \leq n$  into Theorem 3.1 yields

$$\left( \sum_{k=1}^n p_k x_k \right) \left( \sum_{k=1}^n \frac{1}{x_k^{p_k}} \right) = n \left( \sum_{k=1}^n \frac{1}{x_k^{1/x_k}} \right) \geq \left( \sum_{k=1}^n \frac{1}{x_k} \right)^2 = 1$$

completing the proof. □


Finally, we give two inequalities similar to the ones obtained in [6] for the triangle.

**Theorem 3.3.** Let  $a, b$  and  $c$  be positive real numbers such that  $a + b + c = 1$ . Then, the following inequality

$$a^{a(a+2b)} \cdot b^{b(b+2c)} \cdot c^{c(c+2a)} \geq \frac{1}{3}$$

holds.

*Proof.* Since  $a + b + c = 1$ , then  $a^2 + b^2 + c^2 + 2(ab + bc + ca) = 1$ . Therefore, choosing  $p_1 = a^2$ ,  $p_2 = b^2$ ,  $p_3 = c^2$ ,  $p_4 = 2ab$ ,  $p_5 = 2bc$ ,  $p_6 = 2ca$  and  $x_1 = 1/a$ ,  $x_2 = 1/b$ ,  $x_3 = 1/c$ ,  $x_4 = 1/a$ ,  $x_5 = 1/b$ ,  $x_6 = 1/c$ , and applying




---

**Certain Inequalities for Convex Functions**

P.G. Popescu and  
J.L. Diaz-Barrero

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 12

Jensen's inequality to the function  $f(t) = \ln t$  that is concave for all  $t \geq 0$ , we obtain

$$\begin{aligned} \ln \left( a^2 \frac{1}{a} + b^2 \frac{1}{b} + c^2 \frac{1}{c} + 2ab \frac{1}{a} + 2bc \frac{1}{b} + 2ca \frac{1}{c} \right) \\ \geq a^2 \ln \frac{1}{a} + b^2 \ln \frac{1}{b} + c^2 \ln \frac{1}{c} + 2ab \ln \frac{1}{a} + 2bc \ln \frac{1}{b} + 2ca \ln \frac{1}{c}, \end{aligned}$$

from which, we get

$$\ln 3 \geq \ln \left( \frac{1}{a^a(a+2b) \cdot b^b(b+2c) \cdot c^c(c+2a)} \right)$$

and this completes the proof.  $\square$

**Theorem 3.4.** Let  $a, b, c$  be positive numbers such that  $ab + bc + ca = abc$ . Then,

$$\sqrt[b]{a} \sqrt[c]{b} \sqrt[a]{c} (a + b + c) \geq abc$$


holds.

*Proof.* Since  $ab + bc + ca = abc$ , then  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 1$ . So, choosing  $p_1 = \frac{1}{a}$ ,  $p_2 = \frac{1}{b}$ ,  $p_3 = \frac{1}{c}$  and  $x_1 = ab$ ,  $x_2 = bc$ ,  $x_3 = ca$ , and applying Jensen's inequality to  $f(t) = \ln t$  again, we get

$$\ln(a + b + c) \geq \frac{1}{a} \ln ab + \frac{1}{b} \ln bc + \frac{1}{c} \ln ca$$

or

$$a + b + c \geq a^{\frac{1}{a} + \frac{1}{c}} \cdot b^{\frac{1}{b} + \frac{1}{a}} \cdot c^{\frac{1}{c} + \frac{1}{b}}.$$



### Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

Title Page

Contents




Go Back

Close

Quit

Page 10 of 12

Now, taking into account that  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 1$ , we obtain:  $a + b + c \geq a^{1-\frac{1}{b}} \cdot b^{1-\frac{1}{c}} \cdot c^{1-\frac{1}{a}}$ , from which the statement immediately follows and the proof is complete.  $\square$



---

### Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

---

Title Page

Contents



Go Back


Close

Quit

Page 11 of 12

## References

- [1] J.L.W.V. JENSEN, Sur les fonctions convexes et les inégalités entre les valeurs moyennes, *Acta Math.*, **30** (1906), 175–193.
- [2] Th.M. RASSIAS, *Survey on Classical Inequalities*, Kluwer Academic Publishers, Dordrecht, 2000.
- [3] M. BENCZE, *Octagon Mathematical Magazine Collection*, 1993–2004.
- [4] J.F. STEFFENSEN, On certain inequalities and methods of approximation, *J. Institute Actuaries*, **51** (1919), 279–297.
- [5] P.M. VASIĆ AND J.E. PEČARIĆ, On Jensen inequality, *Univ. Beograd. Publ. Elektrotehn Fak. Ser. Mat. Fis.*, **639-677** (1979), 50–54.
- [6] J.L. DÍAZ-BARRERO, Some cyclical inequalities for the triangle, *J. Ineq. Pure and Appl. Math.*, **6**(1) (2005), Art. 20. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=489>]



---

### Certain Inequalities for Convex Functions

P.G. Popescu and  
J.L. Díaz-Barrero

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12