


# Journal of Inequalities in Pure and Applied Mathematics



## APPLICATIONS OF THE EXTENDED HERMITE-HADAMARD INEQUALITY

Z. RETKES

University of Szeged  
Bolyai Institute  
Aradi vértanúk tere 1  
Szeged, H-6720 Hungary.

EMail: [retkes@math.u-szeged.hu](mailto:retkes@math.u-szeged.hu)

volume 7, issue 1, article 24,  
2006.

*Received 28 September, 2005;  
accepted 13 November, 2005.*

*Communicated by: P.S. Bullen*

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



## Abstract

In this paper we give some combinatorial applications according to a new extension of the classical Hermite-Hadamard inequality proved in [1].

*2000 Mathematics Subject Classification:* 26D10, 49J40, 49K40.

*Key words:* Dynamical Systems, Monotone Trajectories, Generalized Jacobian, Variational Inequalities.

## Contents

|   |                               |    |
|---|-------------------------------|----|
| 1 | Introduction .....            | 3  |
| 2 | Asymptotical Formulae .....   | 5  |
| 3 | Formulae for Power Sums ..... | 10 |

### References

---

### Monotone Trajectories of Dynamical Systems and Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 2 of 15**

# 1. Introduction

In the paper [1], we have proved the following generalization of the classical Hermite-Hadamard inequality for convex functions, extended it to  $n$  nodes: Suppose that  $-\infty \leq a < b \leq \infty$ ,  $f : (a, b) \rightarrow \mathbb{R}$  is a strict convex function,  $x_i \in (a, b)$ ,  $i = 1, \dots, n$  such that  $x_i \neq x_j$  if  $1 \leq i < j \leq n$ . Then the following inequality holds:


$$\sum_{i=1}^n \frac{F^{(n-1)}(x_i)}{\Pi_i(x_1, \dots, x_n)} < \frac{1}{n!} \sum_{i=1}^n f(x_i).$$

In the concave case  $<$  sign is changed to  $>$ . Here  $\Pi_i(x_1, \dots, x_n) := \prod_{k=1, k \neq i}^n (x_i - x_k)$  and  $F^{(0)}(s), F^{(1)}(s), \dots, F^{(n-1)}(s), \dots$  is a sequence of functions defined recursively by  $F^{(0)}(s) = f(s)$  and  $\frac{d}{ds} F^{(n)}(s) = F^{(n-1)}(s)$ ,  $n = 1, 2, \dots$ . These sequences of functions of  $f$  are known as the iterated integrals of  $f$ . As an application of this main result we have proved the following identities:

$$(1.1) \quad \sum_{k=1}^n \frac{x_k^n}{\Pi_k(x_1, \dots, x_n)} = \sum_{k=1}^n x_k,$$

$$(1.2) \quad \sum_{k=1}^n \frac{x_k^{n-1}}{\Pi_k(x_1, \dots, x_n)} = 1,$$

$$(1.3) \quad \sum_{k=1}^n \frac{1}{x_k} = (-1)^{n-1} \prod_{k=1}^n x_k \sum_{k=1}^n \frac{1}{x_k^2 \Pi_k(x_1, \dots, x_n)} \quad (x_k \neq 0)$$



---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 3 of 15**

and

$$(1.4) \quad \prod_{k=1}^n \frac{1}{x_k} = (-1)^{n-1} \sum_{k=1}^n \frac{1}{x_k \Pi_k(x_1, \dots, x_n)} \quad (x_k \neq 0).$$

In this paper we apply the formulae above to obtain closed combinatorial formulae and to investigate the asymptotic behaviours of these sums. For the sake of the convenience of the reader we collect the transformational regulations for the quantity  $\Pi_k(x_1, \dots, x_n)$  which plays a significant role in all of the formulae. Note that all of them are simple consequences of the definition of  $\Pi_k(x_1, \dots, x_n)$ .

a) If  $\alpha \in \mathbb{R}$ , then

$$\Pi_k(\alpha + x_1, \dots, \alpha + x_n) = \Pi_k(x_1, \dots, x_n),$$

that is,  $\Pi_k$  is shift invariant.

b)

$$\Pi_k(\alpha x_1, \dots, \alpha x_n) = \alpha^{n-1} \Pi_k(x_1, \dots, x_n).$$

c)

$$\Pi_k(\alpha - x_1, \dots, \alpha - x_n) = (-1)^{n-1} \Pi_k(x_1, \dots, x_n).$$

d) If  $x_k \neq 0$ ,  $k = 1, \dots, n$ , then

$$\Pi_k\left(\frac{1}{x_1}, \dots, \frac{1}{x_n}\right) = (-1)^{n-1} \frac{\Pi_k(x_1, \dots, x_n)}{x_k^{n-2} \prod_{j=1}^n x_j}.$$



---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 4 of 15**

## 2. Asymptotical Formulae

a) Let  $1 < x_1 < \dots < x_n$  be variables and  $s_1 = x_1, s_2 = x_1 + x_2, \dots, s_n = x_1 + \dots + x_n$ . Then  $s_i \neq s_j$  if  $1 \leq i < j \leq n$ , hence formula (1.1) can be applied to  $s_1, \dots, s_n$ , consequently we have

$$(2.1) \quad \sum_{j=1}^n s_j = \sum_{j=1}^n \frac{s_j^n}{\Pi_j(s_1, \dots, s_n)}.$$

For the left-hand side of (2.1)


$$\sum_{j=1}^n s_j = x_1 + (x_1 + x_2) + \dots + (x_1 + \dots + x_n) = \sum_{j=1}^n j \cdot x_{n-j+1}.$$

The definition of  $\Pi_j$  implies

$$\begin{aligned} \Pi_j(s_1, \dots, s_n) &= (s_j - s_1) \cdots (s_j - s_{j-1})(s_j - s_{j+1}) \cdots (s_j - s_n) \\ &= (x_2 + \dots + x_j) \cdots (x_{j-1} + x_j) \cdot x_j (-1)^{n-j} x_{j+1} (x_{j+1} + x_{j+2}) \\ &\quad \cdots (x_{j+1} + \dots + x_n). \end{aligned}$$

Using the above expressions we have the following identities for all  $1 < x_1 < \dots < x_n$ :

$$(2.2) \quad \begin{aligned} \sum_{j=1}^n j \cdot x_{n-j+1} &= \sum_{j=1}^n \frac{(-1)^{n-j} (x_1 + \dots + x_j)^n}{(x_2 + \dots + x_j) \cdots (x_{j-1} + x_j) x_j x_{j+1} \cdots (x_{j+1} + \dots + x_n)}. \end{aligned}$$



---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 5 of 15**

Now, if  $x_k \rightarrow 1$ ,  $k = 1, \dots, n$  then (2.2) gives


$$\binom{n+1}{2} = \sum_{j=1}^n j = \sum_{j=1}^n \frac{(-1)^{n-j} j^n}{(j-1)!(n-j)!}.$$

Multiplying both sides by  $(n-1)!$  leads us to

$$\sum_{j=1}^n (-1)^{n-j} j^n \binom{n-1}{j-1} = \frac{(n+1)!}{2}.$$

b) Apply now formula (1.3) to  $x_k = 1 + \frac{k}{n}$ ,  $k = 1, \dots, n$ . Hence

$$\begin{aligned} \sum_{k=1}^n \frac{1}{1 + \frac{k}{n}} &= (-1)^{n-1} \prod_{k=1}^n \left(1 + \frac{k}{n}\right) \sum_{k=1}^n \frac{1}{(1 + \frac{k}{n})^2 \Pi_k(1 + \frac{1}{n}, \dots, 1 + \frac{n}{n})} \\ &= (-1)^{n-1} \frac{1}{n^n} \prod_{k=1}^n (n+k) \sum_{k=1}^n \frac{n^2}{(n+k)^2 \Pi_k(\frac{1}{n}, \frac{2}{n}, \dots, \frac{n}{n})} \\ &= (-1)^{n-1} \frac{1}{n^n} \prod_{k=1}^n (n+k) \sum_{k=1}^n \frac{n^2}{(n+k)^2 \frac{1}{n^{n-1}} \Pi_k(1, \dots, n)} \\ &= (-1)^{n-1} \prod_{k=1}^n (n+k) \sum_{k=1}^n \frac{n}{(n+k)^2 (k-1)! (-1)^{n-k} (n-k)!} \\ &= n \cdot \binom{2n}{n} \sum_{k=1}^n \frac{(-1)^{k-1}}{(n+k)^2} \binom{n-1}{k-1}. \end{aligned}$$




---

### Monotone Trajectories of Dynamical Systems and Clarke's Generalized Jacobian

Giovanni P. Crespi and  
[mrocca@eco.uninsubria.it](mailto:mrocca@eco.uninsubria.it)

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 6 of 15**

From this equality chain we conclude that

$$(2.3) \quad \binom{2n}{n} \sum_{k=1}^n \frac{(-1)^{k-1}}{(n+k)^2} \binom{n-1}{k-1} = \frac{1}{n} \sum_{k=1}^n \frac{1}{1 + \frac{k}{n}} = \sum_{k=1}^n \frac{1}{n+k}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \frac{1}{1 + \frac{k}{n}} = \int_0^1 \frac{1}{1+u} du = \log 2,$$

hence the left-hand side of (2.3) is convergent and

$$\lim_{n \rightarrow \infty} \binom{2n}{n} \sum_{k=1}^n \frac{(-1)^{k-1}}{(n+k)^2} \binom{n-1}{k-1} = \log 2.$$

Pursuant to the Stirling formula  $\binom{2n}{n} \sim \frac{4^n}{\sqrt{\pi n}}$  ( $n \rightarrow \infty$ ), consequently we have the following asymptotic expression for the sum on the left:

$$\sum_{k=1}^n \frac{(-1)^{k-1}}{(n+k)^2} \binom{n-1}{k-1} \sim \sqrt{\pi} \log 2 \frac{\sqrt{n}}{4^n} \quad (n \rightarrow \infty).$$

c) This example follows along the lines above but choosing  $x_k = 1 + \left(\frac{k}{n}\right)^2$  ( $k = 1, \dots, n$ ). Without the detailed computation we have

$$(2.4) \quad \frac{1}{n} \sum_{k=1}^n \frac{1}{1 + \left(\frac{k}{n}\right)^2} = 2n \frac{\prod_{k=1}^n (n^2 + k^2)}{(n!)^2} \sum_{k=1}^n \frac{(-1)^{k-1} k^2}{(n^2 + k^2)^2} \cdot \frac{\binom{n}{k}}{\binom{n+k}{n}}.$$




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 7 of 15](#)

From this equality the limit of the right-hand side exists and

$$\begin{aligned} \lim_{n \rightarrow \infty} 2n \cdot \frac{\prod_{k=1}^n (n^2 + k^2)}{(n!)^2} \sum_{k=1}^n \frac{(-1)^{k-1} k^2}{(n^2 + k^2)^2} \cdot \frac{\binom{n}{k}}{\binom{n+k}{n}} &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \frac{1}{1 + \left(\frac{k}{n}\right)^2} \\ &= \int_0^1 \frac{1}{1+u^2} du \\ &= \arctan 1 = \frac{\pi}{4}. \end{aligned}$$

Combining these observations we have


$$\sum_{k=1}^n \frac{(-1)^{k-1} k^2}{(n^2 + k^2)^2} \frac{\binom{n}{k}}{\binom{n+k}{n}} \sim \frac{\pi}{8n} \cdot \frac{(n!)^2}{\prod_{k=1}^n (n^2 + k^2)} \quad (n \rightarrow \infty).$$

Moreover  $(n!)^2 \sim 2\pi n \left(\frac{n}{e}\right)^{2n}$  and

$$\prod_{k=1}^n (n^2 + k^2) = n^{2n} \prod_{k=1}^n \left[ 1 + \left( \frac{k}{n} \right)^2 \right].$$

On the other hand

$$\begin{aligned} \frac{1}{n} \log \prod_{k=1}^n \left[ 1 + \left( \frac{k}{n} \right)^2 \right] &= \frac{1}{n} \sum_{k=1}^n \log \left[ 1 + \left( \frac{k}{n} \right)^2 \right] \\ &\rightarrow \int_0^1 \log(1+x^2) dx \quad (n \rightarrow \infty), \end{aligned}$$




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
[mrocca@eco.uninsubria.it](mailto:mrocca@eco.uninsubria.it)

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 8 of 15**

hence

$$\prod_{k=1}^n \left[ 1 + \left( \frac{k}{n} \right)^2 \right] \sim e^{n \int_0^1 \log(1+x^2) dx}.$$

Using integration by parts to evaluate the integral in the exponent gives

$$\int_0^1 \log(1+x^2) dx = \log 2 + \frac{\pi}{2} - 2 = \alpha > 0.$$

Putting together these results gives the following asymptotical expression:

$$\sum_{k=1}^n \frac{(-1)^{k-1} k^2}{(n^2 + k^2)^2} \frac{\binom{n}{k}}{\binom{n+k}{n}} \sim \frac{\pi^2}{4} e^{-n(\log 2 + \frac{\pi}{2})} \quad (n \rightarrow \infty).$$



---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 9 of 15**

### 3. Formulae for Power Sums

Let  $u_1, \dots, u_n \in \mathbb{R}$  such that  $u_i \neq 0$ ,  $|u_i| < 1$ ,  $u_i \neq u_j$  if  $1 \leq i < j \leq n$  and  $x_i = 1 - u_i$  ( $i = 1, \dots, n$ ). Then applying (1.3) gives

$$\begin{aligned} \sum_{i=1}^n \frac{1}{1-u_i} &= (-1)^{n-1} \prod_{k=1}^n (1-u_k) \sum_{i=1}^n \frac{1}{(1-u_i)^2 \prod_i (1-u_1, \dots, 1-u_n)} \\ &= \prod_{k=1}^n (1-u_k) \sum_{i=1}^n \frac{1}{(1-u_i)^2 \prod_i (u_1, \dots, u_n)}. \end{aligned}$$

Under the above conditions  $\frac{1}{1-u_i} = \sum_{\ell=0}^{\infty} u_i^{\ell}$ , hence

$$(3.1) \quad \sum_{i=1}^n \frac{1}{1-u_i} = \sum_{i=1}^n \sum_{\ell=0}^{\infty} u_i^{\ell} = \sum_{\ell=0}^{\infty} \sum_{i=1}^n u_i^{\ell} = \sum_{\ell=0}^{\infty} P(\ell),$$

where we use the denotation  $P(\ell) := u_1^{\ell} + \dots + u_n^{\ell}$  for ( $\ell = 0, 1, 2, \dots$ ), and thus we have the following identities for  $u_1, \dots, u_n$ :

$$(3.2) \quad \sum_{\ell=0}^{\infty} P(\ell) = \prod_{k=1}^n (1-u_k) \sum_{i=1}^n \frac{1}{(1-u_i)^2 \prod_i (u_1, \dots, u_n)}.$$

Apply now (3.2) to  $u_k = \frac{k}{n+1}$  ( $k = 1, \dots, n$ ). For the left-hand side:

$$\sum_{\ell=0}^{\infty} P(\ell) = \sum_{\ell=0}^{\infty} \sum_{k=1}^n \left( \frac{k}{n+1} \right)^{\ell} = \sum_{\ell=0}^{\infty} \frac{1}{(n+1)^{\ell}} \sum_{k=1}^n k^{\ell} = \sum_{\ell=0}^{\infty} \frac{\tilde{P}(\ell)}{(n+1)^{\ell}},$$




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
[mrocca@eco.uninsubria.it](mailto:mrocca@eco.uninsubria.it)

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 10 of 15**

where  $\tilde{P}(\ell) = 1^\ell + 2^\ell + \cdots + n^\ell$  ( $\ell = 0, 1, 2, \dots$ ). On the other hand

$$\prod_{k=1}^n \left(1 - \frac{k}{n+1}\right) = \frac{n!}{(n+1)^n} \quad \text{and}$$

$$\Pi_i \left( \frac{1}{n+1}, \dots, \frac{n}{n+1} \right) = \frac{(-1)^{n-i}(i-1)!(n-i)!}{(n+1)^{n-1}}.$$

Putting these expressions into the right-hand side of (3.2) we obtain

$$(3.3) \quad \sum_{\ell=0}^{\infty} \frac{\tilde{P}(\ell)}{(n+1)^\ell} = n(n+1) \sum_{j=1}^n \frac{(-1)^{j-1}}{j^2} \binom{n-1}{j-1}.$$

A simple rearrangement of (3.3) and using  $\frac{n}{j} \binom{n-1}{j-1} = \binom{n}{j}$  gives


$$\sum_{\ell=0}^{\infty} \frac{\tilde{P}(\ell)}{(n+1)^{\ell+1}} = \sum_{j=1}^n \frac{(-1)^{j-1}}{j} \binom{n}{j},$$

and since it is known that

$$\sum_{j=1}^n \frac{(-1)^{j-1}}{j} \binom{n}{j} = \int_0^1 \frac{1 - (1-s)^n}{s} ds = \sum_{k=1}^n \frac{1}{k},$$

consequently we have the following identity for all fixed  $n$ :

$$\sum_{\ell=0}^{\infty} \frac{\tilde{P}(\ell)}{(n+1)^{\ell+1}} = \sum_{k=1}^n \frac{1}{k}.$$




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 15 of 15

Let us start now with (3.1) substituting  $u_k = \frac{1}{x_k}$ ,  $x_k > 1$  ( $k = 1, \dots, n$ ) and  $x_i \neq x_j$  if  $1 \leq i < j \leq n$ . Using the same technique applied above and the transformational rule d) we get the following identity for  $x_1, \dots, x_n$ :

$$(3.4) \quad \sum_{\ell=0}^{\infty} \sum_{k=1}^n \frac{1}{x_k^\ell} = (-1)^{n-1} \prod_{k=1}^n (x_k - 1) \sum_{i=1}^n \frac{x_i^n}{(x_i - 1)^2 \prod_i (x_1, \dots, x_n)}.$$

Putting  $x_k = k + 1$  ( $k = 1, \dots, n$ ) in (3.4) and simplifying the right-hand side we have the following equality:


$$(3.5) \quad \sum_{\ell=0}^{\infty} \left[ \frac{1}{2^\ell} + \dots + \frac{1}{(n+1)^\ell} \right] = \sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i}.$$

Observe that  $\frac{1}{2^\ell} + \dots + \frac{1}{(n+1)^\ell} = \zeta^{[n+1]}(\ell) - 1$ , where  $\zeta^{[n+1]}(\ell)$  denotes the  $(n+1)$ th partial sum of the series  $\zeta(\ell) = \sum_{k=1}^{\infty} \frac{1}{k^\ell}$ , if  $\ell \geq 2$ . Moreover let  $H_n$  denote the  $n$ th partial sum of the harmonic series, that is  $H_n = \sum_{k=1}^n \frac{1}{k}$ . Using this quantity and separating the summands corresponding to  $\ell = 0$  and  $\ell = 1$  gives

$$(3.6) \quad n + H_{n+1} - 1 + \sum_{\ell=2}^{\infty} [\zeta^{[n+1]}(\ell) - 1] = \sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i}.$$

Rearranging (3.6) and taking the limit as  $n \rightarrow \infty$  yields that:

$$(3.7) \quad \sum_{\ell=2}^{\infty} [\zeta(\ell) - 1] = 1 + \lim_{n \rightarrow \infty} \left[ \sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i} - n - H_{n+1} \right].$$




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 12 of 15](#)

This relation prompted us to investigate the quantities arising in both sides of the equality. On one hand the sum of the series is equal to 1 as an easy computation shows below:

$$\begin{aligned}\sum_{\ell=2}^{\infty} [\zeta(\ell) - 1] &= \sum_{\ell=2}^{\infty} \sum_{k=2}^{\infty} \frac{1}{k^{\ell}} = \sum_{k=2}^{\infty} \frac{1}{k^2} \sum_{j=0}^{\infty} \frac{1}{k^j} \\ &= \sum_{k=2}^{\infty} \frac{1}{k^2} \frac{1}{1 - \frac{1}{k}} \\ &= \sum_{k=2}^{\infty} \frac{1}{k(k-1)} = 1.\end{aligned}$$

Hence


$$\lim_{n \rightarrow \infty} \left[ \sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i} - n - H_{n+1} \right] = 0.$$

In fact we prove more in the following lemma.

**Lemma 3.1.** *For all  $n \geq 1$  the identity below holds*

$$\sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i} = n + H_n.$$

*Proof.* In the paper [1] we proved that if  $x_1, \dots, x_n \in \mathbb{R}$  such that  $x_i \neq x_j$  if




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
[mrocca@eco.uninsubria.it](mailto:mrocca@eco.uninsubria.it)

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 13 of 15

$1 \leq i < j \leq n$  then

$$\sum_{i=1}^n \frac{x_i^j}{\Pi_i(x_i, \dots, x_n)} = \begin{cases} \sum_{k=1}^n x_k & \text{if } j = n \\ 1 & \text{if } j = n - 1 \\ 0 & \text{if } 0 \leq j \leq n - 2. \end{cases}$$

Applying this result to  $x_i = \frac{1}{i}$  ( $i = 1, \dots, n$ ) gives that

$$\sum_{i=1}^n \frac{(-1)^{i-1}}{i} i^{n-j} \binom{n}{i} = \begin{cases} H_n & \text{if } j = n \\ 1 & \text{if } j = n - 1 \\ 0 & \text{if } 0 \leq j \leq n - 2. \end{cases}$$

Moreover

$$\begin{aligned} \sum_{i=1}^n \frac{(-1)^{i-1}}{i} (i+1)^n \binom{n}{i} &= \sum_{i=1}^n \frac{(-1)^{i-1}}{i} \left[ \sum_{k=0}^n i^{n-k} \binom{n}{k} \right] \binom{n}{i} \\ &= \sum_{k=0}^n \left[ \sum_{i=1}^n \frac{(-1)^{i-1}}{i} i^{n-k} \binom{n}{i} \right] \binom{n}{k} \\ &= \binom{n}{n-1} + H_n \binom{n}{n} = n + H_n \end{aligned}$$

as we stated. □




---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
[mrocca@eco.uninsubria.it](mailto:mrocca@eco.uninsubria.it)

---

[Title Page](#)

[Contents](#)



[Go Back](#)


[Close](#)

[Quit](#)

Page 14 of 15

## References

- [1] Z. RETKES, An extension of the Hermite-Hadamard inequality (submitted)
- [2] S.S. DRAGOMIR AND C.E.M. PEARCE, *Selected Topics on Hermite-Hadamard Inequalities*, RGMIA Monographs, Victoria University, 2000.  
[ONLINE: <http://rgmia.vu.edu.au/monographs/index.html>]
- [3] W. RUDIN, *Real and Complex Analysis*, McGraw-Hill Book Co., 1966.
- [4] G.H. HARDY, J.E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge University Press, 1934.



---

Monotone Trajectories of  
Dynamical Systems and  
Clarke's Generalized Jacobian

Giovanni P. Crespi and  
mrocca@eco.uninsubria.it

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 15 of 15