

# Journal of Inequalities in Pure and Applied Mathematics



## SOME INEQUALITIES FOR THE EXPECTATION AND VARIANCE OF A RANDOM VARIABLE WHOSE PDF IS $n$ -TIME DIFFERENTIABLE

NEIL S. BARNETT, PIETRO CERONE, SEVER S. DRAGOMIR AND JOHN ROUMELIOTIS

School of Communications and Informatics  
Victoria University of Technology  
PO Box 14428, Melbourne City MC  
8001 Victoria, Australia

EMail: [Neil.Barnett@vu.edu.au](mailto:Neil.Barnett@vu.edu.au)  
URL: <http://sci.vu.edu.au/staff/neilb.html>

EMail: [Peter.Cerone@vu.edu.au](mailto:Peter.Cerone@vu.edu.au)  
URL: <http://sci.vu.edu.au/staff/peterc.html>

EMail: [Sever.Dragomir@vu.edu.au](mailto:Sever.Dragomir@vu.edu.au)  
URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>

EMail: [John.Roumeliotis@vu.edu.au](mailto:John.Roumeliotis@vu.edu.au)  
URL: <http://www.staff.vu.edu.au/johnr/>

volume 1, issue 2, article 21,  
2000.

Received 25 February, 2000;  
accepted 29 May, 2000.

Communicated by: *S.P. Singh*

Abstract

Contents




Home Page

Go Back

Close

Quit





## Abstract

Some inequalities for the expectation and variance of a random variable whose p.d.f. is  $n$ -time differentiable are given.

*2000 Mathematics Subject Classification:* 60E15, 26D15

*Key words:* Random Variable, Expectation, Variance, Dispersion

## Contents

1	Introduction .....	3
2	Some Preliminary Integral Identities .....	6
3	Some Inequalities .....	12
References		

---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 2 of 29**

# 1. Introduction

Let  $f : [a, b] \rightarrow \mathbb{R}_+$  be the p.d.f. of the random variable  $X$  and

$$E(X) := \int_a^b t f(t) dt$$

its *expectation* and

$$\sigma(X) = \left[ \int_a^b (t - E(X))^2 f(t) dt \right]^{\frac{1}{2}} = \left[ \int_a^b t^2 f(t) dt - [E(X)]^2 \right]^{\frac{1}{2}}$$

its dispersion or standard deviation.

In [1], using the identity

$$(1.1) \quad [x - E(X)]^2 + \sigma^2(X) = \int_a^b (x - t)^2 f(t) dt$$

and applying a variety of inequalities such as: Hölder's inequality, pre-Grüss, pre-Chebychev, pre-Lupaş, or Ostrowski type inequalities, a number of results concerning the expectation and variance of the random variable  $X$  were obtained.



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 3 of 29

For example,

$$(1.2) \quad \sigma^2(X) + [x - E(X)]^2$$

$$\leq \begin{cases} (b-a) \left[ \frac{(b-a)^2}{12} + \left( x - \frac{a+b}{2} \right)^2 \right] \|f\|_{\infty}, & \text{if } f \in L_{\infty}[a, b]; \\ \left[ \frac{(b-x)^{2q+1} + (x-a)^{2q+1}}{2q+1} \right]^{\frac{1}{q}} \|f\|_p, & \text{if } f \in L_p[a, b], \\ \left( \frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right)^2, & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \end{cases}$$

for all  $x \in [a, b]$ , which imply, amongst other things, that

$$(1.3)$$

$$0 \leq \sigma(X)$$

$$\leq \begin{cases} (b-a)^{\frac{1}{2}} \left[ \frac{(b-a)^2}{12} + [E(X) - \frac{a+b}{2}]^2 \right]^{\frac{1}{2}} \|f\|_{\infty}^{\frac{1}{2}}, & \text{if } f \in L_{\infty}[a, b]; \\ \left\{ \frac{[b-E(X)]^{2q+1} + [E(X)-a]^{2q+1}}{2q+1} \right\}^{\frac{1}{2q}} \|f\|_p^{\frac{1}{2}}, & \text{if } f \in L_p[a, b], \\ \frac{b-a}{2} + \left| E(X) - \frac{a+b}{2} \right|, & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \end{cases}$$

and

$$(1.4) \quad 0 \leq \sigma^2(X) \leq [b - E(X)][E(X) - a] \leq \frac{1}{4}(b-a)^2.$$

In this paper more accurate inequalities are obtained by assuming that the p.d.f. of  $X$  is  $n$ -time differentiable and that  $f^{(n)}$  is absolutely continuous on




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 4 of 29](#)

$[a, b]$ . For other recent results on the application of Ostrowski type inequalities in Probability Theory, see [2]-[4].



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 5 of 29

## 2. Some Preliminary Integral Identities

The following lemma, which is interesting in itself, holds.

**Lemma 2.1.** *Let  $X$  be a random variable whose probability distribution function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ . Then*

$$(2.1) \quad \sigma^2(X) + [E(X) - x]^2 = \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3) k!} f^{(k)}(x) + \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt$$

for all  $x \in [a, b]$ .


*Proof.* Is by Taylor's formula with integral remainder. Recall that

$$(2.2) \quad f(t) = \sum_{k=0}^n \frac{(t-x)^k}{k!} f^{(k)}(x) + \frac{1}{n!} \int_x^t (t-s)^n f^{(n+1)}(s) ds$$

for all  $t, x \in [a, b]$ .

Together with

$$(2.3) \quad \sigma^2(X) + [E(X) - x]^2 = \int_a^b (t-x)^2 f(t) dt,$$



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 6 of 29

where  $f$  is the p.d.f. of the random variable  $X$ , we obtain

(2.4)

$$\begin{aligned} \sigma^2(X) + [E(X) - x]^2 &= \int_a^b (t-x)^2 \left[ \sum_{k=0}^n \frac{(t-x)^k}{k!} f^{(k)}(x) + \frac{1}{n!} \int_x^t (t-s)^n f^{(n+1)}(s) ds \right] dt \\ &= \sum_{k=0}^n f^{(k)}(x) \int_a^b \frac{(t-x)^{k+2}}{k!} dt \\ &\quad + \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt \end{aligned}$$

and since

$$\int_a^b \frac{(t-x)^{k+2}}{k!} dt = \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3) k!},$$

the identity (2.4) readily produces (2.1) □

**Corollary 2.2.** *Under the above assumptions, we have*

(2.5)

$$\begin{aligned} \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 &= \sum_{k=0}^n \frac{\left[ 1 + (-1)^k \right] (b-a)^{k+3}}{2^{k+3} (k+3) k!} f^{(k)}\left(\frac{a+b}{2}\right) \\ &\quad + \frac{1}{n!} \int_a^b \left( t - \frac{a+b}{2} \right)^2 \left( \int_{\frac{a+b}{2}}^t (t-s)^n f^{(n+1)}(s) ds \right) dt. \end{aligned}$$




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 7 of 29**

The proof follows by using (2.4) with  $x = \frac{a+b}{2}$ .

**Corollary 2.3.** Under the above assumptions,

$$(2.6) \quad \begin{aligned} \sigma^2(X) + \frac{1}{2} [(E(X) - a)^2 + (E(X) - b)^2] \\ = \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \\ + \frac{1}{n!} \int_a^b \int_a^b K(t, s) (t-s)^n f^{(n+1)}(s) ds dt, \end{aligned}$$

where


$$K(t, s) := \begin{cases} \frac{(t-a)^2}{2} & \text{if } a \leq s \leq t \leq b, \\ -\frac{(t-b)^2}{2} & \text{if } a \leq t < s \leq b. \end{cases}$$

*Proof.* In (2.1), choose  $x = a$  and  $x = b$ , giving

$$(2.7) \quad \begin{aligned} \sigma^2(X) + [E(X) - a]^2 \\ = \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} f^{(k)}(a) + \frac{1}{n!} \int_a^b (t-a)^2 \left( \int_a^t (t-s)^n f^{(n+1)}(s) ds \right) dt \end{aligned}$$

and

$$(2.8) \quad \begin{aligned} \sigma^2(X) + [E(X) - b]^2 \\ = \sum_{k=0}^n \frac{(-1)^k (b-a)^{k+3}}{(k+3)k!} f^{(k)}(b) + \frac{1}{n!} \int_a^b (t-b)^2 \left( \int_b^t (t-s)^n f^{(n+1)}(s) ds \right) dt. \end{aligned}$$




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)

[◀◀](#)

[▶▶](#)

[◀](#)

[▶](#)

[Go Back](#)

[Close](#)

[Quit](#)

**Page 8 of 29**

Adding these and dividing by 2 gives (2.6).  $\square$

Taking into account that  $\mu = E(X) \in [a, b]$ , then we also obtain the following.

**Corollary 2.4.** *With the above assumptions,*

$$(2.9) \quad \sigma^2(X) = \sum_{k=0}^n \frac{(b-\mu)^{k+3} + (-1)^k (\mu-a)^{k+3}}{(k+3) k!} f^{(k)}(\mu) \\ + \frac{1}{n!} \int_a^b (t-\mu)^2 \left( \int_\mu^t (t-s)^n f^{(n+1)}(s) ds \right) dt.$$


*Proof.* The proof follows from (2.1) with  $x = \mu \in [a, b]$ .  $\square$

**Lemma 2.5.** *Let the conditions of Lemma 2.1 relating to  $f$  hold. Then the following identity is valid.*

$$(2.10) \quad \sigma^2(X) + [E(X) - x]^2 \\ = \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} + \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds,$$

where

$$(2.11) \quad K(x, s) = \begin{cases} (-1)^{n+1} \psi_n(s-a, x-s), & a \leq s \leq x \\ \psi_n(b-s, s-x), & x < s \leq b \end{cases}$$




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 9 of 29**

with

$$(2.12) \quad \psi_n(u, v) = \frac{u^{n+1}}{(n+3)(n+2)(n+1)} \cdot [(n+2)(n+1)u^2 + 2(n+3)(n+1)uv + (n+3)(n+2)v^2].$$


*Proof.* From (2.1), an interchange of the order of integration gives

$$\begin{aligned} & \frac{1}{n!} \int_a^b (t-x)^2 dt \int_x^t (t-s)^n f^{(n+1)}(s) ds \\ &= \frac{1}{n!} \left\{ - \int_a^x \int_a^s (t-x)^2 (t-s)^n f^{(n+1)}(s) dt ds \right. \\ & \quad \left. + \int_x^b \int_s^b (t-x)^2 (t-s)^n f^{(n+1)}(s) dt ds \right\} \\ &= \frac{1}{n!} \int_a^b \tilde{K}_n(x, s) f^{(n+1)}(s) ds, \end{aligned}$$

where

$$\tilde{K}_n(x, s) = \begin{cases} p_n(x, s) = - \int_a^s (t-x)^2 (t-s)^n dt, & a \leq s \leq x \\ q_n(x, s) = \int_s^b (t-x)^2 (t-s)^n dt, & x < s < b. \end{cases}$$

To prove the lemma it is sufficient to show that  $K \equiv \tilde{K}$ .




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 10 of 29](#)

Now,

$$\begin{aligned}\tilde{p}_n(x, s) &= - \int_a^s (t-x)^2 (t-s)^n dt = (-1)^{n+1} \int_0^{s-a} (u+x-s)^2 u^n du \\ &= (-1)^{n+1} \int_0^{s-a} [u^2 + 2(x-s)u + (x-s)^2] u^n du \\ &= (-1)^{n+1} \psi_n(s-a, x-s),\end{aligned}$$

where  $\psi(\cdot, \cdot)$  is as given by (2.12). Further,

$$\tilde{q}_n(x, s) = \int_s^b (t-x)^2 (t-s)^n dt = \int_0^{b-s} [u+(s-x)]^2 u^n du = \psi_n(b-s, s-x),$$

where, again,  $\psi(\cdot, \cdot)$  is as given by (2.12). Hence  $K \equiv \tilde{K}$  and the lemma is proved.  $\square$



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 11 of 29

### 3. Some Inequalities

We are now able to obtain the following inequalities.

**Theorem 3.1.** Let  $X$  be a random variable whose probability density function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ , then

$$(3.1) \quad \left| \sigma^2(X) + [E(X) - x]^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3) k!} f^{(k)}(x) \right| \\ \leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{(n+1)!(n+4)} [(x-a)^{n+4} + (b-x)^{n+4}], & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{n!(n+3+\frac{1}{q})} \frac{[(x-a)^{n+3+\frac{1}{q}} + (b-x)^{n+3+\frac{1}{q}}]}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{n!(n+3)} [(x-a)^{n+3} + (b-x)^{n+3}], & \end{cases}$$

for all  $x \in [a, b]$ , where  $\|\cdot\|_p$  ( $1 \leq p \leq \infty$ ) are the usual Lebesgue norms on  $[a, b]$ , i.e.,

$$\|g\|_\infty := \text{ess sup}_{t \in [a, b]} |g(t)| \quad \text{and} \quad \|g\|_p := \left( \int_a^b |g(t)|^p dt \right)^{\frac{1}{p}}, \quad p \geq 1.$$




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
n-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 12 of 29


*Proof.* By Lemma 2.1,

$$\begin{aligned}
 (3.2) \quad & \sigma^2(X) + [E(X) - x]^2 = \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k! (k+3)} f^{(k)}(x) \\
 & = \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt \\
 & := M(a, b; x).
 \end{aligned}$$

Clearly,

$$\begin{aligned}
 |M(a, b; x)| & \leq \frac{1}{n!} \int_a^b (t-x)^2 \left| \int_x^t (t-s)^n f^{(n+1)}(s) ds \right| dt \\
 & \leq \frac{1}{n!} \int_a^b (t-x)^2 \left[ \sup_{s \in [x,t]} |f^{(n+1)}(s)| \left| \int_x^t |t-s|^n ds \right| \right] dt \\
 & \leq \frac{\|f^{(n+1)}\|_\infty}{n!} \int_a^b \frac{(t-x)^2 |t-x|^{n+1}}{n+1} dt \\
 & = \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |t-x|^{n+3} dt \\
 & = \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \left[ \int_a^x (x-t)^{n+3} dt + \int_x^b (t-x)^{n+3} dt \right] \\
 & = \frac{\|f^{(n+1)}\|_\infty [(x-a)^{n+4} + (b-x)^{n+4}]}{(n+1)! (n+4)}
 \end{aligned}$$

and the first inequality in (3.1) is obtained.




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 13 of 29](#)

For the second, we use Hölder's integral inequality to obtain


$$\begin{aligned}
 |M(a, b; x)| &\leq \frac{1}{n!} \int_a^b (t-x)^2 \left| \int_x^t |t-s|^{nq} ds \right|^{\frac{1}{q}} \left| \int_x^t |f^{(n+1)}(s)|^p ds \right|^{\frac{1}{p}} dt \\
 &\leq \frac{1}{n!} \left( \int_a^b |f^{(n+1)}(s)|^p ds \right)^{\frac{1}{p}} \int_a^b (t-x)^2 |t-x|^{\frac{nq+1}{q}} dt \\
 &= \frac{1}{n!} \frac{\|f^{(n+1)}\|_p}{(nq+1)^{\frac{1}{q}}} \int_a^b |t-x|^{n+2+\frac{1}{q}} dt \\
 &= \frac{1}{n!} \frac{\|f^{(n+1)}\|_p}{(nq+1)^{\frac{1}{q}}} \left[ \frac{(b-x)^{n+3+\frac{1}{q}} + (x-a)^{n+3+\frac{1}{q}}}{n+3+\frac{1}{q}} \right].
 \end{aligned}$$

Finally, note that

$$\begin{aligned}
 |M(a, b; x)| &\leq \frac{1}{n!} \int_a^b (t-x)^2 |t-x|^n \left| \int_x^t |f^{(n+1)}(s)| ds \right| dt \\
 &\leq \frac{\|f^{(n+1)}\|_1}{n!} \int_a^b |t-x|^{n+2} dt \\
 &= \frac{\|f^{(n+1)}\|_1}{n!} \left[ \frac{(x-a)^{n+3} + (b-x)^{n+3}}{n+3} \right]
 \end{aligned}$$

and the third part of (3.1) is obtained.  $\square$

It is obvious that the best inequality in (3.1) is when  $x = \frac{a+b}{2}$ , giving Corollary 3.2.




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 14 of 29

**Corollary 3.2.** With the above assumptions on  $X$  and  $f$ ,

(3.3)

$$\begin{aligned} & \left| \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 - \sum_{k=0}^n \frac{\left[ 1 + (-1)^k \right] (b-a)^{k+3}}{2^{k+3} (k+3) k!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ & \leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{2^{n+3}(n+1)!(n+4)} (b-a)^{n+4}, & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{2^{n+2+\frac{1}{q}} n! (n+3+\frac{1}{q})} \frac{(b-a)^{n+3+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], p > 1, \\ & \quad \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{2^{n+2} n! (n+3)} (b-a)^{n+3}. \end{cases} \end{aligned}$$

The following corollary is interesting as it provides the opportunity to approximate the variance when the values of  $f^{(k)}(\mu)$  are known,  $k = 0, \dots, n$ .




---

Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is  $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)


Page 15 of 29

**Corollary 3.3.** With the above assumptions and  $\mu = \frac{a+b}{2}$ , we have

$$(3.4) \quad \left| \sigma^2(X) - \sum_{k=0}^n \frac{(b-\mu)^{k+3} + (-1)^k (\mu-a)^{k+3}}{(k+3) k!} f^{(k)}(\mu) \right| \\ \leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{(n+1)!(n+4)} [(\mu-a)^{n+4} + (b-\mu)^{n+4}], & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{n!(n+3+\frac{1}{q})} \frac{[(\mu-a)^{n+3+\frac{1}{q}} + (b-\mu)^{n+3+\frac{1}{q}}]}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{n!(n+3)} [(\mu-a)^{n+3} + (b-\mu)^{n+3}]. \end{cases}$$

The following result also holds.

**Theorem 3.4.** Let  $X$  be a random variable whose probability density function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 16 of 29](#)

$[a, b]$ , then

$$(3.5) \quad \left| \sigma^2(X) + \frac{1}{2} [(E(X) - a)^2 + (E(X) - b)^2] \right. \\ \left. - \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \right| \\ \leq \begin{cases} \frac{1}{(n+4)(n+1)!} \|f^{(n+1)}\|_\infty (b-a)^{n+4}, & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{2^{1/q-1}}{n!(qn+1)^{1/q}[(n+2)q+2]^{1/q}} \|f^{(n+1)}\|_p \frac{(b-a)^{n+3+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ \frac{1}{2n!} \|f^{(n+1)}\|_1 (b-a)^{n+3}, & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \end{cases}$$

where  $\|\cdot\|_p$  ( $1 \leq p \leq \infty$ ) are the usual Lebesgue  $p$ -norms.

*Proof.* Using Corollary 2.3

$$\left| \sigma^2(X) + \frac{1}{2} [(E(X) - a)^2 + (E(X) - b)^2] \right. \\ \left. - \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \right| \\ \leq \frac{1}{n!} \int_a^b \int_a^b |K(t, s)| |t-s|^n |f^{(n+1)}(s)| ds dt \\ =: N(a, b).$$




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)


[Page 17 of 29](#)

It is obvious that

$$N(a, b)$$

$$\begin{aligned} &\leq \|f^{(n+1)}\|_{\infty} \frac{1}{n!} \int_a^b \int_a^b |K(t, s)| |t - s|^n ds dt \\ &= \|f^{(n+1)}\|_{\infty} \frac{1}{n!} \int_a^b \left( \int_a^t |K(t, s)| |t - s|^n ds + \int_t^b |K(t, s)| |t - s|^n ds \right) dt \\ &= \frac{1}{n!} \|f^{(n+1)}\|_{\infty} \int_a^b \left[ \frac{(t-a)^2}{2} \cdot \frac{(t-a)^{n+1}}{n+1} + \frac{(t-b)^2}{2} \cdot \frac{(b-t)^{n+1}}{n+1} \right] dt \\ &= \frac{1}{2(n+1)!} \|f^{(n+1)}\|_{\infty} \int_a^b [(t-a)^{n+3} + (b-t)^{n+3}] dt \\ &= \frac{1}{2(n+1)!} \|f^{(n+1)}\|_{\infty} \left[ \frac{(b-a)^{n+4}}{n+4} + \frac{(b-a)^{n+4}}{n+4} \right] \\ &= \frac{\|f^{(n+1)}\|_{\infty}}{(n+4)(n+1)!} (b-a)^{n+4} \end{aligned}$$

so the first part of (3.5) is proved.



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)


[Close](#)

[Quit](#)

Page 18 of 29

Using Hölder's integral inequality for double integrals,

$$\begin{aligned}
 N(a, b) &\leq \frac{1}{n!} \left( \int_a^b \int_a^b |f^{(n+1)}(s)|^p ds dt \right)^{\frac{1}{p}} \times \left( \int_a^b \int_a^b |K(t, s)|^q |t - s|^{qn} ds dt \right)^{\frac{1}{q}} \\
 &= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \left[ \int_a^b \left( \int_a^t |K(t, s)|^q |t - s|^{qn} ds \right. \right. \\
 &\quad \left. \left. + \int_t^b |K(t, s)|^q |t - s|^{qn} ds \right) dt \right]^{\frac{1}{q}} \\
 &= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \left[ \int_a^b \left[ \frac{(t-a)^{2q}}{2^q} \int_a^t |t-s|^{qn} ds \right. \right. \\
 &\quad \left. \left. + \frac{(t-b)^{2q}}{2^q} \int_t^b |t-s|^{qn} ds \right] dt \right]^{\frac{1}{q}} \\
 &= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \left[ \int_a^b \left[ \frac{(t-a)^{2q} (t-a)^{qn+1}}{2^q (qn+1)} + \frac{(t-b)^{2q} (b-t)^{qn+1}}{2^q (qn+1)} \right] dt \right]^{\frac{1}{q}} \\
 &= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \cdot \left[ \frac{1}{2^q (qn+1)} \right]^{\frac{1}{q}} \left[ \int_a^b (t-a)^{(n+2)q+1} dt \right. \\
 &\quad \left. + \int_a^b (b-t)^{(n+2)q+1} dt \right]^{\frac{1}{q}}
 \end{aligned}$$




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 19 of 29

$$\begin{aligned}
&= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \cdot \left[ \frac{1}{2^q (qn+1)} \right]^{\frac{1}{q}} \left[ \frac{(b-a)^{(n+2)q+2}}{(n+2)q+2} + \frac{(b-a)^{(n+2)q+2}}{(n+2)q+2} \right]^{\frac{1}{q}} \\
&= \frac{2^{1/q} \|f^{(n+1)}\|_p (b-a)^{n+2+\frac{1}{p}+\frac{2}{q}}}{n! 2 (qn+1)^{\frac{1}{q}} ((n+2)q+2)^{\frac{1}{q}}} \\
&= \frac{2^{1/q-1} \|f^{(n+1)}\|_p \left[ (b-a)^{n+3+\frac{1}{q}} \right]}{n! (qn+1)^{\frac{1}{q}} [(n+2)q+2]^{\frac{1}{q}}}
\end{aligned}$$

and the second part of (3.5) is proved.

Finally, we observe that

$$\begin{aligned}
N(a,b) &\leq \frac{1}{n!} \sup_{(t,s) \in [a,b]^2} |K(t,s)| |t-s|^n \int_a^b \int_a^b |f^{(n+1)}(s)| ds dt \\
&= \frac{1}{n!} \frac{(b-a)^2}{2} \cdot (b-a)^n (b-a) \int_a^b |f^{(n+1)}(s)| ds \\
&= \frac{1}{2n!} (b-a)^{n+3} \|f^{(n+1)}\|_1,
\end{aligned}$$

which is the final result of (3.5).  $\square$

The following particular case can be useful in practical applications. For




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

<a href="#">Title Page</a>	
<a href="#">Contents</a>	
<a href="#"></a>	<a href="#"></a>
<a href="#"></a>	<a href="#"></a>
<a href="#">Go Back</a>	
<a href="#">Close</a>	
<a href="#">Quit</a>	
<a href="#">Page 20 of 29</a>	

$n = 0$ , (3.1) becomes

$$(3.6) \quad \left| \sigma^2(X) + [E(X) - x]^2 - (b - a) \left[ \left( x - \frac{a+b}{2} \right)^2 + \frac{(b-a)^2}{12} \right] f(x) \right| \\ \leq \begin{cases} \frac{\|f'\|_\infty}{4} [(x-a)^4 + (b-x)^4], & \text{if } f' \in L_\infty[a,b]; \\ \frac{q\|f'\|_p}{3q+1} \left[ (x-a)^{3+\frac{1}{q}} + (b-x)^{3+\frac{1}{q}} \right], & \text{if } f' \in L_p[a,b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \|f'\|_1 \left[ \frac{(b-a)^2}{12} + \left( x - \frac{a+b}{2} \right)^2 \right], & \end{cases}$$

for all  $x \in [a, b]$ . In particular, for  $x = \frac{a+b}{2}$ ,

$$(3.7) \quad \left| \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 - \frac{(b-a)^3}{12} f\left(\frac{a+b}{2}\right) \right| \\ \leq \begin{cases} \frac{\|f'\|_\infty}{32} (b-a)^4, & \text{if } f' \in L_\infty[a,b]; \\ \frac{q\|f'\|_p (b-a)^{3+\frac{1}{q}}}{2^{2+\frac{1}{q}}(3q+1)}, & \text{if } f' \in L_p[a,b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f'\|_1}{12} (b-a)^3, & \end{cases}$$

which is, in a sense, the best inequality that can be obtained from (3.6). If in




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 21 of 29

(3.6)  $x = \mu = E(X)$ , then

$$(3.8) \quad \left| \sigma^2(X) - (b-a) \left[ \left( E(X) - \frac{a+b}{2} \right)^2 + \frac{(b-a)^2}{12} \right] f(E(X)) \right| \\ \leq \begin{cases} \frac{\|f'\|_\infty}{4} [(E(X)-a)^4 + (b-E(X))^4], & \text{if } f' \in L_\infty[a,b]; \\ \frac{\|f'\|_p}{(3+\frac{1}{q})} [(E(X)-a)^4 + (b-E(X))^4], & \text{if } f' \in L_p[a,b], p > 1, \\ \|f'\|_1 \left[ \frac{(b-a)^2}{12} + \left( E(X) - \frac{a+b}{2} \right)^2 \right]. \end{cases}$$

In addition, from (3.5),

$$(3.9) \quad \left| \sigma^2(X) + \frac{1}{2} [(E(X)-a)^2 + (E(X)-b)^2] - \frac{(b-a)^3}{3} \left[ \frac{f(a) + f(b)}{2} \right] \right| \\ \leq \begin{cases} \frac{1}{4} \|f'\|_\infty (b-a)^4, & \text{if } f' \in L_\infty[a,b]; \\ \frac{1}{n! 2^{\frac{1}{q}} (q+1)^{\frac{1}{q}}} \|f'\|_p (b-a)^{3+\frac{1}{q}}, & \text{if } f' \in L_p[a,b], p > 1, \\ \frac{1}{2} \|f'\|_1 (b-a)^3, \end{cases}$$

which provides an approximation for the variance in terms of the expectation and the values of  $f$  at the end points  $a$  and  $b$ .





---

Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is  $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---



**Theorem 3.5.** Let  $X$  be a random variable whose p.d.f.  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ . Then

(3.10)

$$\left| \sigma^2(X) + (E(X) - x)^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} \right| \\ \leq \begin{cases} [(x-a)^{n+4} + (b-x)^{n+4}] \frac{\|f^{(n+1)}\|_\infty}{(n+1)!(n+4)}, & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ C^{\frac{1}{q}} \left[ (x-a)^{(n+3)q+1} + (b-x)^{(n+3)q+1} \right]^{\frac{1}{q}} \frac{\|f^{(n+1)}\|_p}{n!}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ p > 1; \\ \left[ \frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right]^{n+3} \cdot \frac{\|f^{(n+1)}\|_1}{n!(n+3)}, & \end{cases}$$


where

$$(3.11) \quad C = \int_0^1 \left[ \frac{u^{n+3}}{n+3} + 2(1-u) \frac{u^{n+2}}{n+2} + (1-u)^2 \frac{u^{n+1}}{n+1} \right]^q du.$$

*Proof.* From (2.10),

(3.12)

$$\left| \sigma^2(X) + (E(X) - x)^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} \right| \\ = \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right|.$$




---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 23 of 29

Now, on using the fact that from (2.11), (2.12),  $\psi_n(u, v) \geq 0$  for  $u, v \geq 0$ ,

$$(3.13) \quad \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \leq \frac{\|f^{(n+1)}\|_\infty}{n!} \left\{ \int_a^x \psi_n(s-a, x-s) ds + \int_x^b \psi_n(b-s, s-x) ds \right\}.$$

Further,

$$(3.14) \quad \psi_n(u, v) = \frac{u^{n+3}}{n+3} + 2v \frac{u^{n+2}}{n+2} + v^2 \frac{u^{n+1}}{n+1}$$

and so


$$(3.15)$$

$$\begin{aligned} & \int_a^x \psi_n(s-a, x-s) ds \\ &= \int_a^x \left[ \frac{(s-a)^{n+3}}{n+3} + 2(x-s) \frac{(s-a)^{n+2}}{n+2} + (x-s)^2 \frac{(s-a)^{n+1}}{n+1} \right] ds \\ &= (x-a)^{n+4} \int_0^1 \left[ \frac{\lambda^{n+3}}{n+3} + 2(1-\lambda) \frac{\lambda^{n+2}}{n+2} + (1-\lambda)^2 \frac{\lambda^{n+1}}{n+1} \right] d\lambda, \end{aligned}$$

where we have made the substitution  $\lambda = \frac{s-a}{x-a}$ .

Collecting powers of  $\lambda$  gives

$$\lambda^{n+3} \left[ \frac{1}{n+3} - \frac{2}{n+2} + \frac{1}{n+1} \right] - \frac{2\lambda^{n+2}}{(n+2)(n+1)} + \frac{\lambda^{n+1}}{n+1}$$




---

### Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)


[Page 24 of 29](#)

and so, from (3.15),

$$(3.16) \quad \begin{aligned} & \int_a^x \psi_n(s-a, x-s) ds \\ &= (x-a)^{n+4} \left\{ \frac{1}{n+4} \left[ \frac{1}{n+3} - \frac{2}{n+2} + \frac{1}{n+1} \right] \right. \\ & \quad \left. - \frac{2}{(n+3)(n+2)(n+1)} + \frac{1}{(n+2)(n+1)} \right\} \\ &= \frac{(x-a)^{n+4}}{(n+4)(n+1)}. \end{aligned}$$

Similarly, on using (3.14),

$$\begin{aligned} & \int_x^b \psi_n(b-s, s-x) ds \\ &= \int_x^b \left[ \frac{(b-s)^{n+3}}{n+3} + 2(s-x) \frac{(b-s)^{n+2}}{n+2} + (s-x)^2 \frac{(b-s)^{n+1}}{n+1} \right] ds \end{aligned}$$



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
*n*-Time Differentiable

---

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 25 of 29

and making the substitution  $\nu = \frac{b-s}{b-x}$  gives

(3.17)

$$\begin{aligned} & \int_x^b \psi_n(b-s, s-x) ds \\ &= (b-x)^{n+4} \int_0^1 \left[ \frac{\nu^{n+3}}{n+3} + 2(1-\nu) \frac{\nu^{n+2}}{n+2} + (1-\nu)^2 \frac{\nu^{n+1}}{n+1} \right] d\nu \\ &= \frac{(b-x)^{n+4}}{(n+4)(n+1)}, \end{aligned}$$

where we have used (3.15) and (3.16). Combining (3.16) and (3.17) gives the first inequality in (3.10). For the second inequality in (3.10), we use Hölder's integral inequality to obtain

(3.18)

$$\left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \leq \frac{\|f^{(n+1)}(s)\|_p}{n!} \left( \int_a^b |K_n(x, s)|^q ds \right)^{\frac{1}{q}}.$$

Now, from (2.11) and (3.14)

$$\begin{aligned} \int_a^b |K_n(x, s)|^q ds &= \int_a^x \psi^q(s-a, x-s) ds + \int_x^b \psi^q(b-s, s-x) ds \\ &= C \left[ (x-a)^{(n+3)q+1} + (b-x)^{(n+3)q+1} \right], \end{aligned}$$

where  $C$  is as defined in (3.11) and we have used (3.15) and (3.16). Substitution into (3.18) gives the second inequality in (3.10).




---

Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is  $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

<a href="#">Title Page</a>	
<a href="#">Contents</a>	
<a href="#"></a>	<a href="#"></a>
<a href="#"></a>	<a href="#"></a>
<a href="#">Go Back</a>	
<a href="#">Close</a>	
<a href="#">Quit</a>	
<a href="#">Page 26 of 29</a>	

Finally, for the third inequality in (3.10). From (3.12),

(3.19)

$$\begin{aligned} & \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \\ & \leq \frac{1}{n!} \left\{ \int_a^x \psi_n(s-a, x-s) |f^{(n+1)}(s)| ds + \int_x^b \psi_n(b-s, s-x) |f^{(n+1)}(s)| ds \right\} \\ & \leq \frac{1}{n!} \left\{ \psi_n(x-a, 0) \int_a^x |f^{(n+1)}(s)| ds + \psi_n(b-x, 0) \int_x^b |f^{(n+1)}(s)| ds \right\}, \end{aligned}$$

where, from (3.14),

$$(3.20) \quad \psi_n(u, 0) = \frac{u^{n+3}}{n+3}.$$


Hence, from (3.19) and (3.20)

$$\begin{aligned} & \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \leq \frac{1}{n!} \max \left\{ \frac{(x-a)^{n+3}}{n+3}, \frac{(b-x)^{n+3}}{n+3} \right\} \|f^{(n+1)}(\cdot)\|_1 \\ & = \frac{1}{n!(n+3)} [\max \{x-a, b-x\}]^{n+3} \|f^{(n+1)}(\cdot)\|_1, \end{aligned}$$

which, on using the fact that for  $X, Y \in \mathbb{R}$

$$\max \{X, Y\} = \frac{X+Y}{2} + \left| \frac{X-Y}{2} \right|$$

gives, from (3.12), the third inequality in (3.10). The theorem is now completely proved.  $\square$




---

Some Inequalities for the Expectation and Variance of a Random Variable whose PDF is  $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

<a href="#">Title Page</a>	
<a href="#">Contents</a>	
<a href="#"></a>	<a href="#"></a>
<a href="#"></a>	<a href="#"></a>
<a href="#">Go Back</a>	
<a href="#">Close</a>	
<a href="#">Quit</a>	
<a href="#">Page 27 of 29</a>	

*Remark 3.1.* The results of Theorem 3.5 may be compared with those of Theorem 3.1. Theorem 3.5 is based on the single integral identity developed in Lemma 2.5, while Theorem 3.1 is based on the double integral identity representation for the bound. It may be noticed from (3.1) and (3.10) that the bounds are the same for  $f^{(n+1)} \in L_\infty [a, b]$ , that for  $f^{(n+1)} \in L_1 [a, b]$  (which is always true since  $f^{(n)}$  is absolutely continuous) the bound obtained in (3.1) is better and for  $f^{(n+1)} \in L_p [a, b]$ ,  $p > 1$ , the result is inconclusive.



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

---

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 28 of 29

## References

- [1] N.S. BARNETT, P. CERONE, S.S. DRAGOMIR AND J. ROUMELIOTIS, Some inequalities for the dispersion of a random variable whose p.d.f. is defined on a finite interval, *J. Inequal. Pure and Appl. Math.*, accepted for publication.
- [2] N.S. BARNETT AND S.S. DRAGOMIR, An inequality of Ostrowski type for cumulative distribution functions, *Kyungpook Math. J.*, **39**(2) (1999), 303–311.  
[ONLINE] (Preprint available) <http://rgmia.vu.edu.au/v1n1.html>
- [3] N.S. BARNETT AND S.S. DRAGOMIR, An Ostrowski type inequality for a random variable whose probability density function belongs to  $L_\infty [a, b]$ , *Nonlinear Anal. Forum*, **5** (2000), 125–135.  
[ONLINE] (Preprint available) <http://rgmia.vu.edu.au/v1n1.html>
- [4] S.S. DRAGOMIR, N.S. BARNETT AND S. WANG, An Ostrowski type inequality for a random variable whose probability density function belongs to  $L_p [a, b]$ ,  $p > 1$ , *Math. Inequal. Appl.*, **2**(4) (1999), 501–508.



---

Some Inequalities for the  
Expectation and Variance of a  
Random Variable whose PDF is  
 $n$ -Time Differentiable

Neil S. Barnett, Pietro Cerone,  
Sever S. Dragomir and  
John Roumeliotis

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 29 of 29