

# Journal of Inequalities in Pure and Applied Mathematics

## REGULARITY RESULTS FOR VECTOR FIELDS OF BOUNDED DISTORTION AND APPLICATIONS

ALBERTO FIORENZA AND FLAVIA GIANNETTI

Dipartimento di Costruzioni e Metodi Matematici in Architettura  
via Monteoliveto 3  
-80134 Napoli, ITALY  
*E*Mail: [fiorenza@unina.it](mailto:fiorenza@unina.it)  
*U*RL: <http://cds.unina.it/~fiorenza/>

Dipartimento di Matematica e Applicazioni "R. Caccioppoli"  
via Cintia  
80126 Napoli, ITALY  
*E*Mail: [giannett@matna2.dma.unina.it](mailto:giannett@matna2.dma.unina.it)



---

volume 1, issue 2, article 14,  
2000.

*Received 17 January, 2000;*  
*accepted 4 April, 2000.*

*Communicated by: S. Saitoh*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

In this paper we prove higher integrability results for vector fields  $B, E, (B, E) \in L^{2-\varepsilon}(\Omega, \mathbb{R}^n) \times L^{2-\varepsilon}(\Omega, \mathbb{R}^n)$ ,  $\varepsilon$  small, such that  $\operatorname{div} B = 0$ ,  $\operatorname{curl} E = 0$  satisfying a “reverse” inequality of the type

$$|B|^2 + |E|^2 \leq \left( K + \frac{1}{K} \right) \langle B, E \rangle + |F|^2$$

with  $K \geq 1$  and  $F \in L^r(\Omega, \mathbb{R}^n)$ ,  $r > 2 - \varepsilon$ . Applications to the theory of quasiconformal mappings and partial differential equations are given. In particular, we prove regularity results for very weak solutions of equations of the type

$$\operatorname{div} a(x, \nabla u) = \operatorname{div} F.$$

If  $|a(x, z)|^2 + |z|^2 \leq (K + 1/K) \langle a(x, z), z \rangle$ , in the homogeneous case, our method provides a new proof of the regularity result

$$u \in W_{loc}^{1,2-\varepsilon}(\Omega) \Rightarrow u \in W_{loc}^{r,2+\varepsilon}(\Omega)$$

where  $\varepsilon$  is sufficiently small. A result of higher integrability for functions verifying a reverse integral inequality is used, and its optimality is proved.

**2000 Mathematics Subject Classification:** 35J60, 26D15

**Key words:** Reverse Inequalities, Finite Distortion Vector Fields, Div-Curl Vector Fields, Elliptic Partial Differential Equations

We wish to thank Prof. T. Iwaniec for stimulating discussions on this subject.

This work has been performed as a part of a National Research Project supported by M.U.R.S.T.



---

### Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back


Close

Quit

Page 2 of 27

# Contents

1	Introduction .....	4
2	Preliminary Results .....	6
3	Reverse Hölder Inequalities and Higher Integrability .....	8
4	A Counterexample .....	12
5	Higher Integrability Results and Applications .....	15
6	Regularity Results for Nonhomogeneous Equations .....	19
	References	



---

## Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 27

# 1. Introduction

The usual way to establish the  $W_{loc}^{1,2+\varepsilon}(\Omega)$ ,  $\varepsilon > 0$ , regularity of solutions  $u \in W_{loc}^{1,2}(\Omega)$  of equations of the type

$$\operatorname{div} a(x, \nabla u) = \operatorname{div} F \quad \text{in } \Omega,$$

where  $|a(x, z)|^2 + |z|^2 \leq (K + 1/K) \langle a(x, z), z \rangle$ , is to combine the Caccioppoli inequality

$$\int_Q |\nabla u|^2 dx \leq c \left[ \int_{2Q} \left| \frac{u - u_{2Q}}{2R} \right|^2 dx + \int_{2Q} |F|^2 dx \right],$$

where  $R$  is the sidelength of the cube  $Q \subset 2Q \subset \Omega$ , with the Poincaré-Sobolev inequality


$$\left( \int_Q \left| \frac{u - u_R}{R} \right|^2 dx \right)^{\frac{1}{2}} \leq \left( \int_Q |\nabla u|^{\frac{2n}{n+2}} dx \right)^{\frac{n+2}{2n}},$$

to obtain the nonhomogeneous reverse Hölder inequality

$$(1.1) \quad \int_Q |\nabla u|^2 dx \leq c \left\{ \left[ \int_{2Q} (|\nabla u|^2)^{\frac{n}{n+2}} dx \right]^{\frac{n+2}{n}} + \int_{2Q} |F|^2 dx \right\}.$$

The higher integrability result then arises by using the well-known Giaquinta-Modica technique [3, 2].

The aim of this paper is to provide a different way to get regularity results, based on inequalities for div-curl vector fields (see Theorem 2.1, [5, 10]). Starting from these inequalities, under the assumption of bounded distortion, we get



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 4 of 27


directly a family of reverse type inequalities, namely

$$(1.2) \quad \int_Q (|\nabla u|^2)^{1-\varepsilon} dx$$

$$\leq c_1 \varepsilon \int_{2Q} (|\nabla u|^2)^{1-\varepsilon} dx + c_2 \left[ \int_{2Q} (|\nabla u|^2)^{(1-\varepsilon)\frac{n}{n+1}} dx \right]^{\frac{n+1}{n}} + c_3 \int_{2Q} (|F|^2)^{(1-\varepsilon)} dx.$$

Notice that, even if inequality (1.2) contains an extra term, by using our method we are able to obtain a higher integrability result also for *very weak* solutions of some nonlinear elliptic equations by just assuming an integrability on the gradient below the natural exponent (see [8]). Let us observe also that if  $\varepsilon = 0$  the exponent  $n/(n + 1)$  in inequality (1.2) is larger than the exponent  $n/(n + 2)$  in inequality (1.1). Actually, inequality (1.2) follows from a more general argument about vector fields of bounded distortion, which includes an analogous result of the theory of quasiregular mappings (with the same exponent we get in (1.2), see [7]).

After recalling known results in Section 2, we prove a higher integrability result for functions verifying a reverse-type inequality (Theorem 3.1) in Section 3. In Section 4 we give a counterexample showing that generally the assumptions in Theorem 3.1 cannot be weakened. In Section 5 we prove a higher integrability result for finite distortion vector fields (see Proposition 5.1), and we give some applications to the theory of quasiconformal mappings and to the theory of regularity for very weak solutions of homogeneous nonlinear elliptic equations in divergence form. Finally, in Section 6, we extend our method to the case of more general vector fields in order to study the case of nonhomogeneous equations (see Theorem 6.1).




---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 27

## 2. Preliminary Results

In the following we will consider div-curl vector fields  $B = (B_1, \dots, B_n) \in L^q(\mathbb{R}^n, \mathbb{R}^n)$ ,  $E = (E_1, \dots, E_n) \in L^p(\mathbb{R}^n, \mathbb{R}^n)$ ,  $1 < p, q < \infty$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , i.e.

$$\operatorname{curl} E = \left( \frac{\partial E_i}{\partial x_j} - \frac{\partial E_j}{\partial x_i} \right)_{i,j=1,\dots,n} = 0 \quad (2.1)$$


$$\operatorname{div} B = \sum_{i=1}^n \frac{\partial B_i}{\partial x_i} = 0$$

in the sense of distributions.

The following basic estimates are established in [5] (see also [10] for the present formulation). We denote by  $Q_0, Q$  open cubes in  $\mathbb{R}^n$  with sides parallel to the coordinate axis, and by  $2Q$  the cube with the same center of  $Q$  and double side-length.

**Theorem 2.1.** *Let  $1 < p, q < \infty$  be a Hölder conjugate pair,  $\frac{1}{p} + \frac{1}{q} = 1$ , and let  $1 < r, s < \infty$  be a Sobolev conjugate pair,  $\frac{1}{r} + \frac{1}{s} = 1 + \frac{1}{n}$ . Then there exists a constant  $c_n = c_n(p, s)$  such that for each cube  $Q$  such that  $2Q \subset Q_0 \subset \mathbb{R}^n$  we have*

$$(2.2) \quad \left| \int_Q \frac{\langle B, E \rangle}{|B|^\varepsilon |E|^\varepsilon} dx \right| \leq c_n \varepsilon \left[ \int_{2Q} |E|^{(1-\varepsilon)p} dx \right]^{\frac{1}{p}} \left[ \int_{2Q} |B|^{(1-\varepsilon)q} dx \right]^{\frac{1}{q}} \\ + c_n \left[ \int_{2Q} |E|^{(1-\varepsilon)s} dx \right]^{\frac{1}{s}} \left[ \int_{2Q} |B|^{(1-\varepsilon)r} dx \right]^{\frac{1}{r}},$$



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and  
Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 6 of 27

whenever  $0 \leq 2\varepsilon \leq \min \left\{ \frac{p-1}{p}, \frac{q-1}{q}, \frac{r-1}{r}, \frac{s-1}{s} \right\}$  and  $\operatorname{div} B = 0$ ,  $\operatorname{curl} E = 0$ .

The following proposition by Giaquinta-Modica [3, 2] will be useful in the sequel.

**Proposition 2.2.** *Let  $g \in L^\alpha(Q_0)$ ,  $\alpha > 1$  and  $f \in L^r(Q_0)$ ,  $r > \alpha$  be two non-negative functions and suppose that for every cube  $Q$  such that  $2Q \subset Q_0$  the following estimate holds*

$$(2.3) \quad \int_Q g^\alpha dx \leq b \left\{ \left( \int_{2Q} g dx \right)^\alpha + \int_{2Q} f^\alpha dx \right\} + \theta \int_{2Q} g^\alpha dx$$

with  $b > 1$ . There exist constants  $\theta_0 = \theta_0(\alpha, n)$ ,  $\sigma_0 = \sigma_0(b, \theta, \alpha, r, n)$  such that if  $\theta < \theta_0$ , then  $g \in L_{loc}^{\alpha+\sigma}(Q_0)$  for all  $0 < \sigma < \sigma_0$  and

$$(2.4) \quad \left( \int_Q g^{\alpha+\sigma} dx \right)^{\frac{1}{\alpha+\sigma}} \leq c \left\{ \left( \int_{2Q} g^\alpha dx \right)^{\frac{1}{\alpha}} + \left( \int_{2Q} f^{\alpha+\sigma} dx \right)^{\frac{1}{\alpha+\sigma}} \right\},$$

where  $c$  is a positive constant depending on  $b, \theta, \alpha, r, n$ .



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 7 of 27

### 3. Reverse Hölder Inequalities and Higher Integrability


This section is concerned with a variant of the result established in Proposition 2.2. We remark that in our assumption (3.1) we will consider a family of inequalities of the type (2.3) in which both the exponent of integrability of the function  $g$  and a coefficient in the right hand side depend on  $\varepsilon$ . Nevertheless, even if Proposition 2.2 cannot be applied *a priori*, in the theorem we will prove that we can obtain a higher integrability result for  $g$  and an estimate of the type (2.4).

**Theorem 3.1.** *Let  $g \in L^{2(1-\varepsilon)}(Q_0)$  and  $f \in L^r(Q_0)$ ,  $0 \leq \varepsilon < \frac{1}{2}$ ,  $r > 2(1 - \varepsilon)$ , be nonnegative functions such that*

$$(3.1) \quad \int_Q g^{2(1-\varepsilon)} dx \leq c_1 \varepsilon \int_{2Q} g^{2(1-\varepsilon)} dx + c_2 \left\{ \left( \int_{2Q} g^{2(1-\varepsilon) \frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} + \left( \int_{2Q} f^{2(1-\varepsilon)} dx \right) \right\}$$

for every cube  $Q \subset 2Q \subset Q_0$ , for some constants  $c_1 \geq 0$ ,  $c_2 > 0$ .

Then there exist  $\bar{\varepsilon} = \bar{\varepsilon}(c_1, n)$  and  $\bar{\eta} = \bar{\eta}(c_1, c_2, r, \varepsilon, n)$  such that if  $0 \leq \varepsilon < \bar{\varepsilon}$ ,



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 8 of 27

then  $g \in L_{loc}^{2(1-\varepsilon)+\eta}(Q_0)$ ,  $\forall 0 \leq \eta < \bar{\eta}$  and

$$\left( \int_Q g^{2(1-\varepsilon)+\eta} dx \right)^{\frac{1}{2(1-\varepsilon)+\eta}} \leq c \left\{ \left( \int_{2Q} g^{2(1-\varepsilon)} dx \right)^{\frac{1}{2(1-\varepsilon)}} + \left( \int_{2Q} f^{2(1-\varepsilon)+\eta} dx \right)^{\frac{1}{2(1-\varepsilon)+\eta}} \right\},$$

where  $c$  is a positive constant depending on  $c_2, r, \varepsilon, n$ .

*Proof.* Since the functions  $g_\varepsilon = g^{2(1-\varepsilon)\frac{n}{n+1}}$ ,  $f_\varepsilon = f^{2(1-\varepsilon)\frac{n}{n+1}}$  verify the inequality


$$(3.2) \quad \int_Q g_\varepsilon^{\frac{n+1}{n}} dx \leq c_2 \left\{ \left( \int_{2Q} g_\varepsilon dx \right)^{\frac{n+1}{n}} + \left( \int_{2Q} f_\varepsilon^{\frac{n+1}{n}} dx \right) \right\} + c_1 \varepsilon \int_{2Q} g_\varepsilon^{\frac{n+1}{n}} dx$$

we can apply Proposition 2.2 with  $\alpha = \frac{n+1}{n}$  and  $b = c_2$ . We get  $\theta_0 = \theta_0(n)$  and  $\sigma_0 = \sigma_0(c_2, r, \varepsilon, n)$  such that, if (3.2) holds with  $c_1 \varepsilon < \frac{\theta_0}{2}$ , then  $g_\varepsilon \in L_{loc}^{\alpha+\sigma}(Q_0)$  for every  $0 < \sigma < \sigma_0$ , i.e.

$$\left[ g^{2(1-\varepsilon)\frac{n}{n+1}} \right]^{\frac{n+1}{n}+\sigma} \in L_{loc}^1(Q_0) \quad \forall 0 < \sigma < \sigma_0$$

and

$$(3.3) \quad \left( \int_Q g_\varepsilon^{\frac{n+1}{n}+\sigma} dx \right)^{\frac{1}{\frac{n+1}{n}+\sigma}} \leq c \left\{ \left( \int_{2Q} g_\varepsilon^{\frac{n+1}{n}} dx \right)^{\frac{n}{n+1}} + \left( \int_{2Q} f_\varepsilon^{\frac{n+1}{n}+\sigma} dx \right)^{\frac{n}{n+1}+\sigma} \right\}$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 9 of 27

with  $c$  depending on  $c_2, r, \varepsilon, n$ .

Set

$$0 < \bar{\varepsilon} < \frac{\theta_0}{2c_1}, \quad 0 < \bar{\eta} < (1 - \bar{\varepsilon}) \frac{2n\sigma_0}{n+1}.$$

If  $0 \leq \varepsilon < \bar{\varepsilon}$  and  $0 \leq \eta < \bar{\eta}$ , we have

$$\varepsilon < \bar{\varepsilon} < 1 - \bar{\eta} \frac{n+1}{2n\sigma_0} < 1 - \eta \frac{n+1}{2n\sigma_0}$$

or, equivalently,

$$2(1 - \varepsilon) + \eta < 2(1 - \varepsilon) \frac{n}{n+1} \left[ \frac{n+1}{n} + \sigma_0 \right].$$


Therefore we get

$$g \in L_{loc}^{2(1-\varepsilon)+\eta}(Q_0)$$

and inequality (3.3) becomes

$$\begin{aligned} & \left( \int_Q g^{2(1-\varepsilon)+\eta} dx \right)^{\frac{1}{2(1-\varepsilon)+\eta}} \\ & \leq c \left\{ \left( \int_{2Q} g^{2(1-\varepsilon)} dx \right)^{\frac{1}{2(1-\varepsilon)}} + \left( \int_{2Q} f^{2(1-\varepsilon)+\eta} dx \right)^{\frac{1}{2(1-\varepsilon)+\eta}} \right\}. \end{aligned}$$

□



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 10 of 27

Let us observe that upon closer inspection of the proof of Theorem 3.1, one can note that the gain of integrability given by  $\sigma_0 = \sigma_0(c_2, r, \varepsilon, n)$  is actually dependent only on  $c_2, \frac{r}{2(1-\varepsilon)}, n$ . Nevertheless, if  $f \equiv 0$  a.e. in  $Q_0$ , the number  $\sigma_0$ , and therefore also  $\bar{\eta}$  and  $c$ , do not depend on  $\varepsilon$ . This remark is crucial to prove the following.

**Corollary 3.2.** *Let  $0 \leq \varepsilon < \frac{1}{2}$  and  $g \in L^{2(1-\varepsilon)}(Q_0)$ ,  $Q_0 \subset \mathbb{R}^n$ , be such that*

$$\int_Q g^{2(1-\varepsilon)} dx \leq c_1 \int_{2Q} g^{2(1-\varepsilon)} dx + c_2 \left( \int_{2Q} g^{2(1-\varepsilon) \frac{n}{n+1}} dx \right)^{\frac{n+1}{n}}$$


for every cube  $Q \subset 2Q \subset Q_0$ .

Then there exists  $\bar{\varepsilon} = \bar{\varepsilon}(c_1, n)$  such that if  $0 \leq \varepsilon < \bar{\varepsilon}$ , then  $g \in L_{loc}^{2+2\varepsilon}(Q_0)$  and

$$(3.4) \quad \left( \int_Q g^{2(1+\varepsilon)} dx \right)^{\frac{1}{2(1+\varepsilon)}} \leq c \left( \int_{2Q} g^{2(1-\varepsilon)} dx \right)^{\frac{1}{2(1-\varepsilon)}}$$

where  $c$  is a positive constant depending on  $c_2, n$ .

*Proof.* Let us apply Theorem 3.1 with  $f \equiv 0$  a.e. in  $Q_0$ . If  $\varepsilon < \min(\bar{\varepsilon}, \frac{\bar{\eta}}{4})$ , choosing  $\eta = 4\varepsilon$ , from inequality (3.1) we get  $g \in L_{loc}^{2+2\varepsilon}(Q_0)$  and inequality (3.4) holds.  $\square$



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 11 of 27

## 4. A Counterexample

Let us consider  $f, g$  non-negative functions on a cube  $Q_0$  satisfying assumptions of the type of Theorem 3.1 with  $c_1 = 0$ , namely,  $f, g$  are such that  $g \in L^\alpha(Q_0)$ ,  $f \in L^{\lambda\alpha}(Q_0)$  for some  $\alpha > 1$ ,  $\lambda > 1$  and

$$(4.1) \quad \left( \int_Q g^\alpha dx \right)^{\frac{1}{\alpha}} \leq a \int_{2Q} g dx + b \left( \int_{2Q} f^\alpha dx \right)^{\frac{1}{\alpha}} \quad \forall Q, 2Q \subset Q_0.$$


In this case it is known, [6], that if  $\lambda$  is sufficiently close to 1,  $g \in L_{loc}^{\lambda\alpha}(Q_0)$  and

$$(4.2) \quad \left( \int_Q g^{\lambda\alpha} dx \right)^{\frac{1}{\lambda\alpha}} \leq a_\lambda \left( \int_{2Q} g^\lambda dx \right)^{\frac{1}{\lambda}} + b_\lambda \left( \int_{2Q} f^{\lambda\alpha} dx \right)^{\frac{1}{\lambda\alpha}},$$

where  $a_\lambda$  and  $b_\lambda$  are constants depending only on  $n, \alpha, a, b$ .

In the following we show that, even if it is still true that  $g \in L_{loc}^{\lambda\alpha}(Q_0)$  for any  $\lambda < 1$  (sufficiently small), one cannot find any  $\lambda < 1$ ,  $a_\lambda > 0$ ,  $b_\lambda > 0$  such that estimate (4.2) holds for any  $g \in L^\alpha(Q_0)$ ,  $f \in L^\alpha(Q_0)$  verifying (4.1). If we consider a function  $f \in \bigcap_{1 \leq p < \alpha} L^p(Q_0)$  such that  $\int_Q f^\alpha dx = +\infty \forall Q \subset Q_0$ , of course we have  $f \in L^{\lambda\alpha}(Q_0)$  for  $\lambda < 1$ , and it is possible to show immediately that there are no  $a_\lambda, b_\lambda > 0$  such that (4.2) holds for any  $g \in L^\alpha(Q_0)$ , for any  $f \in L^{\lambda\alpha}(Q_0)$  verifying (4.1).

We will proceed as follows: by a contradiction argument, we will be able to prove that there exists  $\lambda < 1$  such that *any* function  $g_0 \in L^{\lambda\alpha}(Q_0)$ ,  $g_0 > 0$ , satisfies a certain reverse inequality, which is generally false.



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 12 of 27

Let  $g \in C(\overline{Q_0})$ ,  $g > 0$ . Then there exists  $\delta > 0$  such that

$$\left( \int_{2Q} g^\alpha dx \right)^{\frac{1}{\alpha}} \leq 2 \int_{2Q} g dx \quad \forall Q, 2Q \subset Q_0, \text{diam } 2Q < \delta.$$

In fact,

$$\sup_{x \in 2Q} g(x) \leq \sup_{x, y \in 2Q} |g(x) - g(y)| + g(y) \quad \forall y \in 2Q$$

and then, because of the uniform continuity of  $g$ , we have

$$\sup_{2Q} g \leq 2 \inf_{2Q} g.$$

Let us divide  $Q_0$  into a finite number of disjoint cubes

$$Q_0 = \bigcup_{j=1}^N Q_j$$

such that


$$2Q \subset Q_0, \text{diam } 2Q \geq \delta \Rightarrow \exists Q_j : Q_j \subset 2Q.$$

Now let us point out that if  $f$  is any function in  $L^r(Q_0)$  for every  $1 \leq r < \alpha$ , but

$$\int_{Q_j} f^\alpha dx = +\infty \quad \forall j = 1, \dots, N$$

then there exist

$$f_k = \max_{j=1, \dots, N} \frac{k \min\{f, k\}}{\left( \int_{Q_j} (\min\{f, k\})^\alpha dx \right)^{\frac{1}{2\alpha}}}$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 13 of 27

such that

$$\left( \int_{2Q} \left( \frac{1}{k} f_k \right)^\alpha \right)^{\frac{1}{\alpha}} \geq \max_{Q_0} g \geq \left( \int_{2Q} g^\alpha dx \right)^{\frac{1}{\alpha}} \quad \forall Q, 2Q \subset Q_0, \text{diam } 2Q \geq \delta.$$

Therefore

$$\left( \int_{2Q} g^\alpha dx \right)^{\frac{1}{\alpha}} \leq 2 \int_{2Q} g dx + \left( \int_{2Q} \left( \frac{1}{k} f_k \right)^\alpha \right)^{\frac{1}{\alpha}} \quad \forall Q, 2Q \subset Q_0,$$

i.e.  $g, \frac{1}{k} f_k$  satisfy inequality (4.1).

Let us suppose to the contrary that for some  $\lambda < 1$

$$\left( \int_Q g^{\lambda\alpha} dx \right)^{\frac{1}{\lambda\alpha}} \leq a_\lambda \left( \int_{2Q} g^\lambda dx \right)^{\frac{1}{\lambda}} + b_\lambda \left( \int_{2Q} \left( \frac{1}{k} f_k \right)^{\lambda\alpha} \right)^{\frac{1}{\lambda\alpha}} \quad \forall Q, 2Q \subset Q_0.$$

Letting  $k$  tend to infinity, we have the inequality


$$\left( \int_Q g^{\lambda\alpha} dx \right)^{\frac{1}{\lambda\alpha}} \leq a_\lambda \left( \int_{2Q} g^\lambda dx \right)^{\frac{1}{\lambda}} \quad \forall Q, 2Q \subset Q_0$$

for every continuous function  $g$ .

This inequality, by an approximation argument, extends to every function  $g_0 \in L^{\lambda\alpha}(Q_0), g_0 > 0$

$$\left( \int_Q g_0^{\lambda\alpha} dx \right)^{\frac{1}{\lambda\alpha}} \leq a_\lambda \left( \int_{2Q} g_0^\lambda dx \right)^{\frac{1}{\lambda}} \quad \forall Q, 2Q \subset Q_0,$$

which is absurd, since this inequality implies a higher integrability for  $g_0$ .



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and  
Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 14 of 27

## 5. Higher Integrability Results and Applications

We start with the following regularity result for vector fields of bounded distortion

**Proposition 5.1.** *Let  $\Omega \subset \mathbb{R}^n$ ,  $0 < \epsilon < 1$  and  $\Phi = (E, B) \in L^{2-\epsilon}(\Omega, \mathbb{R}^n) \times L^{2-\epsilon}(\Omega, \mathbb{R}^n)$  be such that  $\operatorname{div} B = 0$ ,  $\operatorname{curl} E = 0$  and*

$$(5.1) \quad |B(x)|^2 + |E(x)|^2 \leq \left( K + \frac{1}{K} \right) \langle B(x), E(x) \rangle \quad \text{a.e. in } \Omega,$$


where  $K \geq 1$ . Then there exists  $\bar{\epsilon} = \bar{\epsilon}(K, n)$  such that  $\Phi \in L_{loc}^{2+\epsilon}(\Omega, \mathbb{R}^n) \times L_{loc}^{2+\epsilon}(\Omega, \mathbb{R}^n)$  for all  $0 < \epsilon < \bar{\epsilon}$  and

$$\left( \int_Q |\Phi|^{2+\epsilon} dx \right)^{\frac{1}{2+\epsilon}} \leq c \left( \int_{2Q} |\Phi|^{2-\epsilon} dx \right)^{\frac{1}{2-\epsilon}} \quad \forall Q, 2Q \subset \Omega,$$

where  $c$  is a positive constant depending on  $K, n$ .

*Proof.* Let us fix the cube  $Q$  such that  $2Q \subset \Omega$ . Applying Theorem 2.1 with  $p = q = 2$  and  $r = s = \frac{2n}{n+1}$ , from inequality (5.1) we get

$$\begin{aligned} & \int_Q (|B|^2 + |E|^2)^{1-\epsilon} dx \\ & \leq c_{n,K} \epsilon \int_{2Q} (|B|^2 + |E|^2)^{1-\epsilon} dx + c_{n,K} \left( \int_{2Q} (|B|^2 + |E|^2)^{(1-\epsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} \end{aligned}$$



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 15 of 27

for  $\varepsilon$  sufficiently small. Substituting  $g^2$  for  $|B|^2 + |E|^2$  in the last inequality gives

$$\int_Q g^{2-2\varepsilon} dx \leq c_{n,K} \varepsilon \int_{2Q} g^{2-2\varepsilon} dx + c_{n,K} \left( \int_{2Q} g^{(2-2\varepsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}}.$$

By Corollary 3.2 there exists  $\bar{\varepsilon} = \bar{\varepsilon}(K, n)$  such that if  $0 \leq \varepsilon < \bar{\varepsilon}$ , then  $g \in L_{loc}^{2+2\varepsilon}(\Omega)$  and

$$\left( \int_Q g^{2+2\varepsilon} dx \right)^{\frac{1}{2+2\varepsilon}} \leq c \left( \int_{2Q} g^{2-2\varepsilon} dx \right)^{\frac{1}{2-2\varepsilon}}$$

and then the assertion. □

Now we consider the equation on a bounded open set  $\Omega \subset \mathbb{R}^n$ ,


$$Au = 0 \quad \text{in } \Omega \subset \mathbb{R}^n,$$

where  $A$  is a differential operator defined by

$$Au = \operatorname{div} a(x, \nabla u).$$

Here  $a : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a mapping such that  $x \rightarrow a(x, z)$  is measurable for all  $z \in \mathbb{R}^n$  and  $z \rightarrow a(x, z)$  is continuous for almost every  $x \in \Omega$ . Furthermore, we assume that there exists  $K \geq 1$  such that for almost every  $x \in \Omega$  we have

$$(5.2) \quad |a(x, z)|^2 + |z|^2 \leq \left( K + \frac{1}{K} \right) \langle a(x, z), z \rangle,$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 16 of 27

where  $x, z$  are arbitrary vectors in  $\mathbb{R}^n$ .

Let us prove the following result (originally proved in [9], in the linear case, by using a duality technique).

**Corollary 5.2.** *Let  $0 < \epsilon < \frac{1}{2}$  and  $u \in W_{loc}^{1,2-2\epsilon}(\Omega)$  be a very weak solution of*

$$\operatorname{div} a(x, \nabla u) = 0.$$

*Then there exists  $\bar{\epsilon} = \bar{\epsilon}(K, n)$  such that  $u \in W_{loc}^{1,2+2\epsilon}(\Omega)$  for all  $0 < \epsilon < \bar{\epsilon}$  and*

$$\left( \int_Q |\nabla u|^{2+2\epsilon} dx \right)^{\frac{1}{2+2\epsilon}} \leq c \left( \int_{2Q} |\nabla u|^{2-2\epsilon} dx \right)^{\frac{1}{2-2\epsilon}},$$

where  $c$  is a positive constant depending on  $K, n$ .


*Proof.* Setting

$$E = \nabla u, \quad B = a(x, \nabla u)$$

we have  $\operatorname{div} B = 0$ ,  $\operatorname{curl} E = 0$  and, by (5.2),

$$|E|^2 + |B|^2 = |\nabla u|^2 + |a(x, \nabla u)|^2 \leq \left( K + \frac{1}{K} \right) \langle a(x, \nabla u), \nabla u \rangle$$

so that  $E, B$  are div-curl fields of bounded distortion. Applying Proposition 5.1 we get the assertion.  $\square$




---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit

Page 17 of 27

Another interesting case to which Proposition 5.1 applies is when one considers a homeomorphism


$$f = (f^1, f^2) : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$f^i \in W^{1,2-\varepsilon}(\Omega) \quad i = 1, 2,$$

$$f \text{ } K\text{-quasiregular, } K \geq 1, \quad \text{i.e.} \quad |Df(x)|^2 \leq \left( K + \frac{1}{K} \right) J(x, f),$$

where  $|Df(x)|$  denotes the norm of the distributional differential  $Df(x)$  and  $J(x, f)$  is the Jacobian determinant  $J(x, f) = \det Df(x)$ .

Then, writing the Jacobian  $J(x, f)$  as  $\langle B, E \rangle$ , where  $E = \nabla f^1 = (f_x^1, f_y^1)$  and  $B = (f_y^2, -f_x^2)$  we have  $\operatorname{div} B = 0$ ,  $\operatorname{curl} E = 0$  and that (5.1) holds. It follows that for  $\varepsilon$  sufficiently small  $f \in W^{1,2+\varepsilon}(\Omega, \mathbb{R}^2)$ , giving back in this way the celebrated theorem by Bojarski [1]. Significant results about the Jacobian determinant are in [7].




---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 27

## 6. Regularity Results for Nonhomogeneous Equations

In this section we consider  $\Phi = (E, B) \in L^{2-2\varepsilon}(\Omega, \mathbb{R}^n) \times L^{2-2\varepsilon}(\Omega, \mathbb{R}^n)$  such that

$$(6.1) \quad \operatorname{div} B = 0, \quad \operatorname{curl} E = 0,$$

$$(6.2) \quad |B(x)|^2 + |E(x)|^2 \leq \left( K + \frac{1}{K} \right) \langle B(x), E(x) \rangle + |F|^2,$$

where  $F$  is a function in  $L^r(\Omega, \mathbb{R}^n)$ ,  $r > 2 - 2\varepsilon$ , for  $\varepsilon$  sufficiently small.


**Theorem 6.1.** *Let  $0 \leq \varepsilon < \frac{1}{2}$  and  $E, B$  vector fields as in (6.1), (6.2). Then there exist  $\bar{\varepsilon} = \bar{\varepsilon}(K, n)$  and  $\bar{\eta} = \bar{\eta}(K, r, \varepsilon, n)$  such that if  $0 \leq \varepsilon < \bar{\varepsilon}$ , then  $\Phi = (E, B) \in L_{loc}^{2-2\varepsilon+\eta}(\Omega, \mathbb{R}^n) \times L_{loc}^{2-2\varepsilon+\eta}(\Omega, \mathbb{R}^n)$  for all  $0 \leq \eta < \bar{\eta}$  and*

$$(6.3) \quad \left( \int_Q |\Phi|^{2-2\varepsilon+\eta} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \leq c \left\{ \left( \int_{2Q} |\Phi|^{2-2\varepsilon} dx \right)^{\frac{1}{2-2\varepsilon}} + \left( \int_{2Q} (|F|^2)^{\frac{2-2\varepsilon+\eta}{2}} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \right\},$$

where  $c$  is a positive constant depending on  $K, r, \varepsilon, n$ .

*Proof.* Let us fix  $Q$  a cube such that  $2Q \subset \Omega$  and set

$$Q^+ = \{x \in Q \mid \langle B, E \rangle \geq 0 \text{ a.e.}\}$$



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 19 of 27

$$Q^- = \{x \in Q \mid \langle B, E \rangle \leq 0 \text{ a.e.}\}$$

Let us observe that by (6.2), replacing  $|F|$  with  $f$ , we have


$$\begin{aligned} \int_{Q^-} \frac{-\langle B, E \rangle}{|B|^\varepsilon |E|^\varepsilon} dx &\leq \int_{Q^-} (|B||E|)^{1-\varepsilon} dx \\ &\leq \int_{Q^-} (|B|^2 + |E|^2)^{1-\varepsilon} dx \\ &\leq \int_{Q^-} \left[ \left( K + \frac{1}{K} \right) \langle B, E \rangle + f^2 \right]^{1-\varepsilon} dx \\ &\leq \int_{Q^-} f^{2-2\varepsilon} dx \leq \int_Q f^{2-2\varepsilon} dx \end{aligned}$$

and therefore

$$\begin{aligned} \int_Q \frac{\langle B, E \rangle}{|B|^\varepsilon |E|^\varepsilon} dx &= \int_{Q^+} \frac{\langle B, E \rangle}{|B|^\varepsilon |E|^\varepsilon} dx + \int_{Q^-} \frac{\langle B, E \rangle}{|B|^\varepsilon |E|^\varepsilon} dx \\ &\geq \int_{Q^+} \frac{\langle B, E \rangle}{(|B|^2 + |E|^2 + f^2)^\varepsilon} dx - \int_Q f^{2-2\varepsilon} dx \end{aligned}$$

Applying Theorem 2.1 with  $p = q = 2$  and  $r = s = \frac{2n}{n+1}$ , for  $\varepsilon$  sufficiently small, we get

$$\begin{aligned} \int_Q \frac{\langle B, E \rangle}{(|B|^2 + |E|^2 + f^2)^\varepsilon} dx &\leq c_n \varepsilon \int_{2Q} (|B|^2 + |E|^2 + f^2)^{1-\varepsilon} dx \\ &\quad + c_n \left( \int_{2Q} (|B|^2 + |E|^2 + f^2)^{(1-\varepsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} + \int_{2Q} f^{2-2\varepsilon} dx. \end{aligned}$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit


Page 20 of 27

By (6.2)

$$\langle B, E \rangle \geq c_K(|B|^2 + |E|^2 - f^2) = c_K(|B|^2 + |E|^2 + f^2) - 2c_K f^2$$

and therefore

$$\begin{aligned} & \int_Q (|B|^2 + |E|^2 + f^2)^{1-\varepsilon} dx \\ & \leq c_{n,K} \varepsilon \int_{2Q} (|B|^2 + |E|^2 + f^2)^{1-\varepsilon} dx \\ & + c_{n,K} \left( \int_{2Q} (|B|^2 + |E|^2 + f^2)^{(1-\varepsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} \\ & + c_K \int_{2Q} \frac{f^2}{(|B|^2 + |E|^2 + f^2)^\varepsilon} dx + \int_{2Q} f^{2-2\varepsilon} dx \\ & \leq c_{n,K} \varepsilon \int_{2Q} (|B|^2 + |E|^2 + f^2)^{1-\varepsilon} dx \\ & + c_{n,K} \left( \int_{2Q} (|B|^2 + |E|^2 + f^2)^{(1-\varepsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} \\ & + (c_K + 1) \int_{2Q} f^{2-2\varepsilon} dx. \end{aligned}$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 21 of 27

Setting  $g^2 = |B|^2 + |E|^2 + f^2$ , the last inequality implies

$$\int_Q g^{2-2\varepsilon} dx \leq c_{n,K} \varepsilon \int_{2Q} g^{2-2\varepsilon} dx + c_{n,K} \left( \int_{2Q} g^{(2-2\varepsilon)\frac{n}{n+1}} dx \right)^{\frac{n+1}{n}} + (c_K + 1) \int_{2Q} f^{2-2\varepsilon} dx.$$

By Theorem 3.1 there exist  $\bar{\varepsilon} = \bar{\varepsilon}(K, n)$  and  $\bar{\eta} = \bar{\eta}(K, r, \varepsilon, n)$  such that if  $0 \leq \varepsilon < \bar{\varepsilon}$ , then  $g \in L_{loc}^{2-2\varepsilon+\eta}(\Omega)$  for all  $0 \leq \eta < \bar{\eta}$  and

$$\left( \int_Q g^{2-2\varepsilon+\eta} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \leq c \left\{ \left( \int_{2Q} g^{2-2\varepsilon} dx \right)^{\frac{1}{2-2\varepsilon}} + \left( \int_{2Q} (f^2)^{\frac{2-2\varepsilon+\eta}{2}} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \right\}$$

and then the assertion.  $\square$

Let us consider now the following equation on a bounded open set  $\Omega \subset \mathbb{R}^n$

$$(6.4) \quad \operatorname{div} a(x, \nabla u) = \operatorname{div} F,$$

where  $F \in L^r(\Omega)$ ,  $r > 2 - 2\varepsilon$ , for  $\varepsilon$  sufficiently small and  $a : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a mapping satisfying the assumptions of Section 5.

**Corollary 6.2.** *Let  $0 \leq \varepsilon < \frac{1}{2}$  and  $u \in W_{loc}^{1,2-2\varepsilon}(\Omega)$  be a very weak solution of the equation (6.4). Then there exist  $\bar{\varepsilon} = \bar{\varepsilon}(K, n)$  and  $\bar{\eta} = \bar{\eta}(K, r, \varepsilon, n)$  such that*



Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 22 of 27

if  $0 \leq \varepsilon < \bar{\varepsilon}$ , then  $u \in W_{loc}^{1,2-2\varepsilon+\eta}(\Omega)$  for all  $0 \leq \eta < \bar{\eta}$  and

$$\begin{aligned} & \left( \int_Q |\nabla u|^{2-2\varepsilon+\eta} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \\ & \leq c \left\{ \left( \int_{2Q} |\nabla u|^{2-2\varepsilon} dx \right)^{\frac{1}{2-2\varepsilon}} + \left( \int_{2Q} |F|^{2-2\varepsilon+\eta} dx \right)^{\frac{1}{2-2\varepsilon+\eta}} \right\} \end{aligned}$$


for all cubes  $Q$  such that  $2Q \subset \Omega$  and where  $c$  is a positive constant depending on  $K, r, \varepsilon, n$ .

*Proof.* Setting

$$E = \nabla u, \quad B = a(x, \nabla u) - F,$$

we have

$$\begin{aligned} |E|^2 + |B|^2 & \leq |\nabla u|^2 + (|a(x, \nabla u)| + |F|)^2 \\ & \leq 2(|a(x, \nabla u)|^2 + |\nabla u|^2) + 2|F|^2 \\ & \leq 2 \left( K + \frac{1}{K} \right) \langle a(x, \nabla u), \nabla u \rangle + 2|F|^2 \\ & = 2 \left( K + \frac{1}{K} \right) \langle a(x, \nabla u) - F, \nabla u \rangle + 2 \left( K + \frac{1}{K} \right) \langle F, \nabla u \rangle + 2|F|^2. \end{aligned}$$



**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

Title Page

Contents



Go Back

Close

Quit

Page 23 of 27

Since by Young's inequality

$$\begin{aligned}
 \langle F, \nabla u \rangle &= 2\langle F, \nabla u \rangle - \langle F, \nabla u \rangle \\
 &\leq 2 \cdot 2 \left( K + \frac{1}{K} \right) |F|^2 + 2 \frac{1}{2 \left( K + \frac{1}{K} \right)} |\nabla u|^2 - \langle F, \nabla u \rangle \\
 &\leq 4 \left( K + \frac{1}{K} \right) |F|^2 + \frac{1}{K + \frac{1}{K}} (|a(x, \nabla u)|^2 + |\nabla u|^2) - \langle F, \nabla u \rangle \\
 &\leq 4 \left( K + \frac{1}{K} \right) |F|^2 + \langle a(x, \nabla u), \nabla u \rangle - \langle F, \nabla u \rangle \\
 &= 4 \left( K + \frac{1}{K} \right) |F|^2 + \langle a(x, \nabla u) - F, \nabla u \rangle
 \end{aligned}$$


we get

$$|E|^2 + |B|^2 \leq 4 \left( K + \frac{1}{K} \right) \langle B, E \rangle + \left[ 8 \left( K + \frac{1}{K} \right)^2 + 2 \right] |F|^2,$$

i.e.  $E, B$  are vector fields satisfying (6.1) and (6.2). From Theorem 6.1 we get the higher integrability for  $|E|^2 + |B|^2$  and then for  $|\nabla u|$ ; the estimate follows directly from (6.3).  $\square$

*Remark 6.1.* Let  $u \in W_{loc}^{1,2}(\Omega)$  be a solution of the equation (6.4). Corollary 6.2 asserts that the function  $g = |\nabla u|$  verifies inequality (4.1) and, surprisingly, satisfies also inequality (4.2) with  $\lambda < 1$  sufficiently small.

*Remark 6.2.* We note that, arguing as in the end of Section 5, our result of higher integrability applies also to  $(K, K')$ -quasiregular mappings (see [4]), i.e.




---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit


Page 24 of 27

functions  $f$  verifying

$$f = (f^1, f^2) : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$f^i \in W^{1,2-\varepsilon}(\Omega) \quad i = 1, 2,$$

$$|Df(x)|^2 \leq \left( K + \frac{1}{K} \right) J(x, f) + K'.$$



---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back


Close

Quit

Page 25 of 27

## References

- [1] B. BOJARSKI, Homeomorphic solutions of Beltrami system, *Dokl. Akad. Nauk. SSSR*, **102** (1955), 661–664.
- [2] E. GIUSTI, Metodi Diretti nel Calcolo delle Variazioni, *Unione Matematica Italiana* (1994).
- [3] M. GIAQUINTA AND G. MODICA, Regularity results for some classes of higher order non linear elliptic systems, *J. Reine Angew. Math.*, **311/312** (1979), 145–169.
- [4] D. GILBARG AND N.S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, Springer (1983).
- [5] T. IWANIEC, Integrability Theory of the Jacobians, *Lipschitz Lectures* (1995).
- [6] T. IWANIEC, The Gehring Lemma, Quasiconformal Mappings and Analysis, *A Collection of Papers Honoring F.W. Gehring P.L. Duren, J.M. Heinonen, B.G. Osgood, B.P. Palka* (1998) Springer-Verlag, 181–204.
- [7] T. IWANIEC AND C. SBORDONE, On the integrability of the Jacobian under minimal hypothesis, *Arch. Rat. Mech. Anal.*, **119** (1992), 129–143.
- [8] T. IWANIEC AND C. SBORDONE, Weak minima of variational integrals, *J. Reine Angew. Math.*, **454** (1994), 143–161.
- [9] N. MEYERS AND A. ELCRAT, Some results on regularity for solutions of nonlinear elliptic systems and quasiregular functions, *Duke Math. J.*, **42**(1) (1975), 121–136.



---

Regularity results for vector fields of bounded distortion and applications

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents




Go Back

Close

Quit

Page 26 of 27

- [10] C. SBORDONE, New estimates for div-curl products and very weak solutions of PDEs, *Annali Scuola Normale Superiore di Pisa (IV)*, **25**(3-4) (1997), 739–756.



---

**Regularity results for vector fields of bounded distortion and applications**

Alberto Fiorenza and  
Flavia Giannetti

---

Title Page

Contents



Go Back

Close

Quit

Page 27 of 27