

Journal of Inequalities in Pure and Applied Mathematics

A PICK FUNCTION RELATED TO AN INEQUALITY FOR THE ENTROPY FUNCTION

CHRISTIAN BERG

Department of Mathematics,
University of Copenhagen,
Universitetsparken 5
DK-2100 Copenhagen, DENMARK.
*E*Mail: berg@math.ku.dk
*U*RL: <http://www.math.ku.dk/~berg/>

©2000 Victoria University
ISSN (electronic): 1443-5756
024-01



volume 2, issue 2, article 26,
2001.

Received 6 November, 2000;
accepted 6 March, 2001.

Communicated by: F. Hansen

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


The function $\psi(z) = 2/(1+z) + 1/(\text{Log}(1-z)/2)$, holomorphic in the cut plane $\mathbb{C} \setminus [1, \infty[$, is shown to be a Pick function. This leads to an integral representation of the coefficients in the power series expansion $\psi(z) = \sum_{n=0}^{\infty} \beta_n z^n$, $|z| < 1$. The representation shows that (β_n) decreases to zero as conjectured by F. Topsøe. Furthermore, (β_n) is completely monotone.

2000 Mathematics Subject Classification: 30E20, 44A60.

Key words: Pick functions, completely monotone sequences.

Contents

1	Introduction and Statement of Results	3
2	Proofs	5
	References	



A Pick Function Related to an Inequality for the Entropy Function

Christian Berg

Title Page

Contents



Go Back

Close

Quit

Page 2 of 8

1. Introduction and Statement of Results

In the paper [2] about bounds for entropy Topsøe considered the function

$$(1.1) \quad \psi(x) = \frac{2}{1+x} + \frac{1}{\ln \frac{1-x}{2}}, \quad -1 < x < 1$$

with the power series expansion

$$(1.2) \quad \psi(x) = \sum_{n=0}^{\infty} \beta_n x^n$$

and conjectured from numerical evidence that (β_n) decreases to zero.

The purpose of this note is to prove the conjecture by establishing the integral representation

$$(1.3) \quad \beta_n = \int_1^{\infty} \frac{dt}{t^{n+1}(\pi^2 + \ln^2 \frac{t-1}{2})}, \quad n \geq 0.$$


This formula clearly shows $\beta_0 > \beta_1 > \dots > \beta_n \rightarrow 0$. Furthermore, by a change of variable we find

$$\beta_n = \int_0^1 s^n \frac{ds}{s(\pi^2 + \ln^2 \frac{1-s}{2s})}, \quad n \geq 0,$$

which shows that (β_n) is a completely monotone sequence, cf. [3].

The representation (1.3) follows from the observation that ψ is the restriction of a Pick function with the following integral representation

$$(1.4) \quad \psi(z) = \int_1^{\infty} \frac{dt}{(t-z)(\pi^2 + \ln^2 \frac{t-1}{2})}, \quad z \in \mathbb{C} \setminus [1, \infty[.$$



A Pick Function Related to an
Inequality for the Entropy
Function

Christian Berg

Title Page

Contents




Go Back

Close

Quit

Page 3 of 8

From (1.4) we immediately get (1.3) since $\beta_n = \psi^{(n)}(0)/n!$.



**A Pick Function Related to an
Inequality for the Entropy
Function**

Christian Berg

Title Page

Contents



Go Back

Close

Quit

Page 4 of 8

2. Proofs

A holomorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ in the upper half-plane is called a Pick function, cf. [1], if $\text{Im } f(z) \geq 0$ for all $z \in \mathbb{H}$. Pick functions are also called Nevanlinna functions or Herglotz functions. They have the integral representation

$$(2.1) \quad f(z) = az + b + \int_{-\infty}^{\infty} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu(t),$$

where $a \geq 0$, $b \in \mathbb{R}$ and μ is a non-negative Borel measure on \mathbb{R} satisfying

$$\int \frac{d\mu(t)}{1+t^2} < \infty.$$


It is known that

$$(2.2) \quad a = \lim_{y \rightarrow \infty} f(iy)/iy, \quad b = \text{Re } f(i), \quad \mu = \lim_{y \rightarrow 0^+} \frac{1}{\pi} \text{Im } f(t + iy) dt,$$

where the limit refers to the vague topology. Finally f has a holomorphic extension to $\mathbb{C} \setminus [1, \infty[$ if and only if $\text{supp } (\mu) \subseteq [1, \infty[$.

Let $\text{Log } z = \ln |z| + i \text{Arg } z$ denote the principal logarithm in the cut plane $\mathbb{C} \setminus]-\infty, 0]$, with $\text{Arg } z \in]-\pi, \pi[$. Hence $\text{Log } \frac{1-z}{2}$ is holomorphic in $\mathbb{C} \setminus [1, \infty[$ with $z = -1$ as a simple zero. It is easily seen that

$$(2.3) \quad \psi(z) = \frac{2}{1+z} + \frac{1}{\text{Log } \frac{1-z}{2}}, \quad z \in \mathbb{C} \setminus [1, \infty[$$



**A Pick Function Related to an
Inequality for the Entropy
Function**

Christian Berg

Title Page

Contents



Go Back

Close

Quit

Page 5 of 8

is a holomorphic extension of (1.1) with a removable singularity for $z = -1$ where $\psi(-1) = 1/2$. To see that $V(z) = \text{Im } \psi(z) \geq 0$ for $z \in \mathbb{H}$ it suffices by the boundary minimum principle for harmonic functions to verify $\liminf_{z \rightarrow x} V(z) \geq 0$ for $x \in \mathbb{R}$ and $\liminf_{|z| \rightarrow \infty} V(z) \geq 0$, where in both cases $z \in \mathbb{H}$.

We find

$$\lim_{z \rightarrow x} \psi(z) = \begin{cases} \psi(x), & x \leq 1 \text{ (with } \psi(1) = 1) \\ \frac{2}{1+x} + \frac{1}{\ln \frac{x-1}{2} - i\pi}, & x > 1 \end{cases}$$

hence


$$\lim_{z \rightarrow x} V(z) = \begin{cases} 0, & x \leq 1 \\ \frac{\pi}{\pi^2 + \ln^2 \frac{x-1}{2}}, & x > 1, \end{cases}$$

whereas $\lim_{|z| \rightarrow \infty} \psi(z) = 0$. This shows that ψ is a Pick function, and from (2.2) we see that $a = 0$ and μ has the following continuous density with respect to Lebesgue measure

$$d(x) = \begin{cases} 0, & x \leq 1 \\ 1 / (\pi^2 + \ln^2 \frac{x-1}{2}), & x > 1. \end{cases}$$

Therefore

$$\psi(z) = b + \int_1^\infty \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) \frac{dt}{\pi^2 + \ln^2 \frac{t-1}{2}}.$$



A Pick Function Related to an Inequality for the Entropy Function

Christian Berg

Title Page

Contents



Go Back

Close

Quit

Page 6 of 8


In this case we can integrate term by term, and since $\lim_{x \rightarrow -\infty} \psi(x) = 0$, we find

$$\psi(z) = \int_1^{\infty} \frac{dt}{(t-z)(\pi^2 + \ln^2 \frac{t-1}{2})}$$

and

$$b = \operatorname{Re} \psi(i) = 1 - \frac{8 \ln 2}{\pi^2 + 4 \ln^2 2} = \int_1^{\infty} \frac{t dt}{(1+t^2)(\pi^2 + \ln^2 \frac{t-1}{2})},$$

which establishes (1.4).



A Pick Function Related to an Inequality for the Entropy Function

Christian Berg

Title Page

Contents



Go Back


Close

Quit

Page 7 of 8

References

- [1] W.F. DONOGHUE, *Monotone Matrix Functions and Analytic Continuation*, Berlin, Heidelberg, New York, 1974.
- [2] F. TOPSØE, Bounds for entropy and divergence for distributions over a two-element set, *J. Ineq. Pure And Appl. Math.*, **2**(2) (2001), Article 25.
http://jipam.vu.edu.au/v2n2/044_00.html
- [3] D.V. WIDDER, *The Laplace Transform*, Princeton 1941.



A Pick Function Related to an Inequality for the Entropy Function

Christian Berg

Title Page

Contents



Go Back

Close

Quit

Page 8 of 8