

Journal of Inequalities in Pure and Applied Mathematics

ON THE UTILITY OF THE TELYAKOVSKII'S CLASS S

L. LEINDLER

Bolyai Institute
University of Szeged
Aradi vértanúk tere 1
6720 Szeged
HUNGARY
EMail: leindler@math.u-szeged.hu

©2000 Victoria University
ISSN (electronic): 1443-5756
008-01



volume 2, issue 3, article 32,
2001.

*Received 29 January, 2001;
accepted 26 April, 2001.*

Communicated by: S.S. Dragomir

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


An illustration is given showing the advantage of the definition given by Telyakovskii for the class introduced by Sidon. It is also verified that if a sequence $\{a_n\}$ belongs to the recently defined subclass S_γ of S , $\gamma > 0$, then the sequence $\{n^\gamma a_n\}$ belongs to the class S , but the converse statement does not hold.

2000 Mathematics Subject Classification: 26D15, 42A20

Key words: Cosine and Sine Series, Fourier Series, Fourier Coefficients, Inequalities, Integrability.

Contents

1	Introduction	3
2	Results	7
3	Proofs	9
	References	



**On the Utility of the
Telyakovskii's Class S**

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

1. Introduction

A great number of mathematicians have studied the question ‘What conditions for a sequence $\{a_n\}$ guarantee that the trigonometric series


$$(1.1) \quad \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$$

and

$$(1.2) \quad \sum_{n=1}^{\infty} a_n \sin nx$$

to be Fourier series, or to converge in L^1 -metric?. We refer only to W.H. Young [13], A.H. Kolmogorov [2], S. Sidon [6], S. A. Telyakovskii [9] and the plentiful references given in [9] and in the excellent monograph by R.P. Boas, Jr. [1]. It is also known that conditions were established with monotone, quasi-monotone, convex and quasi-convex sequences, with null-sequences of bounded variation, and also sequences given by Sidon via a nice special construction.

In 1973 S. A. Telyakovskii [10] introduced a very effective idea, defined a “new” class of coefficient sequences. He denoted this class by S ; the letter S refers to an esteemed result of S. Sidon [6], and to the class defined by him in the same paper. Namely, Telyakovskii also showed that his class and that of Sidon are identical, but to apply his definition is more convenient. This is the reason, in my view, that later most of the authors ([7], [8], [14]), dealing with similar problems, wanted to extend the definition of Telyakovskii.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

In [3] and [4] we showed that some of these “extensions” are equivalent to the class S , and some others are real extensions of S , but they are identical among themselves.

All of these facts show that the class S defined by Telyakovskii plays a very important role in the studies of the problems mentioned above.

The definition of the class S is the following: A null-sequence $\mathbf{a} := \{a_n\}$ belongs to the class S , or briefly $\mathbf{a} \in S$, if there exists a monotonically decreasing sequence $\{A_n\}$ such that $\sum_{n=1}^{\infty} A_n < \infty$ and $|\Delta a_n| \leq A_n$ hold for all n .

The aim of the present note is to give one further illustration which underlies the central position of the class S and the following theorems proved in the same paper where the definition of S was given.

In [10] Telyakovskii, among others, proved the next two theorems.

Theorem 1.1. *Let the coefficients of the series (1.1) belong to the class S . Then the series (1.1) is a Fourier series and*


$$\int_0^{\pi} \left| \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx \right| dx \leq C \sum_{n=0}^{\infty} A_n,$$

where C is an absolute constant.

Theorem 1.2. *Let the coefficients of the series (1.2) belong to the class S . Then for any $p = 1, 2, \dots$*

$$\int_{\frac{\pi}{p+1}}^{\pi} \left| \sum_{n=1}^{\infty} a_n \sin nx \right| dx = \sum_{n=1}^p \frac{|a_n|}{n} + O \left(\sum_{n=1}^{\infty} A_n \right)$$

holds uniformly.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

In particular, the series (1.2) is a Fourier series if and only if

$$\sum_{n=1}^{\infty} \frac{|a_n|}{n} < \infty.$$

Recently Z. Tomovski [12] defined certain subclasses of S , and denoted them by S_r , $r = 1, 2, \dots$ (see also [11] and in [5] the definition of the class $S(\alpha)$). A null-sequence $\{a_n\}$ belongs to the class S_r , if there exists a monotonically decreasing sequence $\{A_n^{(r)}\}$ such that $\sum_{n=1}^{\infty} n^r A_n^{(r)} < \infty$ and $|\Delta a_n| \leq A_n^{(r)}$ for all n . (For $r = 0$ clearly $S_0 = S$ and $A_n^{(0)} = A_n$.)


In [11] Tomovski established, among others, two theorems in connection with the classes S_r as follows:

Theorem 1.3. *Let the coefficients of the series (1.1) belong to the class S_r , $r = 0, 1, \dots$. Then the r -th derivative of the series (1.1) is a Fourier series and if $f^{(r)}(x)$ denotes its sum function we have that*

$$\int_0^{\pi} |f^{(r)}(x)| dx \leq M \sum_{n=0}^{\infty} n^r A_n^{(r)}, \quad M = M(r) > 0.$$

Theorem 1.4. *Let the coefficients of the series (1.2) belong to the class S_r , $r = 0, 1, \dots$, furthermore let $g(x)$ denote the sum function of the series (1.2). Then for any $p = 1, 2, \dots$*

$$\int_{\frac{\pi}{p+1}}^{\pi} |g^{(r)}(x)| dx = \sum_{n=1}^p |a_n| n^{r-1} + O\left(\sum_{n=1}^{\infty} n^r A_n^{(r)}\right).$$



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

In particular, the r -th derivative of the series (1.2) is a Fourier series if and only if

$$\sum_{n=1}^{\infty} |a_n| n^{r-1} < \infty.$$


It is obvious that if $r = 0$ then the Theorems 1.3 and 1.4 reduce to the Theorems 1.1 and 1.2, respectively.

The proof of Theorem 1.3 has not yet appeared, the proof of Theorem 1.4 given in [11] is a constrictive one, follows similar lines as that of Telyakovskii.

Now, we shall verify that if a sequence $\{a_n\}$ belongs to S_r , then the sequence $\{n^r a_n\}$ belongs to S , with such a sequence $\{A_n\}$ which satisfies the inequality

$$(1.3) \quad \sum_{n=1}^{\infty} A_n \leq (r+1) \sum_{n=1}^{\infty} n^r A_n^{(r)}, \quad (A_n \equiv A_n^{(0)}).$$

Thus, this result and the Theorems 1.1 and 1.2 immediately imply the Theorems 1.3 and 1.4, respectively.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

2. Results

We shall deduce our assertion from a somewhat more general result. In the Introduction we have already referred to that in [5], we also defined a certain subclass of S as follows:

Let $\alpha := \{\alpha_n\}$ be a positive monotone sequence tending to infinity. A null-sequence $\{a_n\}$ belongs to the class $S(\alpha)$, if there exists a monotonically decreasing sequence $\{A_n^{(\alpha)}\}$ such that

$$\sum_{n=1}^{\infty} \alpha_n A_n^{(\alpha)} < \infty, \quad \text{and} \quad |\Delta a_n| \leq A_n^{(\alpha)} \quad \text{for all } n.$$

If we denote the class $S(\alpha)$, where $\alpha_n := n^\alpha$, $\alpha > 0$, by S_α , that is, if we introduce the definition $S_\alpha := S(n^\alpha)$, we immediately get the generalization of the classes S_r , $r = 1, 2, \dots$, for any positive α .

We shall prove our result for the classes S_α , $\alpha > 0$.


Theorem 2.1. *Let $\gamma \geq \beta > 0$. If $\{a_n\}$ belongs to the class S_γ , then the sequence $\{n^\beta a_n\}$ belongs to the class $S_{\gamma-\beta}$ and*

$$(2.1) \quad \sum_{n=1}^{\infty} n^{\gamma-\beta} A_n^{(\gamma-\beta)} \leq (\beta + 1) \sum_{n=1}^{\infty} n^\gamma A_n^{(\gamma)}$$

holds.

It is clear that if $\gamma = \beta = r$ then (2.1) gives (1.3). Thus the inequality (1.3), utilizing the assumptions of Theorem 1.3 and 1.4, and the statements of Theorems 1.1 and 1.2, implies the assertions of Theorems 1.3 and 1.4, respectively.

This is a new and short proof for the Theorems 1.3 and 1.4.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13

Remark 2.1. *The statement of the theorem is not reversible in general.*



**On the Utility of the
Telyakovskii's Class S**

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13

3. Proofs

Proof of Theorem 2.1. In order to prove our theorem we have to verify that there exists a monotonically decreasing sequence $\{A_n^{(\gamma-\beta)}\}$ such that (2.1) and

$$(3.1) \quad |\Delta(n^\beta a_n)| \leq A_n^{(\gamma-\beta)}$$

hold. Since $\{a_n\} \in S_\gamma$ thus if $\beta \geq 1$ then

$$(3.2) \quad \begin{aligned} |\Delta(n^\beta a_n)| &= |n^\beta(a_n - a_{n+1}) - a_{n+1}((n+1)^\beta - n^\beta)| \\ &\leq n^\beta |\Delta a_n| + \beta(n+1)^{\beta-1} |a_{n+1}| \\ &\leq n^\beta A_n^{(\gamma)} + \beta(n+1)^{(\beta-1)} \sum_{k=n+1}^{\infty} A_k^{(\gamma)}. \end{aligned}$$

Now define


$$A_n^{(\gamma-\beta)} := n^\beta A_n^{(\gamma)} + \beta \sum_{k=n+1}^{\infty} k^{\beta-1} A_k^{(\gamma)}.$$

By this definition and (3.2) it is clear that (3.1) holds. Next we show that the sequence $\{A_n^{(\gamma-\beta)}\}$ is monotonic, that is

$$A_{n+1}^{(\gamma-\beta)} \leq A_n^{(\gamma-\beta)}.$$

Since $(n+1)^\beta \leq n^\beta + \beta(n+1)^{\beta-1}$ and $A_{n+1}^{(\gamma)} \leq A_n^{(\gamma)}$, thus

$$A_{n+1}^{(\gamma-\beta)} = (n+1)^\beta A_{n+1}^{(\gamma)} + \beta \sum_{k=n+2}^{\infty} k^{\beta-1} A_k^{(\gamma)}$$



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

$$\leq n^\beta A_n^{(\gamma)} + \beta(n+1)^{\beta-1} A_{n+1}^{(\gamma)} + \beta \sum_{k=n+2}^{\infty} k^{\beta-1} A_k^{(\gamma)} = A_n^{(\gamma-\beta)}.$$

Finally we verify (2.1). Since

$$\begin{aligned} \sum_{n=1}^{\infty} n^{\gamma-\beta} A_n^{(\gamma-\beta)} &= \sum_{n=1}^{\infty} n^\gamma A_n^{(\gamma)} + \beta \sum_{n=1}^{\infty} n^{\gamma-\beta} \sum_{k=n+1}^{\infty} k^{\beta-1} A_k^{(\gamma)} \\ &\leq \sum_{n=1}^{\infty} n^\gamma A_n^{(\gamma)} + \beta \sum_{k=2}^{\infty} k^{\beta-1} A_k^{(\gamma)} \sum_{n=1}^k n^{\gamma-\beta} \\ &\leq (\beta+1) \sum_{n=1}^{\infty} n^\gamma A_n^{(\gamma)}. \end{aligned}$$

If $0 < \beta < 1$ then, using the first equality of (3.2), we get that


$$|\Delta(n^\beta a_n)| \leq n^\beta A_n^{(\gamma)} + \beta n^{\beta-1} \sum_{k=n+1}^{\infty} A_k^{(\gamma)}.$$

Henceforth the proof follows the lines given for $\beta \geq 1$ if we define

$$A_n^{(\gamma-\beta)} := n^\beta A_n^{(\gamma)} + \beta n^{\beta-1} \sum_{k=n+1}^{\infty} A_k^{(\gamma)}.$$

Herewith the proof is complete. \square

Proof of Remark 2.1. It suffices to prove the remark for the case $\gamma = \beta = 1$. We know that if $\{a_n\} \in S_1$ then $\{na_n\} \in S$. Our next example will show that



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

there exists a sequence $\{c_n\}$ such that $\{nc_n\} \in S$ but $\{c_n\} \notin S_1$. This verifies that the implication

$$\{a_n\} \in S_1 \Rightarrow \{na_n\} \in S$$

is not reversible.

Put

$$c_n := \frac{1}{n \log(n+1)}, \quad n \geq 1.$$

Then the sequence $\{nc_n\}$ is monotonically decreasing, tends to zero, and thus clearly belongs to the class S .

On the other hand


$$|\Delta c_n| \geq \frac{1}{n(n+1) \log(n+1)},$$

whence

$$\sum_{n=1}^{\infty} n A_n^{(1)} = \infty$$

obviously follows if $A_n^{(1)} \geq |\Delta c_n|$ holds, consequently $\{c_n\}$ does not belong to S_1 .

This proves Remark 2.1. □



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back


Close

Quit

Page 11 of 13

References

- [1] R. P. BOAS JR., *Integrability theorems for trigonometric transforms*, Springer-Verlag, Ergebnisse 38, Berlin, 1967.
- [2] A. N. KOLMOGOROV, Sur l'ordre de grandeur des coefficients de la série de Fourier-Lebesgue, *Bull. Acad. Polon. Sci. (A), Sci. Math.*, (1923), 83–86.
- [3] L. LEINDLER, On the equivalence of classes of Fourier coefficients, *Math. Ineq. & Appl.*, **3** (2000), 45–50.
- [4] L. LEINDLER, On the equivalence of classes of numerical sequences, *Analysis Math.*, **26** (2000), 227–234.
- [5] L. LEINDLER, Classes of numerical sequences, *Math. Ineq & Appl.*, **4**(4) (2001), 515–526.
- [6] S. SIDON, Hinreichende Bedingungen für den Fourier-charakter einer trigonomet-rischen Reihe, *J. London Math. Soc.*, **14** (1939), 158–160.
- [7] N. SING AND K. M. SHARMA, Integrability of trigonometric series, *J. Indian Math. Soc.*, **49** (1985), 31–38.
- [8] Č. V. STANEJEVIČ, and V.B. STANOJEVIČ, Generalizations of Sidon-Telyakovskii, theorem, *Proc. Amer. Math. Soc.*, **101** (1987), 679–684.
- [9] S. A. TELYAKOVSKIĬ, Conditions for integrability of trigonometric series and their application to study linear summability methods of Fourier series, *Izvestiya Akad. Nauk SSSR*, (Russian) **28** (1964), 1209–1236.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents




Go Back

Close

Quit

Page 12 of 13

- [10] S. A. TELYAKOVSKIĪ, On a sufficient condition of Sidon for integrability of trigonometric series, *Math. Zametki*, (Russian) **14** (1973), 317–328.
- [11] Ž. TOMOVSKI, Some results on L^1 -approximation of the r -th derivative of Fourier series, accepted for publication in *J. Inequal. Pure and Appl. Math.* and will appear in volume 3, issue 1, 2002. A pre-print is available on-line at *RGMA Research Report Collection*, **2**(5), article 11, 1999; <http://rgmia.vu.edu.au/v2n5.html>
- [12] Ž TOMOVSKI, An extension of the Sidon-Fomin inequality and applications, *Math. Ineq & Appl.*, **4**(2) (2001), 231–238.
- [13] W. H. YOUNG, On the Fourier series of bounded functions, *Proc. London Math. Soc.*, **12** (1913), 41–70.
- [14] S. Z. A. ZENEI, Integrability of trigonometric series, *Tamkang J. Math.*, **21** (1990), 295–301.



On the Utility of the
Telyakovskii's Class S

L. Leindler

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13