


# Journal of Inequalities in Pure and Applied Mathematics

## ON UNIVALENT HARMONIC FUNCTIONS

METIN ÖZTÜRK AND SIBEL YALÇIN

Uludağ Üniversitesi  
Fen-Ed. Fakültesi  
Matematik Bölümü  
16059 Görükle/BURSA  
TÜRKİYE.  
EMail: [ometin@uludag.edu.tr](mailto:ometin@uludag.edu.tr)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
003-02



---

volume 3, issue 4, article 61,  
2002.

*Received 18 January, 2002;  
accepted 26 June, 2002.*

*Communicated by: H.M. Srivastava*

---

Abstract

Contents




Home Page

Go Back

Close

Quit



## Abstract

Two classes of univalent harmonic functions on unit disc satisfying the conditions  $\sum_{n=2}^{\infty} (n-\alpha)(|a_n|+|b_n|) \leq (1-\alpha)(1-|b_1|)$  and  $\sum_{n=2}^{\infty} n(n-\alpha)(|a_n|+|b_n|) \leq (1-\alpha)(1-|b_1|)$  are given. That the ranges of the functions belonging to these two classes are starlike and convex, respectively. Sharp coefficient relations and distortion theorems are given for these functions. Furthermore results concerning the convolutions of functions satisfying above inequalities with univalent, harmonic and convex functions in the unit disk and with harmonic functions having positive real part.

*2000 Mathematics Subject Classification:* 30C45, 31A05.

*Key words:* Convex harmonic functions, Starlike harmonic functions, Extremal problems.

## Contents

1	Introduction .....	3
2	The Classes $HS(\alpha)$ and $HC(\alpha)$ .....	4
3	Main Results .....	6
	References	

### On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit


Page 2 of 18

# 1. Introduction

Let  $U$  denote the open unit disc and  $S_H$  denote the class of all complex valued, harmonic, orientation-preserving, univalent functions  $f$  in  $U$  normalized by  $f(0) = f_z(0) - 1 = 0$ . Each  $f \in S_H$  can be expressed as  $f = h + \bar{g}$  where  $h$  and  $g$  belong to the linear space  $H(U)$  of all analytic functions on  $U$ .

Firstly, Clunie and Sheil-Small [3] studied  $S_H$  together with some geometric subclasses of  $S_H$ . They proved that although  $S_H$  is not compact, it is normal with respect to the topology of uniform convergence on compact subsets of  $U$ . Meanwhile the subclass  $S_H^0$  of  $S_H$  consisting of the functions having the property  $f_{\bar{z}}(0) = 0$  is compact.

In this article we concentrate on two specific subclasses of univalent harmonic mappings. These classes have corresponding meaning in the class of convex and starlike analytic functions of order  $\alpha$ . The geometrical properties of the functions in these classes together with the neighborhoods in the meaning of Ruschevych and convolution products are considered.



---

On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

## 2. The Classes $HS(\alpha)$ and $HC(\alpha)$

Let  $U_r = \{z : |z| < r, 0 < r \leq 1\}$  and  $U_1 = U$ . Harmonic, complex-valued, orientation-preserving, univalent mappings  $f$  defined on  $U$  can be written as

$$(2.1) \quad f(z) = h(z) + \overline{g(z)},$$

where

$$(2.2) \quad h(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=1}^{\infty} b_n z^n$$

are analytic in  $U$ .


Denote by  $HS(\alpha)$  the class of all functions of the form (2.1) that satisfy the condition

$$(2.3) \quad \sum_{n=2}^{\infty} (n - \alpha)(|a_n| + |b_n|) \leq (1 - \alpha)(1 - |b_1|)$$

and by  $HC(\alpha)$  the subclass of  $HS(\alpha)$  that consists of all functions subject to the condition

$$\sum_{n=2}^{\infty} n(n - \alpha)(|a_n| + |b_n|) \leq (1 - \alpha)(1 - |b_1|),$$

where  $0 \leq \alpha < 1$  and  $0 \leq |b_1| < 1$ . The corresponding subclasses of  $HS(\alpha)$  and  $HC(\alpha)$  with  $b_1 = 0$  will be denoted by  $HS^0(\alpha)$  and  $HC^0(\alpha)$ , respectively. When  $\alpha = 0$ , these classes are denoted by  $HS$  and  $HC$  and have been studied



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 4 of 18

by Y. Avci and E. Zlotkiewicz [2]. If  $|b_1| = 1$  and (2.3) is satisfied, then the mappings  $z + \overline{b_1 z}$  are not univalent in  $U$  and of no interest.

If  $h, g, H, G$  are of the form (2.2) and if

$$f(z) = h(z) + \overline{g(z)} \quad \text{and} \quad F(z) = H(z) + \overline{G(z)}$$

then the convolution of  $f$  and  $F$  is defined to be the function

$$f * F(z) = z + \sum_{n=2}^{\infty} a_n A_n z^n + \overline{\sum_{n=1}^{\infty} b_n B_n z^n}$$

while the integral convolution is defined by


$$f \diamond F(z) = z + \sum_{n=2}^{\infty} \frac{a_n A_n}{n} z^n + \overline{\sum_{n=1}^{\infty} \frac{b_n B_n}{n} z^n}.$$

The  $\delta$ -neighborhood of  $f$  is the set

$$N_{\delta}(f) = \left\{ F : \sum_{n=2}^{\infty} n(|a_n - A_n| + |b_n - B_n|) + |b_1 - B_1| \leq \delta \right\}$$

(see [1], [5]). In this case, let us define the generalized  $\delta$ -neighborhood of  $f$  to be the set

$$N(f) = \left\{ F : \sum_{n=2}^{\infty} (n - \alpha)(|a_n - A_n| + |b_n - B_n|) + (1 - \alpha)|b_1 - B_1| \leq (1 - \alpha)\delta \right\}.$$



### On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 5 of 18

### 3. Main Results

First, let us give the interrelations between the classes  $HS(\alpha)$  and  $HS, HC(\alpha)$  and  $HC$ .

**Theorem 3.1.**  $HS(\alpha) \subset HS$  and  $HC(\alpha) \subset HC$ . Consequently  $HS^0(\alpha) \subset HS^0$  and  $HC^0(\alpha) \subset HC^0$ . In particular if  $0 \leq \alpha_1 \leq \alpha_2 < 1$  then  $HS(\alpha_2) \subset HS(\alpha_1)$  and  $HC(\alpha_2) \subset HC(\alpha_1)$ .

*Proof.* Since

$$(3.1) \quad \sum_{n=2}^{\infty} n(|a_n| + |b_n|) \leq \sum_{n=2}^{\infty} \frac{n - \alpha}{1 - \alpha} (|a_n| + |b_n|) \leq 1 - |b_1|$$

and


$$\sum_{n=2}^{\infty} n^2(|a_n| + |b_n|) \leq \sum_{n=2}^{\infty} \frac{n(n - \alpha)}{1 - \alpha} (|a_n| + |b_n|) \leq 1 - |b_1|$$

we have the proof of theorem.  $\square$

**Corollary 3.2.** (i) Each member of  $HS^0(\alpha)$  maps  $U$  onto a domain starlike with respect to the origin.

(ii) Functions of the class  $HC^0(\alpha)$  maps  $U_r$  onto convex domains.

**Theorem 3.3.** The class  $HS(\alpha)$  consist of univalent sense preserving harmonic mappings.



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 6 of 18

*Proof.* From [2, Theorem 1] for  $f$  in  $HS(\alpha)$  and for  $z_1, z_2 \in U$  with  $z_1 \neq z_2$  we have

$$|h(z_1) - h(z_2)| \geq |z_1 - z_2| \left( 1 - |z_2| \sum_{n=2}^{\infty} n|a_n| \right)$$

and

$$|g(z_1) - g(z_2)| \leq |z_1 - z_2| \left( |b_1| + |z_2| \sum_{n=2}^{\infty} n|b_n| \right).$$

When we consider the relation (3.1), it follows that

$$\begin{aligned} |f(z_1) - f(z_2)| &\geq |z_1 - z_2| \left( 1 - |b_1| - |z_2| \sum_{n=2}^{\infty} n(|a_n| + |b_n|) \right) \\ &= |z_1 - z_2| \left( 1 - |b_1| - |z_2| \sum_{n=2}^{\infty} (n - \alpha)(|a_n| + |b_n|) \right. \\ &\quad \left. - \alpha|z_2| \sum_{n=2}^{\infty} (|a_n| + |b_n|) \right) \\ &\geq |z_1 - z_2| [1 - |b_1| - |z_2|(1 - \alpha)(1 - |b_1|) - \alpha|z_2|(1 - |b_1|)] \\ &= |z_1 - z_2|(1 - |b_1|)(1 - |z_2|) > 0. \end{aligned}$$

So  $f$  is univalent. Since

$$\begin{aligned} J_f(z) &= |h'(z)|^2 - |g'(z)|^2 \\ &\geq (|h'(z)| + |g'(z)|) \left( 1 - |z| \sum_{n=2}^{\infty} n|a_n| - |b_1| - |z| \sum_{n=2}^{\infty} n|b_n| \right) \end{aligned}$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 7 of 18

$$\geq (|h'(z)| + |g'(z)|)(1 - |b_1|)(1 - |z|) > 0$$

$f$  is sense preserving. □

**Remark 3.1.** (i) The functions  $f_n(z) = z + \frac{n-1}{n+1} e^{i\theta} \bar{z}$  are in  $HS(\alpha)$  and the sequence converges uniformly to  $z + e^{i\theta} \bar{z}$ . Thus the class  $HS(\alpha)$  is not compact.

(ii) If  $f \in HS(\alpha)$ , then for each  $r$ ,  $0 < r < 1$ ,  $r^{-1}f(rz) \in HS(\alpha)$ .

(iii) If  $f \in HS(\alpha)$  and  $f_0(z) = \frac{f(z) - \overline{b_1 f(z)}}{1 - |b_1|^2}$  then  $f_0 \in HS^0(\alpha)$ , but  $f(z) = f_0(z) + b_1 \overline{f_0(z)}$  may not be in  $HS(\alpha)$ .

(iv) If  $f = h + \bar{g} \in HS^0(\alpha)$  then the function


$$F(z) = \int_0^1 \frac{f(tz)}{t} dt = \int_0^z \frac{h(u)}{u} du + \overline{\int_0^z \frac{g(u)}{u} du}$$

satisfies (2.3) with  $b_1 = 0$ , hence  $F(z)$  is a convex harmonic mapping. Convexity of  $F(z)$  however, does not imply starlikeness of  $f(z)$  (or even univalence) in general situation.

**Theorem 3.4.** Each function in the class  $HS^0(\alpha)$  maps disks  $U_r$ ,  $r \leq \frac{1-\alpha}{2-\alpha}$  onto convex domains. The constant  $\frac{1-\alpha}{2-\alpha}$  is best possible.

*Proof.* If  $f \in HS^0(\alpha)$  and  $r$ ,  $0 < r < 1$  be fixed, then  $r^{-1}f(rz) \in HS^0(\alpha)$ . We must find an upper bound for  $r$  such that

$$\sum_{n=2}^{\infty} \frac{n(n-\alpha)}{1-\alpha} (|a_n| + |b_n|) r^{n-1} \leq 1.$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 8 of 18

Since  $f \in HS^0$  we have

$$\sum_{n=2}^{\infty} \frac{n(n-\alpha)}{1-\alpha} (|a_n| + |b_n|) r^{n-1} = \sum_{n=2}^{\infty} n(|a_n| + |b_n|) \frac{(n-\alpha)r^{n-1}}{1-\alpha} \leq 1$$

provided  $(n-\alpha)r^{n-1}/(1-\alpha) \leq 1$  which is true if  $r \leq (1-\alpha)/(2-\alpha)$ .  $\square$

**Theorem 3.5.** *If  $f \in HS(\alpha)$ , then*

$$|f(z)| \leq |z|(1 + |b_1|) + \frac{(1-\alpha^2)(1-|b_1|)}{2} |z|^2$$

and

$$|f(z)| \geq (1 - |b_1|) \left( |z| - (1 - \alpha^2) \frac{|z|^2}{2} \right).$$

*Equalities are attained by the functions*

$$f_{\theta}(z) = z + |b_1|e^{i\theta}\bar{z} + \frac{1-|b_1|}{2}(1-\alpha^2)\bar{z}^2$$


*for properly chosen real  $\theta$ .*

*Proof.* We have

$$|f(z)| \leq |z|(1 + |b_1|) + |z|^2 \sum_{n=2}^{\infty} (|a_n| + |b_n|).$$

Since

$$\sum_{n=2}^{\infty} (|a_n| + |b_n|) \leq \frac{1}{2}(1-\alpha)(1-|b_1|) + \frac{\alpha}{2}(|a_2| + |b_2|)$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 9 of 18

$$\begin{aligned}
& -\frac{1}{2} \sum_{n=3}^{\infty} (n - \alpha - 2)(|a_n| + |b_n|) \\
& \leq \frac{1}{2}(1 - \alpha)(1 - |b_1|) + \frac{\alpha}{2}(1 - \alpha)(1 - |b_1|) \\
& = \frac{1}{2}(1 - \alpha^2)(1 - |b_1|)
\end{aligned}$$

it follows that

$$|f(z)| \leq |z|(1 + |b_1|) + \frac{(1 - \alpha^2)(1 - |b_1|)}{2} |z|^2$$

for  $z$  in  $U$ .

Similarly we get

$$\begin{aligned}
|f(z)| & \geq |z|(1 - |b_1|) - |z|^2 \sum_{n=2}^{\infty} (|a_n| + |b_n|) \\
& \geq |z|(1 - |b_1|) - \frac{(1 - \alpha^2)(1 - |b_1|)}{2} |z|^2 \\
& = (1 - |b_1|) \left( |z| - (1 - \alpha^2) \frac{|z|^2}{2} \right).
\end{aligned}$$

The classes  $HS(\alpha)$  and  $HC(\alpha)$  are uniformly bounded, hence they are normal.  $\square$

**Remark 3.2.** We can give a similar result for  $HC(\alpha)$  to that given for  $HS(\alpha)$ . As the proof is similar we shall omit it.



### On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 10 of 18

**Theorem 3.6.** If  $f \in HC(\alpha)$ , then

$$|f(z)| \leq |z|(1 + |b_1|) + \frac{3 - \alpha - 2\alpha^2}{2\alpha}(1 - |b_1|)|z|^2$$

and

$$|f(z)| \geq (1 - |b_1|) \left( |z| - \frac{3 - \alpha - 2\alpha^2}{2\alpha}|z|^2 \right).$$

Equalities are attained by the functions

$$f_\theta(z) = z + |b_1|e^{i\theta}\bar{z} + \frac{3 - \alpha - 2\alpha^2}{2\alpha}\bar{z}^2$$

for properly chosen real  $\theta$ .

**Theorem 3.7.** The extreme points of  $HS^0(\alpha)$  are only the functions of the form:  $z + a_n z^n$  or  $z + \overline{b_m z^m}$  with


$$|a_n| = \frac{1 - \alpha}{n - \alpha}, \quad |b_m| = \frac{1 - \alpha}{m - \alpha}, \quad 0 \leq \alpha < 1.$$

*Proof.* Suppose that

$$f(z) = z + \sum_{n=2}^{\infty} (a_n z^n + \overline{b_n z^n})$$

is such that

$$\sum_{n=2}^{\infty} \frac{n - \alpha}{1 - \alpha} (|a_n| + |b_n|) < 1, \quad a_k > 0.$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 11 of 18

Then, if  $\lambda > 0$  is small enough we can replace  $a_k$  by  $a_k - \lambda$ ,  $a_k + \lambda$  and we obtain two functions  $f_1(z)$ ,  $f_2(z)$  that satisfy the same condition and for which one gets  $f(z) = \frac{1}{2}[f_1(z) + f_2(z)]$ . Hence  $f$  is not a possible extreme point of  $HS^0(\alpha)$ .

Let now  $f \in HS^0(\alpha)$  be such that

$$(3.2) \quad \sum_{n=2}^{\infty} \frac{n - \alpha}{1 - \alpha} (|a_n| + |b_n|) = 1, \quad a_k \neq 0, b_l \neq 0$$

If  $\lambda > 0$  is small enough and if  $\mu, \zeta$  with  $|\mu| = |\zeta| = 1$  are properly chosen complex numbers, then leaving all but  $a_k, b_l$  coefficients of  $f(z)$  unchanged and replacing  $a_k, b_l$  by


$$\begin{aligned} a_k + \lambda \frac{1 - \alpha}{k - \alpha} \mu, & \quad b_l - \lambda \frac{1 - \alpha}{l - \alpha} \zeta \\ a_k - \lambda \frac{1 - \alpha}{k - \alpha} \mu, & \quad b_l + \lambda \frac{1 - \alpha}{l - \alpha} \zeta \end{aligned}$$

we obtain functions  $f_1(z)$ ,  $f_2(z)$  that satisfy (3.2) and such that  $f(z) = \frac{1}{2}[f_1(z) + f_2(z)]$ . In this case  $f$  can not be an extreme point. Thus for  $|a_n| = (1 - \alpha)/(n - \alpha)$ ,  $|b_m| = (1 - \alpha)/(m - \alpha)$ ,  $f(z) = z + a_n z^n$  or  $f(z) = z + \overline{b_m} z^m$  are extreme points of  $HS^0(\alpha)$ .  $\square$

Similarly we can obtain that the following result is true.

**Theorem 3.8.** *The extreme points of  $HC^0(\alpha)$  are only the functions of the form:  $z + a_n z^n$  or  $z + \overline{b_m} z^m$  with*

$$|a_n| = \frac{1 - \alpha}{n(n - \alpha)}, \quad |b_m| = \frac{1 - \alpha}{m(m - \alpha)}, \quad 0 \leq \alpha < 1.$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 12 of 18

Let  $K_H^0$  denote the class of harmonic univalent functions of the form (2.1) with  $b_1 = 0$  that map  $U$  onto convex domains. It is known [3, Theorem 5.10] that the sharp inequalities

$$2|A_n| \leq n + 1, \quad 2|B_n| \leq n - 1$$

are true.

**Theorem 3.9.** *Suppose that*

$$F(z) = z + \sum_{n=2}^{\infty} (A_n z^n + \overline{B_n z^n})$$

*belongs to  $K_H^0$ . Then*

(i) *If  $f \in HC^0(\alpha)$  then  $f * F$  is starlike univalent and  $f \diamond F$  is convex.*


(ii) *If  $f(z)$  satisfies the condition*

$$\sum_{n=2}^{\infty} n^2(n - \alpha)(|a_n| + |b_n|) \leq 1 - \alpha$$

*then  $f * F$  is convex univalent.*

*Proof.* We justify the case (i). Since

$$\begin{aligned} \sum_{n=2}^{\infty} (n - \alpha)(|a_n A_n| + |b_n B_n|) &= \sum_{n=2}^{\infty} n(n - \alpha) \left( |a_n| \left| \frac{A_n}{n} \right| + |b_n| \left| \frac{B_n}{n} \right| \right) \\ &\leq \sum_{n=2}^{\infty} n(n - \alpha)(|a_n| + |b_n|) \leq 1 - \alpha \end{aligned}$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 13 of 18

it follows that  $f * F$  is in  $HS^0(\alpha)$ . Namely  $f * F$  is starlike univalent.

Furthermore, the transformation

$$\int_0^1 \frac{f * F(tz)}{t} dt = f \diamond F(z)$$

now shows that  $f \diamond F \in HC^0(\alpha)$ . □

Let  $S$  denote the class of analytic univalent functions of the form  $F(z) = z + \sum_{n=2}^{\infty} A_n z^n$ . It is well known that the sharp inequality  $|A_n| \leq n$  is true.

**Theorem 3.10.** *If  $f \in HC^0(\alpha)$  and  $F \in S$  then for  $|\varepsilon| \leq 1$ ,  $f * (F + \varepsilon \bar{F})$  is starlike univalent.*

*Proof.* Since

$$\sum_{n=2}^{\infty} (n - \alpha)(|a_n A_n| + |b_n B_n|) \leq \sum_{n=2}^{\infty} n(n - \alpha)(|a_n| + |b_n|) \leq 1 - \alpha$$

it follows that  $f * (F + \varepsilon \bar{F})$  is starlike univalent.


Let  $P_H^0$  denote the class of functions  $F$  complex and harmonic in  $U$ ,  $f = h + \bar{g}$  such that  $\operatorname{Re} f(z) > 0$ ,  $z \in U$  and

$$H(z) = 1 + \sum_{n=1}^{\infty} A_n z^n, \quad G(z) = \sum_{n=2}^{\infty} B_n z^n.$$

It is known [4, Theorem 3] that the sharp inequalities

$$|A_n| \leq n + 1, \quad |B_n| \leq n - 1$$

are true. □



### On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit

Page 14 of 18

**Theorem 3.11.** *Suppose that*

$$F(z) = 1 + \sum_{n=1}^{\infty} (A_n z^n + \overline{B_n z^n})$$

*belongs to  $P_H^0$ . Then*

(i) *If  $f \in HC^0(\alpha)$  then for  $\frac{3}{2} \leq |A_1| \leq 2$ ,  $\frac{1}{A_1} f * F$  is starlike univalent and  $\frac{1}{A_1} f \diamond F$  is convex.*

(ii) *If  $f(z)$  satisfies the condition*


$$\sum_{n=2}^{\infty} n^2(n - \alpha)(|a_n| + |b_n|) \leq 1 - \alpha$$

*then  $\frac{1}{A_1} f * F$  is convex univalent.*

*Proof.* We justify the case (ii). Since

$$\begin{aligned} \sum_{n=2}^{\infty} n(n - \alpha) \left( \left| \frac{a_n A_n}{A_1} \right| + \left| \frac{b_n B_n}{A_1} \right| \right) &\leq \sum_{n=2}^{\infty} n^2(n - \alpha) \left( |a_n| \frac{n+1}{|A_1|n} + |b_n| \frac{n-1}{|A_1|n} \right) \\ &\leq \sum_{n=2}^{\infty} n^2(n - \alpha)(|a_n| + |b_n|) \leq 1 - \alpha \end{aligned}$$

$\frac{1}{A_1} f * F$  is convex univalent. □




---

**On Univalent Harmonic Functions**

Metin Öztürk and Sibel Yalçın

---

Title Page

Contents



Go Back

Close

Quit

Page 15 of 18

**Remark 3.3.** If  $f \in HS^0(\alpha)$  and  $F \in K_H^0$ , then  $f * F$  need not be univalent. For example, if  $f(z) = z + \frac{1-\alpha}{n-\alpha} z^n$  and  $F(z) = \operatorname{Re} \left( \frac{z}{1-z} \right) + i \operatorname{Im} \left( \frac{z}{(1-z)^2} \right)$  then  $f * F(z) = z + \frac{(n+1)(1-\alpha)}{2(n-\alpha)} z^n$  is not univalent in  $U$ . But  $f \diamond F$  is univalent and starlike.


**Theorem 3.12.** Let

$$f(z) = z + \overline{b_1}z + \sum_{n=2}^{\infty} (a_n z^n + \overline{b_n z^n})$$

is a member of  $HC(\alpha)$ . If  $\delta \leq \frac{1-\alpha}{2-\alpha}(1 - |b_1|)$ , then  $N(f) \subset HS(\alpha)$ .

*Proof.* Let  $f \in HC(\alpha)$  and  $F(z) = z + \overline{B_1}z + \sum_{n=2}^{\infty} (A_n z^n + \overline{B_n z^n})$  belong to  $N(f)$ . We have

$$\begin{aligned} & (1-\alpha)|B_1| + \sum_{n=2}^{\infty} (n-\alpha)(|A_n| + |B_n|) \\ & \leq (1-\alpha)|B_1 - b_1| + (1-\alpha)|b_1| \\ & \quad + \sum_{n=2}^{\infty} (n-\alpha)(|A_n - a_n| + |B_n - b_n|) + \sum_{n=2}^{\infty} (n-\alpha)(|a_n| + |b_n|) \\ & \leq (1-\alpha)\delta + (1-\alpha)|b_1| + \frac{1}{2-\alpha} \sum_{n=2}^{\infty} n(n-\alpha)(|a_n| + |b_n|) \\ & \leq (1-\alpha)\delta + (1-\alpha)|b_1| + \frac{1-\alpha}{2-\alpha}(1 - |b_1|) \\ & \leq 1 - \alpha. \end{aligned}$$



On Univalent Harmonic Functions

Metin Öztürk and Sibel Yalçın

Title Page

Contents



Go Back

Close

Quit


Page 16 of 18

Hence, for

$$\delta \leq \frac{1 - \alpha}{2 - \alpha} (1 - |b_1|)$$

$F(z) \in HS(\alpha)$ .

□



---

**On Univalent Harmonic Functions**

Metin Öztürk and Sibel Yalçın

---

Title Page

Contents



Go Back


Close

Quit

Page 17 of 18

## References

- [1] O. ALTINTAŞ, Ö. ÖZKAN AND H.M. SRIVASTAVA, Neighborhoods of a class of analytic functions with negative coefficients, *Appl. Math. Lett.*, **13** (2000), 63–67.
- [2] Y. AVCI AND E. ZLOTKIEWICZ, On harmonic univalent mappings, *Ann. Universitatis Mariae Curie-Sklodowska*, **XLIV**(1), Sectio A (1990), 1–7.
- [3] J. CLUNIE AND T. SHEIL-SMALL, Harmonic univalent functions, *Ann. Acad. Sci. Fen. Series A. I, Math.*, **9** (1984), 3–25.
- [4] Z.J. JAKUBOWSKI, W. MAJCHRZAK AND K. SKALSKA, Harmonic mappings with a positive real part, *Materialy XIV Konferencji z Teorii Zagadnien Ekstremalnych*, Lodz. (1993), 17–24.
- [5] St. RUSCHEWEYH, Neighborhoods of univalent functions, *Proc. Amer. Math. Soc.*, **81** (1981), 521–528.



---

On Univalent Harmonic  
Functions

Metin Öztürk and Sibel Yalçın

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18