

Journal of Inequalities in Pure and Applied Mathematics

SOME INTEGRAL INEQUALITIES INVOLVING TAYLOR'S REMAINDER. II

HILLEL GAUCHMAN

Department of Mathematics,
Eastern Illinois University,
Charleston, IL 61920, USA
EMail: cfhvg@ux1.cts.eiu.edu

©2000 Victoria University
ISSN (electronic): 1443-5756
011-02



volume 4, issue 1, article 1,
2003.

*Received 18 February, 2002;
accepted 12 November, 2002.*

Communicated by: J.E. Pečarić

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this paper, using Grüss' and Chebyshev's inequalities we prove several inequalities involving Taylor's remainder.

2000 Mathematics Subject Classification: 26D15.

Key words: Taylor's remainder, Grüss' inequality, Chebyshev's inequality.

Contents

1	Introduction and Lemma	3
2	Applications of Grüss' Inequality	5
3	Applications of Chebyshev's Inequality	8
	References	



Some Integral Inequalities Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

1. Introduction and Lemma

This paper is a continuation of our paper [4]. As in [4], our goal is to prove several integral inequalities involving Taylor's remainder. Our method is similar to that used in [4]. However, while in [4] we deduced our inequalities from Steffensen's inequality, in the present paper we use Grüss' and Chebyshev's inequalities. We are thankful to Professor S.S. Dragomir who pointed out that Grüss' and Chebyshev's inequalities were used earlier by G.A. Anastassiou and S.S. Dragomir [2], [3] to obtain results on Taylor's remainder different from but related to the results of this paper. The main results of this paper are Theorems 2.1 and 3.1.

In what follows n denotes a non-negative integer. We will denote by $R_{n,f}(c, x)$ the n th Taylor's remainder of function $f(x)$ with center c , i.e.

$$R_{n,f}(c, x) = f(x) - \sum_{k=0}^n \frac{f^{(k)}(c)}{k!} (x - c)^k.$$

Lemma 1.1. *Let f be a function defined on $[a, b]$. Assume that $f \in C^{n+1}([a, b])$. Then one has the representations*

$$(1.1) \quad \int_a^b \frac{(b-x)^{n+1}}{(n+1)!} f^{(n+1)}(x) dx = \int_a^b R_{n,f}(a, x) dx,$$

and

$$(1.2) \quad \int_a^b \frac{(x-a)^{n+1}}{(n+1)!} f^{(n+1)}(x) dx = (-1)^{n+1} \int_a^b R_{n,f}(b, x) dx.$$



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close


Quit

Page 3 of 12

Proof. Observe that:

$$\begin{aligned}
 & \int_a^b \frac{(b-x)^{n+1}}{(n+1)!} f^{(n+1)}(x) dx \\
 &= \int_a^b \frac{(b-x)^{n+1}}{(n+1)!} df^{(n)}(x) \\
 &= f^{(n)}(x) \frac{(b-x)^{n+1}}{(n+1)!} \Big|_{x=a}^{x=b} + \int_a^b \frac{(b-x)^n}{n!} f^{(n)}(x) dx \\
 &= -f^{(n)}(a) \frac{(b-a)^{n+1}}{(n+1)!} + \int_a^b \frac{(b-x)^n}{n!} f^{(n)}(x) dx \\
 &= -f^{(n)}(a) \frac{(b-a)^{n+1}}{(n+1)!} - f^{(n-1)}(a) \frac{(b-a)^n}{n!} + \int_a^b \frac{(b-x)^{n-1}}{(n-1)!} f^{(n-1)}(x) dx \\
 &= \dots \\
 &= -f^{(n)}(a) \frac{(b-a)^{n+1}}{(n+1)!} - f^{(n-1)}(a) \frac{(b-a)^n}{n!} - \dots - f(a) \frac{b-a}{1!} + \int_a^b f(x) dx \\
 &= \int_a^b \left[f(x) - \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k \right] dx \\
 &= \int_a^b R_{n,f}(a, x) dx.
 \end{aligned}$$

The proof of (1.2) is similar to the proof of (1.1) and we omit it. \square



**Some Integral Inequalities
Involving Taylor's Remainder. II**

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

2. Applications of Grüss' Inequality

The following inequality is called Grüss' inequality [5]:

Let $F(x)$ and $G(x)$ be two functions defined and integrable on $[a, b]$. Further let

$$m \leq F(x) \leq M \quad \text{and} \quad \varphi \leq G(x) \leq \Phi$$

for each $x \in [a, b]$, where m, M, φ, Φ are constants. Then


$$\left| \int_a^b F(x)G(x)dx - \frac{1}{b-a} \int_a^b F(x)dx \cdot \int_a^b G(x)dx \right| \leq \frac{b-a}{4}(M-m)(\Phi-\varphi).$$

Theorem 2.1. Let $f(x)$ be a function defined on $[a, b]$ such that $f(x) \in C^{n+1}([a, b])$ and $m \leq f^{(n+1)}(x) \leq M$ for each $x \in [a, b]$, where m and M are constants. Then

$$(2.1) \quad \left| \int_a^b R_{n,f}(a, x)dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!}(b-a)^{n+1} \right| \leq \frac{(b-a)^{n+2}}{4(n+1)!}(M-m)$$

and

$$(2.2) \quad \left| (-1)^{n+1} \int_a^b R_{n,f}(b, x)dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!}(b-a)^{n+1} \right| \leq \frac{(b-a)^{n+2}}{4(n+1)!}(M-m).$$



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents




Go Back

Close

Quit

Page 5 of 12



**Some Integral Inequalities
Involving Taylor's Remainder. II**

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

Proof. Set $F(x) = f^{(n+1)}(x)$, $G(x) = \frac{(b-x)^{n+1}}{(n+1)!}$. Then $m \leq F(x) \leq M$ and $0 \leq G(x) \leq \frac{(b-a)^{n+1}}{(n+1)!}$. By Grüss' inequality,

$$\left| \int_a^b \frac{(b-x)^{n+1}}{(n+1)!} f^{(n+1)}(x) dx - \frac{1}{b-a} \int_a^b f^{(n+1)}(x) dx \cdot \int_a^b \frac{(b-x)^{n+1}}{(n+1)!} dx \right| \leq \frac{b-a}{4} \cdot \frac{(b-a)^{n+1}}{(n+1)!} (M-m).$$

Using Lemma 1.1, we obtain

$$\left| \int_a^b R_{n,f}(a, x) - \frac{1}{b-a} [f^{(n)}(b) - f^{(n)}(a)] \cdot \frac{(b-a)^{n+2}}{(n+2)!} \right| \leq \frac{(b-a)^{n+2}}{4(n+1)!} (M-m).$$

That proves (2.1).

To prove (2.2), we set $F(x) = f^{(n+1)}(x)$, $G(x) = \frac{(a-x)^{n+1}}{(n+1)!}$, and continue as in the proof of (2.1). □

Now we consider the simplest cases of Theorem 2.1, namely the cases when $n = 0$ or 1 .

Corollary 2.2. *Let $f(x)$ be a function defined on $[a, b]$ such that $f(x) \in C^2([a, b])$*

and $m \leq f''(x) \leq M$ for each $x \in [a, b]$, where m and M are constants. Then

$$(2.3) \quad \left| \int_a^b f(x)dx - f(a)(b-a) - \frac{2f'(a) + f'(b)}{6}(b-a)^2 \right| \leq \frac{(b-a)^3}{8}(M-m),$$

$$(2.4) \quad \left| \int_a^b f(x)dx - f(b)(b-a) + \frac{2f'(b) + f'(a)}{6}(b-a)^2 \right| \leq \frac{(b-a)^3}{8}(M-m),$$


$$(2.5) \quad \left| \int_a^b f(x)dx - \frac{f(a) + f(b)}{2}(b-a) + \frac{f'(b) - f'(a)}{12}(b-a)^2 \right| \leq \frac{(b-a)^3}{8}(M-m).$$

Proof. To obtain (2.3) and (2.4) we take $n = 1$ in (2.1) and (2.2) of Theorem 2.1. Taking half the sum of (2.3) and (2.4), we obtain (2.5). \square

Remark 2.1. Taking $n = 0$ in (2.1) and (2.2), we obtain that if $m \leq f'(x) \leq M$ on $[a, b]$, then

$$\left| \int_a^b f(x)dx - \frac{f(a) + f(b)}{2}(b-a) \right| \leq \frac{(b-a)^2}{4}(M-m).$$

This inequality is weaker than a modification of Iyengar's inequality due to Agarwal and Dragomir [1].



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12

3. Applications of Chebyshev's Inequality

The following is Chebyshev's inequality [5]:

Let $F, G : [a, b] \rightarrow \mathbb{R}$ be integrable functions, both increasing or both decreasing. Then

$$\int_a^b F(x)G(x)dx \geq \frac{1}{b-a} \int_a^b F(x)dx \cdot \int_a^b G(x)dx.$$

If one of the functions is increasing and the other decreasing, then the above inequality is reversed.


Theorem 3.1. Let $f(x)$ be a function defined on $[a, b]$ such that $f(x) \in C^{(n+1)}([a, b])$.

If $f^{(n+1)}(x)$ is increasing on $[a, b]$, then,

$$(3.1) \quad -\frac{f^{(n+1)}(b) - f^{(n+1)}(a)}{4(n+1)!}(b-a)^{n+2} \leq \int_a^b R_{n,f}(a, x)dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!}(b-a)^{n+1} \leq 0,$$

and

$$(3.2) \quad 0 \leq (-1)^{(n+1)} \int_a^b R_{n,f}(b, x)dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!}(b-a)^{n+1} \leq \frac{f^{(n+1)}(b) - f^{(n+1)}(a)}{4(n+1)!}(b-a)^{n+2}.$$



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12

If $f^{(n+1)}(x)$ is decreasing on $[a, b]$, then

$$(3.3) \quad 0 \leq \int_a^b R_{n,f}(a, x) dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!} (b-a)^{n+1} \\ \leq \frac{f^{(n+1)}(a) - f^{(n+1)}(b)}{4(n+1)!} (b-a)^{n+2},$$

and


$$(3.4) \quad -\frac{f^{(n+1)}(a) - f^{(n+1)}(b)}{4(n+1)!} (b-a)^{n+2} \\ \leq (-1)^{(n+1)} \int_a^b R_{n,f}(b, x) dx - \frac{f^{(n)}(b) - f^{(n)}(a)}{(n+2)!} (b-a)^{n+1} \\ \leq 0.$$

Proof. Set $F(x) = f^{(n+1)}(x)$ and $G(x) = \frac{(b-x)^{(n+1)}}{(n+1)!}$. Then $F(x)$ is increasing and $G(x)$ decreasing on $[a, b]$. Using Chebyshev's inequality for $F(x)$ and $G(x)$ and (1.1), we obtain right inequality in (3.1). Left inequality in (3.1) follows readily from (2.1), if we take into account that since $f^{(n+1)}(x)$ is increasing on $[a, b]$, $f^{(n+1)}(a) \leq f^{(n+1)}(x) \leq f^{(n+1)}(b)$ for all $x \in [a, b]$.

To prove (3.2), set $F(x) = f^{(n+1)}(x)$ and $G(x) = \frac{(x-a)^{(n+1)}}{(n+1)!}$. The rest of the proof is the same as in the proof of (3.1).

The proofs of (3.3) and (3.4) are similar to those of (3.1) and (3.2) respectively, and we omit them. \square

We now consider the simplest cases of Theorem 3.1, namely the cases when $n = 0$ or 1 .



Some Integral Inequalities Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 9 of 12

Corollary 3.2. Let $f(x)$ be a function defined on $[a, b]$ such that $f(x) \in C^2([a, b])$. If $f''(x)$ is increasing on $[a, b]$, then


$$(3.5) \quad -\frac{f''(b) - f''(a)}{8}(b-a)^2 \leq \frac{1}{b-a} \int_a^b f(x)dx - f(a) - \frac{2f'(a) + f'(b)}{6}(b-a) \leq 0$$

$$(3.6) \quad 0 \leq \frac{1}{b-a} \int_a^b f(x)dx - f(b) + \frac{f'(a) + 2f'(b)}{6}(b-a) \leq \frac{f''(b) - f''(a)}{8}(b-a)^2,$$

$$(3.7) \quad \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{f(a) + f(b)}{2} + \frac{f'(b) - f'(a)}{12}(b-a) \right| \leq \frac{f''(b) - f''(a)}{16}(b-a)^2.$$

Proof. To obtain (3.5) and (3.6) we take $n = 1$ in (3.1) and (3.2) of Theorem 3.1. We obtain (3.7) taking half the sum of (3.5) and (3.6). \square

Remark 3.1. The inequalities similar to (3.5) – (3.7) for the case of decreasing $f''(x)$ can be obtained substituting $-f(x)$ instead of $f(x)$ into inequalities (3.5) – (3.7).



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 10 of 12

Remark 3.2. Taking $n = 0$ in Theorem 3.1, we obtain that if $f'(x)$ is increasing on $[a, b]$, then

$$(3.8) \quad \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{4}(b - a) \leq \frac{1}{b - a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

Let us compare (3.8) with the following Hermite-Hadamard's inequality [6]:

If $f(x)$ is convex on $[a, b]$ (in particular if $f'(x)$ exists and increasing on $[a, b]$), then


$$f\left(\frac{a + b}{2}\right) \leq \frac{1}{b - a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

We see that the right inequality in (3.8) is the same as the right Hermite-Hadamard's inequality. However, it can be easily proved that the left inequality in (3.8) is weaker than the left Hermite-Hadamard's inequality.

Remark 3.3. Taking the difference of (3.5) and (3.6), we obtain that if $f''(x)$ is increasing on $[a, b]$, then

$$0 \leq \frac{f'(a) + f'(b)}{2} - \frac{f(b) - f(a)}{b - a} \leq \frac{f''(b) - f''(a)}{4}(b - a).$$

This inequality follows readily if we take $f'(x)$ instead of $f(x)$ in (3.8).



Some Integral Inequalities
Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back


Close

Quit

Page 11 of 12

References

- [1] R.P. AGARWAL AND S.S. DRAGOMIR, An application of Hayashi's inequality for differentiable functions, *Computers Math. Applic.*, **32**(6) (1996), 95–99.
- [2] G.A. ANASTASSIOU AND S.S. DRAGOMIR, On some estimates of the remainder in Taylor's Formula, *J. Math. Anal. Applic.*, **263** (2001), 246–263.
- [3] S.S. DRAGOMIR, New estimation of the remainder in Taylor's formula using Grüss' type inequalities and applications, *Math. Ineq. and Applics.*, **2**(7) (1999), 183–193.
- [4] H. GAUCHMAN, Some integral inequalities involving Taylor's remainder. I, *J. Inequal. Pure and Appl. Math.*, **3**(2) (2002), Article 26. [ONLINE: http://jipam.vu.edu.au/v3n2/068_01.html]
- [5] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/New York, 1993.
- [6] J.E. PEČARIĆ, F. PROSCHAN AND Y.L. TONG, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, 1992.



Some Integral Inequalities Involving Taylor's Remainder. II

Hillel Gauchman

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12