

COMPARISON OF GREEN FUNCTIONS FOR GENERALIZED SCHRÖDINGER OPERATORS ON $C^{1,1}$ -DOMAINS

LOTFI RIAHI

Department of Mathematics,
Faculty of Sciences of Tunis,
Campus Universitaire 1060
Tunis, TUNISIA.

EMail: Lotfi.Riahi@fst.rnu.tn



volume 4, issue 1, article 22,
2003.

*Received 16 September, 2002;
accepted 10 February, 2003.*

Communicated by: A.M. Fink

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

We establish some inequalities on the $\frac{1}{2}\Delta$ -Green function G on bounded $C^{1,1}$ -domain. We use these inequalities to prove the existence of the $(\frac{1}{2}\Delta - \mu)$ -Green function G_μ and its comparability to G , where μ is in some general class of signed Radon measures. Finally we prove that the choice of this class is essentially optimal.


2000 Mathematics Subject Classification: 34B27, 35J10.

Key words: Green function, Schrödinger operator, 3G-Theorem.

Contents

1	Introduction	3
2	Preliminaries and Notations	5
3	Inequalities on the Green Function G	7
4	The Class $\mathcal{K}(\Omega)$	11
5	The Green Function for $\frac{1}{2}\Delta - \mu$	21

References



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 2 of 31

1. Introduction


The first aim of this paper is to prove some inequalities on the Green function G of $\frac{1}{2}\Delta$ on bounded $C^{1,1}$ -domain Ω in \mathbb{R}^n , $n \geq 3$, where Δ is the Laplacian operator. In particular we give an alternative and shorter proof of the $3G$ -Theorem established in [9] using long and sharp discussions. The $3G$ -Theorem includes the usual one proved in [11], [4] and [3], which was very useful to obtain some potential theoretic results. The second is to prove a comparison theorem between the Green function G and the Green function G_μ of the Schrödinger operator $\frac{1}{2}\Delta - \mu$ on Ω , where μ is allowed to be in some class of signed Radon measures. In contrast to [9], there is no restriction on the sign of μ in this work. This comparison theorem is very important in the sense that it enables us to deduce some potential theoretic results for $\frac{1}{2}\Delta - \mu$ which are known to hold for $\frac{1}{2}\Delta$. This is stated at the end of the paper. Moreover our result covers the case of signed Radon measures with bounded Newtonian potentials i.e,

$$\sup_{x \in \Omega} \int_{\Omega} \frac{1}{|x - y|^{n-2}} |\mu|(dy) < +\infty.$$

The Schrödinger operator $\frac{1}{2}\Delta - f$, with f belonging to the Kato class K_n^{loc} which is studied by several authors (see [1], [3], [4], [11]) is just the special case where μ has the density f with respect to the Lebesgue measure. In particular our results cover the ones proved by Zhao [11]. Finally we show that the choice of this class is essentially optimal.

Our paper is organized as follows.

In Section 2, we give some notations and recall some known results. In Section 3, we prove some inequalities on the Green function of $\frac{1}{2}\Delta$ on bounded



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back


Close

Quit

Page 3 of 31

$C^{1,1}$ -domain. A new and a shorter proof of the $3G$ -Theorem established in [9] is given. In Section 4, we introduce a general class of signed Radon measures on Ω denoted by $\mathcal{K}(\Omega)$ that will be considered in this work. We give some examples and we study some properties of this class. In Section 5, we prove a comparison theorem between the Green functions of $\frac{1}{2}\Delta$ and the Schrödinger operator $\frac{1}{2}\Delta - \mu$, where μ is in the class $\mathcal{K}(\Omega)$. We also show that when μ is nonnegative the condition $\mu \in \mathcal{K}(\Omega)$ is necessary for the comparison theorem to hold.

Throughout the paper the letter C will denote a generic positive constant which may vary in value from line to line.



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 4 of 31

2. Preliminaries and Notations

Throughout the paper Ω denotes a bounded $C^{1,1}$ -domain in \mathbb{R}^n , $n \geq 3$. This means that for each $z \in \partial\Omega$ there exists a ball $B(z, R_0)$, $R_0 > 0$ and a coordinate system of \mathbb{R}^n such that in these coordinates,

$$B(z, R_0) \cap \Omega = B(z, R_0) \cap \{(x', x_n)/x' \in \mathbb{R}^{n-1}, x_n > f(x')\},$$

and

$$B(z, R_0) \cap \partial\Omega = B(z, R_0) \cap \{(x', f(x'))/x' \in \mathbb{R}^{n-1}\},$$

where f is a $C^{1,1}$ -function.

Δ denotes the Laplacian operator on \mathbb{R}^n and G its Green function on Ω . For a signed Radon measure μ on Ω , we denote by G_μ the $(\frac{1}{2}\Delta - \mu)$ -Green function on Ω , when it exists.

For $x \in \Omega$ let $d(x) = d(x, \partial\Omega)$, the distance from x to the boundary of Ω . We denote by $d(\Omega)$ the diameter of Ω .

Since Ω is a bounded $C^{1,1}$ -domain, then it has the following geometrical property:

There exists $r_0 > 0$ depending only on Ω such that for any $z \in \partial\Omega$ and $0 < r \leq r_0$ there exist two balls $B_1^z(r)$ and $B_2^z(r)$ of radius r such that $B_1^z(r) \subset \Omega$, $B_2^z(r) \subset \mathbb{R}^n \setminus \bar{\Omega}$ and $\{z\} = \partial B_1^z(r) \cap \partial B_2^z(r)$.

We recall the following interesting estimates on the Green function G which are due to Grüter and Widman [5], Zhao [11] and Hueber [6].

Theorem 2.1. *There exists a constant $C > 0$ depending on the diameter of Ω ,*



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close


Quit

Page 5 of 31

on the curvature of $\partial\Omega$ and on the dimension n such that

$$C^{-1} \min \left(1, \frac{d(x)d(y)}{|x-y|^2} \right) \frac{1}{|x-y|^{n-2}} \leq G(x,y) \\ \leq C \min \left(1, \frac{d(x)d(y)}{|x-y|^2} \right) \frac{1}{|x-y|^{n-2}}.$$

for all $x, y \in \Omega$.



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 6 of 31

3. Inequalities on the Green Function G

In this section we first give a new and a simple proof of the $3G$ -Theorem established in [9]. We also derive other inequalities on the Green function G that will be used in the next sections.

Theorem 3.1 (3G-Theorem). *There exists a constant $C = C(\Omega, n) > 0$ such that for $x, y, z \in \Omega$, we have*

$$\frac{G(x, z)G(z, y)}{G(x, y)} \leq C \left(\frac{d(z)}{d(x)} G(x, z) + \frac{d(z)}{d(y)} G(z, y) \right).$$

Proof. The inequality of the theorem is equivalent to

$$(3.1) \quad \frac{1}{d(z)G(x, y)} \leq C \left(\frac{1}{d(x)G(z, y)} + \frac{1}{d(y)G(x, z)} \right).$$

On the other hand, since for $a > 0, b > 0$,


$$\frac{ab}{a+b} \leq \min(a, b) \leq 2 \frac{ab}{a+b},$$

then

$$\frac{d(x)d(y)}{|x-y|^2 + d(x)d(y)} \leq \min \left(1, \frac{d(x)d(y)}{|x-y|^2} \right) \leq 2 \frac{d(x)d(y)}{|x-y|^2 + d(x)d(y)},$$

and hence, from Theorem 2.1, we obtain

$$C^{-1}N(x, y) \leq G(x, y) \leq CN(x, y),$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 7 of 31

where

$$N(x, y) = \frac{d(x)d(y)}{|x - y|^{n-2}(|x - y|^2 + d(x)d(y))}.$$

Therefore (3.1) is equivalent to

$$(3.2) \quad \begin{aligned} |x - y|^{n-2}(|x - y|^2 + d(x)d(y)) \\ \leq C(|z - y|^{n-2}(|z - y|^2 + d(z)d(y)) \\ + |x - z|^{n-2}(|x - z|^2 + d(x)d(z))). \end{aligned}$$

Then, we shall prove (3.2). By symmetry we may assume that $|x - z| \leq |y - z|$. We have

$$(3.3) \quad \begin{aligned} |x - y|^{n-2} &\leq (|x - z| + |z - y|)^{n-2} \\ &\leq 2^{n-2}|z - y|^{n-2}, \end{aligned}$$

and


$$(3.4) \quad \begin{aligned} |x - y|^2 + d(x)d(y) &\leq (|x - z| + |z - y|)^2 + (|x - z| + d(z))d(y) \\ &\leq 4|z - y|^2 + |z - y|d(y) + d(z)d(y). \end{aligned}$$

If $|z - y| \leq d(z)$, then

$$(3.5) \quad |z - y|d(y) \leq d(z)d(y).$$

If $|z - y| \geq d(z)$, then

$$(3.6) \quad \begin{aligned} |z - y|d(y) &\leq |z - y|(d(z) + |z - y|) \\ &\leq 2|z - y|^2. \end{aligned}$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 8 of 31

From (3.4), (3.5) and (3.6), we obtain

$$(3.7) \quad |x - y|^2 + d(x)d(y) \leq 6(|z - y|^2 + d(y)d(z)).$$

From (3.3) and (3.7), we obtain

$$|x - y|^{n-2}(|x - y|^2 + d(x)d(y)) \leq 2^{n+1}|z - y|^{n-2}(|z - y|^2 + d(z)d(y)).$$

This proves (3.2) with $C = 2^{n+1}$. \square

Lemma 3.2. *There exists a constant $C = C(\Omega, n) > 0$ such that for all $x, y \in \Omega$, we have*


$$\frac{d(y)}{d(x)}G(x, y) \leq \frac{C}{|x - y|^{n-2}}.$$

Proof. By Theorem 2.1, we have

$$\frac{d(y)}{d(x)}G(x, y) \leq \frac{C}{|x - y|^{n-2}} \min \left(\frac{d(y)}{d(x)}, \frac{d(y)^2}{|x - y|^2} \right).$$

Put $t = \frac{d(y)}{d(x)} > 0$. From the inequality $|x - y| \geq |d(y) - d(x)|$, it follows

$$\begin{aligned} \min \left(\frac{d(y)}{d(x)}, \frac{d(y)^2}{|x - y|^2} \right) &\leq \min \left(\frac{d(y)}{d(x)}, \frac{d(y)^2}{|d(y) - d(x)|^2} \right) \\ &= \min \left(t, \frac{t^2}{(t - 1)^2} \right). \end{aligned}$$



Since $\min\left(t, \frac{t^2}{(t-1)^2}\right) \leq 4$, for all $t > 0$, then we obtain

$$\frac{d(y)}{d(x)}G(x, y) \leq \frac{4C}{|x - y|^{n-2}}.$$

By symmetry we also have


$$\frac{d(x)}{d(y)}G(x, y) \leq \frac{4C}{|x - y|^{n-2}}.$$

This ends the proof. □

The usual $3G$ -Theorem proved in [3, 4, 11] is well known under the following form which is a simple consequence of Theorem 3.1 and Lemma 3.2.

Corollary 3.3. *There exists a constant $C = C(\Omega, n) > 0$ such that for $x, y, z \in \Omega$, we have*

$$\frac{G(x, z)G(z, y)}{G(x, y)} \leq C \left(\frac{1}{|x - z|^{n-2}} + \frac{1}{|z - y|^{n-2}} \right).$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 10 of 31

4. The Class $\mathcal{K}(\Omega)$

Definition 4.1. Let μ be a signed Radon measure on Ω . We say that μ is in the class $\mathcal{K}(\Omega)$ if it satisfies

$$\|\mu\| \equiv \sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) |\mu|(dy) < +\infty,$$

where $|\mu|$ is the total variation of μ .

In the following we study some properties of the class $\mathcal{K}(\Omega)$ and to this end we first need to prove the following lemma.

Lemma 4.1. For $x, y \in \Omega$, we have

If $d(x)d(y) \geq |x - y|^2$, then

$$\frac{1}{3}d(y) \leq d(x) \leq 3d(y).$$

If $d(x)d(y) \leq |x - y|^2$, then


$$\max(d(x), d(y)) \leq 2|x - y|.$$

Proof. If $d(x)d(y) \geq |x - y|^2$, then in view of the inequality $|x - y| \geq |d(x) - d(y)|$, we obtain

$$d(x)d(y) \geq |d(x) - d(y)|^2,$$

which implies

$$3d(x)d(y) \geq d(x)^2 + d(y)^2,$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 11 of 31

and then

$$\frac{1}{3}d(y) \leq d(x) \leq 3d(y).$$

If $d(x)d(y) \leq |x - y|^2$, then in view of the inequality $d(x) \geq d(y) - |x - y|$, we obtain

$$d(y)(d(y) - |x - y|) \leq |x - y|^2,$$

which gives

$$\begin{aligned} d(y)^2 &\leq |x - y|^2 + d(y)|x - y| \\ &\leq \left(|x - y| + \frac{1}{2}d(y) \right)^2. \end{aligned}$$

The last inequality yields

$$\frac{1}{2}d(y) \leq |x - y|.$$


Similarly, we have

$$\frac{1}{2}d(x) \leq |x - y|.$$

□

The following proposition provides some interesting examples of measures in the class $\mathcal{K}(\Omega)$.

Proposition 4.2. *For $\alpha \in \mathbb{R}$, the measure $\frac{1}{d(y)^\alpha} dy$ is in the class $K(\Omega)$ if and only if $\alpha < 2$.*



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 12 of 31

Proof. We first assume $\alpha < 2$ and we will prove that

$$\sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^{\alpha}} dy < +\infty.$$

By Theorem 2.1, we have


$$(4.1) \quad \sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^{\alpha}} dy \leq C \sup_{x \in \Omega} \int_{\Omega} \min \left(\frac{1}{d(x)d(y)}, \frac{1}{|x-y|^2} \right) \frac{d(y)^{2-\alpha}}{|x-y|^{n-2}} dy.$$

On the other hand

$$(4.2) \quad \begin{aligned} & \int_{\Omega} \min \left(\frac{1}{d(x)d(y)}, \frac{1}{|x-y|^2} \right) \frac{d(y)^{2-\alpha}}{|x-y|^{n-2}} dy \\ &= \int_{\Omega \cap (d(x)d(y) \geq |x-y|^2)} \dots dy + \int_{\Omega \cap (d(x)d(y) \leq |x-y|^2)} \dots dy \\ &\equiv I_1 + I_2. \end{aligned}$$

We estimate I_1 . From Lemma 4.1, we have

$$\begin{aligned} I_1 &= \int_{\Omega \cap (d(x)d(y) \geq |x-y|^2)} \frac{d(y)^{1-\alpha}}{d(x)|x-y|^{n-2}} dy \\ &\leq C d(x)^{-\alpha} \int_{|x-y| \leq \sqrt{3}d(x)} \frac{1}{|x-y|^{n-2}} dy \end{aligned}$$



Comparison of Green Functions for Generalized Schrödinger Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 13 of 31

$$\begin{aligned}
 &\leq Cd(x)^{-\alpha} \int_0^{\sqrt{3}d(x)} r dr \\
 &\leq Cd(x)^{2-\alpha} \\
 (4.3) \quad &\leq Cd(\Omega)^{2-\alpha}.
 \end{aligned}$$

Now we estimate I_2 . From Lemma 4.1, we have

$$\begin{aligned}
 I_2 &= \int_{\Omega \cap \{d(x)d(y) \leq |x-y|^2\}} \frac{d(y)^{2-\alpha}}{|x-y|^n} dy \\
 &\leq 2^{2-\alpha} \int_{\Omega} \frac{1}{|x-y|^{n-2+\alpha}} dy \\
 &\leq 2^{2-\alpha} w_{n-1} \int_0^{d(\Omega)} r^{1-\alpha} dr \\
 (4.4) \quad &= \frac{2^{2-\alpha} w_{n-1}}{2-\alpha} d(\Omega)^{2-\alpha},
 \end{aligned}$$


where w_{n-1} is the area of the unit sphere S_{n-1} in \mathbb{R}^n .

Combining (4.1), (4.2), (4.3) and (4.4), we obtain

$$\sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^\alpha} dy \leq Cd(\Omega)^{2-\alpha} < +\infty.$$

Now we assume $\alpha \geq 2$ and we will prove that

$$\sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^\alpha} dy = +\infty.$$



Comparison of Green Functions for Generalized Schrödinger Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 14 of 31

We first remark that when $d(x) \leq \frac{\sqrt{5}-1}{2}|x-y|$, we have

$$d(y) \leq d(x) + |x-y| \leq \frac{\sqrt{5}+1}{2}|x-y|$$

and then $d(x)d(y) \leq |x-y|^2$. By Theorem 2.1, we have


$$\begin{aligned} & \sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^\alpha} dy \\ & \geq C^{-1} \sup_{x \in \Omega} \int_{\Omega} \min \left(\frac{1}{d(x)d(y)}, \frac{1}{|x-y|^2} \right) \frac{d(y)^{2-\alpha}}{|x-y|^{n-2}} dy \\ (4.5) \quad & \geq C^{-1} \sup_{x \in \Omega} \int_{\Omega \cap \{d(x) \leq \frac{\sqrt{5}-1}{2}|x-y|\}} \frac{d(y)^{2-\alpha}}{|x-y|^n} dy. \end{aligned}$$

Let $z_0 \in \partial\Omega$ and put x_0 the center of $B_1^{z_0}(r_0)$. This means $B_1^{z_0}(r_0) = B(x_0, r_0) \subset \Omega$. For $x \in]z_0, x_0]$, we have

$$|y-x| \leq |y-z_0| + d(x),$$

and

$$\left\{ y \in D : d(x) \leq \frac{3-\sqrt{5}}{2}|y-z_0| \right\} \subset \left\{ y \in D : d(x) \leq \frac{\sqrt{5}-1}{2}|y-x| \right\}.$$



Comparison of Green Functions for Generalized Schrödinger Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 15 of 31

Hence for $x \in]z_0, x_0]$, we have

$$(4.6) \quad \int_{\Omega \cap (d(x) \leq \frac{\sqrt{5}-1}{2}|x-y|)} \frac{d(y)^{2-\alpha}}{|x-y|^n} dy \geq \int_{\Omega \cap (d(x) \leq \frac{3-\sqrt{5}}{2}|y-z_0|)} \frac{|y-z_0|^{2-\alpha}}{(|y-z_0| + d(x))^n} dy.$$

From (4.5) and (4.6), we obtain

$$(4.7) \quad \begin{aligned} \sup_{x \in \Omega} \int_{\Omega} \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^\alpha} dy &\geq C^{-1} \int_{\Omega} \frac{1}{|y-z_0|^{n+\alpha-2}} dy \\ &\geq C^{-1} \int_{B_1^{z_0}(r_0)} \frac{1}{|y-z_0|^{n+\alpha-2}} dy \\ &= C^{-1} \int_{|y-x_0| < r_0} \frac{1}{|y-z_0|^{n+\alpha-2}} dy \\ &= C^{-1} \int_{|y| < r_0} \frac{1}{|y-\xi|^{n+\alpha-2}} dy, \end{aligned}$$

where $\xi = z_0 - x_0$ with $|\xi| = r_0$.

We take a spherical coordinate system $(r, \theta_1, \dots, \theta_{n-1})$ such that $\xi = (|\xi|, 0, \dots, 0)$. Then, we have

$$(4.8) \quad \begin{aligned} \int_{|y| < r_0} \frac{1}{|y-\xi|^{n+\alpha-2}} dy &= w_{n-2} \int_0^{r_0} r^{n-1} \int_0^\pi \frac{(\sin \theta_1)^{n-2}}{(r^2 + r_0^2 - 2rr_0 \cos \theta_1)^{\frac{n+\alpha-1}{2}}} d\theta_1 dr. \end{aligned}$$



Comparison of Green Functions for Generalized Schrödinger Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 16 of 31

By making the change of variables $t = \tan \frac{\theta_1}{2}$, we obtain

$$\begin{aligned} & \int_0^\pi \frac{(\sin \theta_1)^{n-2}}{(r^2 + r_0^2 - 2rr_0 \cos \theta_1)^{\frac{n+\alpha}{2}-1}} d\theta_1 \\ &= 2^{n-1} \int_0^{+\infty} \frac{t^{n-2}(1+t^2)^{\frac{\alpha-n}{2}}}{((r+r_0)^2 t^2 + (r_0-r)^2)^{\frac{n+\alpha}{2}-1}} dt \\ &= \frac{2^{n-1}(r_0+r)^{1-n}}{(r_0-r)^{\alpha-1}} \int_0^{+\infty} \frac{s^{n-2} \left(1 + \left(\frac{r_0-r}{r_0+r}\right)^2 s^2\right)^{\frac{\alpha-n}{2}}}{(s^2+1)^{\frac{n+\alpha}{2}-1}} ds \\ &\geq \frac{k}{(r_0-r)^{\alpha-1}}, \end{aligned}$$

where $k = k(r_0, \alpha, n) > 0$.


This implies

$$(4.9) \quad \int_0^{r_0} r^{n-1} \int_0^\pi \frac{(\sin \theta_1)^{n-2}}{(r^2 + r_0^2 - 2rr_0 \cos \theta_1)^{\frac{n+\alpha}{2}-1}} d\theta_1 dr \geq k \int_0^{r_0} \frac{r^{n-1}}{(r_0-r)^{\alpha-1}} dr = +\infty.$$

From (4.7), (4.8) and (4.9), we obtain

$$\sup_{x \in \Omega} \int_\Omega \frac{d(y)}{d(x)} G(x, y) \frac{1}{d(y)^\alpha} dy = +\infty.$$

This ends the proof. □



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 17 of 31

Now we compare the class $\mathcal{K}(\Omega)$ with the class of signed Radon measures with bounded Newtonian potentials. A signed Radon measure μ is said to be of bounded Newtonian potential if $\sup_{x \in \Omega} \int_{\Omega} \frac{1}{|x-y|^{n-2}} |\mu|(dy) < +\infty$.

Proposition 4.3. *The class $\mathcal{K}(\Omega)$ properly contains the class of signed Radon measures with bounded Newtonian potentials.*

Proof. From definitions and Lemma 3.2, it is clear that the class of signed Radon measures with bounded Newtonian potentials is contained in $\mathcal{K}(\Omega)$. In the sequel we will prove that for $1 \leq \alpha$,

$$\int_{\Omega} \frac{1}{d(y)^{\alpha}} dy = +\infty$$

and then

$$\sup_{x \in \Omega} \int_{\Omega} \frac{1}{|x-y|^{n-2}} \frac{1}{d(y)^{\alpha}} dy = +\infty.$$


In particular for $1 \leq \alpha < 2$, $\frac{1}{d(y)^{\alpha}} dy$ does not define a bounded Newtonian potential and by Proposition 4.2, we know that $\frac{1}{d(y)^{\alpha}} dy \in \mathcal{K}(\Omega)$.

Without loss of generality we assume that $0 \in \partial\Omega$. We know that there exists $R_0 > 0$ such that

$$B(0, R_0) \cap \Omega = B(0, R_0) \cap \{(x', x_n) / x' \in \mathbb{R}^{n-1}, x_n > f(x')\},$$

and

$$B(0, R_0) \cap \partial\Omega = B(0, R_0) \cap \{(x', f(x')) / x' \in \mathbb{R}^{n-1}\},$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 18 of 31

where f is a $C^{1,1}$ -function.

By the continuity of f , there exists $\rho_0 \in]0, \frac{R_0}{4}[$ such that for $|y'| < \rho_0$, we have $|f(y')| < \frac{R_0}{2}$.

Hence for all $y = (y', y_n)$ such that $|y'| < \rho_0$ and $0 < y_n - f(y') < \frac{R_0}{4}$, we have $(y', f(y')) \in \partial\Omega$ and $y \in B(0, R_0) \cap \Omega$ which give $d(y) \leq y_n - f(y')$.

Using these observations we have

$$\begin{aligned} \int_{\Omega} \frac{1}{d(y)^\alpha} dy &\geq \int_{\Omega \cap B(0, R_0)} \frac{1}{d(y)^\alpha} dy \\ &\geq \int_{|y'| < \rho_0} \int_{0 < y_n - f(y') < \frac{R_0}{4}} \frac{1}{(y_n - f(y'))^\alpha} dy_n dy' \\ &= \int_{|y'| < \rho_0} dy' \int_0^{\frac{R_0}{4}} \frac{1}{r^\alpha} dr = +\infty. \end{aligned}$$


□

We next prove that the Kato class K_n^{loc} is properly contained in $\mathcal{K}(\Omega)$. For the reader's convenience we recall the definition of the Kato class K_n^{loc} .

Definition 4.2. A Borel measurable function f on Ω is in the Kato class K_n^{loc} if it satisfies

$$\limsup_{r \rightarrow 0} \int_{x \in \Omega} \int_{(|x-y| < r) \cap \Omega} \frac{|f(y)|}{|x-y|^{n-2}} dy = 0.$$

Proposition 4.4. The class $\mathcal{K}(\Omega)$ properly contains the Kato class K_n^{loc} .



Comparison of Green Functions for Generalized Schrödinger Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 19 of 31

Proof. Let f be in K_n^{loc} . We have

$$\limsup_{r \rightarrow 0} \sup_{x \in \Omega} \int_{(|x-y| < r) \cap \Omega} \frac{|f(y)|}{|x-y|^{n-2}} dy = 0.$$

Then, there exists $r > 0$ such that

$$(4.10) \quad \sup_{x \in \Omega} \int_{(|x-y| < r) \cap \Omega} \frac{|f(y)|}{|x-y|^{n-2}} dy \leq 1.$$

This yields

$$\sup_{x \in \Omega} \int_{(|x-y| < r) \cap \Omega} |f(y)| dy \leq r^{n-2}.$$

On the other hand, since $\bar{\Omega}$ is a compact subset then there are $x_1, \dots, x_p \in \Omega$, $p \in \mathbb{N}^*$ such that $\bar{\Omega} = \cup_{i=1}^p B(x_i, r) \cap \bar{\Omega}$. Hence the last inequality gives

$$\int_{\Omega} |f(y)| dy \leq pr^{n-2}.$$


It follows that

$$(4.11) \quad \sup_{x \in \Omega} \int_{(|x-y| \geq r) \cap \Omega} \frac{|f(y)|}{|x-y|^{n-2}} dy \leq p.$$

From (4.10) and (4.11) we obtain

$$\sup_{x \in \Omega} \int_{\Omega} \frac{|f(y)|}{|x-y|^{n-2}} dy \leq p + 1 < +\infty.$$

This means that $f(y)dy$ defines a bounded Newtonian potential and the result holds from Proposition 4.3. \square



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 20 of 31

5. The Green Function for $\frac{1}{2}\Delta - \mu$

In this section we prove that when $\mu \in \mathcal{K}(\Omega)$ the Green function G_μ of the Schrödinger operator $\frac{1}{2}\Delta - \mu$ exists and it is comparable to G . We first prove the following result.

Theorem 5.1. *There exists a constant $C = C(\Omega, n) > 0$ such that for all $\mu \in \mathcal{K}(\Omega)$ and all nonnegative superharmonic function h on Ω , we have*

$$\int_{\Omega} G(x, y)h(y)|\mu|(dy) \leq C\|\mu\|h(x),$$

for all $x \in \Omega$.

Proof. By the 3G-Theorem, we have

$$(5.1) \quad \int_{\Omega} G(x, y)G(y, z)|\mu|(dy) \leq 2C\|\mu\|G(x, z),$$

for all $x, z \in \Omega$.


Now let h be a nonnegative superharmonic function on Ω ; there is an increasing sequence $(h_n)_n$ of nonnegative measurable functions on Ω such that

$$h(x) = \sup_n \int_{\Omega} G(x, z)h_n(z)dz,$$

for all $x \in \Omega$.

From (5.1), we have

$$\int_{\Omega} \int_{\Omega} G(x, y)G(y, z)|\mu|(dy)h_n(z)dz \leq 2C\|\mu\| \int_{\Omega} G(x, z)h_n(z)dz,$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 21 of 31

for all $x \in \Omega$.

By the Fubini's theorem, we obtain

$$\int_{\Omega} G(x, y) \int_{\Omega} G(y, z) h_n(z) dz |\mu|(dy) \leq 2C \|\mu\| \int_{\Omega} G(x, z) h_n(z) dz,$$

for all $x \in \Omega$.

When n tends to $+\infty$, we obtain

$$\int_{\Omega} G(x, y) h(y) |\mu|(dy) \leq 2C \|\mu\| h(x),$$

for all $x \in \Omega$. □

Corollary 5.2. *Let $\mu \in \mathcal{K}(\Omega)$. Then*


$$\sup_{x \in \Omega} \int_{\Omega} G(x, y) |\mu|(dy) < +\infty.$$

Let μ be a signed Radon measure in the class $\mathcal{K}(\Omega)$, i.e. $\|\mu\| < +\infty$. The Jordan decomposition into positive and negative parts says that $\mu = \mu^+ - \mu^-$ and $|\mu| = \mu^+ + \mu^-$. From Corollary 5.2, the functions

$$x \rightarrow \int_{\Omega} G(x, y) \mu^+(dy) \text{ and } x \rightarrow \int_{\Omega} G(x, y) \mu^-(dy)$$

are two continuous potentials on Ω , and the real continuous function

$$x \rightarrow \int_{\Omega} G(x, y) \mu(dy)$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 22 of 31

corresponds to the difference of these two potentials. Hence from the perturbed theory studied in [2], it follows that there exists a Green function G_μ for the Schrödinger operator $\frac{1}{2}\Delta - \mu$ on Ω satisfying the resolvent equation:

$$G(x, y) = G_\mu(x, y) + \int_{\Omega} G(x, z)G_\mu(z, y)d\mu(z),$$

for all $x, y \in \Omega$.

Our main result is the following.

Theorem 5.3. *Assume that $\mu \in \mathcal{K}(\Omega)$ with $\|\mu\|$ sufficiently small. Then the Green functions G and G_μ are comparable, i.e. there is a constant $C = C(\Omega, n, \|\mu\|) > 0$ such that*

$$C^{-1}G \leq G_\mu \leq CG.$$

Proof. We have the resolvent equation:


$$\begin{aligned} G(x, y) &= G_\mu(x, y) + \int_{\Omega} G(x, z)G_\mu(z, y)d\mu(z) \\ &\equiv G_\mu(x, y) + G * G_\mu(x, y). \end{aligned}$$

Then

$$G_\mu = G - G * G_\mu.$$

By iteration we obtain

$$(5.2) \quad G_\mu = G + \sum_{m \geq 1} (-1)^m G^{*m+1},$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 23 of 31

where

$$G^{*2}(x, y) \equiv G * G(x, y) = \int_{\Omega} G(x, z)G(z, y)d\mu(z),$$

and

$$G^{*m+1} = G^{*m} * G.$$

From the 3G-Theorem, we have

$$\begin{aligned} \frac{1}{G(x, y)} \int_{\Omega} G(x, z)G(z, y)|\mu|(dz) \\ \leq C \left(\int_{\Omega} \frac{d(z)}{d(x)}G(x, z)|\mu|(dz) + \int_{\Omega} \frac{d(z)}{d(y)}G(z, y)|\mu|(dz) \right) \\ \leq 2C\|\mu\|. \end{aligned}$$

In particular, we have


$$|G^{*2}| \leq 2C\|\mu\|G.$$

By recurrence, we obtain

$$(5.3) \quad |G^{*m+1}| \leq (2C\|\mu\|)^m G.$$

When $\|\mu\|$ is sufficiently small so that $2C\|\mu\| < \frac{1}{2}$, we obtain, from (5.2) and (5.3),

$$\begin{aligned} |G_{\mu} - G| &\leq \sum_{m \geq 1} (2C\|\mu\|)^m G \\ &= \frac{2C\|\mu\|}{1 - 2C\|\mu\|} G, \end{aligned}$$



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 24 of 31

which yields

$$\left(\frac{1 - 4C\|\mu\|}{1 - 2C\|\mu\|} \right) G \leq G_\mu \leq \frac{1}{1 - 2C\|\mu\|} G.$$

□

Recall that when μ is a nonnegative Radon measure, we know by [8] that the Green function G_μ of $\frac{1}{2}\Delta - \mu$ exists and satisfies the resolvent equation:

$$G(x, y) = G_\mu(x, y) + \int_{\Omega} G(x, z)G_\mu(z, y)d\mu(z),$$

for all $x, y \in \Omega$.

Next we show that in this case, the condition $\mu \in \mathcal{K}(\Omega)$ is necessary and sufficient for the comparability result.

Lemma 5.4. *There exists a constant $C = C(\Omega, n) > 0$ such that*

$$C^{-1}d(x) \leq \int_{\Omega} G(x, y)dy \leq Cd(x),$$

for all $x \in \Omega$.


Proof. From Theorem 2.1, we have

$$G(x, y) \leq \frac{C}{|x - y|^{n-2}} \min \left(1, \frac{d(x)d(y)}{|x - y|^2} \right),$$

for all $x, y \in \Omega$.

If $d(y) \leq 2|x - y|$, then

$$(5.4) \quad G(x, y) \leq 2C \frac{d(x)}{|x - y|^{n-1}}.$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 25 of 31

If $d(y) \geq 2|x - y|$, then $d(x) \geq d(y) - |x - y| \geq |x - y|$, which implies

$$(5.5) \quad G(x, y) \leq \frac{C}{|x - y|^{n-2}} \leq C \frac{d(x)}{|x - y|^{n-1}}.$$

Combining (5.4) and (5.5), we obtain

$$G(x, y) \leq 2C \frac{d(x)}{|x - y|^{n-1}},$$

for all $x, y \in \Omega$.


This yields

$$\begin{aligned} \int_{\Omega} G(x, y) dy &\leq 2Cd(x) \int_{\Omega} \frac{1}{|x - y|^{n-1}} dy \\ &\leq 2Cd(x) \int_{0 \leq |x-y| \leq d(\Omega)} \frac{1}{|x - y|^{n-1}} dy \\ &= 2Cw_{n-1}d(x) \int_0^{d(\Omega)} dr \\ &= C_1d(x). \end{aligned}$$

From Theorem 2.1, we also have

$$\frac{C^{-1}}{|x - y|^{n-2}} \min \left(1, \frac{d(x)d(y)}{|x - y|^2} \right) \leq G(x, y),$$

for all $x, y \in \Omega$.



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 26 of 31

This implies

$$C^{-1} \frac{d(x)d(y)}{d(\Omega)^n} \leq G(x, y),$$

for all $x, y \in \Omega$.

Hence

$$C^{-1} d(\Omega)^{-n} d(x) \int_{\Omega} d(y) dy \leq \int_{\Omega} G(x, y) dy,$$

which means

$$C_2 d(x) \leq \int_{\Omega} G(x, y) dy,$$

for all $x \in \Omega$. □

Theorem 5.5. *Let μ be a nonnegative Radon measure. Then, the Green function G_{μ} of $\frac{1}{2}\Delta - \mu$ on Ω is comparable to G if and only if $\mu \in \mathcal{K}(\Omega)$.*

Proof. We have the integral equation:

$$G(x, y) = G_{\mu}(x, y) + \int_{\Omega} G(x, z)G_{\mu}(z, y)d\mu(z),$$


for all $x, y \in \Omega$.

We first assume that G_{μ} and G are comparable which means that there exists a constant $C \geq 1$ such that

$$C^{-1}G \leq G_{\mu} \leq G.$$

Hence

$$\int_{\Omega} G(x, z)G(z, y)d\mu(z) \leq (C - 1)G(x, y),$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 27 of 31

for all $x, y \in \Omega$.

This implies

$$\int_{\Omega} \int_{\Omega} G(x, z)G(z, y)d\mu(z)dy \leq (C - 1) \int_{\Omega} G(x, y)dy.$$

for all $x \in \Omega$.

Using the Fubini's theorem, it follows that

$$\int_{\Omega} G(x, z) \int_{\Omega} G(z, y)dyd\mu(z) \leq (C - 1) \int_{\Omega} G(x, y)dy.$$

for all $x \in \Omega$.

From Lemma 5.4, we deduce that

$$\int_{\Omega} d(z)G(x, z)d\mu(z) \leq C'd(x),$$

for all $x \in \Omega$.


This means that

$$\sup_{x \in \Omega} \int_{\Omega} \frac{d(z)}{d(x)} G(x, z)d\mu(z) \leq C',$$

and then $\mu \in \mathcal{K}(\Omega)$.

Now let $\mu \in \mathcal{K}(\Omega)$, which means $\|\mu\| < +\infty$. By Theorem 5.3 the Green function $G_{\frac{\mu}{8C\|\mu\|}}$ of the Schrödinger operator $\Delta - \frac{\mu}{8C\|\mu\|}$ is comparable to G . This means that there exists $C > 1$ such that

$$C^{-1}G \leq G_{\frac{\mu}{8C\|\mu\|}} \leq G.$$



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 28 of 31

By Theorem 1 in [10], it follows that

$$C^{-8C\|\mu\|}G \leq G_\mu \leq G,$$


which ends the proof. \square

Remark 5.1. *In view of the paper [7], our results hold also when we replace the Laplace operator by an elliptic operator*

$$L = \sum_{i,j=1}^n a_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^n b_i \frac{\partial}{\partial x_i}$$

which is uniformly elliptic with bounded Hölder continuous coefficients $a_{i,j}$, b_i .

Remark 5.2. *The comparison theorem serves as a main tool to obtain some potential-theoretic results. For example it implies the equivalence of $(\frac{1}{2}\Delta - \mu)$ -potential and $\frac{1}{2}\Delta$ -potential of any measure with support contained in Ω and then the equivalence of $(\frac{1}{2}\Delta - \mu)$ -capacity and $\frac{1}{2}\Delta$ -capacity of any set in Ω . These equivalences say that the fine topology, polar sets, etc. are the same for $\frac{1}{2}\Delta$ and $\frac{1}{2}\Delta - \mu$. Following the argument in [7], the comparison theorem also implies the equivalence of $(\frac{1}{2}\Delta - \mu)$ -harmonic measure and $\frac{1}{2}\Delta$ -harmonic measure on $\partial\Omega$. This gives rise to a boundary Harnack principle and a comparison theorem for nonnegative $(\frac{1}{2}\Delta - \mu)$ -solutions and nonnegative $\frac{1}{2}\Delta$ -solutions vanishing continuously on a part of $\partial\Omega$ (see [4]).*



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents



Go Back


Close

Quit

Page 29 of 31

References

- [1] A. AIZENMAN AND B. SIMON, Brownian motion and Harnack's inequality for Schrödinger's operators, *Comm. Pure Appl. Math.*, **35** (1982), 209–273.
- [2] A. BOUKRICHA, W. HANSEN AND H. HUEBER, Continuous solutions of the generalized Schrödinger equation and perturbation of harmonic spaces, *Expositiones Math.*, **5** (1987), 97–135.
- [3] K.L. CHUNG AND Z. ZHAO, *From Brownian motion to Schrödinger's equation*, Springer Verlag, New York, 1995.
- [4] M. CRANSTON, E.B. FABES AND Z. ZHAO, Conditional gauge and potential theory for the Schrödinger operator, *Trans. Amer. Math. Soc.*, **307**(1) (1988), 171–194.
- [5] M. GRÜTER AND K.O. WIDMAN, The Green function for uniformly elliptic equations, *Manuscripta Math.*, **37** (1982), 303–342.
- [6] H. HUEBER, A uniform estimate for the Green functions on $C^{1,1}$ -domains, *Bibos publication Universität Bielefeld*, (1986).
- [7] H. HUEBER AND M. SIEVEKING, Uniform bounds for quotients of Green functions on $C^{1,1}$ -domains, *Ann. Inst. Fourier*, **32**(1) (1982) 105–117.
- [8] H. MAAGLI AND M. SELMI, Perturbation des résolventes et des semi-groupes par une mesure de Radon positive, *Mathematische Zeitschrift*, **205** (1990), 379–393.



Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains

Lotfi Riahi

Title Page

Contents




Go Back

Close

Quit

Page 30 of 31

- [9] M. SELMI, Comparaison des noyaux de Green sur les domaines $C^{1,1}$, *Revue Roumaine de Mathématiques Pures et Appliquées*, **36** (1991), 91–100.
- [10] M. SELMI, Critere de comparaison de certains noyaux de Green, *Lecture Notes in Math.*, **1235** (1987), 172–193.
- [11] Z. ZHAO, Green function for Schrödinger operator and condioned Feynmann Kac Gauge, *Jour. Math. Anal. Appl.*, **116** (1986), 309–334.



**Comparison of Green Functions
for Generalized Schrödinger
Operators on $C^{1,1}$ -Domains**

Lotfi Riahi

Title Page

Contents



Go Back

Close

Quit

Page 31 of 31