

Journal of Inequalities in Pure and Applied Mathematics

RATE OF CONVERGENCE OF SUMMATION-INTEGRAL TYPE OPERATORS WITH DERIVATIVES OF BOUNDED VARIATION

VIJAY GUPTA, VIPIN VASISHTHA AND M. K. GUPTA

School of Applied Sciences,
Netaji Subhas Institute of Technology,
Sector 3 Dwarka, New Delhi-110045,
India.

E-Mail: vijay@nsit.ac.in

Department of Mathematics,
Hindu College,
Moradabad -244001, India.

Department of Mathematics,
Ch. Charan Singh University,
Meerut-255004,
India.

E-Mail: mkgupta2002@hotmail.com

©2000 Victoria University
ISSN (electronic): 1443-5756
013-03



volume 4, issue 2, article 34,
2003.

*Received 28 January, 2003;
accepted 29 March, 2003.*

Communicated by: A. Fiorenza

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In the present paper, we estimate the rate of convergence of the recently introduced generalized sequence of linear positive operators $G_{n,c}(f, x)$ with derivatives of bounded variation.

2000 Mathematics Subject Classification: 41A25, 41A30.

Key words: Linear positive operators, Bounded variation, Total variation, Rate of convergence.

Contents

1	Introduction	3
2	Auxiliary Results	8
3	Main Result	10



Rate of Convergence of Summation-Integral Type Operators with Derivatives Of Bounded Variation

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 2 of 18

1. Introduction

Let $DB_\gamma(0, \infty)$, $(\gamma \geq 0)$ be the class of all locally integrable functions defined on $(0, \infty)$, satisfying the growth condition $|f(t)| \leq Mt^\gamma$, $M > 0$ and $f' \in BV$ on every finite subinterval of $[0, \infty)$. Then for a function $f \in DB_\gamma(0, \infty)$ we consider the generalized family of linear positive operators which includes some well known operators as special cases. The generalized sequence of operators is defined by


$$(1.1) \quad G_{n,c}(f, x) = n \sum_{k=1}^{\infty} p_{n,k}(x; c) \int_0^{\infty} p_{n+c, k-1}(t; c) f(t) dt + p_{n,0}(x; c) f(0), \quad x \in [0, \infty)$$

where $p_{n,k}(x; c) = (-1)^k \frac{x^k}{k!} \phi_{n,c}^{(k)}(x)$,

- (i) $\phi_{n,c}(x) = e^{-nx}$ for $c = 0$,
- (ii) $\phi_{n,c}(x) = (1 + cx)^{-n/c}$ for $c \in \mathbb{N}$,

and $\{\phi_{n,c}\}_{n \in \mathbb{N}}$ be a sequence of functions defined on an interval $[0, b]$, $b > 0$ having the following properties for every $n \in \mathbb{N}$, $k \in \mathbb{N}_0$:

- (i) $\phi_{n,c} \in C^\infty([a, b])$;
- (ii) $\phi_{n,c}(0) = 1$;
- (iii) $\phi_{n,c}$ is completely monotone $(-1)^k \phi_{n,c}^{(k)}(x) \geq 0$;



Rate of Convergence of Summation-Integral Type Operators with Derivatives Of Bounded Variation

Vijay Gupta, Vipin Vasishtha and M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

(iv) There exists an integer c such that $\phi_{n,c}^{(k+1)} = -n\phi_{n+c,c}^{(k)}$, $n > \max\{0, -c\}$.

Remark 1.1. We may remark here that the functions $\phi_{n,c}$ have various applications in different fields, like potential theory, probability theory, physics and numerical analysis. A collection of most interesting properties of such functions can be found in [10, Ch. 4].

It is easily verified that the operators (1.1) are linear positive operators. Also $G_{n,c}(1, x) = 1$. The generalized new sequence $G_{n,c}$ was recently introduced by Srivastava and Gupta [9].


For $c = 0$ and $\phi_{n,c}(x) = e^{-nx}$ the operators $G_{n,c}$ reduce to the Phillips operators (see e.g. [7], [8]), which are defined by

$$(1.2) \quad G_{n,0}(f, x) = n \sum_{k=1}^{\infty} p_{n,k}(x; 0) \int_0^{\infty} p_{n,k-1}(t; 0) f(t) dt + e^{-nx} f(0), \quad x \in [0, \infty),$$

where $p_{n,k}(x; 0) = \frac{e^{-nx}}{k!} (nx)^k$.

For $c = 1$ and $\phi_{n,c}(x) = (1 + cx)^{-n/c}$ the operators $G_{n,c}$ reduce to the new sequence of summation integral type operators [6], which are defined by

$$(1.3) \quad G_{n,1}(f, x) = n \sum_{k=1}^{\infty} p_{n,k}(x; 1) \int_0^{\infty} p_{n+1,k-1}(t; 1) f(t) dt + (1+x)^{-n} f(0), \quad x \in [0, \infty),$$



Rate of Convergence of Summation-Integral Type Operators with Derivatives Of Bounded Variation

Vijay Gupta, Vipin Vasishtha and M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 4 of 18

where

$$p_{n,k}(x; 1) = \binom{n+k-1}{k} x^k (1+x)^{n-k}.$$


Remark 1.2. It may be noted that for $c = 1$, we get the Baskakov basis functions $p_{n,k}(x; 1)$ which are closely related to the well known Meyer-Konig and Zeller basis functions $m_{n,k}(t) = \binom{n+k-1}{k} t^k (1-t)^n$, $t \in [0, 1]$ because by replacing the variable t with $\frac{x}{1+x}$ in the above MKZ basis functions we get the Baskakov basis functions. Zeng [11] obtained the exact bound for the Meyer Konig Zeller basis functions. Very recently Gupta et al. [6] used the bound of Zeng [11] and estimated the rate of convergence for the operators $G_{n,1}(f, x)$ on functions of bounded variation.

The operators (1.3) are slightly modified form of the operators introduced by Agarwal and Thamer [1], which are defined by

$$(1.4) \quad G_{n,1}^*(f, t) = (n-1) \sum_{k=1}^{\infty} p_{n,k}(x; 1) \int_0^{\infty} p_{n,k-1}(t; 1) f(t) dt + (1+x)^{-n} f(0), \quad x \in [0, \infty),$$

where $p_{n,k}(x; 1)$ is as defined by (1.3) above.

Recently Gupta [5] estimated the rate of approximation for the sequence (1.4) for bounded variation functions. Although the operators defined by (1.3) and (1.4) above are almost the same, but the main advantage to consider the operators in the form (1.3) rather than the form (1.4) is that some approximation properties become simpler in the analysis for the form (1.3) in comparison to the form (1.4). The rate of approximation with derivatives of bounded variation



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 5 of 18

has been studied by several researchers. Bojanic and Cheng ([2], [3]) estimated the rate of convergence with derivatives of bounded variation for Bernstein and Hermite-Fejer polynomials by using different methods.

Alternatively we may rewrite the operators (1.1) as

$$(1.5) \quad G_{n,c}(f, x) = \int_0^\infty K_n(x, t; c) f(t) dt,$$

where

$$K_n(x, t; c) = n \sum_{k=1}^{\infty} p_{n,k}(x; c) p_{n+c, k-1}(t; c) + p_{n,0}(x; c) p_{n,0}(t; c) \delta(t),$$

$\delta(t)$ being the Dirac delta function. Also let


$$(1.6) \quad \beta_n(x, t; c) = \int_0^t K_n(x, s; c) ds$$

then

$$\beta_n(x, \infty; c) = \int_0^\infty K_n(x, s; c) ds = 1.$$

In the present paper we extend the results of [4] and [6] and study the rate of convergence by means of the decomposition technique of functions with derivatives of bounded variation. More precisely the functions having derivatives of bounded variation on every finite subinterval on the interval $[0, \infty)$ be defined as

$$f(x) = f(0) + \int_0^x \psi(t) dt, \quad 0 < a \leq x \leq b,$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close


Quit

Page 6 of 18

where ψ is a function of bounded variation on $[a, b]$ and c is a constant.

We denote the auxiliary function f_x , by

$$f_x(t) = \begin{cases} f(t) - f(x^-), & 0 \leq t < x; \\ 0, & t = x; \\ f(t) - f(x^+), & x < t < \infty. \end{cases}$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 7 of 18

2. Auxiliary Results

In this section we give certain results, which are necessary to prove the main result.

Lemma 2.1. [9]. Let the function $\mu_{n,m}(x)$, $m \in N^0$, be defined as

$$\mu_{n,m}(x; c) = n \sum_{k=1}^{\infty} p_{n,k}(x; c) \int_0^{\infty} p_{n+c,k-1}(t; c) (t-x)^m dt + (-x)^m p_{n,0}(x; c).$$

Then


$$\begin{aligned} \mu_{n,0}(x; c) &= 1, \quad \mu_{n,1}(x; c) = \frac{cx}{(n-c)}, \\ \mu_{n,2}(x; c) &= \frac{x(1+cx)(2n-c) + (1+3cx)cx}{(n-c)(n-2c)}, \end{aligned}$$

and there holds the recurrence relation

$$\begin{aligned} [n - c(m+1)] \mu_{n,m+1}(x; c) \\ = x(1+cx) [\mu_{n,m}^{(1)}(x; c) + 2m\mu_{n,m-1}(x; c)] \\ + [m(1+2cx) + cx] \mu_{n,m}(x; c). \end{aligned}$$

Consequently for each $x \in [0, \infty)$, we have from this recurrence relation that

$$\mu_{n,m}(x; c) = O(n^{-(m+1)/2}).$$



Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 8 of 18

Remark 2.1. In particular, given any number $\lambda > 2$ and $x > 0$ from Lemma 2.1, we have for $c \in N^0$ and n sufficiently large

$$(2.1) \quad G_{n,c}((t-x)^2, x) \equiv \mu_{n,2}(x; c) \leq \frac{\lambda x(1+cx)}{n}.$$

Remark 2.2. It is also noted from (2.1), that

$$(2.2) \quad G_{n,c}(|t-x|, x) \leq (G_{n,c}((t-x)^2, x))^{\frac{1}{2}} \leq \frac{\sqrt{\lambda x(1+cx)}}{\sqrt{n}}.$$

Lemma 2.2. Let $x \in (0, \infty)$ and $K_n(x, t)$ be defined by (1.5). Then for $\lambda > 2$ and for n sufficiently large, we have


$$(i) \quad \beta_n(x, y; c) = \int_0^y K_n(x, t; c) dt \leq \frac{\lambda x(1+cx)}{n(x-y)^2}, \quad 0 \leq y < x,$$

$$(ii) \quad 1 - \beta_n(x, z; c) = \int_z^\infty K_n(x, t; c) dt \leq \frac{\lambda x(1+cx)}{n(z-x)^2}, \quad x < z < \infty.$$

Proof. First, we prove (i). In view of (2.1), we have

$$\begin{aligned} \int_0^y K_n(x, t; c) dt &\leq \int_0^y \frac{(x-t)^2}{(x-y)^2} K_n(x, t; c) dt \leq (x-y)^{-2} \mu_{n,2}(x; c) \\ &\leq \frac{\lambda x(1+cx)}{n(x-y)^2}. \end{aligned}$$

The proof of (ii) is similar. ■



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 9 of 18

3. Main Result

In this section we prove the following main theorem.


Theorem 3.1. *Let $f \in DB_\gamma(0, \infty)$, $\gamma > 0$, and $x \in (0, \infty)$. Then for $\lambda > 2$ and for n sufficiently large, we have*

$$\begin{aligned}
 |G_{n,c}(f, x) - f(x)| &\leq \frac{\lambda(1+cx)}{n} \left(\sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{k}}^{x+\frac{x}{k}} ((f')_x) + \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^{x+\frac{x}{\sqrt{n}}} ((f')_x) \right) \\
 &+ \frac{\lambda(1+cx)}{n} (|f(2x) - f(x) - xf'(x^+)| + |f(x)|) \\
 &+ \frac{\sqrt{\lambda x(1+cx)}}{\sqrt{n}} (M2^\gamma O(n^{-\gamma/2}) + |f'(x^+)|) \\
 &+ \frac{1}{2} \frac{\sqrt{\lambda x(1+cx)}}{\sqrt{n}} |f'(x^+) - f'(x^-)| \\
 &+ \frac{cx}{2(n-c)} |f'(x^+) + f'(x^-)|,
 \end{aligned}$$

where $\bigvee_a^b(f_x)$ denotes the total variation of f_x on $[a, b]$.

Proof. We have

$$\begin{aligned}
 G_{n,c}(f, x) - f(x) &= \int_0^\infty K_n(x, t; c) (f(t) - f(x)) dt \\
 &= \int_0^\infty \left(\int_x^t K_n(x, t; c) f'(u) du \right) dt.
 \end{aligned}$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 10 of 18

Using the identity

$$\begin{aligned}
 f'(u) &= \frac{1}{2} [f'(x^+) + f'(x^-)] \\
 &\quad + (f')_x(u) + \frac{1}{2} [f'(x^+) - f'(x^-)] \operatorname{sgn}(u-x) \\
 &\quad + \left[f'(x) - \frac{1}{2} [f'(x^+) + f'(x^-)] \right] \chi_x(u),
 \end{aligned}$$

it is easily verified that


$$\int_0^\infty \left(\int_x^t f'(x) - \frac{1}{2} [f'(x^+) + f'(x^-)] \chi_x(u) du \right) K(x, t; c) dt = 0.$$

Also

$$\begin{aligned}
 \int_0^\infty \left(\int_x^t \frac{1}{2} [f'(x^+) - f'(x^-)] \operatorname{sgn}(u-x) du \right) K_n(x, t; c) dt \\
 = \frac{1}{2} [f'(x^+) - f'(x^-)] G_{n,c}(|t-x|, x)
 \end{aligned}$$

and

$$\begin{aligned}
 \int_0^\infty \left(\int_x^t \frac{1}{2} [f'(x^+) + f'(x^-)] du \right) K(x, t; c) dt \\
 = \frac{1}{2} [f'(x^+) + f'(x^-)] G_{n,c}((t-x), x).
 \end{aligned}$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit


Page 11 of 18

Thus we have

$$\begin{aligned}
 (3.1) \quad & |G_{n,c}(f, x) - f(x)| \\
 & \leq \left| \int_x^\infty \left(\int_x^t (f')_x(u) du \right) K_n(x, t; c) dt \right. \\
 & \quad \left. - \int_0^x \left(\int_x^t (f')_x(u) du \right) K_n(x, t; c) dt \right| \\
 & \quad + \frac{1}{2} |f'(x^+) - f'(x^-)| G_{n,c}(|t-x|, x) \\
 & \quad + \frac{1}{2} |f'(x^+) + f'(x^-)| G_{n,c}((t-x), x) \\
 & = |A_n(f, x; c) + B_n(f, x; c)| \\
 & \quad + \frac{1}{2} |f'(x^+) - f'(x^-)| G_{n,c}(|t-x|, x) \\
 & \quad + \frac{1}{2} |f'(x^+) + f'(x^-)| G_{n,c}((t-x), x).
 \end{aligned}$$

To complete the proof of the theorem it is sufficient to estimate the terms $A_n(f, x; c)$ and $B_n(f, x; c)$. Applying integration by parts, using Lemma 2.2 and taking $y = x - x/\sqrt{n}$, we have

$$|B_n(f, x; c)| = \left| \int_0^x \left(\int_x^t (f')_x(u) du \right) dt (\beta_n(x, t; c)) \right|,$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 12 of 18


$$\begin{aligned}
& \int_0^x \beta_n(x, t; c) (f')_x(t) dt \\
& \leq \left(\int_0^y + \int_y^x \right) |(f')_x(t)| |\beta_n(x, t; c)| dt \\
& \leq \frac{\lambda x(1+cx)}{n} \int_0^y \bigvee_t^x ((f')_x) \frac{1}{(x-t)^2} dt + \int_y^x \bigvee_t^x ((f')_x) dt \\
& \leq \frac{\lambda x(1+cx)}{n} \int_0^y \bigvee_t^x ((f')_x) \frac{1}{(x-t)^2} dt + \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^x ((f')_x).
\end{aligned}$$

Let $u = x/(x-t)$. Then we have

$$\begin{aligned}
\frac{\lambda x(1+cx)}{n} \int_0^y \bigvee_t^x ((f')_x) \frac{1}{(x-t)^2} dt &= \frac{\lambda x(1+cx)}{n} \int_1^{\sqrt{n}} \bigvee_{x-\frac{x}{u}}^x ((f')_x) du \\
&\leq \frac{\lambda(1+cx)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{u}}^x ((f')_x).
\end{aligned}$$

Thus

$$(3.2) \quad |B_n(f, x; c)| \leq \frac{\lambda(1+cx)}{n} \sum_{k=1}^{[\sqrt{n}]} \bigvee_{x-\frac{x}{u}}^x ((f')_x) + \frac{x}{\sqrt{n}} \bigvee_{x-\frac{x}{\sqrt{n}}}^x ((f')_x).$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back


Close

Quit

Page 13 of 18

On the other hand, we have

$$\begin{aligned}
 (3.3) \quad |A_n(f, x; c)| &= \left| \int_x^\infty \left(\int_x^t (f')_x(u) du \right) K_n(x, t; c) dt \right| \\
 &= \left| \int_{2x}^\infty \left(\int_x^t (f')_x(u) du \right) K_n(x, t; c) dt \right. \\
 &\quad \left. + \int_x^{2x} \left(\int_x^t (f')_x(u) du \right) dt (1 - \beta_n(x, t; c)) \right| \\
 &\leq \left| \int_{2x}^\infty (f(t) - f(x)) K_n(x, t; c) dt \right| \\
 &\quad + |f'(x^+)| \left| \int_{2x}^\infty (t - x) K_n(x, t; c) dt \right| \\
 &\quad + \left| \int_x^{2x} (f')_x(u) du \right| |1 - \beta_n(x, 2x; c)| \\
 &\quad + \int_x^{2x} |(f')_x(t)| |1 - \beta_n(x, t; c)| dt \\
 &\leq \frac{M}{x} \int_{2x}^\infty K_n(x, t; c) t^\gamma |t - x| dt \\
 &\quad + \frac{|f(x)|}{x^2} \int_{2x}^\infty K_n(x, t; c) (t - x)^2 dt \\
 &\quad + |f'(x^+)| \int_{2x}^\infty K_n(x, t; c) |t - x| dt \\
 &\quad + \frac{\lambda(1 + cx)}{nx} |f(2x) - f(x) - xf'(x^+)|
 \end{aligned}$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 14 of 18


$$+ \frac{\lambda(1+cx)}{n} \sum_{k=1}^{\lfloor \sqrt{n} \rfloor} \bigvee_x^{x+\frac{x}{k}} ((f')_x) + \frac{x}{\sqrt{n}} \bigvee_x^{x+\frac{x}{\sqrt{n}}} ((f')_x).$$

Next applying Hölder's inequality, and Lemma 2.1, we proceed as follows for the estimation of the first two terms in the right hand side of (3.3):

$$\begin{aligned} (3.4) \quad & \frac{M}{x} \int_{2x}^{\infty} K_n(x, t; c) t^\gamma |t-x| dt + \frac{|f(x)|}{x^2} \int_{2x}^{\infty} K_n(x, t; c) (t-x)^2 dt \\ & \leq \frac{M}{x} \left(\int_{2x}^{\infty} K_n(x, t; c) t^{2\gamma} dt \right)^{\frac{1}{2}} \left(\int_0^{\infty} K_n(x, t; c) (t-x)^2 dt \right)^{\frac{1}{2}} \\ & \quad + \frac{|f(x)|}{x^2} \int_{2x}^{\infty} K_n(x, t; c) (t-x)^2 dt \\ & \leq M 2^\gamma O(n^{-\gamma/2}) \frac{\sqrt{\lambda x(1+cx)}}{\sqrt{n}} + |f(x)| \frac{\lambda(1+cx)}{nx}. \end{aligned}$$

Also the third term of the right side of (3.3) is estimated as

$$\begin{aligned} & |f'(x^+)| \int_{2x}^{\infty} K_n(x, t; c) |t-x| dt \\ & \leq |f'(x^+)| \int_0^{\infty} K_n(x, t; c) |t-x| dt \\ & \leq |f'(x^+)| \left(\int_0^{\infty} K_n(x, t; c) (t-x)^2 dt \right)^{\frac{1}{2}} \left(\int_0^{\infty} K_n(x, t; c) dt \right)^{\frac{1}{2}} \\ & = |f'(x^+)| \frac{\sqrt{\lambda x(1+cx)}}{\sqrt{n}}. \end{aligned}$$



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 15 of 18

Combining the estimates (3.1) – (3.4), we get the desired result.

This completes the proof of Theorem 3.1. ■


Remark 3.1. For negative values of c , the operators $G_{n,c}$ may be defined in different ways. Here we consider one such example, when $c = -1$ then $\phi_{n,c}(x) = (1-x)^n$, the operator reduces to

$$G_{n,-1}(f, x) = n \sum_{k=1}^n p_{n,k}(x; -1) \int_0^1 p_{n-1,k-1}(t; -1) f(t) dt + (1-x)^n f(0), \quad x \in [0, 1],$$

where

$$p_{n,k}(x; -1) = \binom{n}{k} x^k (1-x)^{n-k}.$$

The rate of convergence for the operators $G_{n-1}(f, x)$ is analogous so we omit the details.



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back


Close

Quit

Page 16 of 18

References

- [1] P.N. AGARWAL AND K.J. THAMAR, Approximation of unbounded function by a new sequence of linear positive operators, *J. Math. Anal. Appl.*, **225** (1998), 660–672.
- [2] R. BOJANIĆ AND F. CHENG, Rate of convergence of Bernstein polynomials for functions with derivatives of bounded variation, *J. Math. Anal Appl.*, **141** (1989), 136–151.
- [3] R. BOJANIĆ AND F. CHENG, Rate of convergence of Hermite Fejer polynomials for functions with derivatives of bounded variation, *Acta Math. Hungar.*, **59** (1992), 91–102.
- [4] V. GUPTA AND G.S. SRIVASTAVA, On the rate of convergence of Phillips operators for functions of bounded variation, *Commentation Math.*, **XXXVI** (1996), 123–130.
- [5] V. GUPTA, Rate of approximation by new sequence of linear positive operators, *Comput. Math. Appl.*, **45**(12) (2003), 1895–1904.
- [6] V. GUPTA, M.K. GUPTA AND V. VASISHTHA, An estimate on the rate of convergence of Bezier type summation integral operators, *Kyungpook Math. J.*, (to appear).
- [7] C.P. MAY, On Phillips operators, *J. Approx. Theory*, **20** (1977), 315–322.
- [8] R.S. PHILLIPS, An inversion formula for semi groups of linear operators, *Ann. Math.*, **59** (1954), 352–356.



Rate of Convergence of Summation-Integral Type Operators with Derivatives Of Bounded Variation

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents




Go Back

Close

Quit

Page 17 of 18

- [9] H.M. SRIVASTAVA AND V. GUPTA, A certain family of summation integral type operators, *Math. Comput. Modelling*, **37** (2003), 1307–1315.
- [10] D.V. WIDDER, *The Laplace Transform*, Princeton University Press, Princeton, 1941.
- [11] X.M. ZENG, Bounds for Bernstein basis functions and Meyer Konig and Zeller basis functions, *J. Math. Anal. Appl.*, **219** (1998), 364–376.



**Rate of Convergence of
Summation-Integral Type
Operators with Derivatives Of
Bounded Variation**

Vijay Gupta, Vipin Vasishtha and
M.K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18