

# Journal of Inequalities in Pure and Applied Mathematics

## REVERSE INEQUALITIES ON CHAOTICALLY GEOMETRIC MEAN VIA SPECHT RATIO, II

MASATOSHI FUJII, JADRANKA MIČIĆ, J.E. PEČARIĆ  
AND YUKI SEO


Department of Mathematics  
Osaka Kyoiku University  
Kashiwara, Osaka 582-8582, Japan.  
*E-Mail:* [mfujii@cc.osaka-kyoiku.ac.jp](mailto:mfujii@cc.osaka-kyoiku.ac.jp)

Technical College Zagreb  
University of Zagreb  
Konavoska 2, 10000 Zagreb, Croatia.  
*E-Mail:* [Jadranka.Micic@public.srce.hr](mailto:Jadranka.Micic@public.srce.hr)

Faculty of Textile Technology  
University of Zagreb  
Pierottijeva 6  
10000 Zagreb, Croatia.  
*E-Mail:* [pecaric@mahazu.hazu.hr](mailto:pecaric@mahazu.hazu.hr)

Tennoji Branch, Senior High School  
Osaka Kyoiku University  
Tennoji, Osaka 543-0054, Japan.  
*E-Mail:* [yukis@cc.osaka-kyoiku.ac.jp](mailto:yukis@cc.osaka-kyoiku.ac.jp)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
028-03



---

volume 4, issue 2, article 40,  
2003.

*Received 24 January, 2003;  
accepted 5 March, 2003.*

*Communicated by: S. Saitoh*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract


In 1967, as a converse of the arithmetic-geometric mean inequality, Mond and Shisha gave an estimate of the difference between the arithmetic mean and the geometric one, which we call it the Mond-Shisha difference. As an application of the Mond-Pečarić method, we show some order relations between the power means of positive operators on a Hilbert space. Among others, we show that the upper bound of the difference between the arithmetic mean and the chaotically geometric one of positive operators coincides with the Mond-Shisha difference.

*2000 Mathematics Subject Classification:* Primary 47A30, 47A63.

*Key words:* Operator concavity, Power mean, Arithmetic mean, Geometric mean.

## Contents

1	Introduction .....	3
2	Preliminary on the Mond-Pečarić Method .....	6
3	Reverse Inequality on Operator Convexity .....	9
4	Comparison Between Arithmetic and Chaotically Geometric Means .....	12
	References	



### Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

# 1. Introduction

In 1960, as a converse of the arithmetic-geometric mean inequality, W. Specht [13] estimated the upper bound of the arithmetic mean by the geometric one for positive numbers: For  $x_1, \dots, x_n \in [m, M]$  with  $0 < m < M$ ,

$$(1.1) \quad \sqrt[n]{x_1 x_2 \cdots x_n} \leq \frac{x_1 + x_2 + \cdots + x_n}{n} \leq M_h(1) \sqrt[n]{x_1 x_2 \cdots x_n},$$

where  $h = \frac{M}{m} (\geq 1)$  is a generalized condition number in the sense of Turing [15] and the Specht ratio  $M_h(1)$  is defined for  $h \geq 1$  as

$$M_h(1) = \frac{(h-1)h^{\frac{1}{h-1}}}{e \log h} \quad (h > 1) \quad \text{and} \quad M_1(1) = 1.$$

On the other hand, Mond and Shisha [11, 12] gave an estimate of the difference between the arithmetic mean and the geometric one: For  $x_1, \dots, x_n \in [m, M]$  with  $0 < m < M$ ,

$$(1.2) \quad 0 \leq \frac{x_1 + x_2 + \cdots + x_n}{n} - \sqrt[n]{x_1 x_2 \cdots x_n} \leq L(m, M) \log M_h(1),$$

where the logarithmic mean  $L(m, M)$  is defined for  $0 < m < M$  as

$$L(m, M) = \frac{M - m}{\log M - \log m} \quad (M \neq m) \quad \text{and} \quad L(m, m) = m.$$

J.I. Fujii and one of the authors [1, 2] showed an operator version of the Mond-Shisha theorem (1.2): Let  $A$  be a positive operator on a Hilbert space  $H$  satisfying  $m \leq A \leq M$  for some scalars  $0 < m < M$ . Then

$$(1.3) \quad (Ax, x) - \exp(\log Ax, x) \leq L(m, M) \log M_h(1)$$



## Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

holds for every unit vector  $x$  in  $H$ . Incidentally, if we put  $A = \text{diag}(x_1, x_2, \dots, x_n)$  and  $x = \frac{1}{\sqrt{n}}(1, 1, \dots, 1)$  in (1.3), then we have (1.2).

Next, we recall the geometric mean in the sense of Kubo-Ando theory [7]: For two positive operators  $A$  and  $B$  on a Hilbert space  $H$ , the geometric mean and arithmetic mean of  $A$  and  $B$  are defined as follows:

$$A \sharp_{\lambda} B = A^{\frac{1}{2}}(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^{\lambda}A^{\frac{1}{2}} \quad \text{and} \quad A \nabla_{\lambda} B = (1 - \lambda)A + \lambda B$$

for  $\lambda \in [0, 1]$ . Like the numerical case, the arithmetic-geometric mean inequality holds:

$$(1.4) \quad A \sharp_{\lambda} B \leq A \nabla_{\lambda} B \quad \text{for all } \lambda \in [0, 1].$$

Tominaga [14] showed the following inequality, as a reverse inequality of the noncommutative arithmetic-geometric mean inequality (1.4) which differs from (1.3): Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$ . Then


$$(1.5) \quad 0 \leq A \nabla_{\lambda} B - A \sharp_{\lambda} B \leq hL(m, M) \log M_h(1) \quad \text{for all } \lambda \in [0, 1],$$

where  $h = \frac{M}{m}$ . It is considered as another operator version of the Mond-Shisha theorem (1.2).

On the other hand, M. Fujii and R. Nakamoto discussed the monotonicity of a family of power means in [4]. For fixed  $A, B > 0$  and  $\lambda \in [0, 1]$ , we put

$$F(r) = (A^r \nabla_{\lambda} B^r)^{\frac{1}{r}} \quad (r \neq 0), \quad = e^{\log A \nabla_{\lambda} \log B} \quad (r = 0).$$

Then the power mean  $F(r)$  is monotone increasing on  $\mathbb{R}$  under the chaotic order  $X \gg Y$ , i.e.,  $\log X \geq \log Y$  for  $X, Y > 0$ , [4, Lemma 2]. In particular,




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit


Page 4 of 20

$A \diamond_{\lambda} B = e^{\log A \nabla_{\lambda} \log B}$  is called the chaotically  $\lambda$ -geometric mean. In general, it does not coincide with  $A \sharp_{\lambda} B$ .

In this note, as a continuation of [3], we consider some order relations between the arithmetic mean and the chaotically geometric one. Among others, we show that if  $A$  and  $B$  are positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $h = \frac{M}{m}$ , then

$$\begin{aligned} -L(m, M) \log M_h(1) &\leq A \nabla_{\lambda} B - A \diamond_{\lambda} B \\ &\leq L(m, M) \log M_h(1) \quad \text{for all } \lambda \in [0, 1]. \end{aligned}$$

Concluding this section, we have to mention that almost all results in this note are based on our previous result [8, Corollary 4] coming from the Mond-Pečarić method [9]. Namely this note might be understood as an application of the Mond-Pečarić method.




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

## 2. Preliminary on the Mond-Pečarić Method

Let  $A$  be a positive operator on a Hilbert space  $H$  satisfying  $m \leq A \leq M$  for some scalars  $0 < m < M$ , and let  $f(t)$  be a real valued continuous convex function on  $[m, M]$ . Mond and Pečarić [9] proved that

$$(2.1) \quad 0 \leq (f(A)x, x) - f((Ax, x)) \leq \beta(m, M, f)$$

holds for every unit vector  $x \in H$ , where

$$(2.2) \quad \beta(m, M, f) = \max \left\{ \frac{f(M) - f(m)}{M - m}(t - m) + f(m) - f(t); t \in [m, M] \right\}.$$


Similarly, we have the following complementary result of (2.1) for a concave function. If  $f(t)$  is concave, then

$$(2.3) \quad \bar{\beta}(m, M, f) \leq (f(A)x, x) - f((Ax, x)) \leq 0$$

holds for every unit vector  $x \in H$ , where

$$(2.4) \quad \bar{\beta}(m, M, f) = \min \left\{ \frac{f(M) - f(m)}{M - m}(t - m) + f(m) - f(t); t \in [m, M] \right\}.$$

The following result is a generalization of (2.1) and based on the idea due to Furuta's work [5, 6]. We cite it here for convenience:



Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II

Masatoshi Fujii, Jadranka Mikić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 6 of 20

**Theorem A ([8]).** Let  $A_j$  ( $j = 1, 2, \dots, k$ ) be positive operators on a Hilbert space  $H$  satisfying  $m \leq A_j \leq M$  for some scalars  $0 < m < M$ . Let  $f(t)$  be a real valued continuous convex function on  $[m, M]$ . Then

$$(2.5) \quad 0 \leq \sum_{j=1}^k (f(A_j)x_j, x_j) - f\left(\sum_{j=1}^k (A_j x_j, x_j)\right) \leq \beta(m, M, f)$$

holds for all  $k$ -tuples  $(x_1, \dots, x_k)$  in  $H$  with  $\sum_{j=1}^k \|x_j\|^2 = 1$ , where  $\beta(m, M, f)$  is defined as in (2.2).


For the power function  $f(t) = t^p$ , we know the following fact, which is a reverse inequality of the Hölder-McCarthy inequality:

**Theorem B.** Let  $A$  be a positive operator on a Hilbert space  $H$  satisfying  $m \leq A \leq M$  for some scalars  $0 < m < M$  and put  $h = \frac{M}{m}$ . For each  $p > 1$

$$(2.6) \quad 0 \leq (A^p x, x) - (Ax, x)^p \leq C(m, M, p)$$

holds for every unit vector  $x \in H$ , where the constant  $C(m, M, p)$  ([8, 16]) is defined as

$$(2.7) \quad C(m, M, p) = \frac{Mm^p - mM^p}{M - m} + (p - 1) \left( \frac{M^p - m^p}{p(M - m)} \right)^{\frac{p}{p-1}} \quad \text{for all } p > 1.$$




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit


Page 7 of 20

We obtain a complement of Theorem B: Under the assumption of Theorem B, for each  $0 < p < 1$

$$(2.8) \quad -\frac{M^p - m^p}{M - m} C\left(m^p, M^p, \frac{1}{p}\right) \leq (A^p x, x) - (Ax, x)^p \leq 0$$

holds for every unit vector  $x \in H$ . It easily can be proved by the fact that

$$\bar{\beta}(m, M, t^p) = -\frac{M^p - m^p}{M - m} C\left(m^p, M^p, \frac{1}{p}\right) \text{ for } 0 < p < 1.$$




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 20

### 3. Reverse Inequality on Operator Convexity

Continuous functions which are convex as real functions need not be operator convex. In this section, we estimate the bounds of the operator convexity for convex functions.

**Lemma 3.1.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$ . If  $f(t)$  is a real valued continuous convex function on  $[m, M]$ , then for each  $\lambda \in [0, 1]$*

$$(3.1) \quad -\beta(m, M, f) \leq f(A)\nabla_\lambda f(B) - f(A\nabla_\lambda B) \leq \beta(m, M, f),$$

where  $\beta(m, M, f)$  is defined as (2.2).

*Proof.* For each  $0 < \lambda < 1$  and unit vector  $x \in H$ , put  $A_1 = A$ ,  $A_2 = B$ ,  $x_1 = \sqrt{1-\lambda}x$  and  $x_2 = \sqrt{\lambda}x$  in Theorem A. Then we have


$$(1-\lambda)(f(A)x, x) + \lambda(f(B)x, x) \leq f((1-\lambda)(Ax, x) + \lambda(Bx, x)) + \beta(m, M, f).$$

Hence it follows that

$$\begin{aligned} (((1-\lambda)f(A) + \lambda f(B))x, x) &\leq f(((1-\lambda)A + \lambda B)x, x) + \beta(m, M, f) \\ &\leq (f((1-\lambda)A + \lambda B)x, x) + \beta(m, M, f) \end{aligned}$$

where the last inequality holds by the convexity of  $f(t)$  [9, Theorem 1] or (2.1). Therefore we have

$$f(A)\nabla_\lambda f(B) \leq f(A\nabla_\lambda B) + \beta(m, M, f).$$



Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 9 of 20

Next, since  $f(t)$  is convex, it follows that

$$\begin{aligned} (1 - \lambda)(f(A)x, x) + \lambda(f(B)x, x) &\geq (1 - \lambda)f((Ax, x)) + \lambda f((Bx, x)) \\ &\geq f((1 - \lambda)(Ax, x) + \lambda(Bx, x)). \end{aligned}$$

Since  $0 < m \leq (1 - \lambda)A + \lambda B \leq M$ , it follows from (2.1) that

$$\begin{aligned} f((1 - \lambda)(Ax, x) + \lambda(Bx, x)) &= f((A \nabla_{\lambda} B)x, x) \\ &\geq (f(A \nabla_{\lambda} B)x, x) - \beta(m, M, f) \end{aligned}$$

holds for every unit vector  $x \in H$ . Therefore we have

$$-\beta(m, M, f) + f(A \nabla_{\lambda} B) \leq f(A) \nabla_{\lambda} f(B).$$

□


We have the following complementary result of Lemma 3.1 for concave functions.

**Lemma 3.2.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$ . If  $f(t)$  is a real valued continuous concave function on  $[m, M]$ , then for each  $\lambda \in [0, 1]$*

$$(3.2) \quad -\bar{\beta}(m, M, f) \geq f(A) \nabla_{\lambda} f(B) - f(A \nabla_{\lambda} B) \geq \bar{\beta}(m, M, f),$$

where  $\bar{\beta}(m, M, f)$  is defined as (2.4).

Next, consider the functions  $f(t) = t^r$  on  $[0, \infty)$ . Then  $f(t)$  is operator concave if  $0 \leq r \leq 1$ , operator convex if  $1 \leq r \leq 2$ , and  $f(t)$  is not operator convex but it is convex if  $r > 2$ . By Lemmas 3.1 and 3.2, we obtain the following reverse inequalities on operator convexity and operator concavity for  $f(t) = t^r$ .



### Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

**Corollary 3.3.** Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $\lambda \in [0, 1]$ .

(i) If  $0 < r \leq 1$ , then

$$-\frac{M^r - m^r}{M - m} C\left(m^r, M^r, \frac{1}{r}\right) \leq A^r \nabla_\lambda B^r - (A \nabla_\lambda B)^r \leq 0.$$

(ii) If  $1 \leq r \leq 2$ , then

$$0 \leq A^r \nabla_\lambda B^r - (A \nabla_\lambda B)^r \leq C(m, M, r).$$

(iii) If  $r > 2$ , then


$$-C(m, M, r) \leq A^r \nabla_\lambda B^r - (A \nabla_\lambda B)^r \leq C(m, M, r),$$

where  $C(m, M, r)$  is defined as (2.7).

*Proof.* Put  $f(t) = t^r$  for  $r > 1$  in Lemma 3.1, then we obtain  $\beta(m, M, f) = C(m, M, r)$ . Also, in the case of  $0 < r \leq 1$ , we have

$$\bar{\beta}(m, M, f) = -\frac{M^r - m^r}{M - m} C\left(m^r, m^r, \frac{1}{r}\right)$$

in Lemma 3.2. □



**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 11 of 20

## 4. Comparison Between Arithmetic and Chaotically Geometric Means

Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  and  $\lambda \in [0, 1]$ . The operator function  $F(r) = (A^r \nabla_\lambda B^r)^{1/r}$  ( $r \in \mathbb{R}$ ) is monotone increasing on  $[1, \infty)$  and not monotone increasing on  $(0, 1]$  under the usual order. Recently, Nakamoto and one of the authors [4] investigated some properties of the chaotically geometric mean  $A \diamond_\lambda B = e^{\log A \nabla_\lambda \log B}$  and showed that the operator function  $F(r)$  is monotone increasing on  $\mathbb{R}$  under the chaotic order and  $F(r)$  converges to  $A \diamond_\lambda B$  as  $r \rightarrow +0$  in the strong operator topology.

In this section, we shall consider some order relations among the chaotically geometric mean, the arithmetic one and the power mean  $F(r)$  by using the results in the previous section. The obtained inequality

$$-L(m, M) \log M_h(1) \leq A \nabla_\lambda B - A \diamond_\lambda B \leq L(m, M) \log M_h(1)$$

is understood as a variant of a reverse Young inequality

$$0 \leq A \nabla_\lambda B - A \sharp_\lambda B \leq hL(m, M) \log M_h(1)$$

due to Tominaga [14], where  $h = \frac{M}{m}$ .

Firstly, by virtue of Corollary 3.3, we see an estimate of the bounds of the difference among the family  $\{F(r) : r > 0\}$ . Incidentally the constant  $C(m, M, r)$  is defined as (2.7).

**Theorem 4.1.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $\lambda \in [0, 1]$ .*



---

Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II

Masatoshi Fujii, Jadranka Micić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 20

(i) If  $0 < r \leq 1 \leq s$ , then

$$\begin{aligned} -C\left(m^r, M^r, \frac{1}{r}\right) &\leq F(s) - F(r) \\ &\leq C\left(m^r, M^r, \frac{1}{r}\right) + \frac{M - m}{M^s - m^s} C(m, M, s). \end{aligned}$$

(ii) If  $0 < 1 \leq r \leq s$ , then

$$0 \leq F(s) - F(r) \leq \frac{M - m}{M^s - m^s} C(m, M, s).$$

(iii) If  $0 < r \leq s \leq 1$ , then


$$|F(s) - F(r)| \leq C\left(m^r, M^r, \frac{1}{r}\right) + C\left(m^s, M^s, \frac{1}{s}\right).$$

*Proof.* Suppose that  $0 < r \leq 1$  or  $1 \leq \frac{1}{r}$ . By (iii) of Corollary 3.3, it follows that

$$-C\left(m, M, \frac{1}{r}\right) \leq A^{\frac{1}{r}} \nabla_{\lambda} B^{\frac{1}{r}} - (A \nabla_{\lambda} B)^{\frac{1}{r}} \leq C\left(m, M, \frac{1}{r}\right).$$

We apply it to  $m^r \leq A^r, B^r \leq M^r$  instead of  $m \leq A, B \leq M$ . That is,

$$(4.1) \quad -C\left(m^r, M^r, \frac{1}{r}\right) \leq A \nabla_{\lambda} B - (A^r \nabla_{\lambda} B^r)^{\frac{1}{r}} \leq C\left(m^r, M^r, \frac{1}{r}\right).$$




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

If  $s \geq 1$ , then  $\frac{1}{s} \leq 1$  and by (i) of Corollary 3.3

$$-\frac{M^{1/s} - m^{1/s}}{M - m} C(m^{1/s}, M^{1/s}, s) \leq A^{\frac{1}{s}} \nabla_{\lambda} B^{\frac{1}{s}} - (A \nabla_{\lambda} B)^{\frac{1}{s}} \leq 0.$$

Since  $m^s \leq A^s, B^s \leq M^s$ , we have also

$$(4.2) \quad -\frac{M - m}{M^s - m^s} C(m, M, s) \leq A \nabla_{\lambda} B - (A^s \nabla_{\lambda} B^s)^{\frac{1}{s}} \leq 0.$$


By using (4.1) and (4.2), it follows that

$$\begin{aligned} -C\left(m^r, M^r, \frac{1}{r}\right) &\leq A \nabla_{\lambda} B - (A^r \nabla_{\lambda} B^r)^{1/r} && \text{by (4.1)} \\ &\leq (A^s \nabla_{\lambda} B^s)^{1/s} - (A^r \nabla_{\lambda} B^r)^{1/r} && \text{by (4.2)} \\ &\leq A \nabla_{\lambda} B + \frac{M - m}{M^s - m^s} C(m, M, s) \\ &\quad - A \nabla_{\lambda} B + C\left(m^r, M^r, \frac{1}{r}\right) && \text{by (4.1) and (4.2)} \\ &= \frac{M - m}{M^s - m^s} C(m, M, s) + C\left(m^r, M^r, \frac{1}{r}\right), \end{aligned}$$

and hence we have (i) in the case of  $0 < r \leq 1 \leq s$ .

In the case of  $0 < 1 \leq r \leq s$ , we have  $1/s \leq 1/r \leq 1$  and by (4.2)

$$-\frac{M - m}{M^s - m^s} C(m, M, s) \leq A \nabla_{\lambda} B - (A^s \nabla_{\lambda} B^s)^{1/s}$$



### Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Micić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 14 of 20

and

$$A\nabla_\lambda B - (A^r \nabla_\lambda B^r)^{1/r} \leq 0.$$

Therefore it follows that

$$\begin{aligned} 0 &\leq (A^s \nabla_\lambda B^s)^{1/s} - (A^r \nabla_\lambda B^r)^{1/r} \\ &\leq A\nabla_\lambda B + \frac{M - m}{M^s - m^s} C(m, M, s) - A\nabla_\lambda B \\ &\leq \frac{M - m}{M^s - m^s} C(m, M, s). \end{aligned}$$

In the case of  $0 < r \leq s \leq 1$ , we have  $1 < 1/s \leq 1/r$  and by (4.1)

$$-C\left(m^r, M^r, \frac{1}{r}\right) \leq A\nabla_\lambda B - (A^r \nabla_\lambda B^r)^{\frac{1}{r}} \leq C\left(m^r, M^r, \frac{1}{r}\right)$$


and

$$-C\left(m^s, M^s, \frac{1}{s}\right) \leq A\nabla_\lambda B - (A^s \nabla_\lambda B^s)^{\frac{1}{s}} \leq C\left(m^s, M^s, \frac{1}{s}\right).$$

Therefore it follows that

$$\begin{aligned} -C\left(m^r, M^r, \frac{1}{r}\right) - C\left(m^s, M^s, \frac{1}{s}\right) &\leq (A^s \nabla_\lambda B^s)^{1/s} - (A^r \nabla_\lambda B^r)^{1/r} \\ &\leq C\left(m^r, M^r, \frac{1}{r}\right) + C\left(m^s, M^s, \frac{1}{s}\right). \end{aligned}$$

□



**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

Though the operator function  $F(r)$  converges to  $A \diamond_{\lambda} B$  as  $r \rightarrow 0$  in the strong operator topology,  $F(s)$  is not generally monotone increasing on  $(0, 1]$  under the usual order. Thus, we have the following estimation of the difference between  $F(r)$  and  $A \diamond_{\lambda} B$ .

**Theorem 4.2.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $\lambda \in [0, 1]$ . Put  $h = \frac{M}{m}$ .*

(i) *If  $0 < s < 1$ , then*


$$\begin{aligned} -C \left( m^s, M^s, \frac{1}{s} \right) - L(m, M) \log M_h(1) \\ \leq F(s) - A \diamond_{\lambda} B \\ \leq C \left( m^s, M^s, \frac{1}{s} \right) + L(m, M) \log M_h(1). \end{aligned}$$

(ii) *If  $1 < s$ , then*

$$\begin{aligned} -L(m, M) \log M_h(1) \leq F(s) - A \diamond_{\lambda} B \\ \leq \frac{M - m}{M^s - m^s} C(m, M, s) + L(m, M) \log M_h(1). \end{aligned}$$

*Proof.* To prove this, we need the following facts on  $C(m, M, r)$  for  $0 < m < M$  and  $r > 1$  by Yamazaki [16]:

(a)  $0 \leq C(m, M, r) \leq M(M^{r-1} - m^{r-1})$  for  $r > 1$



## Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

(b)  $C(m, M, r) \rightarrow 0$  as  $r \rightarrow 1$

(c)  $C(m^r, M^r, \frac{1}{r}) \rightarrow L(m, M) \log M_h(1)$  as  $r \rightarrow +0$ .

In the case of  $0 < r \leq s \leq 1$ , if we put  $r \rightarrow 0$  in (iii) of Theorem 4.1, then  $F(r) \rightarrow A \diamond_{\lambda} B$  and  $C(m^r, M^r, \frac{1}{r}) \rightarrow L(m, M) \log M_h(1)$  as  $r \rightarrow 0$ . Therefore we have (i).

In the case of  $0 < r \leq 1 \leq s$ , if we put  $r \rightarrow 0$  in (i) of Theorem 4.1, then we have (ii).  $\square$

As a result, we obtain an operator version of the Mond-Shisha theorem (1.2):

**Theorem 4.3.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $h = \frac{M}{m}$ . Then*

$$-L(m, M) \log M_h(1) \leq A \nabla_{\lambda} B - A \diamond_{\lambda} B \leq L(m, M) \log M_h(1)$$

hold for all  $\lambda \in [0, 1]$ .


*Proof.* Since  $C(m, M, s) \rightarrow 0$  as  $s \rightarrow 1$ , we have the conclusion by (ii) of Theorem 4.2.  $\square$

By combining Theorem 4.3 and a reverse Young inequality (1.4), we obtain an estimate of the difference between the geometric mean and the chaotically geometric one:

**Corollary 4.4.** *Let  $A$  and  $B$  be positive operators on a Hilbert space  $H$  satisfying  $m \leq A, B \leq M$  for some scalars  $0 < m < M$  and  $h = \frac{M}{m}$ . Then*

$$-(1+h)L(m, M) \log M_h(1) \leq A \sharp_{\lambda} B - A \diamond_{\lambda} B \leq L(m, M) \log M_h(1)$$

holds for all  $\lambda \in [0, 1]$ .



---

Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20


*Proof.* Since  $A\sharp_{\lambda}B \leq A\nabla_{\lambda}B$ , it follows from Theorem 4.3 that

$$A\sharp_{\lambda}B - A\diamond_{\lambda}B \leq A\nabla_{\lambda}B - A\diamond_{\lambda}B \leq L(m, M) \log M_h(1).$$

By Theorem 4.3 and a reverse Young inequality (1.4), it follows that

$$\begin{aligned} -L(m, M) \log M_h(1) &\leq A\nabla_{\lambda}B - A\diamond_{\lambda}B \\ &\leq A\sharp_{\lambda}B + hL(m, M) \log M_h(1) - A\diamond_{\lambda}B. \end{aligned}$$

□




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back


Close

Quit

Page 18 of 20

## References

- [1] J.I. FUJII AND Y. SEO, Determinant for positive operators, *Sci. Math.*, **1**(2) (1998), 153–156.
- [2] J.I. FUJII AND Y. SEO, Characterizations of chaotic order associated with the Mond-Shisha difference, *Math. Inequal. Appl.*, **5** (2002), 725–734.
- [3] M. FUJII, S.H. LEE, Y. SEO AND D. JUNG, Reverse inequalities on chaotically geometric mean via Specht ratio, *Math. Inequal. Appl.*, to appear.
- [4] M. FUJII AND R. NAKAMOTO, A geometric mean in the Furuta inequality, *Sci. Math. Japon.*, **55**(3) (2002), 615–621.
- [5] T. FURUTA, Extensions of Hölder-McCarthy and Kantorovich inequalities and their applications, *Proc. Japan Acad.*, Ser. A, **73** (1997), 38–41.
- [6] T. FURUTA, Operator inequalities associated with Hölder-McCarthy and Kantorovich inequalities, *J. Inequal. Appl.*, **2** (1998), 137–148.
- [7] F. KUBO AND T. ANDO, Means of positive linear operators, *Math. Ann.*, **246**(1980), 205–224.
- [8] J. MIČIĆ, Y. SEO, S.-E. TAKAHASHI AND M. TOMINAGA, Inequalities of Furuta and Mond-Pečarić, *Math. Inequal. Appl.*, **2** (1999), 83–111.
- [9] B. MOND AND J.E. PEČARIĆ, Convex inequalities in Hilbert spaces, *Houston J. Math.*, **19** (1993), 405–420.



---

### Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents




Go Back

Close

Quit

Page 19 of 20

- [10] B. MOND AND J.E. PEČARIĆ, Convex inequalities for several positive operators in Hilbert space, *Indian J. Math.*, **35** (1993), 121–135.
- [11] B. MOND AND O. SHISHA, Difference and ratio inequalities in Hilbert space, *Inequalities II*, (O.Shisha, Ed.), Academic Press, New York, 1970, 241–249.
- [12] O. SHISHA AND B. MOND, Bounds on difference of means, *Inequalities*, (O.Shisha, Ed.). Academic Press, New York, 1967, 293–308.
- [13] W. SPECHT, Zur Theorie der elementaren Mittel, *Math. Z.*, **74** (1960), 91–98.
- [14] M. TOMINAGA, Specht's ratio in the Young inequality, *Sci. Math. Japon.*, **55** (2002), 585–588.
- [15] A.M. TURING, Rounding off-errors in matrix processes, *Quart. J. Mech. Appl. Math.*, **1** (1948), 287–308.
- [16] T. YAMAZAKI, An extension of Specht's theorem via Kantorovich inequality and related results, *Math. Ineq. Appl.*, **3** (2000), 89–96.




---

**Reverse Inequalities on  
Chaotically Geometric Mean via  
Specht Ratio, II**

Masatoshi Fujii, Jadranka Mičić,  
J. Pečarić and Yuki Seo

---

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20