


PREDICTOR-CORRECTOR METHODS FOR GENERALIZED GENERAL MULTIVALUED MIXED QUASI VARIATIONAL INEQUALITIES

CHAO FENG SHI, SAN YANG LIU AND LIAN JUN LI

Department of Applied Mathematics,
Xidian University,
XI'AN, 710071, P.R. CHINA
E-Mail: Shichf@163.com



volume 4, issue 2, article 38,
2003.

*Received 04 April, 2003;
accepted 01 May, 2003.*

Communicated by: *S.S. Dragomir*

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this paper, a class of generalized general mixed quasi variational inequalities is introduced and studied. We prove the existence of the solution of the auxiliary problem for the generalized general mixed quasi variational inequalities, suggest a predictor-corrector method for solving the generalized general mixed quasi variational inequalities by using the auxiliary principle technique. If the bi-function involving the mixed quasi variational inequalities is skew-symmetric, then it is shown that the convergence of the new method requires the partially relaxed strong monotonicity property of the operator, which is a weak condition than cocoercivity. Our results can be viewed as an important extension of the previously known results for variational inequalities.

2000 Mathematics Subject Classification: 11N05, 11N37

Key words: Variational inequalities, Auxiliary principle, Predictor-corrector method, Convergence.

Contents

1	Introduction	3
2	Preliminaries	5
3	Main Results	9



Predictor-Corrector Methods for Generalized General Multivalued Mixed Quasi Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 2 of 17

1. Introduction

In recent years, variational inequalities have been generalized and extended in many different directions using novel and innovative techniques to study wider classes of unrelated problems in mechanics, physics, optimization and control, nonlinear programming, economics, regional, structural, transportation, elasticity, and applied sciences, etc., see [1] – [8] and the references therein. An important and useful generalization of variational inequalities is called the general mixed quasi variational inequality involving the nonlinear bifunction. It is well-known that due to the presence of the nonlinear bifunction, projection method and its variant forms including the Wiener-Hopf equations, descent methods cannot be extended to suggest iterative methods for solving the general mixed quasi variational inequalities. In particular, it has been shown that if the nonlinear bifunction is proper, convex and lower semicontinuous with respect to the first argument, then the general mixed quasi variational inequalities are equivalent to the fixed-point problems. This equivalence has been used to suggest and analyze some iterative methods for solving the general mixed quasi variational inequalities. In this approach, one has to evaluate the resolvent of the operator, which is itself a difficult problem. To overcome these difficulties, Glowinski et al. [6] suggested another technique, which is called the auxiliary principle technique. Recently, Noor [1] extended the auxiliary principle technique to suggest and analyze a new predictor-corrector method for solving general mixed quasi variational inequalities. However, the main results in [1, Algorithm 3.1, Lemma 3.1 and Theorem 3.1] are wrong. Also, Algorithm 3.1 in [1] is based on the assumption that auxiliary problem has a solution, but the author did not show the existence of the solution for this auxiliary problem. On the other hand, in 1999,



Predictor-Corrector Methods for Generalized General Multivalued Mixed Quasi Variational Inequalities

Chao Feng Shi, San Yang Liu and Lian Jun Li

Title Page

Contents




Go Back

Close

Quit

Page 3 of 17

Huang et al. [7] modified and extended the auxiliary principle technique to study the existence of a solution for a class of generalized set-valued strongly nonlinear implicit variational inequalities and suggested some general iterative algorithms. Inspired and motivated by recent research going on in this fascinating and interesting field, in this paper, a class of generalized general mixed quasi variational inequalities is introduced and studied, which includes the general mixed quasi variational inequality as a special case. We prove the existence of the solution of the auxiliary problem for the generalized general mixed quasi variational inequalities, and suggest a predictor-corrector method for solving the generalized general mixed quasi variational inequalities by using the auxiliary principle technique. If the bi-function involving the mixed quasi variational inequalities is skew-symmetric, then it is shown that the convergence of the new method requires the partially relaxed strong monotonicity property of the operator, which is a weaker condition than cocoercivity. Our results extend, improve and modify the main results of Noor [1].



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 4 of 17

2. Preliminaries

Let H be a real Hilbert space whose inner product and norm are denoted by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$, respectively. Let $CB(H)$ be the family of all nonempty closed and bounded sets in H . Let K be a nonempty closed convex set in H . Let $\varphi(\cdot, \cdot) : H \times H \rightarrow H$ be a nondifferentiable nonlinear bifunction. For given nonlinear operators $N(\cdot, \cdot) : H \times H \rightarrow H, g : H \rightarrow H$ and two set-valued operators $T, V : H \rightarrow CB(H)$, consider the problem of finding $u \in H, w \in T(u), y \in V(u)$ such that

$$(2.1) \quad \langle N(w, y), g(v) - g(u) \rangle + \varphi(g(v), g(u)) - \varphi(g(u), g(u)) \geq 0, \forall g(v) \in H.$$

The inequality of type (2.1) is called the generalized general multivalued mixed quasi variational inequality.

For a suitable and appropriate choice of the operators N, g, φ and the space H , one can obtain a wide class of variational inequalities and complementarity problems, see [1]. Furthermore, problem (2.1) has important applications in various branches of pure and applied sciences.


Lemma 2.1. *For all $u, v \in H$, we have*

$$(2.2) \quad 2 \langle u, v \rangle = \|u + v\|^2 - \|u\|^2 - \|v\|^2.$$

Definition 2.1. *For all $u_1, u_2, z \in H, x_1 \in T(u_1), x_2 \in T(u_2)$, an operator $N(\cdot, \cdot)$ is said to be:*

- (i) *g -partially relaxed strongly monotone with respect to the first argument, if there exists a constant $\alpha > 0$ such that*

$$\langle N(x_1, \cdot) - N(x_2, \cdot), g(z) - g(u_2) \rangle \geq -\alpha \|g(u_1) - g(z)\|^2.$$



Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 5 of 17

(ii) g -cocoercive with respect to the first argument, if there exists a constant $\mu > 0$ such that

$$\langle N(x_1, \cdot) - N(x_2, \cdot), g(u_1) - g(u_2) \rangle \geq \mu \|N(x_1, \cdot) - N(x_2, \cdot)\|^2.$$

(iii) T is said to be M -lipschitz continuous, if there exists a constant $\delta > 0$ such that

$$M(T(u_1), T(u_2)) \leq \delta \|u_1 - u_2\|,$$

where $M(\cdot, \cdot)$ is the Hausdorff metric on $CB(H)$.


We remark that if $N(x, \cdot) \equiv Tx$, then Definition 2.1 is exactly the Definition 2.1 of Noor and Memon [1]. If $z = u_1$, $N(x, \cdot) \equiv Tx$, g -partially relaxed strongly monotone with respect to the first argument of $N(\cdot, \cdot)$ is exactly g -monotone of T , and g -cocoercive implies g -partially relaxed strongly monotone [3]. This shows that g -partially relaxed strongly monotone with respect to the first argument of $N(\cdot, \cdot)$ is a weaker condition than g -cocoercive with respect to the first argument of $N(\cdot, \cdot)$.

Definition 2.2. For all $u, v \in H$, the bifunction $\varphi(\cdot, \cdot)$ is said to be skew-symmetric, if

$$\varphi(u, u) - \varphi(u, v) - \varphi(v, u) + \varphi(v, v) \geq 0.$$

Note that if the bifunction $\varphi(\cdot, \cdot)$ is linear in both arguments, then it is nonnegative.

In order to obtain our results, we need the following assumption.



Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 6 of 17

Assumption 1. The mappings $N(\cdot, \cdot) : H \times H \rightarrow H$, $g : H \rightarrow H$ satisfy the following conditions:

- (1) for all $w, y \in H$, there exists a constant $\tau > 0$ such that $\|N(w, y)\| \leq \tau(\|w\| + \|y\|)$;
- (2) for a given $x \in H$, mapping $v \mapsto \langle x, g(v) \rangle$ is convex;
- (3) $\varphi(u, v)$ is bounded, that is, there exists a constant $\gamma > 0$ such that

$$|\varphi(u, v)| \leq \gamma \|u\| \|v\|, \forall u, v \in H;$$


- (4) $\varphi(u, v)$ is linear with respect to u .
- (5) $\varphi(\cdot, \cdot)$ is continuous and $\varphi(g(\cdot), \cdot)$ is convex with respect to the first argument.

Remark 2.1. If $g \equiv I$, it is easy to see that the conditions (2), (5) in Assumption 1 can be easily satisfied.

We also need the following lemma.

Lemma 2.2. [4, 5]. Let X be a nonempty closed convex subset of Hausdorff linear topological space E , $\phi, \psi : X \times X \rightarrow \mathbb{R}$ be mappings satisfying the following conditions:

- (1) $\psi(x, y) \leq \phi(x, y), \forall x, y \in X$;
- (2) for each $x \in X$, $\phi(x, y)$ is upper semicontinuous with respect to y ;
- (3) for each $y \in X$, the set $\{x \in X : \psi(x, y) < 0\}$ is a convex set;



Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents




Go Back

Close

Quit

Page 7 of 17

(4) *there exist a nonempty compact set $K \subset X$ and $x_0 \in K$ such that $\psi(x_0, y) < 0$, for any $y \in X \setminus K$. Then there exists a $\bar{y} \in K$ such that $\phi(x, \bar{y}) \geq 0, \forall x \in X$.*



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 8 of 17

3. Main Results

In this section, we give an existence theorem of a solution of the auxiliary problem for the generalized general set-valued quasi variational inequality (2.1). Based on this existence theorem, we suggest and analyze a new iterative method for solving the problem (2.1).

For given $u \in H$, $w \in Tu$, $y \in Vu$, consider the problem of finding a unique $z \in H$ satisfying the auxiliary general mixed quasi variational inequality

$$(3.1) \quad \langle \rho N(w, y) + g(z) - g(u), g(v) - g(z) \rangle \\ + \rho \varphi(g(v), g(z)) - \rho \varphi(g(z), g(z)) \geq 0,$$


for all $v \in H$, where $\rho > 0$ is a constant.

Remark 3.1. We note that if $z = u$, then clearly z is a solution of (2.1).

Theorem 3.1. If Assumption 1 holds, $g : H \rightarrow H$ is invertible and Lipschitz continuous, and $0 < \rho\gamma < 1$, then $P(u, w, y)$ has a solution.

Proof. Define $\phi, \psi : H \times H \rightarrow H$ by

$$\phi(v, z) = \langle g(v), g(v) - g(z) \rangle \\ - \langle g(u), g(v) - g(z) \rangle + \rho \langle N(w, y), g(v) - g(z) \rangle \\ - \rho \varphi(g(z), g(z)) + \rho \varphi(g(v), g(z))$$



Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 9 of 17

and

$$\begin{aligned} \psi(v, z) = & \langle g(z), g(v) - g(z) \rangle \\ & - \langle g(u), g(v) - g(z) \rangle + \rho \langle N(w, y), g(v) - g(z) \rangle \\ & - \rho\varphi(g(z), g(z)) + \rho\varphi(g(v), g(z)), \end{aligned}$$

respectively. Now we show that the mappings ϕ, ψ satisfy all the conditions of Lemma 2.2.


Clearly, ϕ and ψ satisfy condition (1) of Lemma 2.2. It follows from Assumption 1(5) that $\phi(v, z)$ is upper semicontinuous with respect to z . By using Assumption 1 (2) and (5), it is easy to show that the set $\{v \in H \mid \psi(v, z) < 0\}$ is a convex set for each fixed $z \in H$ and so the conditions (2) and (3) of Lemma 2.2 hold.

Now let

$$\omega = \|g(u)\| + \rho\tau(\|w\| + \|y\|), K = \{z \in H : (1 - \rho\gamma)\|g(z)\| \leq \omega\}.$$

Since $g : H \rightarrow H$ is invertible, K is a weakly compact subset of H . For any fixed $z \in H \setminus K$, take $v_0 \in K$ such that $g(v_0) = 0$. From Assumption 1, we have

$$\begin{aligned} \psi(v_0, z) = & - \langle g(z), g(z) \rangle + \langle g(u), g(z) \rangle + \rho \langle N(w, y), -g(z) \rangle - \rho\varphi(g(z), g(z)) \\ \leq & -\|g(z)\|^2 + \|g(u)\|\|g(z)\| + \rho\tau(\|w\| + \|y\|)\|g(z)\| + \rho\gamma\|g(z)\|^2 \\ = & -\|g(z)\|(\|g(z)\| - \|g(u)\| - \rho\tau(\|w\| + \|y\|) - \rho\gamma\|g(z)\|) \\ < & 0. \end{aligned}$$



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 10 of 17

Therefore, the condition (4) of Lemma 2.2 holds. By Lemma 2.2, there exists a $\bar{z} \in H$ such that $\phi(v, \bar{z}) \geq 0$, for all $v \in H$, that is,

$$(3.2) \quad \langle g(v), g(v) - g(\bar{z}) \rangle - \langle g(u), g(v) - g(\bar{z}) \rangle + \rho \langle N(w, y), g(v) - g(\bar{z}) \rangle \\ - \rho\varphi(g(\bar{z}), g(\bar{z})) + \rho\varphi(g(v), g(\bar{z})) \geq 0, \forall v \in H.$$

For arbitrary $t \in (0, 1)$ and $v \in H$, let $g(x_t) = tg(v) + (1 - t)g(\bar{z})$. Replacing v by x_t in (3.2), we obtain

$$0 \leq \langle g(x_t), g(x_t) - g(\bar{z}) \rangle - \langle g(u), g(x_t) - g(\bar{z}) \rangle + \rho \langle N(w, y), g(x_t) - g(\bar{z}) \rangle \\ - \rho\varphi(g(\bar{z}), g(\bar{z})) + \rho\varphi(g(x_t), g(\bar{z})) \\ = t(\langle g(x_t), g(v) - g(\bar{z}) \rangle - \langle g(u), g(v) - g(\bar{z}) \rangle) + \rho t \langle N(w, y), g(v) - g(\bar{z}) \rangle \\ + \rho t\varphi(g(v) - g(\bar{z}), g(\bar{z})).$$

Hence

$$\langle g(x_t), g(v) - g(\bar{z}) \rangle - \langle g(u), g(v) - g(\bar{z}) \rangle + \rho \langle N(w, y), g(v) - g(\bar{z}) \rangle \\ + \rho\varphi(g(v), g(\bar{z})) - \rho\varphi(g(\bar{z}), g(\bar{z})) \geq 0$$

and so

$$\langle g(x_t), g(v) - g(\bar{z}) \rangle \geq \langle g(u), g(v) - g(\bar{z}) \rangle - \rho \langle N(w, y), g(v) - g(\bar{z}) \rangle \\ - \rho\varphi(g(v), g(\bar{z})) + \rho\varphi(g(\bar{z}), g(\bar{z})).$$

Letting $t \rightarrow 0$, we have

$$\langle g(\bar{z}), g(v) - g(\bar{z}) \rangle \geq \langle g(u), g(v) - g(\bar{z}) \rangle - \rho \langle N(w, y), g(v) - g(\bar{z}) \rangle \\ - \rho\varphi(g(v), g(\bar{z})) + \rho\varphi(g(\bar{z}), g(\bar{z})).$$



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 11 of 17

Therefore, $\bar{z} \in H$ is a solution of the auxiliary problem $P(u, w, y)$. This completes the proof. \square

By using Theorem 3.1, we now suggest the following iterative method for solving the generalized general set-valued quasi variational inequality (2.1).

Algorithm 3.1. For given $u_0 \in H, \xi_0 \in Tu_0, \eta_0 \in Vu_0$, compute the approximate solution u_{n+1} by the iterative scheme

$$x_n \in T(w_n) : \|x_{n+1} - x_n\| \leq M(T(w_{n+1}), T(w_n)),$$

$$y_n \in V(w_n) : \|y_{n+1} - y_n\| \leq M(V(w_{n+1}), V(w_n)),$$

$$(3.3) \quad \langle \rho N(x_n, y_n) + g(u_{n+1}) - g(w_n), g(v) - g(u_{n+1}) \rangle + \rho \varphi(g(v), g(u_{n+1})) - \rho \varphi(g(u_{n+1}), g(u_{n+1})) \geq 0, \quad \forall v \in H,$$

and


$$\xi_n \in T(u_n) : \|\xi_{n+1} - \xi_n\| \leq M(T(u_{n+1}), T(u_n)),$$

$$\eta_n \in V(u_n) : \|\eta_{n+1} - \eta_n\| \leq M(V(u_{n+1}), V(u_n)),$$

$$(3.4) \quad \langle \beta N(\xi_n, \eta_n) + g(w_n) - g(u_n), g(v) - g(w_n) \rangle + \beta \varphi(g(v), g(w_n)) - \beta \varphi(g(w_n), g(w_n)) \geq 0, \quad \forall v \in H,$$

where $\rho > 0, \beta > 0$ are constants.

For the convergence analysis of Algorithm 3.1, we need the following result.



Predictor-Corrector Methods for Generalized General Multivalued Mixed Quasi Variational Inequalities

Chao Feng Shi, San Yang Liu and Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 12 of 17

Lemma 3.2. Let $u \in H$, $x \in Tu$, $y \in Vu$ be the exact solution of (2.1) and u_{n+1} be the approximate solution obtained from Algorithm 3.1. If the operator $N(\cdot, \cdot)$ is g -partially relaxed strongly monotone with respect to the first and second argument with constants $a > 0, b > 0$, respectively, the bifunction $\varphi(\cdot, \cdot)$ is skew-symmetric and the conditions in Theorem 3.1 are satisfied, then

$$(3.5) \quad \|g(u_{n+1}) - g(u)\|^2 \leq \|g(u_n) - g(u)\|^2 - (1 - 2\rho(a + b))\|g(u_{n+1}) - g(u_n)\|^2.$$

Proof. Let $u \in H, x \in Tu, y \in Vu$ be a solution of (2.1). Then


$$(3.6) \quad \langle \rho N(x, y), g(v) - g(u) \rangle + \rho\varphi(g(v), g(u)) - \rho\varphi(g(u), g(u)) \geq 0, \forall v \in H,$$

$$(3.7) \quad \langle \beta N(x, y), g(v) - g(u) \rangle + \beta\varphi(g(v), g(u)) - \beta\varphi(g(u), g(u)) \geq 0, \forall v \in H,$$

where $\rho > 0, \beta > 0$ are constants. Now taking $v = u_{n+1}$ in (3.6) and $v = u$ in (3.3), we have

$$(3.8) \quad \langle \rho N(x, y), g(u_{n+1}) - g(u) \rangle + \rho\varphi(g(u_{n+1}), g(u)) - \rho\varphi(g(u), g(u)) \geq 0,$$

$$(3.9) \quad \langle \rho N(x_n, y_n) + g(u_{n+1}) - g(w_n), g(u) - g(u_{n+1}) \rangle + \rho\varphi(g(u), g(u_{n+1})) - \rho\varphi(g(u_{n+1}), g(u_{n+1})) \geq 0.$$



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 13 of 17

Adding (3.8) and (3.9), we have (3.10)

$$\begin{aligned}
 (3.10) \quad & \langle g(u_{n+1}) - g(w_n), g(u) - g(u_{n+1}) \rangle \\
 & \geq \rho \langle N(x_n, y_n) - N(x, y), g(u_{n+1}) - g(u) \rangle + \rho \{ \varphi(g(u), g(u)) \\
 & \quad - \varphi(g(u), g(u_{n+1})) - \varphi(g(u_{n+1}), g(u)) + \varphi(g(u_{n+1}), g(u_{n+1})) \} \\
 & \geq \rho \langle N(x_n, y_n) - N(x_n, y), g(u_{n+1}) - g(u) \rangle \\
 & \quad + \rho \langle N(x_n, y) - N(x, y), g(u_{n+1}) - g(u) \rangle \\
 & \geq -\rho(a + b) \|g(w_n) - g(u_{n+1})\|^2,
 \end{aligned}$$

where we have used the fact that $N(\cdot, \cdot)$ is g -partially relaxed strongly monotone with respect to the first and second argument with constants $a > 0, b > 0$, respectively, and the bifunction $\varphi(\cdot, \cdot)$ is skew-symmetric. Setting $u = g(u) - g(u_{n+1}), v = g(u_{n+1}) - g(w_n)$ in (2.2), we obtain


$$\begin{aligned}
 (3.11) \quad & \langle g(u_{n+1}) - g(w_n), g(u) - g(u_{n+1}) \rangle \\
 & = \frac{1}{2} \{ \|g(u) - g(w_n)\|^2 - \|g(u_{n+1}) - g(w_n)\|^2 - \|g(u) - g(u_{n+1})\|^2 \}.
 \end{aligned}$$

Combining (3.10) and (3.11), we have

$$\begin{aligned}
 (3.12) \quad & \|g(u_{n+1}) - g(u)\|^2 \\
 & \leq \|g(w_n) - g(u)\|^2 - (1 - 2\rho(a + b)) \|g(u_{n+1}) - g(w_n)\|^2,
 \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 (3.13) \quad & \|g(u) - g(w_n)\|^2 \\
 & \leq \|g(u_n) - g(u)\|^2 - (1 - 2\beta(a + b)) \|g(u_n) - g(w_n)\|^2, \\
 & \leq \|g(u_n) - g(u)\|^2, 0 < \beta < 1/2(a + b).
 \end{aligned}$$



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 14 of 17

and

$$\begin{aligned}
 (3.14) \quad \|g(u_{n+1}) - g(w_n)\|^2 &= \|g(u_{n+1}) - g(u_n) + g(u_n) - g(w_n)\|^2 \\
 &= \|g(u_{n+1}) - g(u_n)\|^2 + \|g(u_n) - g(w_n)\|^2 \\
 &\quad + 2 \langle g(u_{n+1}) - g(u_n), g(u_n) - g(w_n) \rangle .
 \end{aligned}$$

Combining (3.12) – (3.14), we have

$$\|g(u_{n+1}) - g(u)\|^2 \leq \|g(u_n) - g(u)\|^2 - (1 - 2\rho(a + b))\|g(u_{n+1}) - g(u_n)\|^2.$$

The required result. \square


Theorem 3.3. *Let H be finite dimensional, $g : H \rightarrow H$ be invertible, g^{-1} is Lipschitz continuous and $0 < \rho < \frac{1}{2}(a + b)$. Let $\{u_n\}, \{\xi_n\}, \{\eta_n\}$ be the sequences obtained from Algorithm 3.1, $u \in H$ be the exact solution of (2.1) and the conditions in Lemma 3.2 are satisfied, then $\{u_n\}, \{\xi_n\}$, and $\{\eta_n\}$ strongly converge to a solution of (2.1).*

Proof. Let $u \in H$ be a solution of (2.1). Since $0 < \rho < \frac{1}{2}(a + b)$, from (3.5), it follows that the sequence $\{\|g(u) - g(u_n)\|\}$ is nonincreasing and consequently $\{u_n\}$ is bounded. Furthermore, we have

$$\Sigma(1 - 2\rho(a + b))\|g(u_{n+1}) - g(u_n)\|^2 \leq \|g(u_0) - g(u)\|^2,$$

which implies that

$$(3.15) \quad \lim_{n \rightarrow \infty} \|g(u_{n+1}) - g(u_n)\| = 0.$$



**Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities**

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 15 of 17

Let \hat{u} be the cluster point of $\{u_n\}$ and the subsequence $\{u_{n_j}\}$ of the sequence $\{u_n\}$ converge to \hat{u} , which implies $\{u_{n_j}\}$ is a Cauchy sequence in H . By (3.4), we know that both $\{\xi_{n_j}\}$ and $\{\eta_{n_j}\}$ are Cauchy sequences in H . Let $\xi_{n_j} \rightarrow \hat{x}$ and $\eta_{n_j} \rightarrow \hat{y}$. Since

$$d(\hat{x}, T(\hat{u})) \leq \|\hat{x} - \xi_{n_j}\| + M(T(u_{n_j}), T(\hat{u})) \rightarrow 0, \quad n_j \rightarrow \infty.$$

So we can obtain $\hat{x} \in T(\hat{u})$. Similarly, we can obtain $\hat{y} \in V(\hat{u})$. Replacing w_n by u_{n_j} in (3.3) and (3.4), the limit $n_j \rightarrow \infty$ and using (3.14), we have


$$\langle N(\hat{x}, \hat{y}), g(v) - g(\hat{u}) \rangle + \varphi(g(v), g(\hat{u})) - \varphi(g(\hat{u}), g(\hat{u})) \geq 0, \quad \forall v \in H,$$

which implies that $\hat{u} \in H$, $\hat{x} \in T\hat{u}$, $\hat{y} \in V\hat{u}$ is a solution of (2.1), and

$$\|g(u_{n+1} - g(u))\|^2 \leq \|g(u_n) - g(u)\|^2.$$

Thus it follows from the above inequality that the sequence $\{u_n\}$ has exactly one cluster point \hat{u} and $\lim_{n \rightarrow \infty} g(u_n) = g(\hat{u})$. Since g is invertible and g^{-1} is Lipschitz continuous, $\lim_{n \rightarrow \infty} u_n = \hat{u}$. The required result. \square

Remark 3.2. Lemma 3.2 and Theorem 3.3 improve and modify the main results of Noor [1].



Title Page

Contents



Go Back


Close

Quit

Page 16 of 17

References

- [1] M.A. NOOR AND Z.A. MEMON, Algorithms for general mixed quasi variational inequalities, *J. Inequal. Pure and Appl. Math.*, **3**(4) (2002), Art. 59. [ONLINE: http://jipam.vu.edu.au/v3n4/044_02.html]
- [2] M.A. NOOR, Solvability of multivalued general mixed variational inequalities, *J. Math. Anal. Applic.*, **261** (2001), 390–402.
- [3] M.A. NOOR, A class of new iterative methods for general mixed variational inequalities, *Math. Computer Modelling*, **31** (2000), 11–19.
- [4] S.S. CHANG, *Variational Inequality and Complementarity Theory with Application*, Shanghai Sci. Technol., Shanghai, 1991.
- [5] S.S. CHANG AND S.W. XIANG, On the existence of solution for a class of quasi-bilinear variational inequalities, *J. Systems Sci. Math. Sci.*, **16** (1996), 136–140.
- [6] R. GLOWINSKI, J.L. LIONS AND R. TREMOLIERES, *Numerical Analysis of Variational Inequalities*, North -Holland, Amsterdam, 1981.
- [7] N.J. HUANG, Y.P. LIU, Y.Y. TANG AND M.R. BAI, On the generalized set-valued strongly nonlinear implicit variational inequalities, *Comput. Math. Appl.*, **37** (1999), 29–36.
- [8] D. KINDERLEHRER AND G. STAMPACCHIA, *An Introduction to Variational Inequalities and their Applications*, SIAM, Philadelphia, 2000.



Predictor-Corrector Methods for
Generalized General
Multivalued Mixed Quasi
Variational Inequalities

Chao Feng Shi, San Yang Liu and
Lian Jun Li

Title Page

Contents



Go Back

Close

Quit

Page 17 of 17