

# Journal of Inequalities in Pure and Applied Mathematics

## INEQUALITIES RELATED TO REARRANGEMENTS OF POWERS AND SYMMETRIC POLYNOMIALS

CEZAR JOIȚA AND PANTELIMON STĂNICĂ

Department of Mathematics,  
Lehigh University,  
Bethlehem, PA 18015, USA.

E-Mail: [cej3@lehigh.edu](mailto:cej3@lehigh.edu)


URL: <http://www.lehigh.edu/~cej3/cej3.html>

Auburn University Montgomery,  
Department of Mathematics,  
Montgomery,  
AL 36124-4023, USA.

E-Mail: [pstanica@mail.aum.edu](mailto:pstanica@mail.aum.edu)

URL: <http://sciences.aum.edu/~stanica>

©2000 Victoria University  
ISSN (electronic): 1443-5756  
058-03



---

volume 4, issue 2, article 37,  
2003.

*Received 29 April, 2003;  
accepted 19 May, 2003.*

*Communicated by: C. Niculescu*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

In [2] the second author proposed to find a description (or examples) of real-valued  $n$ -variable functions satisfying the following two inequalities:

$$\text{if } x_i \leq y_i, i = 1, \dots, n, \text{ then } F(x_1, \dots, x_n) \leq F(y_1, \dots, y_n),$$

with strict inequality if there is an index  $i$  such that  $x_i < y_i$ ; and for  $0 < x_1 < x_2 < \dots < x_n$ , then,

$$F(x_1^{x_2}, x_2^{x_3}, \dots, x_n^{x_1}) \leq F(x_1^{x_1}, x_2^{x_2}, \dots, x_n^{x_n}).$$

In this short note we extend in a direction a result of [2] and we prove a theorem that provides a large class of examples satisfying the two inequalities, with  $F$  replaced by any symmetric polynomial with positive coefficients. Moreover, we find that the inequalities are not specific to expressions of the form  $x^y$ , rather they hold for any function  $g(x, y)$  that satisfies some conditions. A simple consequence of this result is a theorem of Hardy, Littlewood and Polya [1].


**2000 Mathematics Subject Classification:** 05E05, 11C08, 26D05.

**Key words:** Symmetric Polynomials, Permutations, Inequalities.

Both authors are associated with the Institute of Mathematics of Romanian Academy, Bucharest – Romania.

## Contents

1	Introduction .....	3
2	The Results .....	5



### Inequalities Related to Rearrangements of Powers and Symmetric Polynomials

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back

Close

Quit

Page 2 of 11

# 1. Introduction

In [2], the following problem was proposed: *find examples of functions  $F : \mathbb{R}_+^n \rightarrow \mathbb{R}$  with the properties*

$$(1.1) \quad \text{if } x_i \leq y_i, i = 1, \dots, n, \text{ then } F(x_1, \dots, x_n) \leq F(y_1, \dots, y_n),$$

*with strict inequality if there is an index  $i$  such that  $x_i < y_i$ ,*

and

$$(1.2) \quad \text{for } 0 < x_1 < x_2 < \dots < x_n, \text{ then,}$$

$$F(x_1^{x_2}, x_2^{x_3}, \dots, x_n^{x_1}) \leq F(x_1^{x_1}, x_2^{x_2}, \dots, x_n^{x_n}).$$


In [2], the following result was proved.

**Theorem 1.1.** *Assume that the permutation  $\sigma$  can be written as a product of disjoint circular cycles  $C_1 \times C_2 \times \dots \times C_r$ , where each  $C_i$  is a cyclic permutation, that is  $C_i(j) = j + t_i$ , for some fixed  $t_i$ . For any increasing sequence  $0 < x_1 < \dots < x_n$ , we have*

$$(1.3) \quad \sum_{i=1}^n a_i x_i^{x_{\sigma(i)}} \leq \sum_{i=1}^n a_i x_i^{x_i}, \text{ and}$$

$$\prod_{i=1}^n a_i x_i^{x_{\sigma(i)}} \leq \prod_{i=1}^n a_i x_i^{x_i},$$

where  $a_i \geq 0$  is increasing on the cycles  $C_i$  of  $\sigma$ .



Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back

Close

Quit

Page 3 of 11

(The condition on  $a_i$  was inadvertently omitted in the final version of [2].)

In this short note we extend in a direction the previous result of [2] to any permutation, not only the permutations which are products of circular cycles, by proving (1.1) and (1.2) for symmetric polynomials with positive coefficients. Finally, we prove that these inequalities are not specific only to rearrangements of powers, that is, we find other classes of functions of 2-variables with real values, say  $g(x, y)$ , such that, for any  $\sigma \in S_n$  (the group of permutations), we have

$$(1.4) \quad F(g(x_1, x_{\sigma(1)}), \dots, g(x_n, x_{\sigma(n)})) \leq F(g(x_1, x_1), \dots, g(x_n, x_n)),$$

where  $F$  is any symmetric polynomial with positive coefficients.



---

**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 11

## 2. The Results

**Lemma 2.1.** *If  $f \in \mathbb{R}[X_1, X_2]$  is a symmetric polynomial with positive coefficients and  $(x_1, x_2) \in \mathbb{R}_+^2$  and  $(y_1, y_2) \in \mathbb{R}_+^2$  are such that  $x_1x_2 \leq y_1y_2$  and  $x_1^n + x_2^n \leq y_1^n + y_2^n, \forall n \in \mathbb{N}$ , then  $f(x_1, x_2) \leq f(y_1, y_2)$ .*

*Proof.* We have

$$f(X_1, X_2) = \sum_{i < j} a_{ij} X_1^i X_2^j = \sum_{i < j} (a_{ij} X_1^i X_2^j + a_{ji} X_1^j X_2^i) + \sum_{ii} a_{ii} X_1^i X_2^i,$$

where  $a_{ij} \in \mathbb{R}_+$ . Since  $f$  is symmetric  $a_{ij} = a_{ji}$ , and therefore

$$\begin{aligned} f(X_1, X_2) &= \sum_{i < j} a_{ij} (X_1^i X_2^j + X_1^j X_2^i) + \sum_{ii} a_{ii} X_1^i X_2^i \\ &= \sum_{i < j} a_{ij} X_1^i X_2^i (X_1^{j-i} + X_2^{j-i}) + \sum_{ii} a_{ii} X_1^i X_2^i. \end{aligned}$$

It is clear now that the two conditions imposed on  $(x_1, x_2)$  and  $(y_1, y_2)$  imply that  $f(x_1, x_2) \leq f(y_1, y_2)$ .  $\square$

We will consider  $A \subset \mathbb{R}$  and a function  $g : A \times A \rightarrow [0, \infty)$  with the following property: for all  $x_1, x_2, y_1, y_2 \in \mathbb{R}$  such that  $x_1 \leq x_2$  and  $y_1 \leq y_2$  the following two inequalities are satisfied:

$$(2.1) \quad g(x_1, y_2)g(x_2, y_1) \leq g(x_1, y_1)g(x_2, y_2)$$

$$(2.2) \quad [g(x_1, y_2)]^n + [g(x_2, y_1)]^n \leq [g(x_1, y_1)]^n + [g(x_2, y_2)]^n, \forall n \in \mathbb{N}.$$



**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back

Close

Quit

Page 5 of 11

**Theorem 2.2.** Let  $F(X_1, X_2, \dots, X_n)$  be a symmetric polynomial with positive coefficients and  $g$  as above. Then for any  $\sigma \in S_n$  and any  $x_1, x_2, \dots, x_n \in A$  we have:

$$F(g(x_1, x_{\sigma(1)}), g(x_2, x_{\sigma(2)}), \dots, g(x_n, x_{\sigma(n)})) \leq F(g(x_1, x_1), g(x_2, x_2), \dots, g(x_n, x_n)).$$

*Proof.* Consider  $x_1, x_2, \dots, x_n \in A$  arbitrary and fixed. Without loss of generality we may assume that  $x_1 \leq x_2 \leq \dots \leq x_n$ . Let

$$m = \max\{F(g(x_1, x_{\sigma(1)}), \dots, g(x_n, x_{\sigma(n)})) \mid \sigma \in S_n\}$$


and let

$$P = \{\sigma \in S_n \mid F(g(x_1, x_{\sigma(1)}), \dots, g(x_n, x_{\sigma(n)})) = m\}.$$

We would like to prove that  $e \in P$  where  $e$  is the identity. Let  $\tau \in P$  the permutation that has the minimum number of inversions among all elements of  $P$  and suppose that  $\tau \neq e$ . Since  $e$  is the only increasing permutation it follows that there exists  $i \in \{1, 2, \dots, n-1\}$  such that  $\tau(i) > \tau(i+1)$ . Without loss of generality we may assume that  $i = 1$ . Consider  $\tau' \in S_n$  defined as follows:  $\tau'(1) = \tau(2)$ ,  $\tau'(2) = \tau(1)$  and  $\tau'(j) = \tau(j)$  if  $j \geq 3$ . Then  $\tau'$  has fewer inversions than  $\tau$  and therefore  $\tau' \notin P$ , which implies that:

$$(2.3) \quad F(g(x_1, x_{\tau'(1)}), \dots, g(x_n, x_{\tau'(n)})) < F(g(x_1, x_{\tau(1)}), \dots, g(x_n, x_{\tau(n)})).$$

Consider  $f(X_1, X_2) = F(X_1, X_2, g(x_3, x_{\tau(3)}), \dots, g(x_n, x_{\tau(n)}))$ . It follows that  $f$  is symmetric and has positive coefficients. If we set  $y_1 = x_{\tau'(1)} = x_{\tau(2)}$  and



**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back

Close

Quit

Page 6 of 11

$y_2 = x_{\tau'(2)} = x_{\tau(1)}$  it follows that  $y_1 \leq y_2$ . Using the two properties of  $g$  and Lemma 2.1 we deduce that  $f(g(x_1, y_2), g(x_2, y_1)) \leq f(g(x_1, y_1), g(x_2, y_2))$  and therefore

$$F(g(x_1, x_{\tau(1)}), \dots, g(x_n, x_{\tau(n)})) \leq F(g(x_1, x_{\tau'(1)}), \dots, g(x_n, x_{\tau'(n)})),$$

which contradicts (2.3). □

If  $g(x, y) = x^y$ , then the conditions imposed on  $g$  are

$$\begin{aligned} x_1^{y_2} x_2^{y_1} &\leq x_1^{y_1} x_2^{y_2}, \\ x_1^{ny_2} + x_2^{ny_1} &\leq x_1^{ny_1} + x_2^{ny_2}, \end{aligned}$$


which are equivalent to

$$\begin{aligned} x_1^{y_2-y_1} &\leq x_2^{y_2-y_1}, \\ x_1^{ny_1} (x_1^{n(y_2-y_1)} - 1) &\leq x_2^{ny_1} (x_2^{n(y_2-y_1)} - 1). \end{aligned}$$

The first inequality is certainly true as  $x_1 \leq x_2$  and  $y_1 \leq y_2$ . The second inequality is true if  $1 \leq x_1 \leq x_2$  and  $y_1 \leq y_2$ . Therefore

**Corollary 2.3.** *The inequalities (1.1) and (1.2) are satisfied for all  $n$ -variable symmetric polynomials with positive coefficients, defined on  $[1, \infty)^n$ .*

If  $F(x_1, \dots, x_n) := x_1 + \dots + x_n$ , we can prove a result similar to the one of Theorem 2.2 even if we significantly weaken the assumption on  $g$ .



**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back

Close

Quit

Page 7 of 11

**Theorem 2.4.** Let  $A \subset \mathbb{R}$  and  $g : A \times A \rightarrow \mathbb{R}$  be a function such that  $h_{a,b}(y) = g(a, y) - g(b, y)$  ( $a > b$ ) is increasing. Then for any  $x_1, x_2, \dots, x_n \in A$  and any  $\sigma \in S_n$  we have:

$$F(g(x_1, x_{\sigma(1)}), g(x_2, x_{\sigma(2)}), \dots, g(x_n, x_{\sigma(n)})) \leq F(g(x_1, x_1), g(x_2, x_2), \dots, g(x_n, x_n)).$$

*Proof.* We follow the proof of Theorem 2.2 and the only thing we have to check is that

$$F(g(x_1, x_{\tau(1)}), \dots, g(x_n, x_{\tau(n)})) \leq F(g(x_1, x_{\tau'(1)}), \dots, g(x_n, x_{\tau'(n)})).$$

But this inequality is equivalent to

$$g(x_1, x_{\tau(1)}) + g(x_2, x_{\tau(2)}) \leq g(x_1, x_{\tau'(1)}) + g(x_2, x_{\tau'(2)}).$$


If we set  $y_1 = x_{\tau'(1)} = x_{\tau(2)}$  and  $y_2 = x_{\tau'(2)} = x_{\tau(1)}$ , it follows that  $y_1 \leq y_2$  and the previous inequality can be written as

$$g(x_1, y_2) + g(x_2, y_1) \leq g(x_1, y_1) + g(x_2, y_2),$$

which is equivalent to

$$h_{x_2, x_1}(y_1) \leq h_{x_2, x_1}(y_2).$$

This inequality is satisfied because  $y_1 \leq y_2$  and  $h_{x_2, x_1}$  is increasing. □




---

**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 11

**Corollary 2.5.** Let  $u, v$  be increasing functions on  $\mathbb{R}$  with values in  $[1, \infty)$ . The following inequalities are true for all  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$(2.4) \quad \sum_{i=1}^n u(x_i)v(x_{\sigma(i)}) \leq \sum_{i=1}^n u(x_i)v(x_i),$$

$$(2.5) \quad \sum_{i=1}^n u(x_i)^{v(x_{\sigma(i)})} \leq \sum_{i=1}^n u(x_i)^{v(x_i)},$$

$$(2.6) \quad \prod_{i=1}^n u(x_i)^{v(x_{\sigma(i)})} \leq \prod_{i=1}^n u(x_i)^{v(x_i)}.$$

*Proof.* It suffices to prove that the following functions  $g(x, y) = u(x)v(y)$ ,  $g(x, y) = u(x)^{v(y)}$ , or  $g(x, y) = u(y)^{v(x)}$  have the associated  $h$ 's increasing.

Let  $g(x, y) = u(x)v(y)$ . Then  $h(y) = u(a)v(y) - u(b)v(y) = (u(a) - u(b))v(y)$  which is increasing since  $u(a) \geq u(b)$  and  $v(y)$  is increasing.

Let  $g(x, y) = u(x)^{v(y)}$ . Then  $h(y) = u(a)^{v(y)} - u(b)^{v(y)}$ . Since  $u(a) \geq u(b) \geq 1$ , and  $v(y)$  is increasing, by writing

$$h(y) = u(b)^{v(y)} \left( \left( \frac{u(a)}{u(b)} \right)^{v(y)} - 1 \right),$$

we see that  $h$  is increasing.

We remark that to prove (2.4) we only needed  $u, v$  to have positive values. Using the previous remark, to show the last inequality, apply (2.4) with  $w = \log(u)$  and  $v$  (which are both increasing).  $\square$



**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

Title Page

Contents



Go Back


Close

Quit

Page 9 of 11

**Corollary 2.6.** *If the function  $h$  is decreasing on  $A$ , then all the inequalities are reversed.*

**Remark 2.1.** *We see that Theorem 368 of [1] follows from (2.4) and Corollary 2.6.*



---

**Inequalities Related to  
Rearrangements of Powers and  
Symmetric Polynomials**

Cezar Joița and  
Pantelimon Stănică

---

Title Page

Contents



Go Back


Close

Quit

Page 10 of 11

## References

- [1] G. HARDY, J.E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge Univ. Press, 2001.
- [2] P. STĂNICĂ, *Inequalities on linear functions and circular powers*, *J. Ineq. in Pure and Applied Math.*, **3**(3) (2002), Art.43. [ONLINE: [http://jipam.vu.edu.au/v3n3/015\\_02.html](http://jipam.vu.edu.au/v3n3/015_02.html)]



---

### Inequalities Related to Rearrangements of Powers and Symmetric Polynomials

Cezar Joița and  
Pantelimon Stănică

---

Title Page

Contents



Go Back

Close

Quit

Page 11 of 11