

ON AN INTEGRAL INEQUALITY WITH A KERNEL SINGULAR IN TIME AND SPACE

NASSER-EDDINE TATAR

King Fahd University of Petroleum and Minerals
Department of Mathematical Sciences
Dhahran 31261, Saudi Arabia.
EMail: tatarn@kfupm.edu.sa



volume 4, issue 4, article 82,
2003.

*Received 07 March, 2003;
accepted 01 August, 2003.*

Communicated by: D. Bainov

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

In this paper we deal with a nonlinear singular integral inequality which arises in the study of partial differential equations. The integral term is non local in time and space and the kernel involved is also singular in both the time and the space variable. The estimates we prove may be used to establish (global) existence and asymptotic behavior results for solutions of the corresponding problems.

2000 Mathematics Subject Classification: 42B20, 26D07, 26D15.

Key words: Nonlinear integral inequality, Singular kernel.

The author would like to thank King Fahd University of Petroleum and Minerals for its financial support.

Contents

1	Introduction	3
2	Preliminaries	6
3	Estimation	8
	References	

On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. Introduction

We consider the following integral inequality

$$(1.1) \quad \varphi(t, x) \leq k(t, x) + l(t, x) \int_{\Omega} \int_0^t \frac{F(s)\varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} dy ds, \\ x \in \Omega, t > 0,$$

where Ω is a domain in \mathbb{R}^n ($n \geq 1$) (bounded or possibly equal to \mathbb{R}^n), the functions $k(t, x)$, $l(t, x)$ and $F(t)$ are given positive continuous functions in t . The constants $0 < \alpha < n$, $0 < \beta < 1$ and $m > 1$ will be specified below.

This inequality arises in the theory of partial differential equations, for example, when treating the heat equation with a source of polynomial type

$$\begin{cases} u_t(t, x) = \Delta u(t, x) + u^m(t, x), & x \in \mathbb{R}^n, t > 0, m > 1 \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n. \end{cases}$$


If we write the (weak) solution using the fundamental solution $G(t, x)$ of the heat equation

$$u_t(t, x) = \Delta u(t, x),$$

namely,

$$u(t, x) = \int_{\mathbb{R}^n} G(t, x-y)u_0(y)dy + \int_0^t \int_{\mathbb{R}^n} G(t-s, x-y)u^m(s, y)dy ds$$

and take into account the Solonnikov estimates of this fundamental solution (see [14] for instance), then one is led to an inequality of type (1.1).



On an Integral Inequality with a
Kernel Singular in Time and
Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

The features of this inequality, which make it difficult to deal with, are the singularities of the kernel in both the space and time variables. It is also non integrable with respect to the time variable. The standard methods one can find in the literature (see the recent books by Bainov and Simeonov [1] and Pachpatte [12] and the references therein) concerning regular and/or summable kernels cannot be applied in our situation. Indeed, these methods are based on estimates involving the value of the kernels at zero and/or some L^p -norms of the kernels. In contrast, there are very few papers dealing explicitly with singular kernels similar to ours. Let us point out, however, some works concerning integral equations with singularities in time. In Henry [4, Lemmas 7.1.1 and 7.1.2], a similar inequality to (1.1) with only the integral with respect to time, namely


$$\psi(t) \leq a(t) + b \int_0^t (t-s)^{\beta-1} s^{\gamma-1} \psi(s) ds, \quad \beta > 0, \quad \gamma > 0$$

i.e. the linear case ($m = 1$) has been treated. The case $m > 1$ has been considered by Medved in [9], [10]. More precisely, the following inequality

$$\psi(t) \leq a(t) + b(t) \int_0^t (t-s)^{\beta-1} s^{\gamma-1} F(s) \psi^m(s) ds, \quad \beta > 0, \quad \gamma > 0$$

was discussed. The result was used to prove a global existence and an exponential decay result for a parabolic Cauchy problem with a source of power type and a time dependent coefficient, namely

$$\begin{cases} u_t + Au = f(t, u), & u \in X, \\ u(0) = u_0 \in X \end{cases}$$



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 4 of 20

with $\|f(t, u)\| \leq t^\kappa \eta(t) \|u\|_\alpha^m$, $m > 1$, $\kappa \geq 0$, where A is a sectorial operator (see [4]) and $\|\cdot\|_\alpha$ stands for the norm of the fractional space X^α associated to the operator A (see also [11]). This, in turn, has been improved and extended to integro-differential equations and functional differential equations by M. Kirane and N.-E. Tatar in [6] (see also N.-E. Tatar [13] and S. Mazouzi and N.-E. Tatar [7, 8] for more general abstract semilinear evolution problems).

Here, we shall combine the techniques in [9, 10], based on the application of Lemma 2.1 and the use of Lemma 2.3 below, with the Hardy-Littlewood-Sobolev inequality (see Lemma 2.2) to prove our result. We will give sufficient conditions yielding boundedness by continuous functions, exponential decay and polynomial decay of solutions to the integral inequality (1.1).

The paper is organized as follows. In the next section we prepare some notation and lemmas needed in the proofs of our results. Section 3 contains the statement and proof of our result and two corollaries giving sufficient conditions for the exponential decay and the polynomial decay. Finally we point out that our results hold (*a fortiori*) for *weakly* singular kernels in time.



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

2. Preliminaries

In this section we introduce some material necessary for our results. We will use the usual L^p -space with its norm $\|\cdot\|_p$.

Lemma 2.1. (Young inequality) We have, for $\frac{1}{r} = \frac{1}{\rho} + \frac{1}{q}$, the inequality

$$\|fg\|_r \leq \|f\|_\rho \cdot \|g\|_q.$$

Lemma 2.2. Let $\alpha \in [0, 1)$ and $\beta \in \mathbb{R}$. There exists a positive constant $C = C(\alpha, \beta)$ such that


$$\int_0^t s^{-\alpha} e^{\beta s} ds \leq \begin{cases} Ce^{\beta t}, & \text{if } \beta > 0; \\ C(t+1), & \text{if } \beta = 0; \\ C, & \text{if } \beta < 0. \end{cases}$$

Lemma 2.3. (Hardy-Littlewood-Sobolev inequality)

Let $u \in L^p(\mathbb{R}^n)$ ($p > 1$), $0 < \gamma < n$ and $\frac{\gamma}{n} > 1 - \frac{1}{p}$, then $(1/|x|^\gamma) * u \in L^q(\mathbb{R}^n)$ with $\frac{1}{q} = \frac{\gamma}{n} + \frac{1}{p} - 1$. Also the mapping from $u \in L^p(\mathbb{R}^n)$ into $(1/|x|^\gamma) * u \in L^q(\mathbb{R}^n)$ is continuous.

See [5, Theorem 4.5.3, p. 117].

Lemma 2.4. Let $a(t)$, $b(t)$, $K(t)$, $\psi(t)$ be nonnegative, continuous functions on the interval $I = (0, T)$ ($0 < T \leq \infty$), $\omega : (0, \infty) \rightarrow \mathbb{R}$ be a continuous,



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 6 of 20

nonnegative and nondecreasing function, $\omega(0) = 0$, $\omega(u) > 0$ for $u > 0$ and let $A(t) = \max_{0 \leq s \leq t} a(s)$, $B(t) = \max_{0 \leq s \leq t} b(s)$. Assume that

$$\psi(t) \leq a(t) + b(t) \int_0^t K(s)\omega(\psi(s))ds, \quad t \in I.$$

Then

$$\psi(t) \leq W^{-1} \left[W(A(t)) + B(t) \int_0^t K(s)ds \right], \quad t \in (0, T_1),$$

where

$$W(v) = \int_{v_0}^v \frac{d\sigma}{\omega(\sigma)}, \quad v \geq v_0 > 0,$$

W^{-1} is the inverse of W and $T_1 > 0$ is such that

$$W(A(t)) + B(t) \int_0^t K(s)ds \in D(W^{-1})$$

for all $t \in (0, T_1)$.


See [2] (or [5]) for the proof.

Lemma 2.5. If $\delta, \nu, \tau > 0$ and $z > 0$, then

$$z^{1-\nu} \int_0^z (z - \zeta)^{\nu-1} \zeta^{\delta-1} e^{-\tau\zeta} d\zeta \leq M(\nu, \delta, \tau),$$

where $M(\nu, \delta, \tau) = \max(1, 2^{1-\nu}) \Gamma(\delta) \left(1 + \frac{\delta}{\nu}\right) \tau^{-\delta}$.

See [6] for the proof of this lemma.



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

3. Estimation

In this section we state and prove our result on boundedness and also present an exponential and a polynomial decay result.

Theorem 3.1. *Assume that the constants α , β and m are such that $0 < \alpha < \beta n$, $0 < \beta < 1$ and $m > 1$.*

(i) *If $\Omega = \mathbb{R}^n$, then for any r satisfying $\max\left(\frac{(m-1)n}{\alpha}, \frac{m}{\beta}\right) < r < \frac{mn}{\alpha}$, we have*


$$\|\varphi(t, x)\|_r \leq U_{p,r,\rho}(t)$$

with

$$U_{p,r,\rho}(t) = 2^{\frac{m(p-1)}{r}} K(t)^{\frac{1}{p}} \times \left[1 - 2^{m(p-1)} (m-1) C_1^{p-1} C_2^p K(t)^{m-1} L(t) e^{\varepsilon p t} \times \int_0^t e^{-\varepsilon p s} F^p(s) ds \right]^{\frac{m}{(1-m)r}},$$

where $K(t) = \max_{0 \leq s \leq t} \|k(s, \cdot)\|_r^p$, $L(t) = \max_{0 \leq s \leq t} \|l(s, \cdot)\|_\rho^p$, $p = \frac{r}{m}$ and $\rho = \frac{nr}{\alpha r - (m-1)n}$ for some $\varepsilon > 0$. Here C_1 and C_2 are the best constants in Lemma 2.2 and Lemma 2.3, respectively. The estimation is valid as long as

$$(3.1) \quad K(t)^{m-1} L(t) e^{\varepsilon p t} \int_0^t e^{-\varepsilon p s} F^p(s) ds \leq \frac{1}{2^{m(p-1)}} (m-1) C_1^{p-1} C_2^p.$$



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 8 of 20

(ii) If Ω is bounded, then

$$\|\varphi(t, x)\|_{\tilde{r}} \leq U_{p,r,\rho}(t)$$

for any $\tilde{r} \leq r$ where p, r and ρ are as in (i). If moreover, $r < \frac{n}{n\beta-\alpha}$ (but not necessarily $r > \frac{(m-1)n}{\alpha}$) that is $\frac{m}{\beta} < r < \min\left(\frac{mn}{\alpha}, \frac{n}{n\beta-\alpha}\right)$, then this estimation holds for any $\frac{1}{\beta} < p \leq \frac{r}{m}$ provided that $\rho > \frac{nr}{n-(n\beta-\alpha)r}$.

Proof. (i) Suppose that $\Omega = \mathbb{R}^n$. By the Minkowski inequality and the Young inequality (Lemma 2.1), we have


$$(3.2) \quad \|\varphi(t, x)\|_r \leq \|k(t, x)\|_r + \|l(t, x)\|_\rho \left\| \int_{\mathbb{R}^n} \int_0^t \frac{F(s)\varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} ds dy \right\|_q$$

for r, ρ and q such that $\frac{1}{r} = \frac{1}{\rho} + \frac{1}{q}$.

Let p be such that $\frac{1}{p} = \frac{1}{q} + \frac{\alpha}{n}$ and p' its conjugate i.e. $\frac{1}{p} + \frac{1}{p'} = 1$. Using the Hölder inequality, we see that

$$(3.3) \quad \int_0^t (t-s)^{\beta-1} F(s)\varphi^m(s, y) ds \leq \left(\int_0^t (t-s)^{(\beta-1)p'} e^{\varepsilon p' s} ds \right)^{\frac{1}{p'}} \left(\int_0^t e^{-\varepsilon p s} F^p(s)\varphi^{mp}(s, y) ds \right)^{\frac{1}{p}},$$

for any positive constant ε . We have multiplied by $e^{\varepsilon s} \cdot e^{-\varepsilon s}$ before applying Hölder inequality.



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents




Go Back

Close

Quit

Page 9 of 20



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

Choosing $q = \frac{nr}{(nm-\alpha r)}$, we see that $p = \frac{r}{m}$. By our assumption on r it is easy to see that $q > 1$, $p > 1$ and $1 + (\beta - 1)p' > 0$. Therefore, we may apply Lemma 2.2 to get

$$\int_0^t (t-s)^{(\beta-1)p'} e^{\varepsilon p' s} ds < C_1 e^{\varepsilon p' t}.$$

Hence, inequality (3.3) becomes


$$\int_0^t (t-s)^{\beta-1} F(s) \varphi^m(s, y) ds \leq C_1^{\frac{1}{p'}} e^{\varepsilon t} \left(\int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}}.$$

It follows that

$$(3.4) \quad \left\| \int_{\mathbb{R}^n} \int_0^t \frac{F(s) \varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} ds dy \right\|_q \leq C_1^{\frac{1}{p'}} e^{\varepsilon t} \left\| \int_{\mathbb{R}^n} \left(\int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}} \frac{dy}{|x-y|^{n-\alpha}} \right\|_q.$$

As $r < \frac{mn}{\alpha}$, we may apply the results in Lemma 2.3 to obtain

$$(3.5) \quad \left\| \int_{\mathbb{R}^n} \left(\int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}} \frac{dy}{|x-y|^{n-\alpha}} \right\|_q \leq C_2 \left\| \left(\int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, \cdot) ds \right)^{\frac{1}{p}} \right\|_p,$$



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 11 of 20

with p as above and C_2 is the best constant in the result of Lemma 2.3. From (3.2), (3.4) and (3.5) it appears that

$$\begin{aligned} & \|\varphi(t, x)\|_r \\ & \leq \|k(t, x)\|_r + C_3 e^{\varepsilon t} \|l(t, x)\|_\rho \left\| \left(\int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, \cdot) ds \right)^{\frac{1}{p}} \right\|_p \end{aligned}$$

or


$$(3.6) \quad \begin{aligned} & \|\varphi(t, x)\|_r \\ & \leq \|k(t, x)\|_r + C_3 e^{\varepsilon t} \|l(t, x)\|_\rho \left(\int_{\mathbb{R}^n} \int_0^t e^{-\varepsilon ps} F^p(s) \varphi^{mp}(s, x) ds dx \right)^{\frac{1}{p}}, \end{aligned}$$

where $C_3 = C_1^{\frac{1}{p'}} C_2$. Inequality (3.6) can also be written as

$$\begin{aligned} & \|\varphi(t, x)\|_r \\ & \leq \|k(t, x)\|_r + C_3 e^{\varepsilon t} \|l(t, x)\|_\rho \left(\int_0^t e^{-\varepsilon ps} F^p(s) \|\varphi(s, x)\|_{mp}^{mp} ds \right)^{\frac{1}{p}}. \end{aligned}$$

Observe that by our choice of p we have $r = mp$. It follows that

$$(3.7) \quad \begin{aligned} & \|\varphi(t, x)\|_r \\ & \leq \|k(t, x)\|_r + C_3 e^{\varepsilon t} \|l(t, x)\|_\rho \left(\int_0^t e^{-\varepsilon ps} F^p(s) \|\varphi(s, x)\|_r^r ds \right)^{\frac{1}{p}}. \end{aligned}$$



Applying the algebraic inequality

$$(a + b)^p \leq 2^{p-1}(a^p + b^p), \quad a, b \geq 0, \quad p > 1,$$

we deduce from (3.7) that

$$(3.8) \quad \|\varphi(t, x)\|_r^p \leq 2^{p-1} \|k(t, x)\|_r^p + C_4 e^{\varepsilon pt} \|l(t, x)\|_\rho^p \int_0^t e^{-\varepsilon ps} F^p(s) \|\varphi(s, \cdot)\|_r^r ds,$$

where $C_4 = 2^{p-1} C_3^p$. Let us put $\psi(t) = \|\varphi(t, x)\|_r^{r/m}$, then (3.8) takes the form

$$\psi(t) \leq 2^{p-1} \|k(t, \cdot)\|_r^p + C_4 e^{\varepsilon pt} \|l(t, x)\|_\rho^p \int_0^t e^{-\varepsilon ps} F^p(s) \psi^m(s) ds.$$

By the Lemma 2.4 with $\omega(u) = u^m$, $W(v) = \frac{1}{1-m}(v^{1-m} - v_0^{1-m})$ and $W^{-1}(z) = [(1-m)z + v_0^{1-m}]^{\frac{1}{1-m}}$, we conclude that

$$\begin{aligned} \psi(t) &\leq W^{-1} \left[W(2^{p-1} K(t)) + C_4 e^{\varepsilon pt} L(t) \int_0^t e^{-\varepsilon ps} F^p(s) ds \right] \\ &\leq 2^{p-1} K(t) \left[1 - (m-1) C_4 (2^{p-1} K(t))^{m-1} e^{\varepsilon pt} L(t) \right. \\ &\quad \left. \times \int_0^t e^{-\varepsilon ps} F^p(s) ds \right]^{\frac{1}{1-m}}, \end{aligned}$$

where $K(t)$ and $L(t)$ are as in the statement of the theorem.

(ii) If Ω is bounded, then the first part in the assertion (ii) of the theorem follows from the argument in (i) by an extension procedure and the application of the embedding $L^r(\Omega) \subset L^{\tilde{r}}(\Omega)$ for $\tilde{r} \leq r$. Now if $r < \frac{n}{n\beta-\alpha}$ we choose q such that $r < q < \frac{n}{n\beta-\alpha}$ and $\rho > \frac{nr}{n-(n\beta-\alpha)r}$. The Young inequality is therefore applicable. Also as $1 < q < \frac{nr}{n-\alpha p}$, the Hardy-Littlewood-Sobolev inequality (Lemma 2.3) applies (see also [3, p. 660] when Ω is bounded). □

In what follows, in order to simplify the statement of our next results, we define $v := r$ or \tilde{r} according to the cases (i) or (ii) in Theorem 3.1, respectively.


Corollary 3.2. *Suppose that the hypotheses of Theorem 3.1 hold. Assume further that $k(t, x)$ and $l(t, x)$ decay exponentially in time, that is $k(t, x) \leq e^{-\tilde{k}t}\tilde{k}(x)$ and $l(t, x) \leq e^{-\tilde{l}t}\tilde{l}(x)$ for some positive constants \tilde{k} and \tilde{l} . Then $\varphi(t, x)$ is also exponentially decaying to zero i.e.,*

$$(3.9) \quad \|\varphi(t, x)\|_v \leq C_5 e^{-\mu t}, \quad t > 0$$

for some positive constants C_5 and μ provided that

$$\|\tilde{k}(x)\|_r^{m-1} \|\tilde{l}(x)\|_\rho \int_0^\infty F^p(s) ds \leq \frac{1}{2^{m(p-1)}} (m-1) C_6^{p-1} C_2^p,$$

where C_6 is the best constant in Lemma 2.2 (third estimation) and the other constants are as in (i) and (ii) of Theorem 3.1.



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

Proof. From the inequality (1.1) we have

$$(3.10) \quad \varphi(t, x) \leq e^{-\tilde{k}t} \bar{k}(x) + e^{-\tilde{l}t} \bar{l}(x) \int_{\Omega} \int_0^t \frac{F(s) \varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} ds dy.$$

After multiplying by $e^{-m\mu t} \cdot e^{m\mu t}$, where $\mu = \min\{\tilde{k}, \tilde{l}\}$, we use the Hölder inequality to get

$$\begin{aligned} & \int_0^t (t-s)^{\beta-1} F(s) \varphi^m(s, y) ds \\ & \leq \left(\int_0^t (t-s)^{(\beta-1)p'} e^{-m\mu p' t} ds \right)^{\frac{1}{p'}} \left(\int_0^t F^{p'}(s) e^{m\mu p' t} \varphi^{mp'}(s, y) ds \right)^{\frac{1}{p'}}. \end{aligned}$$


As in the proof of Theorem 3.1, $0 < (1-\beta)p' < 1$. If C_6 is the best constant in Lemma 2.2, then we may write

$$(3.11) \quad \int_0^t (t-s)^{\beta-1} F(s) \varphi^m(s, y) ds \leq C_6 \left(\int_0^t F^{p'}(s) \varphi^{mp'}(s, y) ds \right)^{\frac{1}{p'}}.$$

From (3.10) and (3.11) it appears that

$$e^{\mu t} \varphi(t, x) \leq \bar{k}(x) + C_6 \bar{l}(x) \int_{\Omega} \left(\int_0^t F^{p'}(s) e^{m\mu p' t} \varphi^{mp'}(s, y) ds \right)^{\frac{1}{p'}} \frac{dy ds}{|x-y|^{n-\alpha}}.$$

Taking the L^r -norm and applying the Minkowski inequality and then the Young



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 14 of 20

inequality (Lemma 2.1), we find

$$e^{\mu t} \|\varphi(t, x)\|_r \leq \|\bar{k}(x)\|_r + C_6 \|\bar{l}(x)\|_\rho \times \left\| \int_{\Omega} \left(\int_0^t F^p(s) e^{mp\mu t} \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}} \frac{dy ds}{|x-y|^{n-\alpha}} \right\|_q,$$

with $\frac{1}{r} = \frac{1}{\rho} + \frac{1}{q}$. Applying Lemma 2.3, we arrive at


$$e^{\mu t} \|\varphi(t, x)\|_r \leq \|\bar{k}(x)\|_r + C_2 C_6 \|\bar{l}(x)\|_\rho \left\| \left(\int_0^t F^p(s) e^{mp\mu t} \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}} \right\|_p$$

or

$$(3.12) \quad e^{\mu t} \|\varphi(t, x)\|_r \leq \|\bar{k}(x)\|_r + C_2 C_6 \|\bar{l}(x)\|_\rho \left(\int_0^t F^p(s) e^{mp\mu t} \|\varphi(s, x)\|_r^{mp} ds \right)^{\frac{1}{p}}.$$

Taking both sides of (3.12) to the power p , we obtain

$$(3.13) \quad e^{\mu p t} \|\varphi(t, x)\|_r^p \leq 2^{p-1} \|\bar{k}(x)\|_r^p + C_7 \|\bar{l}(x)\|_\rho^p \int_0^t F^p(s) e^{mp\mu t} \|\varphi(s, x)\|_r^{mp} ds.$$



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

Next, putting

$$\chi(t) := e^{\mu pt} \|\varphi(t, x)\|_r^p,$$

the inequality (3.13) may be written as

$$\chi(t) \leq 2^{p-1} \|\bar{k}(x)\|_r^p + C_7 \|\bar{l}(x)\|_\rho^p \int_0^t F^p(s) \chi^m(s) ds.$$

The rest of the proof is essentially the same as that of Theorem 3.1. □

In the following corollary we consider the somewhat more general inequality

$$(3.14) \quad \varphi(t, x) \leq k(t, x) + l(t, x) \int_{\Omega} \int_0^t \frac{s^\delta F(s) \varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} dy ds,$$

$x \in \Omega, t > 0$

for some specified δ .


Corollary 3.3. *Suppose that the hypotheses of Theorem 3.1 hold. Assume further that $k(t, x) \leq t^{-\hat{k}} \bar{k}(x)$ and $1 + \delta p' - m p' \min\{\hat{k}, 1 - \beta\} > 0$. Then any $\varphi(t, x)$ satisfying (3.14) is also polynomially decaying to zero*

$$\|\varphi(t, x)\|_v \leq C_8 t^{-\omega}, \quad C_8, \omega > 0$$

provided that

$$\|\bar{k}(x)\|_r^{m-1} L(t) \int_0^t e^{\varepsilon ps} F^p(s) ds \leq \frac{1}{2^{m(p-1)}} (m-1) C_9^{p-1} C_2^p$$

where C_9 is the best constant in Lemma 2.5.



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

Proof. Let us consider the inequality

$$(3.15) \quad \varphi(t, x) \leq t^{-\hat{k}} \bar{k}(x) + l(t, x) \int_{\Omega} \int_0^t \frac{s^{\delta} F(s) \varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} dy ds.$$

Multiplying by $s^{-m \min\{\hat{k}, 1-\beta\}} e^{-\varepsilon s} \cdot s^{m \min\{\hat{k}, 1-\beta\}} e^{\varepsilon s}$, we obtain


$$(3.16) \quad \int_0^t (t-s)^{\beta-1} s^{\delta} F(s) \varphi^m(s, y) ds \\ \leq \left(\int_0^t (t-s)^{(\beta-1)p'} s^{\delta p' - mp' \min\{\hat{k}, 1-\beta\}} e^{-\varepsilon p' s} ds \right)^{\frac{1}{p'}} \\ \times \left(\int_0^t s^{mp \min\{\hat{k}, 1-\beta\}} e^{p\varepsilon s} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}}.$$

As $1 + \delta p' - mp' \min\{\hat{k}, 1-\beta\} > 0$, we may apply Lemma 2.5 to the first term in the right hand side of (3.16) to get

$$(3.17) \quad \int_0^t (t-s)^{\beta-1} s^{\delta} F(s) \varphi^m(s, y) ds \\ \leq M t^{\beta-1} \left(\int_0^t s^{mp \min\{\hat{k}, 1-\beta\}} e^{p\varepsilon s} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}}.$$

Using (3.17) we infer from inequality (3.15) that

$$t^{\min\{\hat{k}, 1-\beta\}} \varphi(t, x) \leq \bar{k}(x) + M l(t, x) \\ \times \int_{\Omega} \left(\int_0^t s^{mp \min\{\hat{k}, 1-\beta\}} e^{p\varepsilon s} F^p(s) \varphi^{mp}(s, y) ds \right)^{\frac{1}{p}} \frac{dy}{|x-y|^{n-\alpha}}.$$



On an Integral Inequality with a Kernel Singular in Time and Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20

Next, after using the Hardy-Littlewood-Sobolev inequality and defining


$$\phi(t, x) := t^{p \min\{\hat{k}, 1-\beta\}} \|\varphi(t, x)\|_r^p,$$

we proceed as in Theorem 3.1 to find a (uniform) bound for $\phi(t, x)$. □

Remark 3.1. *The investigation of (1.1) with a weakly singular kernel in time, that is*

$$\varphi(t, x) \leq k(t, x) + l(t, x) \int_{\Omega} \int_0^t \frac{e^{-\gamma(t-s)} F(s) \varphi^m(s, y)}{(t-s)^{1-\beta} |x-y|^{n-\alpha}} dy ds, \quad \gamma > 0$$

is simpler since this kernel is summable (in time).



On an Integral Inequality with a
Kernel Singular in Time and
Space

Nasser-eddine Tatar

Title Page

Contents



Go Back


Close

Quit

Page 18 of 20

References

- [1] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, New York, 1992.
- [2] G. BUTLER AND T. ROGERS, A generalization of a lemma of Bihari and applications to pointwise estimates for integral equations, *J. Math. Anal. Appl.* **33**(1) (1971), 77–81.
- [3] R.E. EDWARDS, *Functional Analysis: Theory and Applications*, New York, Holt, Rinehart and Winston, 1965.
- [4] D. HENRY, *Geometric Theory of Semilinear Parabolic Equations*, Springer-Verlag, Berlin, Heidelberg, New York, 1981.
- [5] L. HÖRMANDER, *The Analysis of Linear Partial Differential Operator*, Vol. I, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1983.
- [6] M. KIRANE AND N.-E. TATAR, Global existence and stability of some semilinear problems, *Arch. Math. (Brno)*, **36**(1) (2000), 33–44.
- [7] S. MAZOUZI AND N.-E. TATAR, Global existence for some integrodifferential equations with delay subject to nonlocal conditions, *Zeit. Anal. Anw.*, **21**(1) (2002), 249–256.
- [8] S. MAZOUZI AND N.-E. TATAR, An improved exponential decay result for some semilinear integrodifferential equations, to appear in *Arch. Math.*



On an Integral Inequality with a
Kernel Singular in Time and
Space

Nasser-eddine Tatar

Title Page

Contents




Go Back

Close

Quit

Page 19 of 20

- [9] M. MEDVED', A new approach to an analysis of Henry type integral inequalities and their Bihari type versions, *J. Math. Anal. Appl.*, **214** (1997), 349–366.
- [10] M. MEDVED', Singular integral inequalities and stability of semilinear parabolic equations, *Arch. Math. (Brno)*, **24** (1998), 183–190.
- [11] M. MEDVED', Integral inequalities and global solutions of semilinear evolution equations, *J. Math. Anal. Appl.*, **267** (2002), 643–650.
- [12] B.G. PACHPATTE, *Inequalities for Differential and Integral Equations*, in Mathematics in Science and Engineering Vol. **197**, Academic Press, London, 1998.
- [13] N.-E. TATAR, Exponential decay for a semilinear problem with memory, *Arab J. Math. Sc.*, **7**(1) (2001), 29–45.
- [14] V.A. SOLONNIKOV, Existence of solutions of non-stationary linearized systems of Navier-Stokes equations, *Trudy Mat. Inst. Steklov.* **70** (1964), 213–317 [English trans: *AMS Trans. Ser. II* (1975), 1–116].



On an Integral Inequality with a
Kernel Singular in Time and
Space

Nasser-eddine Tatar

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20