

Journal of Inequalities in Pure and Applied Mathematics

ON GENERALIZED MONOTONE MULTIFUNCTIONS WITH APPLICATIONS TO OPTIMALITY CONDITIONS IN GENERALIZED CONVEX PROGRAMMING

A. HASSOUNI AND A. JADDAR

Département de Mathématiques et d'Informatique,
Faculté des Sciences,
B.P. 1014, Rue Ibn Batouta,
Rabat, Maroc.

*E*Mail: hassounia@hotmail.com

©2000 Victoria University
ISSN (electronic): 1443-5756
062-02



volume 4, issue 4, article 67,
2003.

*Received 29 May, 2002;
accepted 16 March, 2003.*

Communicated by: Z. Nashed

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

Characterization of quasiconvexity and pseudoconvexity of lower semicontinuous functions on Banach spaces are presented in terms of abstract subdifferentials relying on a Mean Value Theorem. We give some properties of the normal cone to the lower level set of f . We also obtain necessary and sufficient optimality conditions in quasiconvex and pseudoconvex programming via variational inequalities.


2000 Mathematics Subject Classification: 90C26, 26B25, 47H04.

Key words: Generalized monotone multifunction, Generalized convex function, Quasiconvex, Pseudoconvex, Generalized subdifferentials, Normal cone, Level set, Local minimum, Global minimum, Variational inequalities.

Contents

1	Introduction	3
2	Preliminaries	4
3	Generalized Convex Functions and Generalized Monotone Multifunctions	8
3.1	Quasiconvex Functions and Quasimonotone Multifunctions	8
3.2	Pseudoconvexity and Subdifferential Properties	13
4	Optimality Conditions and Variational Inequalities	16
4.1	Quasiconvex Programming	16
4.2	Pseudoconvex Programming	20

References



On Generalized Monotone Multifunctions with Applications to Optimality Conditions in Generalized Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 2 of 26


1. Introduction

It is natural in convex analysis to search for characterizations of generalized convex functions in terms of some kind of generalized derivatives or subdifferentials. Several contributions to this question has been made recently. The reader may consult for example [3, 5, 11, 13, 16, 20] for quasiconvex functions and [2, 8, 21, 23, 25] for pseudoconvex functions.

In this paper, we shall define an abstract subdifferential as in [1, 23] which allows us to extend some results in [1, 2, 8, 23] and to give some properties of the normal cone to lower level sets of a given function f .

Notice that the condition $0 \in \partial f(\bar{x})$ for $\bar{x} \in X$, is known to be a necessary but not a sufficient optimality condition in quasiconvex programming for some subdifferentials. We give, using some variational inequalities, a necessary and sufficient condition for a point to be either a local or a global minimum.

After the introduction of some notations and definitions in Section 2, we present in Section 3 some properties of the abstract subdifferential and normal cone to lower level sets of quasiconvex and pseudoconvex functions. Then, in Section 4, we give some optimality conditions involving variational inequalities. This should extend our previous results stated for quasiconvex lower semicontinuous functions on Banach spaces with the Clarke-Rockafellar subdifferential in [13].



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 3 of 26

2. Preliminaries

Let X be a real Banach space, X^* its dual and $\langle \cdot, \cdot \rangle$ the duality pairing between X^* and X . The segment $[a, b]$ is the set $\{a + t(b - a); \quad t \in [0, 1]\}$ while $[a, b[$ is the set $[a, b] \setminus \{b\}$. The open ball with center x and radius r in X is denoted by $B(x, r)$, and the polar cone of a nonempty subset A of X is

$$A^\circ = \{x^* \in X^*; \quad \langle x^*, a \rangle \leq 0, \quad \forall a \in A\}.$$

For an extended real valued function $f : X \mapsto \mathbb{R} \cup \{+\infty\}$, the effective domain is defined by

$$\text{dom}(f) = \{x \in X; \quad f(x) < \infty\}.$$

We write l.s.c. for lower semicontinuous, and $x_n \rightarrow_f x$ when $x_n \rightarrow x$ and $f(x_n) \rightarrow f(x)$.

The abstract subdifferential we consider here is defined as follows:


Definition 2.1. An operator ∂ that associates to any l.s.c. function $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ and a point $x \in X$ a subset $\partial f(x)$ of X^* is a subdifferential if the following assertions hold:

(P1) $\partial f(x) = \{x^* \in X^*; \quad f(y) \geq f(x) + \langle x^*, y - x \rangle \quad \forall y \in X\}$ when f is convex.

(P2) If $x \in \text{dom } f$ is a local minimum of f , then $0 \in \partial f(x)$.

(P3) $\partial f(x) = \partial g(x)$, for any $g : X \mapsto \mathbb{R} \cup \{+\infty\}$ such that $f - g$ is constant in a neighborhood of x .

(P4) $\partial f(x) = \emptyset$, for any $x \in X$ such that $f(x) = +\infty$.



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 4 of 26

It is well known that the Clarke-Rockafellar subdifferential $\partial^{CR} f$ satisfies Zagrodny's Mean value theorem [27]. In order to extend this theorem to our subdifferential, we shall deal with a particular space associated with ∂ called ∂ -reliable.

Definition 2.2. [23]. A Banach space X is ∂ -reliable if for each l.s.c. function $f : X \mapsto \mathbb{R} \cup \{+\infty\}$, for any Lipschitz convex function g and any $x \in \text{dom } f$ such that $f + g$ achieves its minimum in X and each $\varepsilon > 0$ we have:

$$0 \in \partial f(u) + \partial g(v) + \varepsilon B_1^*(0),$$

for some $u, v \in B_\varepsilon(x)$ such that $|f(u) - f(v)| < \varepsilon$.

In the case of the Clarke-Rockafellar subdifferential ∂^{CR} [26] or Iofee subdifferential ∂^I [7], any Banach space is ∂ -reliable.

In the sequel, we will restrict ourselves to subdifferentials that are included in the dag subdifferential


$$\partial^\dagger f(x) = \{x^* \in X^*; \langle x^*, v \rangle \leq f^\dagger(x, v) \quad \forall v \in X\},$$

where

$$f^\dagger(x, v) = \limsup_{(t,y) \rightarrow (0+,x)} t^{-1}(f(y + t(v + x - y)) - f(y)).$$

This subdifferential was introduced by Penot (see [22]), it is large enough to contain the Clarke-Rockafellar ∂^{CR} and Upper Dini ∂^{D+} subdifferentials and still has good properties.

Our results rely on the following mean value theorem.



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 5 of 26

Theorem 2.1. [23]. Let X be a ∂ -reliable space and $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ a l.s.c. function. For any $a \in \text{dom } f$, $b \in X \setminus \{a\}$, $\beta \leq b$, there exists a sequence c_n in X converging to some $c \in [a, b)$ and a sequence $c_n^* \in \partial f(c_n)$ such that for any $b' = c + t(b - a)$, with $t > 0$ we have:

- i) $\liminf_n \langle c_n^*, b - a \rangle \geq \beta - f(a)$,
- ii) $\liminf_n \langle c_n^*, c - c_n \rangle \geq 0$,
- iii) $\liminf_n \left\langle c_n^*, \frac{\|b-a\|}{\|b'-c\|} (b' - c_n) \right\rangle \geq \beta - f(a)$.

Following the methods of [1, 16, 20], we get a similar lemma for our abstract subdifferential, which is immediate by Theorem 2.1.

Lemma 2.2. Let X be a Banach ∂ -reliable space, f a l.s.c. function. Let $a, b \in X$ with $f(a) < f(b)$ then there exists $c \in [a, b[$ and two sequences $c_n \rightarrow c$, $c_n^* \in \partial f(c_n)$ with

$$\langle c_n^*, x - c_n \rangle > 0,$$


for any $x = c + t(b - a)$ with $t > 0$.

Proof. Let $a, b \in X$ with $f(a) < f(b)$, then we can find by Theorem 2.1, $c \in [a, b[$ and two sequences $c_n \rightarrow c$, $c_n^* \in \partial f(c_n)$ with

$$\liminf_n \langle c_n^*, c - c_n \rangle \geq 0,$$

and

$$\liminf_n \langle c_n^*, b - a \rangle \geq f(b) - f(a) > 0.$$



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 6 of 26

For $x = c + t(b - a)$ with $t > 0$, we have

$$\langle c_n^*, x - c_n \rangle = \langle c_n^*, c - c_n \rangle + t \langle c_n^*, b - a \rangle.$$


It follows that

$$\liminf_n \langle c_n^*, x - c_n \rangle > 0.$$

Hence, for n large enough, we have that

$$\langle c_n^*, x - c_n \rangle > 0.$$

□



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 7 of 26

3. Generalized Convex Functions and Generalized Monotone Multifunctions

3.1. Quasiconvex Functions and Quasimonotone Multifunctions

We recall the characterization of quasiconvex functions of [22, 23]. It will allow us to extend and generalize some properties of the normal cone to the lower level set given in [12, 13] to a more general setting.

Indeed, for $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ a l.s.c. function, f is said to be quasiconvex if for every $x, y \in X$ and $\lambda \in [0, 1]$ one has

$$f(\lambda x + (1 - \lambda)y) \leq \max\{f(x), f(y)\}.$$

And denoting by

$$S_f(\lambda) = \{x \in X; f(x) \leq \lambda\}.$$


Quasiconvexity is geometrically equivalent to the fact that $S_f(\lambda)$ is a convex set for all $\lambda \in \mathbb{R}$. In the above one could use the strict level sets as well.

Recall that a multifunction $A : X \rightarrow X^*$ is said to be quasimonotone if for every pair of distinct points $x, y \in X$:

$$\exists x^* \in A(x), \text{ such that } \langle x^*, y - x \rangle > 0 \text{ then, } \forall y^* \in A(y), \langle y^*, y - x \rangle \geq 0.$$

Theorem 3.1. [22, 23] Let X be a Banach space and $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ a l.s.c. function. And consider the following assertions

i) f is quasiconvex.



On Generalized Monotone Multifunctions with Applications to Optimality Conditions in Generalized Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 8 of 26

ii) ∂f is quasimonotone.

Then **i)** implies **ii)** if $\partial f \subset \partial^\dagger f$. And **ii)** implies **i)** if X is ∂ -reliable.

For $x_0 \in X$, set

$$L(x_0) = \{x \in X; \quad f(x) = f(x_0)\}.$$

Then we have

Proposition 3.2. *Let X be a Banach ∂ -reliable space, and f a l.s.c. quasiconvex function such that $\partial f \subset \partial^\dagger f$. If for $x_0 \in X$ there exists $r > 0$ with*

$$0 \notin \partial f(x), \quad \text{for all } x \in B(x_0, r) \cap L(x_0),$$

then we have

$$[\partial f(x_0)]^{\circ\circ} \subset N(S_f(f(x_0)); x_0),$$


where $N(S_f(f(x_0)); x_0)$ is the normal cone to the lower level set $S_f(f(x_0))$ at the point x_0 .

Proof. Suppose by contradiction that there exists v such that

$$v \in [\partial f(x_0)]^{\circ\circ} \quad \text{and} \quad v \notin N(S_f(f(x_0)); x_0).$$

We can check that

$$Cl(\mathbb{R}_+co(\partial f(x_0))) = [\partial f(x_0)]^{\circ\circ}.$$



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 9 of 26

So, we can suppose without loss of generality that $v = x_0^* \in \partial f(x_0)$. Then, we can find some $x_1 \in S_f(x_0)$ such that

$$(3.1) \quad \langle x_0^*, x_1 - x_0 \rangle > 0.$$

We claim that $f(x_0) = f(x_1)$. Otherwise by Lemma 2.2, there exists $c \in [x_1, x_0[$ and two sequences $c_n \rightarrow_f c$ and $c_n^* \in \partial f(c_n)$ with

$$\langle c_n^*, x_0 - c_n \rangle > 0.$$

By using the quasimonotonicity of ∂f we have:

$$\langle x_0^*, x_0 - c_n \rangle \geq 0.$$

Then, letting $n \rightarrow +\infty$ we get

$$\langle x_0^*, x_0 - c \rangle \geq 0.$$

It follows that

$$\langle x_0^*, x_0 - x_1 \rangle \geq 0.$$


A contradiction with (3.1), thus $f(x_0) = f(x_1)$.

Now, set $V_{x_1} = \{x \in X : \langle x_0^*, x - x_0 \rangle > 0\}$.

V_{x_1} is an open neighborhood of x_1 and using the same argument as above we can check that x_1 is a minimum of f on V_{x_1} , and that

$$x_\lambda = x_0 + \lambda(x_1 - x_0) \in V_{x_1} \text{ and } f(x_\lambda) = f(x_0) \text{ for any } \lambda \in]0, 1[.$$

Then there exists $r > 0$ and $\bar{\lambda} \in]0, 1[$ such that $x_{\bar{\lambda}}$ is a global minimum of f on $B(x_0, r) \cap V_{x_1}$. Therefore $0 \in \partial f(x_{\bar{\lambda}})$, which is impossible. \square



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 10 of 26

The former proposition extends some already known results for differentiable functions (see for instance [5]). If we denote by $T(S_f(f(x)); x)$, the tangent cone of the lower level convex set $S_f(f(x))$ at the point $x \in X$, then

$$T(S_f(f(x)); x) = [N(S_f(f(x)); x)]^\circ.$$

A sufficient condition that allows us to obtain the equality in Proposition 3.2 is stated in the following proposition

Proposition 3.3. *Under the hypothesis of Proposition 3.2 and if*

$$[\partial f(x)]^\circ \subset T(S_f(f(x)); x).$$

Then we have

$$N(S_f(f(x)); x) = [\partial f(x)]^{\circ\circ}.$$

Proof. By the bipolar theorem [4] one has


$$[\partial f(x)]^{\circ\circ} \supset N(S_f(f(x)); x).$$

And from Proposition 3.2, the equality immediately holds. □

The following condition

$$N(S_f(f(x)); x) = [\partial f(x)]^{\circ\circ},$$

is in fact a certain kind of regularity condition, which holds only for a subclass of quasiconvex functions. Another abstract approach was developed in [15] based on Crouzeix's representation theorem [6] who obtained a similar equality for his quasi-subdifferential.



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 11 of 26

Consider the multifunction Γ from X to X^* defined by

$$\Gamma(x) = N(S_f(f(x)); x), \quad \text{for } x \in X.$$

Then by using Proposition 3.3, we obtain

Proposition 3.4. *Let X be a Banach ∂ -reliable space, f a l.s.c. quasiconvex function. If for any $x \in X$, $\partial f(x)$ is nonempty such that*

$$(\partial f(x))^\circ \subset T(S_f(f(x)); x).$$

Then, the multifunction Γ is quasimonotone.


Proof. Since f is quasiconvex, by Theorem 3.1 ∂f is quasimonotone. Using Proposition 2.8 of [12], it follows easily that the multifunction $x \mapsto [\partial f(x)]^{\circ\circ}$ is quasimonotone. Then by Proposition 3.3, Γ is also quasimonotone.

It follows that Γ is quasimonotone. □

A particular case of this proposition when ∂ coincides with the Clarke-Rockafellar subdifferential ∂^{CR} , was treated in [13], whose exact statement is the following.

Proposition 3.5. *Let X be a Banach space, f a l.s.c. function from X to $\mathbb{R} \cup \{+\infty\}$ such that $\partial^{CR} f(x)$ is nonempty and $0 \notin \partial^{CR} f(x)$ for all $x \in X$.*

If f is quasiconvex then the multifunction Γ is quasimonotone.



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 12 of 26

3.2. Pseudoconvexity and Subdifferential Properties

The original definition of pseudoconvexity was introduced by Mangazarian in [21] for differentiable functions. This concept was extended later by many authors (see for instance [17, 22, 24]) for arbitrary functions. We will here use the following form:

A function f is said to be pseudoconvex for the subdifferential ∂ if for any $x, y \in X$:

$$\exists x^* \in \partial f(x) : \langle x^*, y - x \rangle \geq 0 \implies f(x) \leq f(y).$$

A multifunction $A : X \rightarrow X^*$ is said to be pseudomonotone if for every pair of distinct points $x, y \in X$


$$\exists x^* \in A(x) : \langle x^*, y - x \rangle > 0 \quad \text{then,} \quad \forall y^* \in A(y), \langle y^*, y - x \rangle > 0.$$

As in the differentiable case, every pseudoconvex function satisfies the fundamental properties:

- every local minimum of f is global.
- $0 \in \partial f(x)$ implies that x is a global minimum of f .

Another interesting property extending a result of [8] where it was stated for the Clarke-Rockafellar subdifferential is the following.

Proposition 3.6. *Let X be a Banach ∂ -reliable and $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ be a l.s.c. function and pseudoconvex function such that $\partial f \subset \partial^\dagger f$, let $x, y \in X$. Then the existence of $x^* \in \partial f(x)$ verifying $\langle x^*, y - x \rangle > 0$ implies $f(x) < f(y)$.*



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 13 of 26

Proof. Let $x, y \in X$ such that $\langle x^*, y - x \rangle > 0$ for some $x^* \in \partial f(x)$, then there exists $\varepsilon > 0$ such that

$$\langle x^*, y' - x \rangle > 0, \quad \forall y' \in B(y, \varepsilon).$$

By the pseudoconvexity of f , we have $f(y') \geq f(x)$.

In particular, $f(y) \geq f(x)$. If we suppose by contradiction that $f(y) = f(x)$, then y must be a global minimum. On the other hand, since $f^\dagger(x, y - x) > 0$ then, there exist two sequences $x_n \rightarrow x, t_n \rightarrow 0^+$ such that

$$t_n^{-1}[f(x_n + t_n(y - x_n)) - f(x_n)] > 0.$$

By the quasiconvexity of the function f (see for instance the proof of Proposition 2.2 in [8]), we get $f(y) > f(x_n)$ which is impossible. \square

We use this proposition to prove a known result for the Clarke-Rockafellar subdifferential for bigger subdifferentials

Theorem 3.7. *Let X be a ∂ -reliable space and $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ a l.s.c. function such that $\partial f \subset \partial^\dagger f$. And consider the following assertions*

- i) f is pseudoconvex.*
- ii) ∂f is pseudomonotone.*

Then, i) implies ii). And ii) implies i) if f is radially continuous.

Proof. The implication **ii)** \implies **i)** is in [23]. For **i)** \implies **ii)**, suppose by contradiction that there exist $x, y \in X$, such that there exist $x^* \in \partial f(x)$ and $y^* \in \partial f(y)$ verifying

$$\langle x^*, y - x \rangle > 0 \quad \text{and} \quad \langle y^*, y - x \rangle \leq 0.$$



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 14 of 26

Then, from Proposition 3.5 and the pseudoconvexity of f we have

$$f(x) < f(y) \quad \text{and} \quad f(y) \leq f(x).$$


A contradiction. □

Now, we state a similar result to Proposition 3.2 for pseudoconvex functions.

Proposition 3.8. *Let X be a Banach ∂ -reliable space with $\partial \subset \partial^\dagger$, f a l.s.c. and pseudoconvex function from X to $\mathbb{R} \cup \{+\infty\}$. Then we have*

$$[\partial f(x)]^{\circ\circ} \subset N(S_f(f(x)); x).$$

Proof. Let $x^* \in \partial f(x)$ and suppose by contradiction that $x^* \notin N(S_f(f(x)); x)$. Then, there exists $y \in S_f(f(x))$ such that $\langle x^*, y - x \rangle > 0$ for some $x^* \in \partial f(x)$. It follows then by Proposition 3.6 that $f(y) > f(x)$, which is impossible. □



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 15 of 26

4. Optimality Conditions and Variational Inequalities

4.1. Quasiconvex Programming

We recall the Minty variational inequality (we use the terminology of Giannessi [9]) that we shall use for our subdifferential. It will be exploited to give some conditions of optimality in nonlinear programming and necessary and sufficient conditions for optimality in quasiconvex programming.

Let Γ be a multifunction from X to X^* , $S \subset X$ and $\bar{x} \in S$.

A point \bar{x} is a Minty equilibrium of Γ if the following variational inequality holds

$$(D) \quad \forall x \in S, \quad \langle \gamma(x), x - \bar{x} \rangle \geq 0, \quad \forall \gamma(x) \in \Gamma(x).$$


Suppose that f is a l.s.c. function from X to $\mathbb{R} \cup \{+\infty\}$ and consider the following minimisation problem

$$(4.1) \quad \text{minimize } f(x), \text{ subject to } x \in C.$$

Then we have

Proposition 4.1. *Let X be a Banach ∂ -reliable space. If \bar{x} is a Minty equilibrium point of ∂f , then we have*

- i) *If $S = X$, then \bar{x} is a global minimum of f .*
- ii) *If $S = N$, where N is a convex open neighborhood of \bar{x} then \bar{x} is a local minimum of f .*



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 16 of 26

Proof. It is enough to prove (ii). Suppose by contradiction that \bar{x} is not a solution of the program (4.1), then there exists $x \in S$ such that $f(x) < f(\bar{x})$. By Lemma 2.2, there exists $c \in [x, \bar{x}[$ and two sequences $c_n \rightarrow_f c$, $c_n^* \in \partial f(c_n)$ with

$$\langle c_n^*, d - c_n \rangle > 0,$$

for any $d = c + t(\bar{x} - x)$ where $t > 0$.

Since S is a convex open neighborhood of \bar{x} then $[x, \bar{x}] \subset S$. Furthermore, for n large enough $c_n \in S$.

In the particular case where $d = \bar{x}$, we have:

$$\langle c_n^*, \bar{x} - c_n \rangle > 0.$$


A contradiction with the variational inequality (D), thus \bar{x} is a local minimum of f . \square

This proposition extends Theorem 2.2 of [18] for nondifferentiable optimization problems.

If in the problem (4.1), the function f to be minimized is l.s.c. and quasi-convex, then we have

Theorem 4.2. *Let X be a Banach ∂ -reliable, and f be a l.s.c. and quasiconvex function such that $\partial f \subset \partial^\dagger f$, and $\bar{x} \in S$. If $S = N$, where N is an open and convex neighborhood of \bar{x} or $S = X$, then the following assertions are equivalent*

- i) \bar{x} is an optimal solution of (4.1).
- ii) \bar{x} is a Minty equilibrium point of ∂f .



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 17 of 26

Proof. **ii) \implies i)** is obtained from Proposition 4.1. Let us show that **i) \implies ii).** Assume that \bar{x} is a strict minimum of (4.1), then for all $x \in S$ such that $x \neq \bar{x}$ we have $f(x) > f(\bar{x})$.

According to Lemma 2.2, there exist $c \in [\bar{x}, x]$, $c_n \rightarrow_f c$ and $c_n^* \in \partial f(c_n)$ such that

$$\langle c_n^*, d - c_n \rangle > 0,$$

for all $d = c + t(x - \bar{x})$ where $t > 0$.

When $d = x$, we obtain that

$$\langle c_n^*, x - c_n \rangle > 0.$$

f being quasiconvex, by Theorem 2.1, ∂f is quasimonotone. It follows then that

$$\text{for all } x^* \in \partial f(x), \quad \langle x^*, x - \bar{x} \rangle \geq 0.$$


Hence, ∂f satisfies the variational inequality (D).

Suppose that we are in the case where \bar{x} is not a strict minimum of (4.1) and let us consider the function $f_{\bar{x}}$ defined by

$$f_{\bar{x}}(x) = \max\{f(x), f(\bar{x})\},$$

and define h by

$$(4.2) \quad h(x) = \begin{cases} f_{\bar{x}}(x) & \text{for } x \neq \bar{x} \\ \nu & \text{for } x = \bar{x} \end{cases}$$



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 18 of 26

where $\nu < f(\bar{x})$. We see easily that h is l.s.c. and quasiconvex and that \bar{x} is a strict local minimum of h . Then, we have

$$\forall x \neq \bar{x} \quad \langle x^*, x - \bar{x} \rangle \geq 0, \quad \forall x^* \in \partial h(x).$$

From (P3), we get $\partial f(x) = \partial h(x)$. □

In the case when 0 is in the interior of $\partial f(\bar{x})$, i.e. $0 \in \text{int}(\partial f(\bar{x}))$, we have the more precise result

Proposition 4.3. *Let X be a ∂ -reliable space and $f : X \mapsto \mathbb{R} \cup \{+\infty\}$ a l.s.c. and quasiconvex function. If $0 \in \text{int}(\partial f(\bar{x}))$ then \bar{x} is a Minty equilibrium point of ∂f . Moreover \bar{x} is a global minimum of f .*

Proof. Assume that $0 \in \text{int}(\partial f(x))$ then

there exists $\varepsilon > 0$ such that $B_{X^*}(0, \varepsilon) \subset \partial f(x)$,

where

$$B_{X^*}(0, \varepsilon) = \{x^* \in X^* : \|x^*\| < \varepsilon\}.$$


Let $d \in X$ such that $d \neq 0$ and consider the linear mapping ℓ_d defined by

$$\ell_d(x^*) = \langle x^*, d \rangle, \quad \text{for } x^* \in X^*.$$

By the open mapping Theorem [4] one has

$$\langle B_{X^*}(0, \varepsilon), d \rangle \subset \langle \partial f(x), d \rangle.$$

Since f is quasiconvex, then ∂f is quasimonotone.



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 19 of 26

We already know by Definition 2.1 of [12] that the multifunction $\partial f_{x,d}$ defined by

$$\partial f_{x,d}(\lambda) = \langle \partial f(x + \lambda d), d \rangle,$$

is quasimonotone, and we can see easily that

$$\langle \lambda d, \partial f(x + \lambda d) \rangle \subset \mathbb{R}_+,$$

for all $\lambda \in \mathbb{R}$ and $d \in X \setminus \{0\}$, thus (D) holds for ∂f . □

4.2. Pseudoconvex Programming

For the pseudoconvex function f , we shall get necessary and sufficient conditions for a point \bar{x} to be a global extremum of f over a convex set C .

First consider the problem (4.1), with f is pseudoconvex, l.s.c. and radially continuous, then we have


Theorem 4.4. *Let X be a Banach space ∂ -reliable, and f a pseudoconvex l.s.c. such that $\partial f \subset \partial^\dagger f$, and let $\bar{x} \in C$. Then the following assertions are equivalent*

i) \bar{x} is an optimal solution of (4.1).

ii) (D) holds.

Proof. Suppose that \bar{x} is a solution of (4.1), then by Proposition 3.6, if $f(\bar{x}) \leq f(x)$, then we must have

$$\forall x^* \in \partial f(x), \quad \langle x^*, \bar{x} - x \rangle \leq 0.$$



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 20 of 26

This means that the variational inequality (D) holds.

Conversely, let $x \in C$ such that $x \neq \bar{x}$ then for some $y \in (\bar{x}, x)$, we have

$$\forall y^* \in \partial f(y), \quad \langle y^*, \bar{x} - y \rangle \leq 0.$$

It follows that

$$\forall y^* \in \partial f(y), \quad \langle y^*, x - y \rangle \leq 0.$$

Since $\partial f(y)$ is nonempty and from the pseudoconvexity of f we have

$$f(y) \leq f(x), \quad \forall y \in (\bar{x}, x).$$

But since f is s.c.i., then $f(\bar{x}) \leq f(x)$. □

We now proceed to the maximisation problem

$$(4.3) \quad \text{maximize } f(x), \text{ subject to } x \in C.$$

For $z \in C$, we denote by

$$C_z = \{x \in C; \quad f(x) = f(z)\}.$$


Then we have

Theorem 4.5. *Let X be a ∂ -reliable space and f a pseudoconvex, l.s.c. and radially continuous such that for any x in C , $\partial f(x)$ is nonempty and $\partial f(x) \subset \partial^+ f(x)$. Let $\bar{x} \in C$ such that*

$$-\infty \leq \inf_C f < f(\bar{x}).$$

Then \bar{x} is a maximum of f on C if and only if

$$\text{for all } x \in C_{\bar{x}}, \quad \partial f(x) \subset N(C, x).$$



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 21 of 26

Proof. Suppose that

$$f(y) \leq f(\bar{x}); \quad \forall y \in C.$$

By Proposition 3.6 we have:

$$\text{for all } x \in C_{\bar{x}}, \quad \partial f(x) \subset N(C, x).$$

Conversely, by contradiction assume that there exists $\bar{z} \in C$ such that

$$f(\bar{z}) > f(\bar{x}).$$

Since by hypothesis, we can find some $z \in C$ with $f(z) < f(\bar{x})$.

By the radial continuity of f , there exists some $x_0 \in (z, \bar{z})$ such that

$$f(x_0) = f(\bar{x}).$$

It follows then that

$$\text{for all } x_0^* \in \partial f(x_0), \quad \langle x_0^*, z - x_0 \rangle = 0.$$

Since f is pseudoconvex then, $f(x_0) \leq f(z)$, a contradiction. □



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back


Close

Quit

Page 22 of 26

References

- [1] D. AUSSEL, *Théorème de la Valeur Moyenne et Convexité Generalisée en Analyse Non Régulière*, Université Blaise Pascal, PhD. Thesis, (1994).
- [2] D. AUSSEL, Subdifferential properties of quasiconvex and pseudoconvex functions: A unified approach, *J. Optim. Th. Appl.*, **97** (1998), 29–45.
- [3] D. AUSSEL, J.-N. CORVELLEC AND M. LASSONDE, Subdifferential characterization of quasiconvexity and convexity, *J. Conv. Analysis*, **1** (1994), 195–201.
- [4] H. BREZIS, *Analyse Fonctionnelle Théorie et Applications*, Masson (1980).
- [5] J.-P. CROUZEIX, About differentiability of order one of quasiconvex functions on \mathbb{R}^n , *J. Optim. Theory Appl.*, **36** (1982), 367–385.
- [6] J.-P. CROUZEIX, Continuity and differentiability properties of quasiconvex functions on \mathbb{R}^n , *Generalized Concavity in Optimization and Economics*, Academic Press, New York, 109–130, (1981).
- [7] A.D. IOFEE, Proximal analysis and approximate subdifferentials, *J. London Math. Soc.*, **41** (1990), 261–268.
- [8] A. DANIILIDIS AND N. HADJISAVVAS, On the subdifferentials of quasiconvex and pseudoconvex functions and cyclic monotonicity, *J. Math. Anal. Appl.*, **237** (1999), 30–42.



On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents




Go Back

Close

Quit

Page 23 of 26

- [9] F. GIANNESI, On Minty variational principle, in: *New Trends in Mathematical Programming*, F.Giannessi, S. Komlósi and T. Rapcsák (Eds.), Kluwer Scientific Publishers, Dordrecht, 93–99, (1998).
- [10] N. HADJISAVVAS AND S. SCHAIBLE, Quasimonotone variational inequalities in Banach spaces, *J. Optim. Th. Appl.*, **90**(1) (1996), 95–111.
- [11] A. HASSOUNI, Sous-différentiels des fonctions quasiconvexes, *Thèse de troisième cycle*, Université Paul sabatier, pp. 68, (1983).
- [12] A. HASSOUNI, Quasimonotone multifunctions; Applications to optimality conditions in quasiconvex programming, *Numer. Funct. Anal. Optim.*, **13**(3-4) (1992), 267–275.
- [13] A. HASSOUNI AND A. JADDAR, Quasiconvex functions and applications to optimality conditions in nonlinear programming, *Appl. Math. Lett.*, **14** (2001), 241–244.
- [14] J.P. HIRIARTY–URRUTY, Tangent cones generalized gradients and mathematical programming in Banach spaces, *Math. Oper. Res.*, **4** (1979).
- [15] S. KOMLÓSI, Quasiconvex first-order approximations and Kuhn-Tucker optimality conditions, *Electron. J. Oper. Res.*, **65** (1993), 327–335.
- [16] S. KOMLÓSI, Monotonicity and quasimonotonicity in nonsmooth analysis, *Recent advances in Nonsmooth Optimization*, World Sci. Publishing, River Edge, NJ, (1995), 193–214.
- [17] S. KOMLÓSI, Generalized monotonicity and generalized convexity, *J. Optim. Appl.*, **84**(2) (1995), 361–376.



On Generalized Monotone Multifunctions with Applications to Optimality Conditions in Generalized Convex Programming

A. Hassouni and A. Jaddar

Title Page

Contents




Go Back

Close

Quit

Page 24 of 26

- [18] S. KOMLÓSI, On the Stampacchia and Minty variational inequality, in: *Generalized Convexity and Optimization for Economics and Financial Decisions*, G. Giorgi and F. Rossi (Eds.), Pitarora editrice, Bogota, 231–260, (1999).
- [19] I.V. KONNOV AND J.C. YAO, On the generalized vector variational inequality problem, *J. Math. Anal. Appl.*, **206** (1997), 42–58.
- [20] D.T. LUC, Characterizations of quasiconvex functions, *Bull. Austral. Math. Soc.*, **48** (1993), 393–405.
- [21] O.L. MANGAZARIAN, Pseudoconvex functions, *SIAM J. Control*, **3** (1965), 281–290.
- [22] J.-P. PENOT, Generalized convex functions in the light of non smooth analysis, *Lecture notes in Economics and Math. Systems*, **429** (1995), Springer Verlag, 269–291.
- [23] J.-P. PENOT, Are generalized derivatives useful for generalized convex functions? , in *Generalized Convexity, Generalized Monotonicity: Recent results*, Crouzeix (Eds.), 3–59, (1998).
- [24] J.-P. PENOT AND P.H. SACH, Generalized monotonicity of subdifferentials and generalized convexity, *J. Optim. Th. Appl.*, **92** (1997).
- [25] J.-P. PENOT AND P.H. QUANG, Generalized convexity of functions and generalized monotonicity of set-valued maps, *J. Optim. Th. Appl.*, **92** (1997), 343–365.



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents




Go Back

Close

Quit

Page 25 of 26

- [26] R.T. ROCKAFELLAR, Generalized directional derivatives and subgradients of nonconvex functions, *Canad. J. Math.*, **32** (1980), 257–280.
- [27] D. ZAGRODNY, Approximate mean value theorem for upper derivatives, *Nonlinear Anal.*, **12** (1988), 1413–1428.



**On Generalized Monotone
Multifunctions with
Applications to Optimality
Conditions in Generalized
Convex Programming**

A. Hassouni and A. Jaddar

Title Page

Contents



Go Back

Close

Quit

Page 26 of 26