

Journal of Inequalities in Pure and Applied Mathematics

ITERATIVE SOLUTION OF NONLINEAR EQUATIONS OF HAMMERSTEIN TYPE

H. ZEGEYE

Bahir Dar University,
P. O. Box. 859,
Bahir Dar, Ethiopia.
EMail: habz@ictp.trieste.it

©2000 Victoria University
ISSN (electronic): 1443-5756
125-03



volume 4, issue 5, article 92,
2003.

*Received 10 August, 2003;
accepted 05 September, 2003.*

Communicated by: R. Verma

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

Suppose X is a real Banach space and $F, K : X \rightarrow X$ are accretive maps. Under different continuity assumptions on F and K such that the inclusion $0 = u + KFu$ has a solution, iterative methods are constructed which converge strongly to such a solution. No invertibility assumption is imposed on K and the operators K and F need not be defined on compact subsets of X . Our method of proof is of independent interest.

2000 Mathematics Subject Classification: 47H04, 47H06, 47H30, 47J05, 47J25

Key words: Accretive operators, Uniformly smooth spaces, Duality maps.

Contents

1	Introduction	3
2	Preliminaries	7
3	Main Results	10
	3.1 Convergence Theorems for Lipschitz Maps	18
	3.2 Convergence Theorems for Uniformly Continuous ϕ -Strongly Accretive Maps	20
	3.3 Explicit Algorithms	23
	References	

Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 2 of 27

1. Introduction

Let X be a real normed linear space with dual X^* . For $1 < q < \infty$, we denote by J_q the generalized duality mapping from X to 2^{X^*} defined by

$$J_q(x) := \{f^* \in X^* : \langle x, f^* \rangle = \|x\| \|f^*\|, \|f^*\| = \|x\|^{q-1}\},$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. If $q = 2$, $J_q = J_2$ and is denoted by J . If X^* is strictly convex, then J_q is single-valued (see e.g., [25]).

A map A with domain $D(A) \subseteq X$ is said to be *accretive* if for every $x, y \in D(A)$ there exists $j(x - y) \in J(x - y)$ such that

$$\langle Ax - Ay, j(x - y) \rangle \geq 0.$$


A is said to be *m-accretive* if it is accretive and $\mathcal{R}(I + \lambda A)$ (range of $(I + \lambda A)$) = X , for all $\lambda > 0$, where I is the identity mapping. A is said to be *ϕ -strongly accretive* if for every $x, y \in D(A)$ there exist $j(x - y) \in J(x - y)$ and a strictly increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$, $\phi(0) = 0$ such that

$$\langle Ax - Ay, j(x - y) \rangle \geq \phi(\|x - y\|) \|x - y\|,$$

and it is *strongly accretive* if for each $x, y \in D(A)$, there exist $j(x - y) \in J(x - y)$ and a constant $k \in (0, 1)$ such that

$$\langle Ax - Ay, j(x - y) \rangle \geq k \|x - y\|^2.$$

Clearly, every strongly accretive map is ϕ -strongly accretive and every ϕ -strongly accretive map is accretive. Closely related to the class of accretive mappings is



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 3 of 27


the class of pseudocontractive mappings. A mapping $T : X \rightarrow X$ is said to be *pseudocontractive* if and only if $A := I - T$ is accretive. One can easily show that the fixed point of pseudocontractive mapping T is the zero of accretive mapping $A := I - T$. If X is a Hilbert space, accretive operators are also called *monotone*. The accretive mappings were introduced independently in 1967 by Browder [3] and Kato [20]. Interest in such mappings stems mainly from their firm connection with equations of evolution. It is known (see e.g., [28]) that many physically significant problems can be modelled by initial-value problems of the form

$$(1.1) \quad x'(t) + Ax(t) = 0, \quad x(0) = x_0,$$

where A is an accretive operator in an appropriate Banach space. Typical examples where such evolution equations occur can be found in the heat, wave or Schrödinger equations. One of the fundamental results in the theory of accretive operators, due to Browder [4], states that if A is locally Lipschitzian and accretive then A is m -accretive which immediately implies that the equation $x + Ax = h$ has a solution $x^* \in D(A)$ for any $h \in X$. This result was subsequently generalized by Martin [22] to the continuous accretive operators. If in (1.1), $x(t)$ is independent of t , then (1.1) reduces to

$$(1.2) \quad Au = 0,$$

whose solutions correspond to the equilibrium points of the system (1.1). Consequently, considerable research efforts have been devoted, especially within the past 20 years or so, to methods of finding approximate solutions (when they exist) of equation (1.2). One important generalization of equation (1.2) is the



**Iterative Solution of Nonlinear
Equations of Hammerstein Type**

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 4 of 27

so-called *equation of Hammerstein type* (see e.g., [18]), where a nonlinear integral equation of Hammerstein type is one of the form:

$$(1.3) \quad u(x) + \int_{\Omega} K(x, y)f(y, u(y))dy = h(x),$$

where dy is a σ -finite measure on the measure space Ω ; the real kernel K is defined on $\Omega \times \Omega$, f is a real-valued function defined on $\Omega \times \mathfrak{R}$ and is, in general, nonlinear and h is a given function on Ω . If we now define an operator K by

$$Kv(x) := \int_{\Omega} K(x, y)v(y)dy; \quad x \in \Omega,$$

and the so-called *superposition* or *Nemytskii* operator by $Fu(y) := f(y, u(y))$ then, the integral equation (1.3) can be put in operator theoretic form as follows:

$$(1.4) \quad u + KFu = 0,$$

where, without loss of generality, we have taken $h \equiv 0$. We note that if K is an arbitrary accretive map (not necessarily the identity), then $A := I + KF$ need not be accretive. Interest in equation (1.4) stems mainly from the fact that several problems that arise in differential equations, for instance, elliptic boundary value problems whose linear parts possess Greens functions can, as a rule, be transformed into the form (1.4) (see e.g., [23, Chapter IV]). Equations of Hammerstein type play a crucial role in the theory of optimal control systems (see e.g., [17]). Several existence and uniqueness theorems have been proved for equations of the Hammerstein type (see e.g., [2, 5, 6, 8, 15]).



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back


Close

Quit

Page 5 of 27

For the iterative approximation of solutions of *equation (1.2)*, the *accretivity/monotonicity* of A is crucial. The Mann iteration scheme (see e.g., [21]) and the Ishikawa iteration scheme (see e.g., [19]) have successfully been employed (see e.g., [7, 10, 11, 12, 13, 14, 16, 19, 21, 24, 27]). Attempts to apply these schemes to equation (1.4) have not provided satisfactory results. In particular, the recursion formulas obtained involved K^{-1} and this is not convenient in applications. Part of the difficulty is, as has already been noted, the fact that the composition of two accretive operators need not be accretive. In the special case in which the operators are defined on subsets D of X which are compact (or more generally, *angle-bounded* (see e.g., [1]), Brèzis and Browder [1] have proved the strong convergence of a suitably defined Galerkin approximation to a solution of (1.4).

It is our purpose in this paper to use the method introduced in [12] which contains an auxiliary operator, defined in terms of K and F in an arbitrary real Banach space which, under certain conditions, is accretive whenever K and F are, and whose zeros are solutions of equation (1.4). Moreover, the operators K and F need not be defined on a compact or angle-bounded subset of X . Furthermore, our method which does not involve K^{-1} provides an explicit algorithm for the computation of solutions of equation (1.4).



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 6 of 27

2. Preliminaries

Let X be a real normed linear space of dimension ≥ 2 . The *modulus of smoothness* of X is defined by:

$$\rho_X(\tau) := \sup \left\{ \frac{\|x+y\| + \|x-y\|}{2} - 1 : \|x\| = 1, \|y\| = \tau \right\}; \quad \tau > 0.$$

It is well known that $\rho_X(\tau) \leq \tau \forall \tau > 0$ (see e.g., [26]). If $\rho_X(\tau) > 0 \forall \tau > 0$, then X is said to be *smooth*. If there exist a constant $c > 0$ and a real number $1 < q < \infty$, such that $\rho_X(\tau) \leq c\tau^q$, then X is said to be *q-uniformly smooth*. A Banach space X is called *uniformly smooth* if $\lim_{\tau \rightarrow 0} \frac{\rho_X(\tau)}{\tau} = 0$. If E is a real uniformly smooth Banach space, then

$$(2.1) \quad \|x+y\|^2 \leq \|x\|^2 + 2\langle y, j(x) \rangle + D \max \left\{ \|x\| + \|y\|, \frac{c}{2} \right\} \rho_X(\|y\|),$$


for every $x, y \in X$, where D and c are positive constants (see e.g., [26]). Typical examples of such uniformly smooth spaces are the Lebesgue L_p , the sequence ℓ_p and the Sobolev W_p^m spaces for $1 < p < \infty$. Moreover, we have

$$(2.2) \quad \rho_{L_p}(\tau) = \rho_{L^p}(\tau) = \rho_{W_p^m}(\tau) \leq \begin{cases} \frac{1}{p}\tau^p, & \text{if } 1 < 2 < p; \\ \frac{p-1}{2}\tau^2, & \text{if } p \geq 2, \end{cases}$$

$\forall \tau > 0$ (see e.g., [26]).

In the sequel we shall need the following results.

Theorem 2.1. [25]. *Let $q > 1$ and X be a real smooth Banach space. Then the following are equivalent.*



Title Page

Contents



Go Back

Close

Quit

Page 7 of 27

1. X is uniformly smooth.

2. There exists a continuous, strictly increasing and convex function $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that for every $x, y \in B_r$ for some $r > 0$ we get

$$(2.3) \quad \|x + y\|^q \leq \|x\|^q + q \langle y, j_q(x) \rangle + g(\|y\|).$$


Lemma 2.2. (see, e.g., [13]). Let X be a normed linear space and J be the normalized duality map on E . Then for any given $x, y \in X$, the following inequality holds:

$$\|x + y\|^2 \leq \|x\|^2 + 2 \langle y, j(x + y) \rangle, \quad \forall j(x + y) \in J(x + y).$$

Theorem 2.3. [9]. Let X be a real Banach space, $A : X \rightarrow X$ be a Lipschitz and strongly accretive map with Lipschitz constant $L > 0$ and strong accretivity constant $\lambda \in (0, 1)$. Assume that $Ax = 0$ has a solution $x^* \in X$. Define $A_\varepsilon : X \rightarrow X$ by $A_\varepsilon x := x - \varepsilon Ax$ for $x \in X$ where $\varepsilon := \frac{1}{2} \left\{ \frac{\lambda}{1+L(3+L-\lambda)} \right\}$. For arbitrary $x_0 \in X$, define the Picard sequence $\{x_n\}$ in X by $x_{n+1} = A_\varepsilon x_n, n \geq 0$. Then, $\{x_n\}$ converges strongly to x^* with $\|x_{n+1} - x^*\| \leq \delta^n \|x_1 - x^*\|$ where $\delta := (1 - \frac{1}{2}\lambda\varepsilon) \in (0, 1)$. Moreover, x^* is unique.

Theorem 2.4. [13] Let X be a real normed linear space. Let $A : X \rightarrow X$ be uniformly continuous ϕ -strongly accretive mapping. Assume $0 = Ax$ has a solution $x^* \in X$. Then, there exists a real number $\gamma_0 > 0$ such that if the real sequence $\{\alpha_n\} \subset [0, \gamma_0]$ satisfies the following conditions:

(i) $\lim \alpha_n = 0$;



Title Page

Contents



Go Back

Close

Quit

Page 8 of 27


$$(ii) \sum \alpha_n = \infty,$$

then for arbitrary $x_0 \in X$ the sequence $\{x_n\}$, defined by

$$x_{n+1} := x_n - \alpha_n Ax_n, \quad n \geq 0,$$

converges strongly to x^* , the unique solution of $0 = Ax$.

We note that Theorem 2.4 is Theorem 3.6 of [13] with A ϕ -strongly accretive mapping.



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 9 of 27

3. Main Results

Lemma 3.1. For $q > 1$, let X be a real uniformly smooth Banach space. Let $E := X \times X$ with norm

$$\|z\|_E := \left(\|u\|_X^q + \|v\|_X^q \right)^{\frac{1}{q}},$$

for arbitrary $z = [u, v] \in E$. Let $E^* := X^* \times X^*$ denote the dual space of E . For arbitrary $x = [x_1, x_2] \in E$ define the map $j_q^E : E \rightarrow E^*$ by

$$j_q^E(x) = j_q^E[x_1, x_2] := [j_q^X(x_1), j_q^X(x_2)],$$

so that for arbitrary $z_1 = [u_1, v_1], z_2 = [u_2, v_2]$ in E the duality pairing $\langle \cdot, \cdot \rangle$ is given by


$$\langle z_1, j_q^E(z_2) \rangle = \langle u_1, j_q^X(u_2) \rangle + \langle v_1, j_q^X(v_2) \rangle.$$

Then

- (a) E is uniformly smooth;
- (b) j_q^E is a single-valued duality mapping on E .

Proof. (a) Let $x = [x_1, x_2], y = [y_1, y_2]$ be arbitrary elements of E . It suffices to show that x and y satisfy condition (2) of Theorem 2.1. We compute as follows:

$$\begin{aligned} \|x + y\|_E^q &= \|[x_1 + y_1, x_2 + y_2]\|_E^q \\ &= \|x_1 + y_1\|_X^q + \|x_2 + y_2\|_X^q \\ &\leq \|x_1\|_X^q + \|x_2\|_X^q + g(\|y_1\|) + g(\|y_2\|) \\ &\quad + q \left\{ \langle y_1, j_q^X(x_1) \rangle + \langle y_2, j_q^X(x_2) \rangle \right\}, \end{aligned}$$



Title Page

Contents



Go Back

Close

Quit

Page 10 of 27



Title Page

Contents



Go Back

Close

Quit

Page 11 of 27

where g is continuous, strictly increasing and a convex function (using (2) of Theorem 2.1, since X is uniformly smooth). It follows that

$$\|x + y\|_E^q \leq \|x\|_E^q + q \langle y, j_q^E(x) \rangle + g'(\|y\|),$$

where $g'(\|y\|) := g(\|y_1\|) + g(\|y_2\|)$. So, the result follows from Theorem 2.1.

(b) For arbitrary $x = [x_1, x_2] \in E$, let $j_q^E(x) = j_q^E[x_1, x_2] = \psi_q$. Then $\psi_q = [j_q^X(x_1), j_q^X(x_2)]$ in E^* . Observe that for $p > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$,

$$\begin{aligned} \|\psi_q\|_{E^*} &= \left(\|[j_q^X(x_1), j_q^X(x_2)]\| \right)^{\frac{1}{p}} \\ &= \left(\|j_q(x_1)\|_{X^*}^p + \|j_q(x_2)\|_{X^*}^p \right)^{\frac{1}{p}} \\ &= \left(\|x_1\|_X^{(q-1)p} + \|x_2\|_X^{(q-1)p} \right)^{\frac{1}{p}} \\ &= \left(\|x_1\|_X^q + \|x_2\|_X^q \right)^{\frac{q-1}{q}} = \|x\|_X^{q-1}. \end{aligned}$$

Hence, $\|\psi_q\|_{E^*} = \|x\|_E^{q-1}$. Furthermore,

$$\begin{aligned} \langle x, \psi_q \rangle &= \langle [x_1, x_2], [j_q^X(x_1), j_q^X(x_2)] \rangle \\ &= \langle x_1, j_q^X(x_1) \rangle + \langle x_2, j_q^X(x_2) \rangle \\ &= \|x_1\|_X^q + \|x_2\|_X^q \\ &= \left(\|x_1\|_X^q + \|x_2\|_X^q \right)^{\frac{1}{q}} \left(\|x_1\|_X^q + \|x_2\|_X^q \right)^{\frac{q-1}{q}} \\ &= \|x\|_E \cdot \|\psi\|_{E^*}^{q-1}. \end{aligned}$$

Hence, j_q^E is a single-valued (since E is uniformly smooth) duality mapping on E . □

Lemma 3.2. *Suppose X is a real normed linear space. Let $F, K : X \rightarrow X$ be maps such that the following conditions hold:*

(i) *For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_1 : [0, \infty) \rightarrow [0, \infty)$, $\phi_1(0) = 0$ such that*

$$\langle Fu_1 - Fu_2, j(u_1 - u_2) \rangle \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(ii) *For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_2 : [0, \infty) \rightarrow [0, \infty)$, $\phi_2(0) = 0$ such that*


$$\langle Ku_1 - Ku_2, j(u_1 - u_2) \rangle \geq \phi_2(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(iii) $\phi_i(t) \geq (2 + r_i)t$ for all $t \in (0, \infty)$ and for some $r_i > 0$, $i = 1, 2$.

Let $E := X \times X$ with norm $\|z\|_E^2 = \|u\|_X^2 + \|v\|_X^2$ for $z = (u, v) \in E$ and define a map $T : E \rightarrow E$ by $Tz := T(u, v) = (Fu - v, u + Kv)$. Then for each $z_1, z_2 \in E$ there exist $j^E(z_1 - z_2) \in J^E(z_1 - z_2)$ and a strictly increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = 0$ such that

$$\langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \geq \phi(\|z_1 - z_2\|)\|z_1 - z_2\|.$$

Proof. Define $\phi : [0, \infty) \rightarrow [0, \infty)$ by $\phi(t) := \min\{r_1, r_2\}t$ for each $t \in [0, \infty)$. Clearly, ϕ is a strictly increasing function with $\phi(0) = 0$. Furthermore,



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 12 of 27

observe that for $z_1 = (u_1, v_1)$ and $z_2 = (u_2, v_2)$ arbitrary elements in E we have $\langle z_1, j^E(z_2) \rangle = \langle u_1, j(u_2) \rangle + \langle v_1, j(v_2) \rangle$. Thus we have the following estimates:

$$\begin{aligned}
 & \langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \\
 &= \langle Fu_1 - Fu_2 - (v_1 - v_2), j(u_1 - u_2) \rangle \\
 & \quad + \langle Kv_1 - Kv_2 + (u_1 - u_2), j(v_1 - v_2) \rangle \\
 &= \langle Fu_1 - Fu_2, j(u_1 - u_2) \rangle - \langle v_1 - v_2, j(u_1 - u_2) \rangle \\
 & \quad + \langle Kv_1 - Kv_2, j(v_1 - v_2) \rangle + \langle u_1 - u_2, j(v_1 - v_2) \rangle \\
 &\geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\
 (3.1) \quad & \quad - \langle v_1 - v_2, j(u_1 - u_2) \rangle + \langle u_1 - u_2, j(v_1 - v_2) \rangle.
 \end{aligned}$$

Since X is an arbitrary real normed linear space, for each $x, y \in X$ and $j(x + y) \in J(x + y)$ (by Lemma 2.2) we have that

$$\begin{aligned}
 \|x + y\|^2 &\leq \|x\|^2 + 2\langle y, j(x + y) - j(x) \rangle + 2\langle y, j(x) \rangle \\
 &\leq \|x\|^2 + 2\|y\|\|j(x + y) - j(x)\| + 2\langle y, j(x) \rangle \\
 &\leq \|x\|^2 + 2\|y\|(\|x + y\| + \|x\|) + 2\langle y, j(x) \rangle \\
 &\leq \|x\|^2 + 2\left(\frac{\|y\|^2}{2} + \frac{\|x + y\|^2}{2} + \frac{\|y\|^2}{2} + \frac{\|x\|^2}{2}\right) + 2\langle y, j(x) \rangle \\
 &= 2\|x\|^2 + 2\|y\|^2 + \|x + y\|^2 + 2\langle y, j(x) \rangle.
 \end{aligned}$$

Thus we get $\langle y, j(x) \rangle \geq -\|x\|^2 - \|y\|^2$.

Replacing y by $-y$ we obtain $-\langle y, j(x) \rangle \geq -\|x\|^2 - \|y\|^2$. Therefore,

$$\begin{aligned}
 -\langle v_1 - v_2, j(u_1 - u_2) \rangle &\geq -\|u_1 - u_2\|^2 - \|v_1 - v_2\|^2 \text{ and} \\
 \langle u_1 - u_2, j(v_1 - v_2) \rangle &\geq -\|v_1 - v_2\|^2 - \|u_1 - u_2\|^2.
 \end{aligned}$$



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 13 of 27

Thus (3.1) and the above estimates give that

$$\begin{aligned}
 & \langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \\
 & \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\
 & \quad - 2\|u_1 - u_2\|^2 - 2\|v_1 - v_2\|^2 \\
 & \geq \left(\phi_1(\|u_1 - u_2\|) - 2\|u_1 - u_2\| \right) \|u_1 - u_2\| \\
 & \quad + \left(\phi_2(\|v_1 - v_2\|) - 2\|v_1 - v_2\| \right) \|v_1 - v_2\| \\
 & \geq r_1\|u_1 - u_2\|^2 + r_2\|v_1 - v_2\|^2 \\
 & \geq \min\{r_1, r_2\} \left\{ \|u_1 - u_2\|^2 + \|v_1 - v_2\|^2 \right\} \\
 & = \min\{r_1, r_2\} \|z_1 - z_2\|^2 \\
 & = \phi(\|z_1 - z_2\|)\|z_1 - z_2\|,
 \end{aligned}$$

completing the proof of Lemma 3.2. □


Lemma 3.3. *Suppose X is a real uniformly smooth Banach space. Let $F, K : X \rightarrow X$ be maps such that the following conditions hold:*

(i) *For each $u_1, u_2 \in X$ there exists a strictly increasing function $\phi_1 : [0, \infty) \rightarrow [0, \infty)$, $\phi_1(0) = 0$ such that*

$$\langle Fu_1 - Fu_2, j(u_1 - u_2) \rangle \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(ii) *For each $u_1, u_2 \in X$ there exists a strictly increasing function $\phi_2 : [0, \infty) \rightarrow [0, \infty)$, $\phi_2(0) = 0$ such that*

$$\langle Ku_1 - Ku_2, j(u_1 - u_2) \rangle \geq \phi_2(\|u_1 - u_2\|)\|u_1 - u_2\|;$$



Title Page

Contents



Go Back

Close

Quit

Page 14 of 27

(iii) $\phi_i(t) \geq (D + r_i)t + \frac{acD}{4}t^{q-1}$, $\rho_X(t) \leq at^q$ for all $t \in (0, \infty)$ and for some $q > 1$, $a > 0$ and $r_i > 0$, $i = 1, 2$, where c and D are the constants appearing in inequality (2.1).

Let E and T be defined as in Lemma 3.2. Then for each $z_1, z_2 \in E$ there exists a strictly increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = 0$ such that


$$\langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \geq \phi(\|z_1 - z_2\|)\|z_1 - z_2\|.$$

Proof. Define $\phi : [0, \infty) \rightarrow [0, \infty)$ by $\phi(t) := \min\{r_1, r_2\}t$ for each $t \in [0, \infty)$. Thus as in the proof of Lemma 3.2 we have that ϕ is a strictly increasing function with $\phi(0) = 0$ and for $z_1 = (u_1, v_1)$ and $z_2 = (u_2, v_2)$ arbitrary elements in E we have the following estimate:

$$\begin{aligned} (3.2) \quad & \langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \\ &= \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\ &\quad - \langle v_1 - v_2, j(u_1 - u_2) \rangle + \langle u_1 - u_2, j(v_1 - v_2) \rangle. \end{aligned}$$

Since X is uniformly smooth for each $x, y \in X$ by (2.1) we have that

$$\begin{aligned} \|x + y\|^2 &\leq \|x\|^2 + 2\langle y, j(x) \rangle + D \max\left\{\|x\| + \|y\|, \frac{c}{2}\right\} \rho_X(\|y\|) \\ &\leq \|x\|^2 + 2\langle y, j(x) \rangle + D \left\{\|x\| + \|y\| + \frac{c}{2}\right\} \rho_X(\|y\|) \\ &\leq \|x\|^2 + 2\langle y, j(x) \rangle + D \left(\|x\|\|y\| + \|y\|^2 + \frac{c}{2}\rho_X(\|y\|)\right) \\ &\text{(since } \rho_X(\|y\|) \leq \|y\|) \end{aligned}$$



Title Page

Contents



Go Back

Close

Quit

Page 15 of 27

$$\leq \|x\|^2 + 2\langle y, j(x) \rangle + D \left(\frac{\|x\|^2}{2} + \frac{\|y\|^2}{2} + \|y\|^2 + \frac{ac}{2}\|y\|^q \right)$$

(since $\rho_X(\|y\|) \leq a\|y\|^q$ by assumption for $q > 1$ and $a > 0$)

$$\leq \left(1 + \frac{D}{2}\right) \|x\|^2 + \frac{3D}{2}\|y\|^2 + \frac{acD}{2}\|y\|^q + 2\langle y, j(x) \rangle,$$

and hence


$$\langle y, j(x) \rangle \geq \frac{1}{2}\|x + y\|^2 - \frac{1}{2} \left(\left(1 + \frac{D}{2}\right) \|x\|^2 + \frac{3D}{2}\|y\|^2 + \frac{acD}{2}\|y\|^q \right).$$

Replacing y by $-y$ we obtain

$$-\langle y, j(x) \rangle \geq \frac{1}{2}\|x - y\|^2 - \frac{1}{2} \left(\left(1 + \frac{D}{2}\right) \|x\|^2 + \frac{3D}{2}\|y\|^2 + \frac{acD}{2}\|y\|^q \right).$$

Thus (3.2) and the above estimates give that

$$\begin{aligned} & \langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle \\ & \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\ & \quad + \frac{1}{2} \left(\|u_1 - u_2 - (v_1 - v_2)\|^2 - \left(1 + \frac{D}{2}\right) \|u_1 - u_2\|^2 \right. \\ & \quad \left. - \frac{3D}{2}\|v_1 - v_2\|^2 - \frac{acD}{2}\|v_1 - v_2\|^q \right) \\ & \quad + \frac{1}{2} \left(\|u_1 - u_2 + v_1 - v_2\|^2 - \left(1 + \frac{D}{2}\right) \|v_1 - v_2\|^2 \right. \\ & \quad \left. - \frac{3D}{2}\|u_1 - u_2\|^2 - \frac{acD}{2}\|u_1 - u_2\|^q \right) \end{aligned}$$



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents




Go Back

Close

Quit

Page 16 of 27



Title Page

Contents



Go Back

Close

Quit

Page 17 of 27

$$\begin{aligned}
 &\geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\
 &\quad + \frac{1}{2} (\|u_1 - u_2 - (v_1 - v_2)\|^2 + \|u_1 - u_2 + v_1 - v_2\|^2) \\
 &\quad - \frac{1}{2} \left((1 + 2D)\|u_1 - u_2\|^2 + \frac{acD}{2}\|u_1 - u_2\|^q \right) \\
 (3.3) \quad &\quad - \frac{1}{2} \left((1 + 2D)\|v_1 - v_2\|^2 + \frac{acD}{2}\|v_1 - v_2\|^q \right).
 \end{aligned}$$

Since for all $x, y \in X, x \neq y$,

$$\left\| \frac{x + y}{2} \right\|^2 \leq \frac{1}{2} (\|x\|^2 + \|y\|^2)$$

we have that

$$\| (u_1 - u_2) - (v_1 - v_2) \|^2 + \| (u_1 - u_2) + (v_1 - v_2) \|^2 \geq \|u_1 - u_2\|^2 + \|v_1 - v_2\|^2.$$

Then (3.3) becomes

$$\begin{aligned}
 \langle Tz_1 - Tz_2, j^E(z_1 - z_2) \rangle &\geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\| - \left(D\|u_1 - u_2\|^2 \right. \\
 &\quad \left. + \frac{acD}{4}\|u_1 - u_2\|^q \right) + \phi_2(\|v_1 - v_2\|)\|v_1 - v_2\| \\
 &\quad - \left(D\|v_1 - v_2\|^2 + \frac{acD}{4}\|v_1 - v_2\|^q \right) \\
 &\geq r_1\|u_1 - u_2\|^2 + r_2\|v_1 - v_2\|^2 \\
 &\geq \min\{r_1, r_2\} \{ \|u_1 - u_2\|^2 + \|v_1 - v_2\|^2 \} \\
 &= \min\{r_1, r_2\} \|z_1 - z_2\|^2 = \phi(\|z_1 - z_2\|)\|z_1 - z_2\|,
 \end{aligned}$$

completing the proof of Lemma 3.3. □

3.1. Convergence Theorems for Lipschitz Maps

Remark 3.1. *If K and F are Lipschitz single-valued maps with Lipschitz constants L_K and L_F respectively, then T is a Lipschitz map with constant $L := \left(d \max\{L_F^2 + 1, L_K^2 + 1\}\right)^{\frac{1}{2}}$ for some constant $d > 0$. Indeed, if $z_1 = (u_1, v_1)$, $z_2 = (u_2, v_2)$ in E then we have that*

$$\begin{aligned} & \|Tz_1 - Tz_2\|^2 \\ &= \|(Fu_1 - Fu_2) - (v_1 - v_2)\|^2 + \|u_1 - u_2 + Kv_1 - Kv_2\|^2 \\ &\leq \left(L_F\|u_1 - u_2\| + \|v_1 - v_2\|\right)^2 + \left(\|u_1 - u_2\| + L_K\|v_1 - v_2\|\right)^2 \\ &\leq d\left(L_F^2\|u_1 - u_2\|^2 + \|v_1 - v_2\|^2 + \|u_1 - u_2\|^2 + L_K^2\|v_1 - v_2\|^2\right) \\ &\text{for some } d > 0 \\ &\leq d \max\{L_F^2 + 1, L_K^2 + 1\} \left(\|u_1 - u_2\|^2 + \|v_1 - v_2\|^2\right) \\ &= d \max\{L_F^2 + 1, L_K^2 + 1\} \|z_1 - z_2\|^2. \end{aligned}$$

Thus $\|Tz_1 - Tz_2\| \leq L\|z_1 - z_2\|$. Consequently, we have the following theorem.

Theorem 3.4. *Let X be real Banach space. Let $F, K : X \rightarrow X$ be Lipschitzian maps with Lipschitz constants L_K and L_F , respectively such that the following conditions hold:*



Iterative Solution of Nonlinear
Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 18 of 27

(i) For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_1 : [0, \infty) \rightarrow [0, \infty)$, $\phi_1(0) = 0$ such that

$$\langle Fu_1 - Fu_2, j(u_1 - u_2) \rangle \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\|;$$


(ii) For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_2 : [0, \infty) \rightarrow [0, \infty)$, $\phi_2(0) = 0$ such that

$$\langle Ku_1 - Ku_2, j(u_1 - u_2) \rangle \geq \phi_2(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(iii) $\phi_i(t) \geq (2 + r_i)t$ for all $t \in (0, \infty)$ and for some $r_i > 0$, $i = 1, 2$ and let $\gamma := \min\{r_1, r_2\}$.

Assume that $u + KF u = 0$ has a solution u^* in X and let $E := X \times X$ and $\|z\|_E^2 = \|u\|_X^2 + \|v\|_X^2$ for $z = (u, v) \in E$ and define the map $T : E \rightarrow E$ by $Tz := T(u, v) = (Fu - v, Kv + u)$. Let L denote the Lipschitz constant of T and $\varepsilon := \frac{1}{2} \left(\frac{\gamma}{1+L(3+L-\gamma)} \right)$. Define the map $A_\varepsilon : E \rightarrow E$ by $A_\varepsilon z := z - \varepsilon Tz$ for each $z \in E$. For arbitrary $z_0 \in E$, define the Picard sequence $\{z_n\}$ in E by $z_{n+1} := A_\varepsilon z_n$, $n \geq 0$. Then $\{z_n\}$ converges strongly to $z^* = [u^*, v^*]$ the unique solution of the equation $Tz = 0$ with $\|z_{n+1} - z^*\| \leq \delta^n \|z_1 - z^*\|$, where $\delta := (1 - \frac{1}{2}\gamma\varepsilon) \in (0, 1)$.

Proof. Observe that u^* is a solution of $u + KF u = 0$ if and only if $z^* = [u^*, v^*]$ is a solution of $Tz = 0$. Hence $Tz = 0$ has a solution $z^* = [u^*, v^*]$ in E . Since T is Lipschitz and by Lemma 3.2 it is strongly accretive with constant γ (which, without loss of generality, we may assume is in $(0, 1)$), the conclusion follows from Theorem 2.3. \square



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 19 of 27

Following the method of the proof of Theorem 3.4 and making use of Lemma 3.3 instead of Lemma 3.2 we obtain the following theorem.

Theorem 3.5. *Let X be a real uniformly smooth Banach space. Let $F, K : X \rightarrow X$ be Lipschitzian maps with Lipschitz constants L_K and L_F , respectively such that conditions (i)-(iii) of Lemma 3.3 are satisfied and let $\gamma := \min\{r_1, r_2\}$. Assume that $u + KF u = 0$ has the solution u^* and set E and T as in Theorem 3.4. Let $L, \varepsilon, A_\varepsilon$, and $\{z_n\}$ be defined as in Theorem 3.4. Then the conclusion of Theorem 3.4 holds.*

3.2. Convergence Theorems for Uniformly Continuous ϕ - Strongly Accretive Maps

Theorem 3.6. *Let X be a real normed linear space. Let $F, K : X \rightarrow X$ be uniformly continuous maps such that the following conditions hold:*


(i) *For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_1 : [0, \infty) \rightarrow [0, \infty)$, $\phi_1(0) = 0$ such that*

$$\langle F u_1 - F u_2, j(u_1 - u_2) \rangle \geq \phi_1(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(ii) *For each $u_1, u_2 \in X$ there exist $j(u_1 - u_2) \in J(u_1 - u_2)$ and a strictly increasing function $\phi_2 : [0, \infty) \rightarrow [0, \infty)$, $\phi_2(0) = 0$ such that*

$$\langle K u_1 - K u_2, j(u_1 - u_2) \rangle \geq \phi_2(\|u_1 - u_2\|)\|u_1 - u_2\|;$$

(iii) $\phi_i(t) \geq (2 + r_i)t$ for all $t \in (0, \infty)$ and for some $r_i > 0, i = 1, 2$.



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents




Go Back

Close

Quit

Page 20 of 27



Title Page

Contents



Go Back

Close

Quit

Page 21 of 27

Assume that $0 = u + KF u$ has a solution u^* in X . Let $E := X \times X$ and $\|z\|_E^2 = \|u\|_X^2 + \|v\|_X^2$ for $z = (u, v) \in E$ and define the map $T : E \rightarrow E$ by $Tz := T(u, v) = (Fu - v, u + Kv)$. Then there exists a real number $\gamma_0 > 0$ such that if the real sequence $\{\alpha_n\} \subset [0, \gamma_0]$ satisfies the following conditions

(a) $\lim_{n \rightarrow \infty} \alpha_n = 0$;

(b) $\sum \alpha_n = \infty$,

then for arbitrary $z_0 \in E$ the sequence $\{z_n\}$, defined by

$$z_{n+1} := z_n - \alpha_n T z_n, \quad n \geq 0,$$

converges strongly to $z^* = [u^*, v^*]$, where u^* is the unique solution of $0 = u + KF u$.

Proof. Since K and F are uniformly continuous maps we have that T is a uniformly continuous map. Observe also that u^* is the solution of $0 = u + KF u$ in X if and only if $z^* = [u^*, v^*]$ is a solution of $0 = Tz$ in E . Thus we obtain that $N(T)$ (null space of T) $\neq \emptyset$. Also by Lemma 3.2, T is ϕ -strongly accretive. Therefore the conclusion follows from Theorem 2.4. \square

Following the method of proof of Theorem 3.6 and making use of Lemma 3.3 instead of Lemma 3.2 we obtain the following theorem.

Theorem 3.7. *Let X be a real uniformly smooth Banach space. Let $F, K : X \rightarrow X$ be uniformly continuous maps such that conditions (i)-(iii) of Theorem 3.5 are satisfied. Assume that $0 = u + KF u$ has a solution u^* in X . Let E, T and $\{z_n\}$ be defined as in Theorem 3.6. Then, the conclusion of Theorem 3.6 holds.*

Remark 3.2. We note that for the special case in which the real Banach space X is q -uniformly smooth using the above method, the author and Chidume [12] proved the following theorem.

Theorem 3.8. [12]. Let X be a real q -uniformly smooth Banach space. Let $F, K : X \rightarrow X$ be Lipschitzian maps with positive constants L_K and L_F respectively with the following conditions:

(i) There exists $\alpha > 0$ such that

$$\langle Fu_1 - Fu_2, j_q(u_1 - u_2) \rangle \geq \alpha \|u_1 - u_2\|^q, \quad \forall u_1, u_2 \in D(F);$$


(ii) There exists $\beta > 0$ such that

$$\langle Ku_1 - Ku_2, j_q(u_1 - u_2) \rangle \geq \beta \|u_1 - u_2\|^q, \quad \forall u_1, u_2 \in D(K);$$

(iii) $\alpha, \beta > d := q^{-1}(1 + d_q - c^{-1}2^{q-1})$ and $\gamma := \min\{\alpha - d, \beta - d\}$ where d_q and c are as in (3.2) and (2.1) of [12], respectively.

Assume that $u + KF u = 0$ has solution u^* and set E and T as in Theorem 3.4. Let L be a Lipschitz constant of T and ε , A_ε and z_n be defined as in Theorem 3.4. Then $\{z_n\}$ converges strongly to $z^* = [u^*, v^*]$ the unique solution of the equation $Tz = 0$ with $\|z_{n+1} - z^*\| \leq \delta^n \|z_1 - z^*\|$, where u^* is the solution of the equation $u + KF u = 0$ and $\delta := (1 - \frac{1}{2}\gamma\varepsilon) \in (0, 1)$.

The cases for Hilbert spaces and L_p spaces ($1 < p < \infty$) are easily deduced from Theorem 3.8. The theorems proved in this paper are analogues of the theorems in [12] for the more general real Banach spaces considered here.



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 22 of 27

3.3. Explicit Algorithms

The method of our proofs provides the following explicit algorithms for computing the solution of the inclusion $0 = u + KF u$ in the space X .

- (a) For Lipschitz operators (Theorem 3.4 and Theorem 3.5) with initial values $u_0, v_0 \in X$, define the sequences $\{u_n\}$ and $\{v_n\}$ in X as follows:


$$u_{n+1} = u_n - \varepsilon (F u_n - v_n);$$
$$v_{n+1} = v_n - \varepsilon (K v_n + u_n).$$

Then $u_n \rightarrow u^*$ in X , the unique solution u^* of $0 = u + KF u$, where ε is as defined in Theorem 3.4.

- (b) For uniformly continuous operators (Theorem 3.6 and Theorem 3.7) with initial values $u_0, v_0 \in X$, define the sequences $\{u_n\}$ and $\{v_n\}$ in X as follows:

$$u_{n+1} = u_n - \alpha_n (F u_n - v_n);$$
$$v_{n+1} = v_n - \alpha_n (K v_n + u_n).$$

Then $u_n \rightarrow u^*$ in X , the unique solution u^* of $0 = u + KF u$, where α_n is as defined in Theorem 3.6.



Iterative Solution of Nonlinear
Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back


Close

Quit

Page 23 of 27

References

- [1] H. BREŽIS AND F.E. BROWDER, Some new results about Hammerstein equations, *Bull. Amer. Math. Soc.*, **80** (1974), 567–572.
- [2] H. BREŽIS AND F.E. BROWDER, Existence theorems for nonlinear integral equations of Hammerstein type, *Bull. Amer. Math. Soc.*, **81** (1975), 73–78.
- [3] F.E. BROWDER, Nonlinear mappings of nonexpansive and accretive type in Banach spaces, *Bull. Amer. Math. Soc.*, **73** (1967), 875–882.
- [4] F.E. BROWDER, Nonlinear monotone and accretive operators in Banach space, *Proc. Nat. Acad. Sci. U.S.A.*, **61** (1968), 388–393.
- [5] F.E. BROWDER, D.G. De FIGUEIREDO AND P. GUPTA, Maximal monotone operators and a nonlinear integral equations of Hammerstein type, *Bull. Amer. Math. Soc.*, **76** (1970), 700–705.
- [6] F.E. BROWDER AND P. GUPTA, Monotone Operators and nonlinear integral equations of Hammerstein type, *Bull. Amer. Math. Soc.*, **75** (1969), 1347–1353.
- [7] R.E. BRUCK, The iterative solution of the equation $f \in x + Tx$ for a monotone operator T in Hilbert space, *Bull. Amer. Math. Soc.*, **79** (1973), 1258–1262.
- [8] R.S. CHEPANOVICH, Nonlinear Hammerstein equations and fixed points, *Publ. Inst. Math. (Beograd) N. S.*, **35**(49) (1984), 119–123.



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 24 of 27

- [9] C.E. CHIDUME, Nonexpansive mappings, generalizations and iterative algorithms, Accepted to appear in *Nonlinear Analysis and Applications*.
- [10] C.E. CHIDUME AND H. ZEGEYE, Approximation of the zeros of nonlinear m -Accretive operators, *Nonlinear Analysis*, **37** (1999), 81–96.
- [11] C.E. CHIDUME AND H. ZEGEYE, Global Iterative Schemes for Accretive Operators, *J. Math. Anal. Appl.*, **257** (2001), 364–377.
- [12] C.E. CHIDUME AND H. ZEGEYE, Iterative approximation of solutions of nonlinear equations of Hammerstein type, *Abstract and Applied Analysis*, **6** (2003), 353–365.
- [13] C.E. CHIDUME AND H. ZEGEYE, Approximation methods for nonlinear operator equations, *Proc. Amer. Math. Soc.*, **131** (2003), 2467–2478.
- [14] M.G. CRANDALL AND A. PAZY, On the range of accretive operators, *Israel J. Math.*, **27**(3-4) (1977), 235–246.
- [15] D.G. De FIGUEIREDO AND C.P. GUPTA, On the variational method for the existence of solutions to nonlinear equations of Hammerstein type, *Proc. Amer. Math. Soc.*, **40** (1973), 470–476.
- [16] L. DENG AND X.P. DING, Iterative approximation of Lipschitz strictly pseudocontractive mappings in uniformly smooth Banach spaces, *Nonlinear Analysis*, **24**(7) (1995), 981–987.
- [17] V. DOLEZALE, Monotone Operators and its Applications in Automation and Network Theory, *Studies in Automation and Control* 3, Elsevier Science Publ. New York, 1979.



Iterative Solution of Nonlinear
Equations of Hammerstein Type

H. Zegeye

Title Page

Contents




Go Back

Close

Quit

Page 25 of 27

- [18] A. HAMMERSTEIN, Nichtlineare integralgleichungen nebst anwendungen, *Acta Math. Soc.*, **54** (1930), 117–176.
- [19] S. ISHIKAWA, Fixed points by a new iteration method, *Proc. Amer. Math. Soc.*, **44**(1) (1974), 147–150.
- [20] T. KATO, Nonlinear semi groups and evolution equations, *J. Math. Soc. Japan*, **19** (1967), 508–520.
- [21] W.R. MANN, Mean value methods in iteration, *Proc. Amer. Math. Soc.*, **4** (1953), 506–510.
- [22] R.H. MARTIN Jr., A global existence theorem for autonomous differential equations in Banach spaces, *Proc. Amer. Math. Soc.*, **26** (1970), 307–314.
- [23] D. PASCALI AND SBURLAN, *Nonlinear Mappings of Monotone Type*, Editura Academiei, Bucuresti, Romania (1978).
- [24] B.E. RHOADES, Fixed point iterations for certain nonlinear mappings, *J. Math. Anal. Appl.*, **183** (1994), 118–120.
- [25] H.K. XU, Inequalities in Banach spaces with applications, *Nonlinear Analysis, TMA*, **16**(2) (1991), 1127–1138.
- [26] Z.B. XU AND G.F. ROACH, Characteristic inequalities for uniformly convex and uniformly smooth Banach spaces, *J. Math. Anal. Appl.*, **157** (1991), 189–210.
- [27] Z.B. XU AND G.F. ROACH, A necessary and sufficient condition for convergence of a steepest descent approximation to accretive operator equations, *J. Math. Anal. Appl.*, **167** (1992), 340–354.



**Iterative Solution of Nonlinear
Equations of Hammerstein Type**

H. Zegeye

Title Page

Contents




Go Back

Close

Quit

Page 26 of 27

[28] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications, Part II: Monotone Operators*, Springer-Verlag, Berlin/New York, 1985.



Iterative Solution of Nonlinear Equations of Hammerstein Type

H. Zegeye

Title Page

Contents



Go Back

Close

Quit

Page 27 of 27