

# Journal of Inequalities in Pure and Applied Mathematics


## A NOTE ON INTEGRAL INEQUALITIES AND EMBEDDINGS OF BESOV SPACES

MORITZ KASSMANN

University of Connecticut  
Department of Mathematics  
196 Auditorium Road  
Storrs, 06269, Connecticut  
USA.

*E*Mail: [kassmann@math.uconn.edu](mailto:kassmann@math.uconn.edu)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
152-03



---

volume 4, issue 5, article 107,  
2003.

*Received 21 October, 2003;*  
*accepted 8 November, 2003.*

*Communicated by: S. Saitoh*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

It is shown that certain known integral inequalities imply directly a well-known embedding theorem of Besov spaces.

*2000 Mathematics Subject Classification:* 60G17, 46E35.

*Key words:* Integral Inequalities, GRR-lemma, Besov spaces, Embeddings.

In [7] the authors prove a theorem which links estimates on the modulus of continuity of a real-valued function to the finiteness of a certain integral. Their result reads as follows:

**Theorem 1.** *Let  $\Psi : \mathbb{R} \rightarrow \mathbb{R}_+$  satisfy  $\Psi(\xi) = \Psi(-\xi)$ ,  $\Psi(\infty) = \infty$  and  $\Psi$  non-decreasing for  $\xi \geq 0$ . Let  $p : [-1, 1] \rightarrow \mathbb{R}_+$  be continuous and satisfy  $p(\xi) = p(-\xi)$ ,  $p(0) = 0$  and  $p$  non-decreasing for  $\xi \geq 0$ . Set*


- (1)  $\Psi^{-1}(\xi) = \sup\{\eta, \Psi(\eta) \leq \xi\}$  for  $\xi \geq \Psi(0)$  and
- (2)  $p^{-1}(\xi) = \max\{\eta, p(\eta) \leq \xi\}$  for  $0 \leq \xi \leq p(1)$ .

*If one has for a function  $f \in C([0, 1])$  that*

$$\int_0^1 \int_0^1 \Psi \left( \frac{f(x) - f(y)}{p(x - y)} \right) dx dy \leq B < \infty,$$

*then one has for all  $s, t \in [0, 1]$ :*

$$|f(s) - f(t)| \leq 8 \int_0^{|s-t|} \Psi^{-1} \left( \frac{4B}{\xi^2} \right) dp(\xi).$$



### A Note on Integral Inequalities and Embeddings of Besov Spaces

Moritz Kassmann

Title Page

Contents



Go Back

Close

Quit

Page 2 of 7

As shown in [6] Theorem 1 can be improved in certain boundary cases. Here we aim to show that Theorem 1, known as the GRR-lemma, can be used to derive directly an embedding theorem from certain Besov spaces into the spaces of Hölder continuous functions. Although this observation is straightforward it is remarkable since the proof of the GRR-lemma is not very complicated. Moreover one gets a better understanding of the arising constant.

Originally, the authors of [7] apply Theorem 1 to study continuity of Gaussian processes. Another application of this theorem is an easy derivation of the Kolmogorov-Prohorov criterion for weak compactness of probability measures or an extension of the Burkholder-Davis-Gundy inequality, see [4]. A generalized version of Theorem 1 has been obtained in [2] and used in [8] where upper bounds for the growth of the diameter of a given set exposed in a diffusive stochastic flow are proved. The  $n$ -dimensional version of Theorem 1 reads in one of its possible forms as follows <sup>1</sup>:


**Theorem 2.** *Let  $(X, d)$  and  $(Y, \rho)$  be metric spaces. Let  $f : X \rightarrow Y$  be a continuous function and let  $m$  be a nonnegative Radon measure. Let further  $\Psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a right-continuous function, nondecreasing satisfying  $\Psi(0) = 0$  and  $\Psi(x) > 0$  for all  $x > 0$ . Define  $\Psi^{-1}$  as in (1). Assume that:*

$$V := \int \int \Psi \left( \frac{\rho(f(x), f(y))}{d(x, y)} \right) m(dx) m(dy) < \infty.$$

Then one has for all  $x, y \in X$ :

$$\rho(f(x), f(y)) \leq 6 \int_0^{d(x,y)} \left\{ \Psi^{-1} \left( \frac{4V}{m(B_r(x))^2} \right) + \Psi^{-1} \left( \frac{4V}{m(B_r(y))^2} \right) \right\} dr.$$

<sup>1</sup>The author thanks M. Scheutzow for providing him with his notes on the GRR-lemma.




---

A Note on Integral Inequalities  
and Embeddings of Besov  
Spaces

Moritz Kassmann

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 7

Note that here the assumptions on  $\Psi$  vary slightly from the ones made in Theorem 1. Let us define Sobolev-spaces of fractional order. For a given open connected set  $\Omega \subset \mathbb{R}^n$ , and parameters  $s \in (0, 1)$ ,  $p \geq 1$  the Banach-space  $W^{(s,p)}(\Omega)$  is defined as the set of all functions  $f \in L^2(\Omega)$  for which the norm

$$\|f\|_{s,p,\Omega}^p := \int_{\Omega} |f|^p dx + \int_{\Omega} \int_{\Omega} \frac{|f(x) - f(y)|^p}{|x - y|^{n+sp}} dy dx$$


is finite. These spaces are called Sobolev-Slobodecki spaces and form a special case of the so called Besov spaces. They appear naturally as trace spaces of Sobolev spaces of integer order of differentiation and in the study of boundary value problems for partial differential equations. The monographs [1, 10, 3, 11, 12, 9] are a good choice out of the broad literature on Besov spaces and embedding theorems.

The following well-known embedding theorem follows from Theorem 2.

**Theorem 3.** Assume that  $s \in (\frac{1}{2}, 1)$  and  $\frac{n}{s} < p \leq \frac{n}{1-s}$ . Consider a open connected set  $\Omega \subset \mathbb{R}^n$ . If  $\Omega$  satisfies the property that  $C(\Omega)$  is dense in  $W^{(s,p)}(\Omega)$  then bounded sets of  $W^{(s,p)}(\Omega)$  are also bounded sets in  $C^\alpha(\Omega)$  with  $\alpha \leq s - \frac{n}{p}$ .

The assumption that  $C(\Omega)$  is dense in  $W^{(s,p)}(\Omega)$  is satisfied for nice sets like  $\Omega = \mathbb{R}^n$ . The assumption stays valid for a wide class of domains, for this subtle matter the reader is referred to [5, 1, 9, 11, 10]. The theorem holds without the restriction  $p \leq \frac{n}{1-s}$ . This is the only concession to the use of Theorem 2.

*Proof of Theorem 3.* Choose  $k = n + sp > 2n$ . Choose  $\gamma = \frac{p}{n+sp}$ . Note that  $\gamma \leq 1$ . Choose in Theorem 2  $X = Y = \mathbb{R}^n$ ,  $d(x, y) = |x - y|$ ,  $\rho(x, y) = |x - y|^\gamma$ . Let  $m$  be the Lebesgue measure supported on  $\Omega$ . Choose  $\Psi(z) = z^k$




---

## A Note on Integral Inequalities and Embeddings of Besov Spaces

Moritz Kassmann

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 7

and note that  $\Psi$  satisfies all assumptions in Theorem 2. Let  $S$  be a set in  $W^{s,p}(\Omega)$  such that  $\|f\|_{s,p,\Omega} \leq K$ . Since  $k\gamma = p$  the assumptions yield that for  $f \in S$ :

$$\begin{aligned} V &= \int \int \Psi \left( \frac{\rho(f(x), f(y))}{d(x, y)} \right) m(dx) m(dy) \\ &= \int_{\Omega} \int_{\Omega} \left( \frac{|f(x) - f(y)|^{\gamma}}{|x - y|} \right)^k dy dx < C. \end{aligned}$$


Theorem 2 now states that for any  $x, y \in \Omega$ :

$$\begin{aligned} |f(x) - f(y)|^{\gamma} &\leq 12 \int_0^{|x-y|} \Psi^{-1} \left( \frac{4V}{C(n)r^{2n}} \right) dr \\ &\leq C(n, k)(4V)^{\frac{1}{k}} \left( \frac{k}{k - 2n} \right) |x - y|^{\left(\frac{k-2n}{k}\right)}. \end{aligned}$$

This leads to:

$$|f(x) - f(y)| \leq C(n, k, V) |x - y|^{\left(\frac{k-2n}{k\gamma}\right)} = C(s, p, V) |x - y|^{s - \frac{n}{p}}.$$

The theorem is thus proved. □




---

**A Note on Integral Inequalities  
and Embeddings of Besov  
Spaces**

Moritz Kassmann

---

Title Page

Contents



Go Back


Close

Quit

Page 5 of 7

## References

- [1] R.A. ADAMS, *Sobolev Spaces*, Academic Press, New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] L. ARNOLD AND P. IMKELLER, Stratonovich calculus with spatial parameters and anticipative problems in multiplicative ergodic theory, *Stochastic Process. Appl.*, **62**(1) (1996), 19–54.
- [3] O.V. BESOV, V.P. IL'IN AND S.M. NIKOL'SKIĭ, *Integralnye predstavleniya funktsii i teoremy vlozheniya*, Fizmatlit “Nauka”, Moscow, second edition, 1996.
- [4] M.T. BARLOW AND M. YOR, Semimartingale inequalities via the Garsia-Rodemich-Rumsey lemma, and applications to local times, *J. Funct. Anal.*, **49**(2) (1982), 198–229.
- [5] A.M. CAETANO, Approximation by functions of compact support in Besov-Triebel-Lizorkin spaces on irregular domains, *Studia Math.*, **142**(1) (2000), 47–63.
- [6] C.A. GREENHALL, Growth and continuity of functions satisfying quadratic integral inequalities, *Indiana Univ. Math. J.*, **21** (1971/1972), 157–175.
- [7] A.M. GARSIA, E. RODEMICH AND H. RUMSEY, Jr., A real variable lemma and the continuity of paths of some Gaussian processes, *Indiana Univ. Math. J.*, **20** (1970/1971), 565–578.



---

A Note on Integral Inequalities  
and Embeddings of Besov  
Spaces

Moritz Kassmann

---

Title Page

Contents




Go Back

Close

Quit

Page 6 of 7

- [8] P. IMKELLER AND M. SCHEUTZOW, On the spatial asymptotic behavior of stochastic flows in Euclidean space, *Ann. Probab.*, **27**(1) (1999), 109–129.
- [9] V.G. MAZ'YA AND S.V. POBORCHI, *Differentiable Functions on Bad Domains*, World Scientific Publishing Co. Inc., River Edge, NJ, 1997.
- [10] T. RUNST AND W. SICKEL, Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations, Volume 3 of *de Gruyter Series in Nonlinear Analysis and Applications*, Walter de Gruyter & Co., Berlin, 1996.
- [11] H. TRIEBEL, Theory of function spaces. II, Volume 84 of *Monographs in Mathematics*, Birkhäuser Verlag, Basel, 1992.
- [12] J. WLOKA, *Partial Differential Equations*, Cambridge University Press, Cambridge, 1987. Translated from the German by C.B. Thomas and M.J. Thomas.



---

**A Note on Integral Inequalities  
and Embeddings of Besov  
Spaces**

Moritz Kassmann

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 7