

A FUNCTIONAL MODEL FOR A FAMILY OF OPERATORS INDUCED BY LAGUERRE OPERATOR

HATAMLEH RA'ED

ABSTRACT. The paper generalizes the instruction, suggested by B. Sz.-Nagy and C. Foias, for operatorfunction induced by the Cauchy problem

$$T_t : \begin{cases} th''(t) + (1-t)h'(t) + Ah(t) = 0 \\ h(0) = h_0(th')(0) = h_1 \end{cases}$$

A unitary dilatation for T_t is constructed in the present paper. then a translational model for the family T_t is presented using a model construction scheme, suggested by Zolotarev, V., [3]. Finally, we derive a discrete functional model of family T_t and operator A applying the Laguerre transform

$$f(x) \rightarrow \int_0^\infty f(x) P_n(x) e^{-x} dx$$

where $P_n(x)$ are Laguerre polynomials [6, 7]. We show that the Laguerre transform is a straightening transform which transfers the family T_t (which is not semigroup) into discrete semigroup e^{-itn} .

INTRODUCTION

Functional models for contraction semigroups $Z_t = \exp(itA)$ and T^n , ($t \geq 0$, $n \in \mathbb{Z}^+$) have been constructed by B. Sz.-Nagy and C. Foias [2] at the beginning of 70-s. The bases of this method is a significant concept of dilatation of contraction semigroup. A spectral realization of the dilatation and subsequent narrowing upon the original space leads to a functional model of the contraction semigroup. As a result an operator $A(T)$ in this case is realized by operators which carry out multiplication by independent variable in a specific functional space. The basis of the concept is the Fourier transform of space L^2 .

2000 *Mathematics Subject Classification*: Primary: 47D06, 47A40; Secondary: 47A50, 47A48, 42A50.

Key words and phrases: Laguerre operator, semigroup, Hilbert space, functional model.

Received November 15, 1999.

1. PRELIMINARY INFORMATION ON THE FUNCTIONAL
MODEL IN A FOURIER REPRESENTATION

1.1. We recall [1] that operator collegation Δ ,

$$(1) \quad \Delta = (A, H, \phi, E, \sigma)$$

is a collection of Hilbert spaces H and E and of linear operators $A : H \rightarrow H$, $\phi : H \rightarrow E$, $\sigma : E \rightarrow E$ ($\sigma^* = \sigma$) where the collegation condition holds:

$$(2) \quad A - A^* = i\phi^* \sigma \phi.$$

It is customary to associate with the collegation (1) an open system [1] which is defined by relations

$$(3) \quad \begin{cases} i \frac{d}{dt} h(t) + Ah(t) = \phi^* \sigma u(t); \\ h(0) = h_0, \quad (t \geq 0); \\ v(t) = u(t) - i\phi h(t) \end{cases}$$

where $h(t)$, $u(t)$, $v(t)$ are vector functions from Hilbert spaces H and E respectively. An important role in the further construction of the model representation plays the conservation Law [1].

Theorem 1.1. *For the open system (3) associated with the collegation Δ (1) the conservation Law holds*

$$(4) \quad \|h_0\|^2 + \int_0^T \langle \sigma u(\zeta), u(\zeta) \rangle d\zeta = \|h(T)\|^2 + \int_0^T \langle \sigma v(\zeta), v(\zeta) \rangle d\zeta$$

for any T , $0 \leq T \leq \infty$.

If operator A is selfadjoint then $\phi = 0$, $\sigma = 0$, and Cauchy problem (3) in induced by the semigroup

$$Z_t = \exp(itA), \quad \text{i.e.} \quad h(t) = Z_t h_0$$

and the conservation Law (4) yields Z_t .

1.2. Let us consider a contractive semigroup $Z_t = \exp(itA)$ ($t \geq 0$), which has a property $\|Z_t h\| \leq \|h\|$ for all $h \in H$.

A unitary dilatation of contractive semigroup Z_t in H is said to be a unitary semigroup U_t in \mathcal{H} [2] such that the following relation holds:

$$(5) \quad \mathcal{H} \supseteq H; \quad P_H U_t|_H = Z_t \quad (t \geq 0)$$

where P_H is an orthoprojector on H . The dilatation U_t in H is said to be minimal if

$$(6) \quad \mathcal{H} = \text{span}\{U_t h; t \in \mathbb{R}, h \in H\}$$

where span in (6) denotes a closed linear span of the vectors $U_t h$ for any $t \in \mathbb{R}$ and any $h \in H$.

A significant role in the theory of dilatation of contractive semigroup Z_t plays the following Theorem 1.2.

Theorem 1.2. *Any contracting semigroup Z_t in H has a unitary dilatation U_t in H . Moreover the minimal dilatation U_t is defined up to isomorphism.*

We present a construction of the dilatation U_t according to the paper [3]. A contractibility of the semigroup Z_t means [2, 3] that A is dissipative, i.e. $-i(A - A^*) \geq 0$. Consequently including A into the collocation Δ (1) we can assume that $\sigma = I$. Therefore the conservation law (4) has the form

$$(7) \quad \|h_0\|^2 + \int_0^T \|u(\zeta)\|^2 d\zeta = \|h(T)\|^2 + \int_0^T \|v(\zeta)\|^2 d\zeta$$

We defined [3] a dilatation space \mathcal{H} , which forms vector-functions $f(\zeta) = (u_+(\zeta), h, u_-(\zeta))$ so that $u_{\pm}(\zeta) \in E$ and $\text{Supp } u_{\pm}(\zeta) \in \mathbb{R}_{\mp}$ for a finite norm

$$(8) \quad \|f\|^2 = \int_{-\infty}^0 \|u_+(\zeta)\|^2 d\zeta + \|h\|^2 + \int_0^{\infty} \|u_-(\zeta)\|^2 d\zeta < \infty.$$

We define a dilatation U_t in \mathcal{H} by the formula

$$(9) \quad (U_t f)(\zeta) = (u_+(t, \zeta), h_t, u_-(t, \zeta))$$

where $u_-(t, \zeta) = P_{\mathbb{R}_+} u_-(\zeta + t)$; $h_t = y_t(0)$, and $y_t(\zeta)$ is a solution of the Cauchy problem

$$\begin{cases} i \frac{d}{d\zeta} y_t(\zeta) + A y_t(\zeta) = \phi^* u_-(\zeta + t); \\ y_t(-t) = 0, \quad \zeta \in (-t, 0); \end{cases}$$

and at last $u_+(t, \zeta) = u_+(t + \zeta) + P_{(-t, 0)} \{u_-(\zeta + t) - i\phi y_t(\zeta)\}$ where $P_{\mathbb{R}_+}$ and $P_{(-t, 0)}$ are operators of narrowing (projection operators at set \mathbb{R}_+ and $(-t, 0)$ respectively), $t \geq 0$.

It is not difficult to show that unitarity of U_t (9) in \mathcal{H} is a consequence of the conservation law (1). By the dilatation construction U_t one can see that the space \mathcal{H} has the form

$$(10) \quad \mathcal{H} = D_+ \oplus H \oplus D_-$$

where the subspace D_+ is formed by vector-function of the form $(u_+(\zeta), 0, 0) \in \mathcal{H}$ and the subspace D_- is formed by vector-function $(0, 0, u_-(\zeta))$ from \mathcal{H} , respectively.

The subspaces D_{\pm} have the following properties:

$$(11) \quad \begin{aligned} U_t D_+ &\subseteq D_+ & (t \geq 0), \\ U_t D_- &\subseteq D_- & (t \leq 0). \end{aligned}$$

Thus D_+ is outgoing subspace and D_- is incoming subspace in the sense of P. D. Lax and R. S. Phillips [4]. In accordance with the paper [3], we define a free unitary group V_t in the space $L_{\mathbb{R}}^2(E)$, which will act as

$$(12) \quad (V_t g)(\zeta) = g(\zeta + t)$$

and vector-function $g(\zeta) \in E$, $\zeta \in \mathbb{R}$ is such that

$$\int_{-\infty}^{\infty} \|g(\zeta)\|^2 d\zeta < \infty.$$

It is evidently that D_{\pm} after identification belongs to $L^2_{\mathbb{R}}(E)$ also.

Wave operators W_{\pm} play a significant role in the scattering theory. They are defined [3, 4] as

$$(13) \quad W_{\pm} = s - \lim_{t \rightarrow \mp\infty} U_+ P_{D_{\pm}} V_{-t}$$

where $P_{D_{\pm}}$ are orthoprojectors on subspaces D_{\pm} . The following theorem holds [3].

Theorem 1.3. *The wave operators W_{\pm} exist as strong limits (13) are isometries from $L^2_{\mathbb{R}}(E)$ to \mathcal{H} , and the relations*

$$(14) \quad W_{\pm} V_t = U_t W_{\pm}, \quad (\forall t), \quad W_{\pm} P_{D_{\pm}} = P_{D_{\pm}}$$

are valid.

The scattering operator S is defined by the wave operator W_{\pm} in a conventional way [3, 4]:

$$(15) \quad S = W_+^* W_-.$$

From Theorem 1.3 there follows a proposition.

Theorem 1.4. *The operator S (15) is a contraction, i.e. $\|S\| \leq 1$ and has the properties:*

$$(16) \quad \begin{aligned} S V_t &= V_t S; \quad S L^2_{\mathbb{R}^+} \subseteq L^2_{\mathbb{R}^+}(E); \\ \overline{S L^2_{\mathbb{R}}(E)} &= L^2_{\mathbb{R}}(E) \end{aligned}$$

1.3. We recall that the collegation Δ (1) is simple [1-3] if $H = \text{span}\{A^n \phi^* g; n \in \mathbb{Z}_+, g \in E\}$. Let us define the following subspaces in \mathcal{H} ,

$$\mathfrak{R}_{\pm} = \overline{W_{\pm} L^2_{\mathbb{R}}(E)}.$$

The following theorem gives a sufficient condition for the completeness of the wave operators W_{\pm} , [3].

Theorem 1.5. *If the collegation Δ is simple then the relation $\mathcal{H} = \text{span}\{f_+ + f_-; f_{\pm} \in \mathfrak{R}_{\pm}\}$ holds.*

Now we construct a translational model [3]. Let $f_k(\zeta) \in L^2_{\mathbb{R}}(E)$, ($k = 1, 2$). We define a mapping

$$\begin{pmatrix} f_1(\zeta) \\ f_2(\zeta) \end{pmatrix} \rightarrow \Psi_p(\zeta) = W_- f_1(\zeta) + W_+ f_2(\zeta) \in \mathcal{H}.$$

Then using isometry of W_{\pm} and the form of operator S (15) it is not difficult to show that

$$(17) \quad \|\Psi_p(\zeta)\|^2 = \int_{-\infty}^{\infty} \left\langle \begin{bmatrix} I & S^* \\ S & I \end{bmatrix} \begin{pmatrix} f_1(\zeta) \\ f_2(\zeta) \end{pmatrix}, \begin{pmatrix} f_1(\zeta) \\ f_2(\zeta) \end{pmatrix} \right\rangle d\zeta,$$

Using Theorem 1.5 we may assert, that space H is isomorphic to the space $L^2 \begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix}$ which is formed by vector-functions $f(\zeta) = \begin{pmatrix} f_1(\zeta) \\ f_2(\zeta) \end{pmatrix}$ for which the norm (17) is finite. By virtue of conditions (14) the dilatation U_t on Ψ_p will act as a shift. Therefore if $f(\zeta) \in L^2 \begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix}$ then the dilatation U_t is transformed into

$$(18) \quad \widehat{U}_t f(\zeta) = f(\zeta + t).$$

Applying again (14), one can easily deduce that the spaces D_{\pm} are realized now in the form

$$(19) \quad \widehat{D}_- = \begin{pmatrix} L^2_{\mathbb{R}_+}(E) \\ 0 \end{pmatrix}, \quad \widehat{D}_+ = \begin{pmatrix} 0 \\ L^2_{\mathbb{R}_-}(E) \end{pmatrix}.$$

Thus the initial space H acquires such model form

$$(20) \quad \begin{aligned} \widehat{H}_p &= L^2 \begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix} \ominus \begin{pmatrix} L^2_{\mathbb{R}_+}(E) \\ L^2_{\mathbb{R}_-}(E) \end{pmatrix} \\ &= f = \left(\begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \in L^2 \begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix}; \begin{matrix} f_1 + S^* f_2 \in L^2_{\mathbb{R}_-}(E) \\ S f_1 + f_2 \in L^2_{\mathbb{R}_+}(E) \end{matrix} \right) \end{aligned}$$

and in the virtue of the dilatation the action of semigroup Z_t is transformed to the shift semigroup

$$(21) \quad \widehat{Z}f(\zeta) = P_{\widehat{H}_p} f(\zeta + t)$$

where $f(\zeta) \in \widehat{H}_p$ (20). Thus the following theorem is proved.

Theorem 1.6. *A minimal unitary dilatation U_t in \mathcal{H} of the contraction semigroup $Z_t = \exp(itA)$ in H , where A is dissipative operator of a simple collegation Δ is unitary equivalent to a translation group \widehat{U}_t (18) in the space $L^2 \begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix}$, and the contraction semigroup Z_t is unitary equivalent to the shift semigroup \widehat{Z}_t (21) in the space \widehat{H}_p respectively.*

The Fourier transform by formula

$$(22) \quad \widetilde{f}(\lambda) = \int_{-\infty}^{\infty} f(\zeta) e^{-i\lambda\zeta} d\zeta$$

in the virtue of Plancherel theorem [2, 3] is a unitary operator in $L^2_{\mathbb{R}}(E)$. By the virtue of Wiener-Paley theorem

$$\tilde{L}^2_{\mathbb{R}_+}(E) = H^2_-(E); \quad \tilde{L}^2_{\mathbb{R}_-}(E) = H^2_+(E)$$

where $H^2_{\pm}(E)$ are Hardy spaces of E -value function from $L^2_{\mathbb{R}}(E)$ which are holomorphically continued into lower (upper) half-plane. Let us apply the Fourier transform (22) to translational model (18) – (21) and take advantage of the following Theorem 1.7

Theorem 1.7. *The Fourier transform of the scattering operator S (15) transfers the operator S into operator performing multiplication by characteristic function*

$$(23) \quad \begin{aligned} S_{\Delta}(\lambda) &= I - \phi(A - \lambda I)^{-1} \phi^*, \quad i.e. \\ (\tilde{S}f)(\lambda) &= S_{\Delta}(\lambda) \tilde{f}(\lambda). \end{aligned}$$

As it is known $\tilde{f}(\lambda + t) = e^{i\lambda t} \tilde{f}(\lambda)$, therefore we derive such functional model.

Theorem 1.8. *A minimal unitary dilatation U_t in H of the contraction semigroup $Z_t = \exp(itA)$ in H , where A is dissipative operator of a simple collegation Δ is unitary equivalent to the group*

$$(24) \quad \tilde{U}_t f(\lambda) = e^{i\lambda t} f(\lambda)$$

where $f(\lambda) \in L^2 \left(\begin{array}{cc} I & S_{\Delta}^*(\lambda) \\ S_{\Delta}(\lambda) & I \end{array} \right)$ and contraction semigroup Z_t is unitary equivalent to semigroup $\tilde{Z}_t f(\lambda) = P_{\tilde{H}_p} e^{i\lambda t} f(\lambda)$, where $f(\lambda)$ belongs to the space

$$\tilde{H}_p = \left\{ f = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}(\lambda) \in \begin{pmatrix} I & S_{\Delta}^*(\lambda) \\ S_{\Delta}(\lambda) & I \end{pmatrix} ; \begin{array}{l} f_1 + S_{\Delta}^*(\lambda) f_2 \in H^2_+(E) \\ S_{\Delta}(\lambda) f_1 + f_2 \in H^2_-(E) \end{array} \right\}$$

Here the main operator \tilde{A} in \tilde{H}_p act as multiplication operator by independent variable

$$(26) \quad \tilde{A}f(\lambda) = P_{\tilde{H}_p} \lambda f(\lambda), \quad f(\lambda) \in \tilde{H}_p.$$

In the next section we will generalize this construction on the case of the Laguerre transform.

2. A FUNCTIONAL MODEL FOR THE LAGUERRE REPRESENTATION

2.1. Let us consider a differential operator

$$(27) \quad \ell = t \frac{d^2}{dt^2} + (1-t) \frac{d}{dt}$$

in what follows called the Laguerre operator; it acts on functions form $C^2 = (\mathbb{R}_+)$. We denote by $L^2_{\mathbb{R}_+}(e^{-t} dt)$ the following space:

$$(28) \quad L^2_{\mathbb{R}_+}(e^{it} dt) = \left\{ f(t), t \in \mathbb{R}_+; \int_0^{\infty} |f(t)|^2 e^{-t} dt < \infty \right\}$$

Proposition 2.1. *An operator ℓ is symmetric in the space $L^2_{\mathbb{R}_+}(e^{-t}dt)$ under the self-adjoint boundary conditions, i.e. $\langle \ell x, y \rangle = \langle y, \ell y \rangle$ for all $x, y \in \mathbb{C}^2(\mathbb{R}_+)$ such that $tx(t)|_{t=0} = 0$, $ty(t)|_{t=0} = 0$ and $ty'(t)|_{t=0} < \infty$, $tx'(t)|_{t=0} < \infty$.*

Proof. We calculate

$$\begin{aligned} \langle \ell x, y \rangle - \langle x, \ell y \rangle &= \int_0^\infty \{ (tx'' + (1-t)x')\bar{y} - x(t\bar{y}'' + (1-t)\bar{y}') \} e^{-t} dt \\ &= \int_0^\infty \{ te^{-t}(x'\bar{y} - \bar{y}'x) \}' dt = \{ te^{-t}(x'\bar{y} - \bar{y}'x) \} \Big|_0^\infty = 0 \end{aligned}$$

by virtue of the boundary conditions. \square

Let us consider now an open system of special form, generated by the Laguerre operator (27) and corresponding to the collegation Δ (1):

$$(29) \quad \begin{cases} \ell h(t) + Ah(t) = \phi^* \sigma u(t); \\ h(0) = h_0(th')(0) = h_1; \\ v(t) = u(t) - i\phi h(t). \end{cases}$$

The following assertion is valid, similar to Theorem (1.1).

Theorem 2.1. *For the open system (29) associated with collegation Δ the law of conservation of energy is valid, i.e.*

$$(30) \quad \begin{aligned} &\int_0^T \langle \sigma u(\zeta), u(\zeta) \rangle e^{-\zeta} d\zeta + \langle I\hat{h}_0, \hat{h}_0 \rangle \\ &= \int_0^T \langle \sigma v(\zeta), v(\zeta) \rangle e^{-\zeta} d\zeta + \langle I\hat{h}_T, \hat{h}_T \rangle \end{aligned}$$

where $I = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ and $h_0 = \begin{pmatrix} h_0 \\ h_t \end{pmatrix}$, $h_T = \begin{pmatrix} h(T) \\ e^{-T}Th'(T) \end{pmatrix}$ for any finite $T > 0$.

Proof. We calculate

$$\begin{aligned} \langle \ell h, h \rangle - \langle h, \ell h \rangle &= \langle \phi^* \sigma u - Ah, h \rangle - \langle h, \psi^* \sigma u - Ah \rangle \\ &= \langle \sigma u, \frac{u-v}{i} \rangle - \langle \frac{u-v}{i}, \sigma u \rangle - \langle (A - A^*)h, h \rangle \\ &= i\langle \sigma u, u-v \rangle + i\langle u-v, \sigma u \rangle - i\langle \phi^* \sigma \phi h, h \rangle \\ &= i\langle \sigma u, u-v \rangle + i\langle u-v, \sigma u \rangle - i\langle \sigma(u-v), u-v \rangle \\ &= i\langle \sigma u, u \rangle - i\langle \sigma v, v \rangle. \end{aligned}$$

Now we integrate the derived equality:

$$\begin{aligned} &\int_0^T \langle \sigma v, v \rangle e^{-t} dt - \int_0^T \langle \sigma u, u \rangle e^{-t} dt \\ &= i \int_0^T [\langle \ell h, h \rangle - \langle h, \ell h \rangle] e^{-t} dt \\ &= i \{ e^{-t} t [\langle h'', h \rangle - \langle h, h' \rangle] \} \Big|_0^T \\ &= \langle I\hat{h}_0, \hat{h}_0 \rangle - \langle I\hat{h}_T, \hat{h}_T \rangle \end{aligned}$$

which proves our assertion. \square

2.2. Let us make use of the energy conservation law (30) to construct a dilatation for operator T_t generated by the Cauchy problem

$$(31) \quad \begin{cases} \ell h(t) + Ah(t) = 0; \\ h(0) = h_0; (th')(0) = h_1; \end{cases}$$

where $T_t(h_0, h_1) = (h(t), th'(t))$. We will call an unitary operator-function U_t in \mathcal{H} a dilatation of family T_t in H , if $\mathcal{H} \supseteq H$, $T_t = P_H U_t|_H$.

Here we do not suppose that T_t and U_t is semigroup. Moreover, the unitary property of U_t may hold not necessarily in Hilbert metric but in indefinite one. The following analog of Theorem 1.2 is valid.

Theorem 2.2. *The operator-function T_t generated by the Cauchy problem (31) with dissipative operator A of collegation Δ (1) (i.e. $\sigma = I$) possesses the unitary (in indefinite metric) dilatation U_t , where the minimal dilatation is determined up to isomorphism.*

Proof. To prove the theorem we bring a construction of dilatation U_t by analog with (8), (9).

Let us consider a Hilbert space

$$(32) \quad \begin{aligned} \mathcal{H} = \{ & f = (u(\zeta), \widehat{h}, v(\zeta)); u(\zeta), v(\zeta) \in E, \text{supp } v \in \mathbb{R}_-, \text{supp } u \in \mathbb{R}_+, \\ & \widehat{h} = \begin{pmatrix} h_0 \\ h_1 \end{pmatrix}, h_k \in H; \|f\|^2 = \int_{-\infty}^0 \|v(\zeta)\|^2 e^{-\zeta} d\zeta + \|\widehat{h}\|^2 \\ & + \int_0^{\infty} \|u(\zeta)\|^2 e^{-\zeta} d\zeta < \infty \}. \end{aligned}$$

We set indefinite metric \mathcal{H}

$$(33) \quad \langle f \rangle_I^2 = \int_{-\infty}^0 \|v(\zeta)\|^2 e^{-\zeta} d\zeta + \langle I\widehat{h}, \widehat{h} \rangle + \int_0^{\infty} \|u(\zeta)\|^2 e^{-\zeta} d\zeta$$

where I has the form indicated in Theorem 2.1.

We construct the dilatation U_t in \mathcal{H} ,

$$(34) \quad U_t f = f_t(u(t, \zeta), \widehat{h}_t, v(t, \zeta)).$$

Let us consider further the Cauchy problem

$$(35) \quad \begin{cases} (i\frac{\partial}{\partial t} + \ell_\zeta) \widehat{u}(t, \zeta) = 0; \\ \widehat{u}(0, \zeta) = u(\zeta); \zeta \in \mathbb{R}_+; \end{cases}$$

where ℓ_ζ is operator ℓ (27) with respect to ζ .

Solution of the problem is easily obtained. In fact, let

$$\widehat{u}(t, \zeta) = \sum_{n \in \mathbb{Z}_+} e^{-itn} C_n g_n(\zeta)$$

where $g_n(\zeta)$ are the Laguerre polynomials [5] which are the solutions of equation $\ell_\zeta g_n(\zeta) + n g_n(\zeta) = 0$ and have the form

$$g_n(\zeta) = \frac{1}{n!} e^\zeta \frac{d^n}{d\zeta^n} (\zeta e^{-\zeta})$$

and make a complete system of orthogonal polynomials in $L^2_{\mathbb{R}_+}(e^{-\zeta} d\zeta)$. The coefficients C_n are obtained from the initial condition $\sum C_n g_n(\zeta) = u(\zeta)$.

Therefore $\widehat{u}(t, \zeta)$ possesses the property $\text{supp } \widehat{u}(t, \zeta) = \text{supp } \widehat{u}(\zeta) \subseteq \mathbb{R}_+$. Now we determine $u(t, \zeta)$ in (34) by the formula

$$(36) \quad u(t, \zeta) = P_{\mathbb{R}_+} \widehat{u}(t, \zeta + t) e^{-\frac{\zeta}{2}}.$$

To set \widehat{h}_t (34), we consider the following Cauchy problem

$$(37) \quad \begin{cases} \ell_\zeta y(\zeta) + Ay(\zeta) = \phi^* \widehat{u}(t, \zeta + t) e^{-\frac{\zeta}{2}}; & \zeta \in (-t, 0); \\ y(-t) = h_0; \\ (-t)e^t y(-t) = h_1; \end{cases}$$

and put $\widehat{h}_t = \begin{pmatrix} y(0) \\ (ty')(0) \end{pmatrix}$.

Finally, to set $v(t, \zeta)$ (34) we consider the similar equation

$$(38) \quad \begin{cases} (i\frac{\partial}{\partial t} + \ell_\zeta) \widehat{v}(t, \zeta) = 0; \\ \widehat{v}(0, \zeta) = v(\zeta); & \zeta \in \mathbb{R}_-; \end{cases}$$

and put $v(t, \zeta) = e^{-\frac{\zeta}{2}} \widehat{v}(t, \zeta + t) + P_{\mathbb{R}_-} \{ \widehat{u}(t, \zeta + t) e^{-\frac{\zeta}{2}} - i\phi y(\zeta) \}$. We show that U_t (34) has property of isometry in the metric (33). To this end we calculate,

$$\begin{aligned} \langle f_t \rangle_I^2 &= \int_{-\infty}^0 \|v(t, \zeta)\|^2 e^{-\zeta} d\zeta + \langle I\widehat{h}_t, \widehat{h}_t \rangle + \int_0^\infty \|u(t, \zeta)\|^2 e^{-\zeta} d\zeta \\ &= \int_{-\infty}^{-t} \|\widehat{v}(t, \zeta + t)\|^2 e^{-\zeta - t} d\zeta + \int_{-t}^0 \|\widehat{u}(t, \zeta + t) e^{-\frac{\zeta}{2}} - i\phi y(\zeta)\|^2 e^\zeta d\zeta \\ &\quad + \langle I\widehat{h}_t, \widehat{h}_t \rangle + \int_0^\infty \|u(t, \zeta + t)\|^2 e^{-\zeta - t} d\zeta \\ &= \int_{-\infty}^{-t} \|\widehat{v}(t, \zeta + t)\|^2 e^{-\zeta - t} d\zeta + \langle I\widehat{h}_0, \widehat{h}_0 \rangle + \int_{-t}^\infty \|\widehat{u}(t, \zeta + t)\|^2 e^{-\zeta - t} d\zeta \\ &= \int_{-\infty}^0 \|\widehat{v}(t, \zeta)\|^2 e^{-\zeta} d\zeta + \langle I\widehat{h}_0, \widehat{h}_0 \rangle + \int_0^\infty \|\widehat{u}(t, \zeta)\|^2 e^{-\zeta} d\zeta \\ &= \langle f \rangle_I^2 \end{aligned}$$

In this calculation we have made use of the conservation law (30) and of the fact that norms of solutions of Cauchy problems $\widehat{u}(t, \zeta)$, $\widehat{v}(t, \zeta)$ (35) and (38) coincide with norms of initial data $u(\zeta)$ and $v(\zeta)$ in the spaces $L^2_{\mathbb{R}_+}(e^{-t} dt)$ and $L^2_{\mathbb{R}_-}(e^{-t} dt)$ by virtue of selfadjointness of operators ℓ_ζ in the spaces.

In order to prove that U_t has the property of being unitary, it is necessary to ascertain that from $U_t^* f = 0$ implies $f = 0$. It is easy to show that U_t^* will act by the formula

$$(39) \quad U_t^* f = (u(t, \zeta), \widehat{h}_t, v(t, \zeta)).$$

Here $v(t, \zeta) = P_{\mathbb{R}_-} \widehat{v}(t, \zeta - t) e^{\frac{t}{2}}$ where $\widehat{v}(t, \zeta)$ is a solution of problem (38).

In order to obtain \widehat{h}_t , it is necessary to consider dual to (37) problem

$$(40) \quad \begin{cases} \ell_\zeta y(\zeta) + A^* y(\zeta) = \phi^* \widehat{v}(\zeta, \zeta - t) e^{\frac{t}{2}}; \\ y(t) = h_0; \\ e^{-t} t y'(t) = h_1; \end{cases}$$

and put $\widehat{h}_t = \begin{pmatrix} y(0) \\ (t y')(0) \end{pmatrix}$. Finally,

$$u(t, \zeta) = \widehat{u}(t, \zeta - t) e^{\frac{t}{2}} + P_{\mathbb{R}_+} \{ \widehat{v}(t, \zeta - t) e^{\frac{t}{2}} + i \phi y(\zeta) \},$$

where $\widehat{u}(t, \zeta)$ is the solution of Cauchy problem (35).

Thus let $U_t^* f = 0$, then $\widehat{u}(t, \zeta) = 0$ and so $\widehat{v}(t, \zeta) = 0$ and $\widehat{v}(t, \zeta - t) e^{\frac{t}{2}} + i \phi y(\zeta) = 0$ therefore $u(\zeta) \equiv 0$. Now, by substituting $\widehat{v}(t, \zeta - t) = -i \phi y(\zeta) e^{-\frac{t}{2}}$ in (40) we obtain a homogeneous equation

$$\ell_\zeta y + A^* y + i \phi^* \phi y = 0$$

with zero condition in the origin $\widehat{h}_t = 0$. By virtue of uniqueness of Cauchy problem solution, this yields that $y(\zeta) \equiv 0$, therefore $\widehat{v}(t, \zeta - t) = 0$ on interval $(0, t)$. Accounting that $\widehat{v}(t, \zeta - t) = 0$ with $(-\infty, 0)$, finally we conclude that $v(\zeta) = 0$. Thus $f = 0$. This proves the property of being unitary for U_t (34) and completes the proof of the theorem. \square

2.3. Let us pass to constructing wave operators. To this end we define a “free” group by analogy with (38)

$$(41) \quad V_t g(\zeta) = g(t, \zeta),$$

where $g(t, \zeta)$ is a solution of Cauchy problem

$$(42) \quad \begin{cases} (i \frac{\partial}{\partial t} + \ell_\zeta) g(t, \zeta) = 0; \\ g(0, \zeta) = g(\zeta) \in L^2_{\mathbb{R}}(e^{-\zeta} d\zeta). \end{cases}$$

It is evident that V_t (41) is unitary. Now we define the operators

$$(43) \quad \begin{aligned} W_- &= s - \lim_{t \rightarrow +\infty} U_t P_{\mathbb{R}_+} V_{-t}, \\ W_+ &= s - \lim_{t \rightarrow -\infty} U_t^* P_{\mathbb{R}_-} V_{-t}^*. \end{aligned}$$

By analogy with Theorem 1.3 we have

Theorem 2.3. *The wave operators W_{\pm} exist as strong limits (43), are isometries from $L^2_{\mathbb{R}}(e^{-\zeta} d\zeta)$ to \mathcal{H} , and the following relations are valid:*

$$(44) \quad \begin{aligned} U_t W_- &= W_- V_t, U_t^* W_+ = W_+ V_t^*, \quad (t \geq 0) \\ W_{\pm} P_{\mathbb{R}_{\mp}} &= P_{\mathbb{R}_{\mp}} \end{aligned}$$

Proof. We prove the assertion of the theorem for W_- (for W_+ the proof is similar). The main matter of the theorem consists of existence proof of W_- since the relation (44) is proved by analogy with arguments given in Section 1; see [2, 3]. Let

$$f_t = U_t P_{\mathbb{R}_+} V_{-t} g = (v(t, \zeta), h_t, u(t, \zeta))$$

then $u(t, \zeta) = P_{\mathbb{R}_+} g(\zeta)$. We consider the Cauchy problem

$$(45) \quad \begin{cases} \ell_{\zeta} y(\zeta) + Ay(\zeta) = \phi^* g(\zeta); \\ y(-t) = 0; y'(-t) = 0, \zeta \in (-t, 0). \end{cases}$$

Then $\widehat{h}_t = \begin{pmatrix} y(0) \\ (ty')(0) \end{pmatrix}$.

We denote by $K(\zeta, \eta)$ a Cauchy function of the problem (45) (i.e. $K(\zeta, \zeta) = 0$, $K'(\zeta, \zeta) = I$), then a solution $y(\zeta)$ of (45) has the form

$$y_t(\zeta) = \int_{-t}^{\zeta} K(\zeta, \eta) \phi^* g(\eta) d\eta.$$

Therefore $V(t, \zeta)$ has the form

$$V(t, \zeta) = P_{(-t, 0)} \{g(\zeta) - i\phi y(\zeta)\}.$$

Thus,

$$f_t = \left(P_{(-t, 0)} \{g(\zeta) - i\phi \int_{-t}^0 K(\zeta, \eta) \phi^* g(\eta) d\eta\}, \begin{pmatrix} \int_{-t}^0 K(0, \eta) \phi^* g(\eta) d\eta \\ \int_{-t}^0 K'(0, \eta) \phi^* g(\eta) d\eta \end{pmatrix}, P_{\mathbb{R}_+} g(\zeta) \right).$$

We show that f_t is a Cauchy sequence, i.e. $\|f_{t+\Delta} - f_t\|^2 \rightarrow 0$ as $t \rightarrow \infty$. Since

$$(46) \quad \|f_{t+\Delta} - f_t\|^2 = \int_{-\infty}^0 \|v_t(t+\Delta, \zeta) - v_t(t, \zeta)\|^2 e^{-\zeta} d\zeta + \|\widehat{h}_{t+\Delta} - \widehat{h}_t\|^2.$$

It is sufficient to show that each summand approaches to zero as $t \rightarrow \infty$. We show that $\|\widehat{h}_{t+\Delta} - \widehat{h}_t\| \rightarrow 0$ when $t \rightarrow \infty$ and we will prove this property component by component. It is obvious that

$$\begin{aligned} \|\widehat{h}_{t+\Delta} - \widehat{h}_t\|^2 &= \left\| \int_{(-t-\Delta)}^{-t} K(0, \eta) \phi^* g(\eta) d\eta \right\|^2 \\ &\leq \int_{-t-\Delta}^{-t} \|K(0, \eta)\|^2 e^{\eta} d\eta \cdot \int_{-t-\Delta}^{-t} e^{-\eta} \|\phi^*\|^2 \|g(\eta)\|^2 d\eta \end{aligned}$$

and since the function $K(0, \eta)e^\eta$ is bounded (see [6, 7]), we obtain that

$$\|h_{t+\Delta} - h_t\|^2 \leq \Delta C \|\phi^*\|^2 \int_{-t-\Delta}^{-t} \|g(\eta)\|^2 e^{-\eta} d\eta \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

since $g(\eta) \in L_{\mathbb{R}}^2(e^{-\eta} d\eta)$.

The convergence of second components $\widehat{h}_{t+\Delta} - \widehat{h}_t$ to zero is proved in a similar way. We show that the first summand in (46) approaches to zero too.

In fact,

$$\begin{aligned} A &= \int_{-\infty}^0 \|P_{(-t-\Delta, -t)}g(\zeta) - iP_{(-t-\Delta, 0)}\phi \int_{-t-\Delta}^{\zeta} K(\zeta, \eta)\phi^*g(\eta) d\eta \\ &\quad + i \int_{-t}^{\zeta} \phi K(\zeta, \eta)\phi^*g(\eta) d\eta\|^2 e^{-\zeta} d\zeta \\ &= \int_{-t-\Delta}^{-t} \|g(\zeta)\|^2 e^{-\zeta} d\zeta + \int_{-\infty}^0 \|P_{(-t-\Delta, 0)}\phi y_{t+\Delta}(\zeta) - P_{(-t, 0)}\phi y_t(\zeta)\|^2 e^{-\zeta} d\zeta \\ &\quad + 2\text{Im} \int_{-t-\Delta}^{-t} \langle g(\zeta), P_{(-t-\Delta, 0)}\phi y(\zeta) - P_{(-t, 0)}\phi y(\zeta) \rangle e^{-\zeta} d\zeta \end{aligned}$$

It is obvious that the first and third summands in the given sum approaches to zero as $t \rightarrow \infty$ because $g(\zeta) \in L_{\mathbb{R}}^2(e^{-\zeta} d\zeta)$. We evaluate the second summand:

$$\begin{aligned} B &= \int_{-\infty}^0 \|P_{(-t-\Delta, 0)}\phi y_{t+\Delta}(\zeta) - P_{(-t, 0)}\phi y_t(\zeta)\|^2 e^{-\zeta} d\zeta \\ &= \int_{-\infty}^0 \langle \phi \Delta y, \phi \Delta y \rangle e^{-\zeta} d\zeta, \end{aligned}$$

where

$$\Delta y = P_{(-t-\Delta, 0)}y_{t+\Delta}(\zeta) - P_{(-t, 0)}y_t(\zeta).$$

Then

$$\begin{aligned} A &= \int_{-\infty}^0 \langle \phi^* \phi \Delta y, \Delta y \rangle e^{-\zeta} d\zeta = \int_{-\infty}^0 \left\langle \frac{A - A^*}{i} \Delta y, \Delta y \right\rangle e^{-\zeta} d\zeta \\ &= 2\text{Im} \int_{-\infty}^0 \langle \phi^* g - \ell \Delta y, \Delta y \rangle e^{-\zeta} d\zeta \\ &= 2\text{Im} \int_{-\infty}^0 \langle \phi^* g, \Delta y \rangle e^{-\zeta} d\zeta + 2\text{Im} \int_{-\infty}^0 \langle \ell \Delta y, \Delta y \rangle e^{-\zeta} d\zeta \end{aligned}$$

the first summand approaches to zero again on account of $g(\zeta) \in L_{\mathbb{R}}^2(e^{-\zeta} d\zeta)$, and the second one yields after integration by parts

$$\|\zeta e^{-\zeta} \Delta y\|_{|\zeta=0} \rightarrow 0 \quad (t \rightarrow \infty)$$

since $\widehat{\Delta h}_t \rightarrow 0$. The theorem is proved. \square

As before, we define the operator S by the formula (15). Then the following theorem holds.

Theorem 2.4. *The operator S (15) is a contraction from $L_{\mathbb{R}}^2(e^{-\zeta} d\zeta)$ to $L_{\mathbb{R}}^2(e^{-\zeta} d\zeta)$ and possesses the following properties:*

$$\begin{aligned} SV_t &= V_t S; \quad SL_{\mathbb{R}_+}^2(e^{-\zeta} d\zeta) \subset L_{\mathbb{R}_+}^2(e^{-\zeta} d\zeta); \\ \overline{SL_{\mathbb{R}}^2(e^{-\zeta} d\zeta)} &= L_{\mathbb{R}}^2(e^{-\zeta} d\zeta). \end{aligned}$$

2.4. Further we suppose that the collegation Δ (1) is simple and as in subsection 1.3 we set a mapping

$$\Psi_p(\zeta = W_- f_1(\zeta)) + W_+ f_2(\zeta)$$

from $L_{\mathbb{R}}^2(e^{-\zeta} d\zeta) + L_{\mathbb{R}}^2(e^{-\zeta} d\zeta)$ to \mathcal{H} . It is obvious that

$$\Psi_p(\zeta) \in L^2 \left(\begin{pmatrix} I & S^* \\ S & I \end{pmatrix}, e^{-\zeta} d\zeta \right)$$

Action of dilatation in this space again reduces to a translation

$$(47) \quad \widehat{U}_t f(\zeta) = f(\zeta + t),$$

since

$$\begin{aligned} U_t \Psi_p(\zeta) &= W_- f_1(\zeta + t) + U_t W_+ f_2(\zeta) \\ &= W_- f_1(\zeta + t) + U_t W_+ V_t^* V_t f_2(\zeta) \\ &= W_- f_1(\zeta + t) + U_t U_t^* W_+ V_t f_2(\zeta) = \Psi_P(\zeta + t). \end{aligned}$$

As earlier, it is obvious that

$$D_- = \begin{pmatrix} L_{\mathbb{R}_+}^2(e^{-\zeta} d\zeta) \\ 0 \end{pmatrix}, \quad D_+ = \begin{pmatrix} 0 \\ L_{\mathbb{R}_-}^2(e^{-\zeta} d\zeta) \end{pmatrix}$$

and the model space H_p has the form

$$(48) \quad H_p = L^2 \left(\begin{pmatrix} 1 & S^* \\ S & 1 \end{pmatrix} e^{-\zeta} d\zeta \right) \ominus \begin{pmatrix} L_{\mathbb{R}_+}^2(e^{-\zeta} d\zeta) \\ L_{\mathbb{R}_-}^2(e^{-\zeta} d\zeta) \end{pmatrix}$$

and in addition T_t passes to shift semigroup

$$(49) \quad \widehat{T}_t f(\zeta) = f(\zeta + t).$$

Now we consider a Laguerre transform

$$(50) \quad L_n = \int_0^\infty e^{-x} P_n(x) f(x) dx$$

where $P_n(x) = \frac{1}{n!} e^{-x} \frac{d^n}{dx^n} (x e^{-x})$ are a Laguerre polynomials, and $f(x) \in L_{\mathbb{R}_+}^2(e^{-x} dx)$. The transform (50) ascertains isomorphism between $L_{\mathbb{R}_+}^2(e^{-x} dx)$ and ℓ^2 .

We extend the Laguerre transform (50) on \mathbb{R}_- in a symmetric way. Then an image of this map yields a space ℓ_-^2 . Let $\ell_{\mathbb{Z}}^2 = \ell_-^2 + \ell_+^2$ is a space of square summable two-sided sequences. Just as for the case of Fourier transform (see Theorem 1.7 in Section 1) a theorem the proof of which repeats the reasonings brought out in [3] holds.

Theorem 2.5. *The Laguerre transform of scattering operator S transfers the operator S into an operator of multiplication by a characteristic function $S_\Delta(n) = I - i\phi(A - nI)^{-1}\phi^*$, $n \in \mathbb{Z}$, i.e.*

$$(51) \quad L_n(Sg) = S_\Delta(n)g_n$$

where $g_n = L_n(g)$.

After realizing the Laguerre transform, the space $L^2\left(\left(\begin{smallmatrix} I & S^* \\ S & I \end{smallmatrix}\right) e^{-\zeta} d\zeta\right)$ passes into the space $\ell_{\mathbb{Z}}^2\left(\begin{smallmatrix} I & S_\Delta^*(n) \\ S_\Delta(n) & I \end{smallmatrix}\right)$ and dilatation \widehat{U}_t (47) is converted into

$$(52) \quad \widehat{U}_t(n)f_n = e^{-itn}f_n.$$

Supspaces D_\pm will have the form

$$D_- = \begin{pmatrix} \ell_-^2 \\ 0 \end{pmatrix}, \quad D_+ = \begin{pmatrix} 0 \\ \ell_+^2 \end{pmatrix}.$$

Therefore H_p is converted to the form

$$(53) \quad \widetilde{H}_p = \left\{ f_n = \begin{pmatrix} f_n^1 \\ f_n^2 \end{pmatrix} \in \ell_{\mathbb{Z}}^2 \begin{pmatrix} I & S_\Delta^*(n) \\ S_\Delta(n) & I \end{pmatrix}; \begin{matrix} f_n^1 + S_\Delta^*(n)f_n^2 \in \ell_+^2 \\ S_\Delta(n)f_n^1 + f_n^2 \in \ell_-^2 \end{matrix} \right\}$$

and a “semigroup” T_t will have the form

$$(54) \quad \widetilde{T}_t(n)f_n = P_{\widetilde{H}_p} e^{-itn}f_n.$$

Thus the following theorem is proved.

Theorem 2.6. *The minimal unitary dilatation U_t (34) in \mathcal{H} (32) of the family of operators T_t (31) with a scattering operator A of collocation Δ (1) is unitary equivalent to $\widehat{U}_t(n)$ (52) in the space $\ell_{\mathbb{Z}}^2\left(\begin{smallmatrix} I & S_\Delta^*(n) \\ S_\Delta(n) & I \end{smallmatrix}\right)$, and the family T_t (31) is unitary equivalent to $\widetilde{T}_t(n)$ (54) in the space \widetilde{H}_p .*

Acknowledgement. The author is grateful to V. A. Zolotarev for setting of the problem and his permanent interest in the work.

REFERENCES

- [1] Livshits, M. S., Yantsevich, A. A., *Theory of operator colligation in Hilbert space*, J. Wiley, N. Y. 1979, Eng. translation.
- [2] Sz.-Nagy, B., Foias, C., *Analyse harmonique des operateurs de l'espace de Hilbert*, Mason, Paris and Akad. Kiado, Budapest 1967; Eng. translation North-Holland, Amsterdam and Akad. Kiado, Budapest 1970.
- [3] Zolotarev, V. A., *Time cones and a functional model on a Riemann surface*, Mat. Sb. **181** (1990), 965–995; Eng. translation in Math. USSR sb. **70** (1991).
- [4] Lax, P., Philips R. S., *Scattering theory*, Academic Press, New York 1967.
- [5] Pavlov, B. S., *Dilatation theory and spectral analysis of nonselfadjoint operators*, Math. programming and Related Questions (Proc. Sevent Winter School, Drogolych, 1994); Theory of Operators in Linear Spaces, Tsentral. Ekonom.-Math. Inst. Akad. Nauk SSSR, Moscow 1976, 3–69; Eng. translation in Amer. Math. Soc. Transl. (2) **115** (1980).
- [6] Mc. Cully J., *The operational calculus of the Laguerre transform*, Ph.D. University of Michigan (1957).
- [7] Kamke, E., *Differentialgleichungen. Lösungsmethoden und Lösungen*, Leipzig 1974.

DEPARTMENT OF MATHEMATICS, IRBID NATIONAL UNIVERSITY
P.O.Box 2600, IRBID, JORDAN
E-mail: raedhat@yahoo.com