

# Counterexample to boundary regularity of a strongly pseudoconvex CR submanifold: An addendum to the paper of Harvey-Lawson

By HING SUN LUK and STEPHEN S.-T. YAU\*

The purpose of this paper is to give a counterexample of Theorem 10.4 in [Ha-La]. In the Harvey-Lawson paper, a global result is claimed, but only a local result is proven. This theorem has had a big impact on CR geometry for almost a quarter of a century because one can use the theory of isolated singularities to study the theory of CR manifolds and vice versa.

*Example.* Consider the following holomorphic map:

$$\begin{aligned} F : \mathbf{C}^2 &\longrightarrow \mathbf{C}^3 \\ (u, v) &\longrightarrow (x, y, z) = (u(u-1), v, u^2(u-1)). \end{aligned}$$

Clearly for any  $c$ ,  $F$  restricted on the line  $\{v = c\}$  is an embedding outside the two points  $(0, c)$  and  $(1, c)$ .  $F$  sends  $(0, t)$  and  $(1, t)$  to  $(0, t, 0)$  for all  $t$ . Now take  $S$ , which is the boundary of a ball  $B = \{(u, v) \in \mathbf{C}^2 : \|(u, v)\| \leq 2\}$ . It is easy to see that the mapping  $F$  restricted on  $S$  is still an embedding. The image of  $S$  under  $F$  is a strongly pseudoconvex CR manifold in  $\mathbf{C}^3$ . The variety that  $F(S)$  bounds is  $F(B)$ . Observe that  $F(B)$  has curve singularities along the line  $(0, t, 0)$ . We remark that  $F(\mathbf{C}^2)$  is a hypersurface  $\{(x, y, z) \in \mathbf{C}^3 : z^2 - zx - x^3 = 0\}$  in  $\mathbf{C}^3$ .

Theorem 10.4 of [Ha-La] was so powerful that it has been used by many researchers. Fortunately, we can replace it by the following theorem, the proof of which will appear elsewhere [Lu-Ya].

**THEOREM.** *Let  $X$  be a strongly pseudoconvex CR manifold of dimension  $2n - 1$ ,  $n \geq 2$ . If  $X$  is contained in the boundary of a bounded strictly pseudoconvex domain  $D$  in  $\mathbf{C}^N$ , then there exists a complex analytic subvariety  $V$  of dimension  $n$  in  $D - X$  such that the boundary of  $V$  is  $X$ . Moreover,  $V$  has boundary regularity at every point of  $X$ , and  $V$  has only isolated singularities in  $V|X$ .*

---

\*Yau's research supported by NSF. Luk's research partially supported by RGC Hong Kong.

*Acknowledgement.* We thank Professor Lempert who first suggested to us that Theorem 10.4 of [Ha-La] may be wrong. In fact it was Lempert who first told the second author a concrete geometric description of how to construct a counterexample to the boundary regularity theorem of Harvey-Lawson (Theorem 10.4 of [Ha-La]) in the higher codimension case. His ideas were realized by two simple examples by us in [Lu-Ya].

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T.,  
HONG KONG

*E-mail address:* hsluk@math.cuhk.edu.hk

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS  
AT CHICAGO, 851 S. MORGAN STREET, CHICAGO, IL, 60607-7045

*E-mail address:* yau@uic.edu

#### REFERENCES

- [Ha-La] F. R. Harvey and H. B. Lawson, Jr., On boundaries of complex analytic varieties, I, *Ann. of Math.* **102** (1975), 223–290.
- [Lu-Ya] H.-S. Luk and S. S.-T. Yau, Kohn-Rossi cohomology, holomorphic de Rham cohomology and the complex plateau problem, preprint.

(Received May 20, 1997)