

On Parallel Vector Fields of Subbundles of Vector Bundles with Linear and Non - Linear Connections

K.Trenčevski and J.Stefanovski

Abstract

In this paper are considered the compatibility conditions for existing of a parallel vector field in vector subbundle of a vector bundle, and further if they are satisfied, the field is found. The compatibility conditions and solutions are given with respect to the initial conditions. The results are generalized for non-linear connections.

Mathematics Subject Classification: 53B15, 35C10.

Key words and Phrases: Parallel vector fields, systems of partial differential equations, non-linear connections.

1 Introduction

This paper is a continuation of the papers [2-6], where [2],[3] and [4] due to the solving of systems of partial differential equations and [5],[6] due to their application in the differential geometry.

In [4] are considered linear and non-linear systems of partial differential equations. The compatibility conditions are found, and if they are satisfied, the solutions are found. All the solutions are given as functional series. The results in [4] are used in [5] for studying the non-linear connections. In [6] are considered parallel vector fields in vector bundles. Indeed the existence of such parallel fields depends on the choice of the initial conditions. The main features in [6] is that the compatibility conditions are found in dependence of the the initial conditions, and if they are satisfied, the vector field is found. It is generalized for non-linear connections and vector bundles with d -connection. In [2] and [3] are considered systems of partial differential equations with linear homogeneous algebraic constraints. It is given an algorithm for reducing to system of lower dimension without constraints and moreover the compatibility conditions for the reduced system are satisfied. This algorithm can also be applied in this paper.

This paper is of interest not only from viewpoint of the differential geometry, but also for differential equations or partial differential equations. Indeed the compatibility

conditions and the solutions for a large class of partial differential equations are given. In this paper we convenient to use the summation convention.

2 On parallel vector fields of subbundles of vector bundles with linear connection

Suppose that the vector field z^i may not be prolonged in general case on the whole manifold M of dimension k , but can be prolonged on distribution η , i.e. on a subbundle of the tangent bundle $T(M)$. Let the rank of η be m ($m \leq k$), and let $Y_{(1)}, \dots, Y_{(m)}$ be a local basis for η . Let ζ be the Lie subalgebra of $T(M)$ generated by $Y_{(1)}, \dots, Y_{(m)}$, and let $Y_{(1)}, \dots, Y_{(p)}$, ($m \leq p \leq k$) be basis of ζ . Then there exist functions $C_{\alpha\beta}^\gamma$ ($\alpha, \beta, \gamma \in \{1, \dots, p\}$) such that

$$(2.1) \quad [Y_{(\alpha)}, Y_{(\beta)}] = C_{\alpha\beta}^\gamma Y_{(\gamma)},$$

i.e. ζ is an integrable distribution. There exist also another basis $X_{(\alpha)} = p_\alpha^\delta Y_{(\delta)}$, $\alpha, \delta \in \{1, \dots, p\}$, such that

$$(2.2) \quad [X_{(\alpha)}, X_{(\beta)}] = 0,$$

i.e.

$$(2.3) \quad p_\alpha^\delta (Y_{(\delta)} p_\beta^\theta) - p_\beta^\delta (Y_{(\delta)} p_\alpha^\theta) + p_\alpha^\delta p_\beta^\gamma C_{\delta\gamma}^\theta = 0.$$

Then there exist uniquely coordinate functions $u^i(x^1, \dots, x^k)$, $i \in \{1, \dots, p\}$ such that

$$(2.4) \quad X_{(\alpha)} u^\beta = \delta_\alpha^\beta$$

and $u^\alpha(0, \dots, 0) = 0$ for $\alpha, \beta \in \{1, \dots, p\}$. We will prove that the system

$$(2.5) \quad \nabla_{Y_{(\alpha)}} Y = 0, \quad (1 \leq \alpha \leq m),$$

is equivalent to

$$(2.6) \quad \nabla_{X_{(\alpha)}} Y = 0, \quad (1 \leq \alpha \leq p).$$

Indeed, the system (2.6) is equivalent with $\nabla_{Y_{(\alpha)}} Y = 0$ for $1 \leq \alpha \leq p$, and it implies (2.5). Conversely, let the system (2.5) is satisfied. Since $R(Y_{(\alpha)}, Y_{(\beta)})Y = 0$, we obtain

$$\nabla_{[Y_{(\alpha)}, Y_{(\beta)}]} Y = R(Y_{(\alpha)}, Y_{(\beta)})Y + \nabla_{Y_{(\alpha)}} (\nabla_{Y_{(\beta)}} Y) - \nabla_{Y_{(\beta)}} (\nabla_{Y_{(\alpha)}} Y) = 0$$

for each $\alpha, \beta \in \{1, \dots, m\}$, and finally we obtain that $\nabla_T Y = 0$ for each vector $T \in \zeta$ and the system (2.6) is satisfied.

Indeed the system (2.6) can be obtained from (2.5) by generating new equations $\nabla_{X_{(\alpha)}} Y = 0$, ($m + 1 \leq \alpha \leq p$) using the Jacobi brackets and then changing the basis $\{Y_1, \dots, Y_p\}$ by $\{X_1, \dots, X_p\}$ such that (2.2) holds. Having this in mind, similarly to the theorem 2.1 [6], the following theorem can be proven.

Theorem 2.1. *Let the analytical system (2.6) with the initial conditions*

$$(2.7) \quad Y(0, \dots, 0) = Z,$$

be given.

(i) *The system (2.6) with (2.7) is integrable if and only if the following system of linear equations*

$$(2.8) \quad \begin{aligned} R(X_{(\alpha)}, X_{(\beta)})T &= 0, \\ (\nabla_{X_{(\gamma_1)}} R(X_{(\alpha)}, X_{(\beta)}))T &= 0, \\ (\nabla_{X_{(\gamma_2)}} \nabla_{X_{(\gamma_1)}} R(X_{(\alpha)}, X_{(\beta)}))T &= 0, \\ &\dots\dots\dots \\ (\nabla_{X_{(\gamma_{n-1})}} \dots \nabla_{X_{(\gamma_1)}} R(X_{(\alpha)}, X_{(\beta)}))T &= 0, \end{aligned}$$

where $\alpha, \beta, \gamma_1, \dots, \gamma_{n-1} \in \{1, \dots, p\}$, has an analytic solution T , such that $T(0, \dots, 0) = Z$.

(ii) *Let the compatibility conditions (2.8) be satisfied and let T be an arbitrary analytical solution of the system (2.8). Then there exist vector fields $Q^{<v_1, \dots, v_p>}$ such that*

$$(2.9) \quad Q^{<0, \dots, 0>} = T,$$

$$(2.10) \quad Q^{<v_1, \dots, v_u+1, \dots, v_p>} = \nabla_{X_{(u)}} Q^{<v_1, \dots, v_p>},$$

and the solution of (2.6) with (2.7) is given by

$$(2.11) \quad Y = \sum_{v_1=0}^{\infty} \sum_{v_2=0}^{\infty} \dots \sum_{v_p=0}^{\infty} \frac{(-u^1)^{v_1}}{v_1!} \frac{(-u^2)^{v_2}}{v_2!} \dots \frac{(-u^p)^{v_p}}{v_p!} \cdot Q^{<v_1, \dots, v_p>}.$$

We notice that $Q^{<v_1, \dots, v_p>}$ is a vector field and u^1, \dots, u^p are scalars, invariant of the coordinate system, and hence the solution (2.11) has covariant form.

In the theorem 2.1 we found the solution of (2.6) on the distribution η , and the solution was unique. Now if there exist r parametric families of such distributions ($r \leq k - p$), then we obtain r parametric solution Y in (2.11), except if the additional restrictions on the initial conditions are required. Specially, if $m = 1$, the compatibility condition vanish, and it is $p = 1$ and $r = k - 1$ in this case.

3 On parallel vector fields of subbundles of vector bundles with non-linear connections

In this section we will consider non-linear connection such that

$$(3.1) \quad \nabla_{X+Y} \neq \nabla_X + \nabla_Y$$

in general case.

Let $X = (X^1, \dots, X^k)$ be a vector. Then we define an operator ∇_X by

$$(3.2) \quad \nabla_X y^j = X^i \frac{\partial}{\partial x^i} y^j + G_s^j(X) y^s.$$

The operator ∇_X acts over the tensor fields, and it holds

$$(3.3) \quad \nabla_X(T_1 \otimes T_2) = (\nabla_X T_1) \otimes T_2 + T_1 \otimes (\nabla_X T_2).$$

Now we will introduce the curvature tensor for the connection ∇_X .

Using (3.2), by direct calculation one obtains

$$(3.4) \quad (\nabla_Y \circ \nabla_X - \nabla_X \circ \nabla_Y - \nabla_{[Y,X]}) y^j = R_q^j(X, Y) y^q,$$

where

$$(3.5) \quad R_q^j(X, Y) = Y^r \frac{\partial}{\partial x^r} G_q^j(X) - X^r \frac{\partial}{\partial x^r} G_q^j(Y) + G_s^j(Y) G_q^s(X) - G_s^j(X) G_q^s(Y) - G_q^j([Y, X]).$$

Note that

$$\begin{aligned} R_q^j(X, Y) &\neq Y^r \frac{\partial}{\partial x^r} G_q^j(X) - X^r \frac{\partial}{\partial x^r} G_q^j(Y) + \\ &+ G_s^j(Y) G_q^s(X) - G_s^j(X) G_q^s(Y) + G_q^j([X, Y]), \\ R_q^j(X, Y) &\neq -R_q^j(Y, X), \end{aligned}$$

and $R_q^j(X, Y)$ is not linear with respect to X and Y . But using the properties (3.3) and (3.4), one can generalize the theorem 2.1 in this case. We will not do that, but we will obtain a further generalization considering a non-linear connection such that $\nabla(X+Y) \neq \nabla X + \nabla Y$ in general case also. Now instead of (3.2) we have

$$(3.6) \quad \nabla_X y^j = X^i \frac{\partial}{\partial x^i} y^j + F^j(x^1, \dots, x^k, X^1, \dots, X^k, y^1, \dots, y^n).$$

Let $Y_{(\alpha)}$ ($1 \leq \alpha \leq m$) be given m vector fields on M . In order to find vector field parallel along $Y_{(\alpha)}$ ($1 \leq \alpha \leq m$) we have to solve the following system

$$(3.7) \quad Y_{(\alpha)}^i \frac{\partial}{\partial x^i} y^j + F_{\alpha}^j(x^1, \dots, x^k, y^1, \dots, y^n) = 0$$

$$(1 \leq \alpha \leq m, 1 \leq j \leq n)$$

of partial differential equations.

We will find the compatibility conditions and the solution of (3.7). In order to do that, first we should transform the given system as follows. The fields $Y_{(\alpha)}$ generate

an integrable distribution ζ of rank p ($m \leq p \leq k$). If $p > m$, then for arbitrary α and β ($\alpha \neq \beta$) we obtain new equations from (3.7) using the Jacobi brackets. Indeed it follows from (3.7) that

$$Y_{(\beta)}^r \frac{\partial}{\partial x^r} \left(Y_{(\alpha)}^i \frac{\partial}{\partial x^i} y^j + F_{(\alpha)}^j \right) - Y_{(\alpha)}^r \frac{\partial}{\partial x^r} \left(Y_{(\beta)}^i \frac{\partial}{\partial x^i} y^j + F_{(\beta)}^j \right) = 0, \text{ i.e.}$$

$$[Y_{(\beta)}, Y_{(\alpha)}]^i \frac{\partial}{\partial x^i} y^j + F^j = 0,$$

where

$$\begin{aligned} F^j &= Y_{(\beta)}^r \frac{\partial}{\partial x^r} F_{(\alpha)}^j - Y_{(\alpha)}^r \frac{\partial}{\partial x^r} F_{(\beta)}^j \\ &= Y_{(\beta)}^r \left[\frac{\partial}{\partial x^r} F_{(\alpha)}^j(x^1, \dots, x^k, y^1, \dots, y^n) + \frac{\partial y^s}{\partial x^r} \frac{\partial}{\partial y^s} F_{(\alpha)}^j(x^1, \dots, x^k, y^1, \dots, y^n) \right] \\ &\quad - Y_{(\alpha)}^r \left[\frac{\partial}{\partial x^r} F_{(\beta)}^j(x^1, \dots, x^k, y^1, \dots, y^n) + \frac{\partial y^s}{\partial x^r} \frac{\partial}{\partial y^s} F_{(\beta)}^j(x^1, \dots, x^k, y^1, \dots, y^n) \right] \\ &= Y_{(\beta)}^r \frac{\partial}{\partial x^r} F_{(\alpha)}^j(x^1, \dots, x^k, y^1, \dots, y^n) - \\ &\quad - F_{(\beta)}^s(x^1, \dots, x^k, y^1, \dots, y^n) \frac{\partial}{\partial y^s} F_{(\alpha)}^j(x^1, \dots, x^k, y^1, \dots, y^n) - \\ &\quad - Y_{(\alpha)}^r \frac{\partial}{\partial x^r} F_{(\beta)}^j(x^1, \dots, x^k, y^1, \dots, y^n) + \\ &\quad + F_{(\alpha)}^s(x^1, \dots, x^k, y^1, \dots, y^n) \frac{\partial}{\partial y^s} F_{(\beta)}^j(x^1, \dots, x^k, y^1, \dots, y^n). \end{aligned}$$

Continuing this procedure, we obtain the following system

$$(3.8) \quad Y_{(\alpha)}^i \frac{\partial}{\partial x^i} y^j + F_{(\alpha)}^j(x^1, \dots, x^k, y^1, \dots, y^n) = 0$$

$$(1 \leq \alpha \leq p, 1 \leq j \leq n)$$

which is equivalent to (3.7), and where $Y_{(\alpha)}$ ($1 \leq \alpha \leq p$) is basis of ζ . Since ζ is an integrable distribution, there exist p vectors $X_{(\alpha)}$ ($1 \leq \alpha \leq p$), $X_{(\alpha)} = C_{\alpha}^{\beta} Y_{(\beta)}$ such that $[X_{(\alpha)}, X_{(\beta)}] = 0$, and moreover there exist p functions $u^i(x^1, \dots, x^k)$ ($1 \leq i \leq p$) such that $X_{(\alpha)} u^i = \delta_{\alpha}^i$. Indeed u^1, \dots, u^k are coordinates on the integrable distribution ζ , and $X_{(\alpha)} = \partial / \partial u^{\alpha}$. Now from (3.8) we obtain the following system

$$X_{(\alpha)}^i \frac{\partial}{\partial x^i} y^j + C_{\alpha}^{\beta} F_{(\beta)}^j(x^1, \dots, x^k, y^1, \dots, y^n) = 0, \text{ i.e.}$$

$$(3.9) \quad X_{(\alpha)}^i \frac{\partial}{\partial x^i} y^j + \Phi_{\alpha}^j(x^1, \dots, x^k, y^1, \dots, y^n) = 0$$

$$(1 \leq \alpha \leq p, 1 \leq j \leq n)$$

which is equivalent to (3.7). In order to consider the system (3.9), assume that Φ_{β}^j are analytical functions, and let

$$(3.10) \quad \Phi_{\beta}^j = \sum_{t \geq 0} \frac{1}{t!} \Gamma_{q_1 \dots q_t \beta}^j(x^1, \dots, x^k) y^{q_1} \dots y^{q_t}.$$

Considering a covariant derivative analogous to (3.19) from [6] and also the curvature tensor, similarly to the theorem 3.2 [6] we have the following

Theorem 3.1. *Let the analytical system (3.9) with the initial conditions*

$$(3.11) \quad Y(0, \dots, 0) = Z$$

be given.

(i) *The system (3.9) with (3.11) is integrable if and only if the following system of non-linear equations*

$$(3.12) \quad \begin{aligned} & \sum_{s \geq 0} \frac{1}{s!} R_{r_1 \dots r_s \alpha \beta}^i t^{r_1} \dots t^{r_s} = 0 \\ & \sum_{s \geq 0} \frac{1}{s!} R_{r_1 \dots r_s \alpha \beta; \gamma_1}^i t^{r_1} \dots t^{r_s} = 0 \\ & \sum_{s \geq 0} \frac{1}{s!} R_{r_1 \dots r_s \alpha \beta; \gamma_1; \gamma_2}^i t^{r_1} \dots t^{r_s} = 0 \\ & \dots\dots\dots \\ & \sum_{s \geq 0} \frac{1}{s!} R_{r_1 \dots r_s \alpha \beta; \gamma_1; \dots; \gamma_{n-1}}^i t^{r_1} \dots t^{r_s} = 0 \end{aligned}$$

has an analytical solution t^j , such that

$$t^j(0, \dots, 0) = z^j \quad (1 \leq j \leq n),$$

where

$$\alpha, \beta, \gamma_1, \dots, \gamma_{n-1} \in \{1, \dots, p\}.$$

(ii) *Let the compatibility conditions for the system (3.9) be satisfied, and let t^j be an arbitrary analytical solution of the system (3.12) with the initial conditions (3.11). Then there exist functions*

$$Q^{r_1 \dots r_s < v_1, \dots, v_p >} \quad (s \geq 0, r_1, \dots, r_s \in \{1, \dots, n\}, v_1, \dots, v_p \in N_0)$$

such that

$$(3.13) \quad \begin{aligned} Q^{<0, \dots, 0>} &= 1, & Q^{r_1 \dots r_s < 0, \dots, 0 >} &= t^{r_1} \cdot t^{r_2} \dots t^{r_s}, \\ Q^{<v_1, \dots, v_u+1, \dots, v_p >} &= 0, \end{aligned}$$

$$(3.14) \quad Q^{r_1 \dots r_s < v_1, \dots, v_u+1, \dots, v_p >} = X_{(u)}^i \frac{\partial}{\partial x^i} Q^{r_1 \dots r_s < v_1, \dots, v_p >} +$$

$$+ \sum_{t \geq 0} \sum_{i=1}^s \frac{1}{t!} \Gamma_{j_1 \dots j_i}^{r_i} Q^{j_1 \dots j_i r_1 \dots r_i \dots r_s \langle v_1, \dots, v_p \rangle}$$

and the solution of (3.9) with (3.11) is given by

$$(3.15) \quad y^r = \sum_{v_1=0}^{\infty} \sum_{v_2=0}^{\infty} \dots \sum_{v_p=0}^{\infty} \frac{(-u^1)^{v_1}}{v_1!} \frac{(-u^2)^{v_2}}{v_2!} \dots \frac{(-u^p)^{v_p}}{v_p!} \times \\ \times Q^{r \langle v_1, \dots, v_p \rangle} \quad (1 \leq r \leq n).$$

The theorem 3.1 can be generalized for parallel tensor fields.

References

- [1] E.Goursat, *Lessons sur le Probleme de Pfaff*, Librairie Scientifique J. Hermann, Paris, 1922.
- [2] J.Stefanovski, K.Trenčevski and V.Covachev: *Charpit system of partial equations with algebraic constraints*, submitted for publication.
- [3] J.Stefanovski, K.Trenčevski and V.Covachev: *Cauchy problem and reduction of system of partial differential equations*, Tamkang Journal of Mathematics, to appear.
- [4] K.Trenčevski, *Solutions of analytical systems of partial differential equations*, Serdica Math. J. 21 (1995), 171-184.
- [5] K.Trenčevski, *A contribution to the non-linear connections*, Tensor N.S. 57 (1996), 226-236.
- [6] K.Trenčevski, *On the parallel vector fields in vector bundles*, Tensor N.S., to appear.

Institute of Mathematics,
P.O.Box 162, 91000 Skopje, Macedonia
e-mail: kostatre@iunona.pmf.ukim.edu.mk

JVP HS Streževo, bul. 1 Maj b.b.
97000 Bitola, Macedonia
e-mail: strezevo@lotus.mpt.com.mk