

Banach J. Math. Anal. 1 (2007), no. 2, 245–251

BANACH JOURNAL OF MATHEMATICAL ANALYSIS ISSN: 1735-8787 (electronic) http://www.math-analysis.org

ON THE STABILITY OF THE QUADRATIC FUNCTIONAL EQUATION IN TOPOLOGICAL SPACES

M. ADAM^{1*} AND S. CZERWIK²

This paper is dedicated to Professor Themistocles M. Rassias.

Submitted by T. Reidel

ABSTRACT. In this paper we investigate the problem of the Hyers–Ulam stability of the generalized quadratic functional equation

f(x+y) + f(x-y) = g(x) + g(y),

where f, g are functions defined on a group with values in a linear topological space.

 1 School of Labour Safety Management, Bankowa 8, PL-40-007 Katowice, Poland.

E-mail address: madam@wszop.edu.pl

 2 Institute of Mathematics, Silesian University of Technology, Kaszubska 23, PL-44-100 Gliwice, Poland.

E-mail address: Stefan.Czerwik@polsl.pl

Date: Received: 20 September 2007; Accepted: 5 November 2007.

^{*} Corresponding author.

²⁰⁰⁰ Mathematics Subject Classification. Primary 39B82; Secondary 54A20.

Key words and phrases. Quadratic functional equation, Hyers–Ulam–Rassias stability in topological spaces.