

Vol. 1 (1996) Paper no. 1, pages 1–18.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

Paper URL

<http://www.math.washington.edu/~ejpecp/EjpVol1/paper1.abs.html>

LÉVY CLASSES AND SELF-NORMALIZATION

Davar Khoshnevisan

Department of Mathematics, University of Utah, Salt Lake City, UT 84112,
davar@math.utah.edu

Abstract: We prove a Chung’s law of the iterated logarithm for recurrent linear Markov processes. In order to attain this level of generality, our normalization is random. In particular, when the Markov process in question is a diffusion, we obtain the integral test corresponding to a law of the iterated logarithm due to Knight.

Keywords: Self-normalization, Lévy classes

AMS subject classification: 60F15, 60J15, 60J45, 60J55

Research supported in part by NSF grant DMS-95-03290.

Submitted to EJP on October 6, 1995. Final version accepted on October 24, 1995.

Lévy Classes and Self-Normalization*

BY DAVAR KHOSHNEVISAN
Department of Mathematics
University of Utah
Salt Lake City, UT. 84112
davar@math.utah.edu

ABSTRACT. We prove a Chung's law of the iterated logarithm for recurrent linear Markov processes. In order to attain this level of generality, our normalization is random. In particular, when the Markov process in question is a diffusion, we obtain the integral test corresponding to a law of the iterated logarithm due to Knight.

A.M.S. SUBJECT 1990 CLASSIFICATION. Primary. 60F15; Secondary. 60J15, 60J45, 60J55
KEY WORDS AND PHRASES. Self-normalization, Lévy classes
SHORT TITLE. Self-normalization

§1. Introduction. Suppose $(B_t; t \geq 0)$ is a one-dimensional Brownian motion. Let $(l_t; t \geq 0)$ denote its local time process evaluated at the level 0. Amongst other results, KNIGHT [Kn, Theorem 3] has proven the following Chung type of iterated logarithm law: almost surely,

$$(1.0) \quad \liminf_{t \rightarrow \infty} \frac{\ln \ln t}{l_t} \sup_{0 \leq s \leq t} |B_s| = 1.$$

Here and throughout, $\ln x \triangleq \log_e(x \vee e)$ where \log_e is the natural logarithm. (1.0) is achieved by demonstrating that for any $\varepsilon > 0$, almost surely $\sup_{0 \leq s \leq t} |B_s| \geq (1 - \varepsilon)l_t / \ln \ln t$, for all t large enough, while there is (a.s.) a random sequence $t_n \rightarrow \infty$, such that for all n , $\sup_{0 \leq s \leq t_n} |B_s| \leq (1 + \varepsilon)l_{t_n} / \ln \ln t_n$. In other words, $(1 + \varepsilon)l_t / \ln \ln t$ is in the upper Lévy class of $\sup_{0 \leq s \leq t} |B_s|$ if $\varepsilon > 0$ and in the lower Lévy class of $\sup_{0 \leq s \leq t} |B_s|$ if $\varepsilon < 0$. It is worth mentioning that (1.0) extends to other diffusions. Furthermore, the results of [Kn] are local, i.e., they hold for $t \rightarrow 0^+$. However, the proofs translate to the case $t \rightarrow \infty$ with no essential changes. Finally, the results of [Kn] are stated for the maximal unreflected process $\sup_{0 \leq s \leq t} B_s$ and due to the choice of the speed measure, the local times in [Kn] are twice ours. The proofs go through with no essential changes.

The main goal of this paper is to extend (1.0) to a broad class of strong Markov processes while providing an exact integral test determining when a suitable function is in the upper (or lower) Lévy class of the maximal process. To this end, let $(X_t; t \geq 0)$ denote a recurrent irreducible linear strong Markov process in the sense of BLUMENTHAL AND GETTOOR [BG]. We shall work under the regime $X_0 = 0$ although this is not necessary. Furthermore, we assume that X possesses local times $(L_t; t \geq 0)$ at 0 (say); for details we refer the reader to [BG], GETTOOR AND KESTEN [GK] and SHARPE [Sh]. In brief, this means that $(L_t; t \geq 0)$ is a continuous additive functional of X whose Revuz measure is proportional to the point mass at 0. (The constant of proportionality does not play a rôle in our main results.)

* Research partially supported by NSF grant DMS-95-03290

It is well known that the recurrence of X implies that $\lim_{t \rightarrow \infty} L_t = \infty$, almost surely (cf. the remarks after Lemma (A.4) below.) Hence, even at this level of generality, it is plausible to try to gauge the lower envelope of $\sup_{s \leq t} |X_s|$ by the random function L_t .

As a consequence of our work, we prove in Theorem (4.2) that when X is a β -symmetric stable Lévy process for $\beta \in (1, 2)$, almost surely,

$$(1.1) \quad \liminf_{t \rightarrow \infty} \left(\frac{\ln \ln t}{L_t} \right)^{1/(\beta-1)} \sup_{0 \leq s \leq t} |X_s| = \left(2\pi^{-1/2} \chi \Gamma\left(\frac{\beta}{2}\right) \Gamma\left(\frac{3-\beta}{2}\right) \Gamma(\beta) \left| \cos\left(\frac{\pi\beta}{2}\right) \right| \right)^{1/(\beta-1)},$$

where the constant χ comes from the characteristic exponent of X as: $\mathbb{P} \exp(i\xi X_t) = \exp(-t\chi|\xi|^\beta)$. Moreover, the process L is normalized as follows:

$$\mathbb{P}\left(\int_0^\infty e^{-s} dL_s \mid X_0 = a\right) = v^1(a).$$

See (4.6) below, together with GETOOR AND KESTEN [GK] for more remarks on normalization of local times. Note also that upon letting $\chi \triangleq \frac{1}{2}$ and $\beta \triangleq 2$, one gets back (1.0). This is not surprising, since in this case X is standard Brownian motion.

The above is part of the motivation behind this paper. Indeed, it is known (cf. DUPUIS [Du], FRISTEDT [F] and WEE [We1, We2]) that for some constant $c = c(\beta) \in (0, \infty)$,

$$\liminf_{t \rightarrow \infty} \left(\frac{\ln \ln t}{t} \right)^{1/\beta} \sup_{0 \leq s \leq t} |X_s| = c.$$

However, c is the principle eigenvalue of corresponding to the Dirichlet problem for Δ^β on $[-1, 1]$ with a 0 exterior condition (cf. WIDOM [Wi]). As such, the value of c is unknown. One of the interesting aspects of (1.1) is that the corresponding problem in self-normalization is quite calculable. Furthermore, together with the LIL of MARCUS AND ROSEN [MR], one can get reasonably good lower bounds on c via $K(\beta)$.

We start with some notation. For any linear Borel set A , we let

$$T(A) \triangleq \inf(s > 0 : X_s \in A).$$

As is sometimes customary, we write \mathbb{P} the probability measure as well as the expectation operator. Furthermore, almost surely (with no reference to the underlying measure) means \mathbb{P} -almost surely. Finally, define for all $x > 0$,

$$(1.2) \quad m(x) \triangleq \frac{1}{\mathbb{P}L_{T(F_x)}},$$

where

$$(1.3) \quad F_x = F(x) \triangleq [x, \infty) \cup (-\infty, -x].$$

For future reference, note that $F(x) \subseteq F(x')$ whenever $x \geq x'$. This implies that $x \mapsto m(x)$ is a decreasing function on $(0, \infty)$. Furthermore, by the strong Markov property, $L_{T(F_x)}$ is an exponential random variable and hence $m(x) \in (0, \infty)$. We are ready to state our main result:

(1.4) **Theorem.** *Suppose $h : (0, \infty) \mapsto (0, \infty)$ is nondecreasing so that $x \mapsto x(m \circ h)(x)$ is also nondecreasing. Then*

$$\mathbb{P}\left(\sup_{0 \leq s \leq t} |X_s| \geq h(L_t), \text{ eventually}\right) = \begin{cases} 1, & \text{if } \int_1^\infty m(h(x))e^{-xm(h(x))} dx < \infty \\ 0, & \text{otherwise} \end{cases}.$$

For the sake of completeness, we also provide the following which generalizes [Kn, Theorem 2]; see MOTOO [Mo] and ERICKSON [Er] for related results.

(1.5) **Theorem.** *Let $h : (0, \infty) \mapsto (0, \infty)$ be a nondecreasing function such that for all x large enough, $x(m \circ h)(x) \leq 1$. Then,*

$$\mathbb{P}\left(\sup_{0 \leq s \leq t} |X_s| \leq h(L_t), \text{ eventually}\right) = \begin{cases} 1, & \text{if } \int_1^\infty m(h(t))dt < \infty \\ 0, & \text{otherwise} \end{cases}.$$

(1.6) **Remark.** Suppose X is a Hunt process on a locally compact space E with a countable basis for which there exists a distinguished point $b \in E$, such that b is regular for $\{b\}$ and is recurrent. Let L_t denote the process of local times at b (cf. [BG] and [Sh] for details). By the Appendix, $L_\infty = \infty$. An inspection of the proofs of Theorems (1.4) and (1.5) show that they immediately extend to results about $(\sup_{s \leq t} g(X_s); t \geq 0)$ where $g : E \mapsto [0, \infty)$ with $g^{-1}(\{0\}) = b$ and F_x is replaced by $g^{-1}([x, \infty))$.

(1.7) **Remark.** Working at the level of generality of Remark (1.6), it is possible to prove local version of Theorems (1.4) and (1.5), i.e., where $t \downarrow 0$. In this case, X need not have a recurrent point; only that L_t exists which is the same as b being regular for $\{b\}$.

Let us finish by discussing what happens when X is a one-dimensional diffusion with scale function S and speed measure M ; see REVUZ AND YOR [RY] for the necessary background. Define for any $x > 0$,

$$g_x(a, b) \triangleq \begin{cases} \frac{(S(a)-S(-x))(S(b)-S(x))}{S(x)-S(-x)}, & \text{if } -x \leq a \leq b \leq x \\ \frac{(S(b)-S(-x))(S(x)-S(a))}{S(x)-S(-x)}, & \text{if } -x \leq b \leq a \leq x \\ 0, & \text{otherwise} \end{cases}.$$

This is the Green's function for the interval $[-x, x]$. One can normalize local times as in [RY, Chapter 7] so that for all $x > 0$,

$$\begin{aligned} \mathbb{P}L_{T(F_x)} &= \mathbb{P}L_{T(\{x, -x\})} \\ &= \int g_x(0, y)M(dy) \\ &= \frac{S(0) - S(-x)}{S(x) - S(-x)} \int_0^x (S(y) - S(x))M(dy) + \frac{S(x) - S(0)}{S(x) - S(-x)} \int_{-x}^0 (S(y) - S(-x))M(dy). \end{aligned}$$

In particular, note that when X is symmetric (i.e., X and $-X$ have the same finite-dimensional distributions), $m(x) = (\int_0^x SdM)^{-1}$.

The organization of this paper is as follows. In Sections 2 and 3, we prove Theorems (1.4) and (1.5), respectively. In Section 4, we show how one can make the necessary computations and use these results when (X_t) is a symmetric stable Lévy process of index $\beta \in (1, 2]$. When $\beta = 2$, we have Brownian motion. In this case, Theorem (1.4) provides the integral test for the Lévy class corresponding to (1.0) above. In Section 5, we discuss the discrete-time analogues of (1.4) and (1.5). More precisely, we discuss self-normalized laws of the iterated logarithms for recurrent walks in \mathbb{Z}^d . For a special class of such walks, another self-normalization appears in the literature. See for example, GRIFFIN AND KUELBS [GK1, GK2] and SHAO [S]; see also LEWIS [Le] and the references

in [S]. The advantage of self-normalizing by local times is apparent in that the class of analyzable processes is quite large. Finally, we provide an appendix which contains several general remarks about NBU random variables and local times.

§2. The Proof of Theorem (1.4). The strategy for proving (1.4) is to time-change

$$(2.1) \quad X_t^* \triangleq \sup_{0 \leq s \leq t} |X_s|,$$

by

$$(2.2) \quad \tau(t) = \tau_t \triangleq \inf \{s > 0 : L_s = t\}.$$

We will exploit the fact that $(X_{\tau(t)}^*; t \geq 0)$ is an extreme-value process. This would in turn allow us to use facts about such processes: see BARNDORFF-NIELSEN [BN], ROBBINS AND SIEGMUND [RS] and KLASS [KL]. Indeed, embedded in our proofs, one can find a streamlined derivation of some of these results. Let us define

$$(2.3) \quad E(t) \triangleq \left\{ \omega : X_{\tau(t)}^* < h(t) \right\},$$

where h is given by (1.4). We begin with some technical lemmas.

$$(2.4) \quad \textbf{Lemma.} \quad (i) \text{ For every } t \geq 0, \mathbb{P}(E(t)) = e^{-tm(h(t))}.$$

$$(ii) \text{ For any } 0 < t < T, \mathbb{P}(E(t) \cap E(T)) = \exp(- (T-t)m(h(T))).$$

Proof. Let $N_x(t)$ denote the total number of excursions before t which hit F_x . (Recall (1.3) as well as the fact that excursions are always counted in the clock τ .) By [It], $(N_x(t); t \geq 0)$ is a Poisson process with mean $\mathbb{P}N_x(t) = t\mathbb{P}N_x(1) = tm(x)$. Part (i) follows since

$$\begin{aligned} \mathbb{P}(E(t)) &= \mathbb{P}(N_{h(t)}(t) = 0) \\ &= e^{-tm(h(t))}, \end{aligned}$$

as desired. To prove part (ii), apply the strong Markov property at time $\tau(t)$. Since $X_{\tau(t)} = X_{\tau(t)^+} = 0$, and since h is increasing,

$$\begin{aligned} \mathbb{P}(E(t) \cap E(T)) &= \mathbb{P}(E(t), \sup_{\tau(t) \leq r \leq \tau(T)} |X_r| < h(T)) \\ &= \mathbb{P}(E(t)) \mathbb{P}\left(\sup_{0 \leq r \leq \tau(T-t)} |X_r| < h(T)\right) \\ &= \mathbb{P}(E(t)) \mathbb{P}(N_{h(T)}(T-t) = 0) \\ &= \mathbb{P}(E(t)) \exp(- (T-t)m(h(T))). \end{aligned}$$

We are done by part (i). ◇

Following ERDŐS [E], define $t_n \triangleq \exp(n/\ln n)$ for all $n \geq 1$. Let us begin with some useful combinatorial properties of $(t_n; n \geq 1)$.

(2.5) **Lemma.** *There exists $c_1 > 1$ such that*

(i) for all $n \geq 1$,

$$c_1^{-1} \frac{t_n}{\ln n} \leq t_{n+1} - t_n \leq c_1 \frac{t_n}{\ln n};$$

(ii) for all $k, n \geq 1$,

$$k \leq c_1 \cdot \frac{t_{n+k} - t_n}{t_n} \cdot \ln n.$$

Proof. By Taylor's expansion, it is easy to see that

$$\lim_{n \rightarrow \infty} \ln n \cdot \frac{t_{n+1} - t_n}{t_n} = 1.$$

Part (i) follows. To prove part (ii), notice that $n \mapsto t_n / \ln n$ is increasing. Hence, by (i),

$$\begin{aligned} t_{n+k} - t_n &= \sum_{j=1}^k (t_{n+j} - t_{n+j-1}) \\ &\geq c_1^{-1} \sum_{j=1}^k \frac{t_{n+j-1}}{\ln(n+j-1)} \\ &\geq c_1^{-1} k \frac{t_n}{\ln n}. \end{aligned}$$

The result follows. ◇

We need some more notation. For positive integers $n < N$, let us define,

$$(2.6) \quad \begin{aligned} \mathcal{G}_n^N &\triangleq [2 \ln n \cdot \ln \ln n, N], \\ \mathcal{B}_n^N &\triangleq [\ln n, 2 \ln n \cdot \ln \ln n), \\ \mathcal{U}_n^N &\triangleq [1, \ln n). \end{aligned}$$

Note that the sets \mathcal{G}_n^N (for good), \mathcal{B}_n^N (for bad) and \mathcal{U}_n^N (for ugly) form an integer partition of $[1, N]$. For integers $0 < n < N$ and $j \geq 1$, define the discrete annuli,

$$(2.7) \quad \mathcal{A}_n^N(j) \triangleq \left\{ 1 \leq k \leq N : \frac{j}{\ln n} \leq 1 - \frac{t_n}{t_{n+k}} \leq \frac{j+1}{\ln n} \right\}.$$

Finally, define

$$(2.8) \quad \mathcal{O}_n^N \triangleq \mathcal{B}_n^N \cup \mathcal{U}_n^N.$$

(2.9) **Lemma.** *There exists a $c_2 > 0$ such that for all integers $n, k, j \geq 1$,*

$$\#(\mathcal{O}_n^N \cap \mathcal{A}_n^N(j)) \leq c_2(j+1)^3.$$

Proof. Without loss of generality, we may suppose that $\mathcal{B}_n^N \cap \mathcal{A}_n^N(j) \neq \emptyset$ and $\mathcal{U}_n^N \cap \mathcal{A}_n^N(j) \neq \emptyset$, for otherwise there is nothing to prove. Let $k \in \mathcal{B}_n^N \cap \mathcal{A}_n^N(j)$. By (2.6) and Lemma (2.5)(ii),

$$(2.10) \quad \begin{aligned} k &\leq c_1 \cdot \frac{t_{n+k} - t_n}{t_n} \cdot \ln n \\ &= c_1 \cdot \left(1 - \frac{t_n}{t_{n+k}}\right) \frac{t_{n+k}}{t_n} \cdot \ln n \\ &\leq c_1(j+1) \frac{t_{n+k}}{t_n}. \end{aligned}$$

By the definition of \mathcal{B}_n^N , $t_{n+k} \leq t_{n+[2 \ln n \cdot \ln \ln n]+1}$. By Taylor's expansion, as $n \rightarrow \infty$,

$$t_{n+[2 \ln n \cdot \ln \ln n]+1} \sim (\ln n)^2 \cdot t_n.$$

Therefore, the following is finite:

$$c_3 \triangleq \sup_n \frac{t_{n+[2 \ln n \cdot \ln \ln n]+1}}{(\ln n)^2 t_n}.$$

To recapitulate, whenever $k \in \mathcal{B}_n^N \cap \mathcal{A}_n^N(j)$,

$$(2.11) \quad k \leq c_3(j+1)(\ln n)^2.$$

On the other hand, for our values of n, N, k and j ,

$$\frac{j+1}{\ln n} \geq 1 - \frac{t_n}{t_{n+k}} \geq 1 - \frac{t_n}{t_{n+[\ln n]}}.$$

Since $\lim_{n \rightarrow \infty} t_{n+[\ln n]}/t_n = e > 1$, it follows that for some $c_4 \in (0, \infty)$, $(j+1) \geq c_4 \ln n$. By (2.11), this means that whenever $k \in \mathcal{B}_n^N \cap \mathcal{A}_n^N(j)$, $k \leq c_3 c_4^{-2}(j+1)^3$. In other words,

$$(2.12) \quad \#(\mathcal{B}_n^N \cap \mathcal{A}_n^N(j)) \leq c_3 c_4^{-2}(j+1)^3.$$

Next, suppose $k \in \mathcal{U}_n^N \cap \mathcal{A}_n^N(j)$. By (2.7) and (2.10),

$$k \leq c_1(j+1) \frac{t_{n+[\ln n]+1}}{t_n}.$$

Since $\lim_{n \rightarrow \infty} t_{n+[\ln n]+1}/t_n = e$, we have shown that for some $c_5 \in (0, \infty)$, $k \leq c_5(j+1)$. In other words, $\#(\mathcal{U}_n^N \cap \mathcal{A}_n^N(j)) \leq c_5(j+1) \leq c_5(j+1)^3$. Together with (2.8) and (2.12), we obtain the lemma. \diamond

We are ready to prove Theorem (1.4). Define $\psi(x) \triangleq xm(h(x))$. Recalling (2.3) and that $t_n = \exp(n/\ln n)$, let $\psi_n \triangleq \psi(t_n)$, $E_n \triangleq E(t_n)$ and $\mathbb{P}_n \triangleq \mathbb{P}(E_n)$, for simplicity. By the argument of ERDŐS [E], it suffices to prove (1.4) when

$$(2.13) \quad c_6 \ln n \leq \psi_n \leq c_7 \ln n,$$

for some $0 < c_6 < c_7 < \infty$ and all $n \geq 1$. With this in mind, it is easy to see that

$$\int_1^\infty m(h(x))e^{-xm(h(x))} dx < \infty \quad \text{if and only if} \quad \sum_n e^{-\psi_n} < \infty.$$

By Lemma (2.4)(i) and the definition of \mathbb{P}_n ,

$$(2.14) \quad \int_1^\infty m(h(x))e^{-xm(h(x))} dx < \infty \quad \text{if and only if} \quad \sum_n \mathbb{P}_n < \infty.$$

Suppose $\int_1^\infty m(h(x))e^{-xm(h(x))} dx = \infty$. It clearly suffices to prove that $\mathbb{P}(E_n, \text{ i.o.}) = 1$. By (2.14), it follows that $\sum_n \mathbb{P}_n = \infty$. By KOCHEN AND STONE [KS], in turn, it suffices to show that

$$\limsup_{N \rightarrow \infty} \frac{\sum_{n=1}^N \sum_{k=1}^{N-n} \mathbb{P}(E_n \cap E_{n+k})}{(\sum_n \mathbb{P}_n)^2} \leq \frac{1}{2}.$$

Recalling the integral partition of $[1, N]$ given by (2.6) and (2.8), it suffices to show the following:

$$(2.15) \quad \mathbb{P}(E_n \cap E_{n+k}) \leq \exp(c_8/\ln n) \mathbb{P}_n \mathbb{P}_{n+k} \quad \forall n < N \text{ and } \forall k \in \mathcal{G}_n^N,$$

$$(2.16) \quad \sum_{n=1}^N \sum_{k \in \mathcal{O}_n^N} \mathbb{P}(E_n \cap E_{n+k}) \leq c_9 \sum_{k=1}^N \mathbb{P}_n.$$

Note that by Lemma (2.4)(i) and (ii),

$$\mathbb{P}(E_n \cap E_{n+k}) = \mathbb{P}_n \mathbb{P}_{n+k} \exp(\psi_{n+k} t_n / t_{n+k}).$$

Thus, to prove (2.15), it suffices to show that there exists some $c_8 > 0$ (independent of $N \geq 1$), such that

$$(2.17) \quad \sup_{k \in \mathcal{G}_n^N} \exp(\psi_{n+k} t_n / t_{n+k}) \leq \exp(c_8 / \ln n).$$

However, for any $k \in \mathcal{G}_n^N$, $t_{n+k} \geq t_{n+2\lfloor \ln n \cdot \ln \ln n \rfloor} \sim t_n (\ln n)^2$. By (2.13) and monotonicity, there exist some $c'_8, c''_8 \in (0, \infty)$, such that

$$\psi_{n+k}^{-1} t_{n+k} / t_n \geq c'_8 (\ln n)^2 \psi_{n+2\lfloor \ln n \cdot \ln \ln n \rfloor}^{-1} \geq c''_8 \ln n.$$

This implies (2.17) with $c_8 \triangleq (c''_8)^{-1}$. As mentioned before, (2.15) follows. To prove (2.16), use Lemmas (2.4)(ii) and (2.9) in the following manner:

$$\begin{aligned} \sum_{n=1}^N \sum_{k \in \mathcal{O}_n^N} \mathbb{P}(E_n \cap E_{n+k}) &= \sum_{n=1}^N \sum_{j=0}^{\infty} \sum_{k \in \mathcal{O}_n^N \cap \mathcal{A}_n^N(j)} \mathbb{P}(E_n \cap E_{n+k}) \\ &= \sum_{n=1}^N \sum_{j=0}^{\infty} \sum_{k \in \mathcal{O}_n^N \cap \mathcal{A}_n^N(j)} \mathbb{P}_n \cdot \exp(-(1 - (t_n/t_{n+k}))\psi_{n+k}) \\ &\leq \sum_{n=1}^N \sum_{j=0}^{\infty} \#(\mathcal{O}_n^N \cap \mathcal{A}_n^N(j)) \cdot \mathbb{P}_n \cdot \exp(-j\psi_n / \ln n), \end{aligned}$$

since $n \mapsto \psi_n$ is increasing. By (2.13), $\psi_n \geq c_7 \ln n$. Hence, by Lemma (2.9),

$$\begin{aligned} \sum_{n=1}^N \sum_{k \in \mathcal{O}_n^N} \mathbb{P}(E_n \cap E_{n+k}) &\leq c_2 \sum_{n=1}^N \sum_{j=0}^{\infty} (j+1)^3 e^{-c_7 j} \mathbb{P}_n \\ &= c_9 \sum_{n=1}^N \mathbb{P}_n, \end{aligned}$$

where $c_9 \triangleq c_2 \sum_{j=0}^{\infty} (j+1)^3 e^{-c_7 j} < \infty$. This proves (2.16) and hence the divergence half of Theorem (1.4).

To prove the convergence half, suppose $\int_1^{\infty} m(h(x)) e^{-xm(h(x))} dx < \infty$. Define

$$\tilde{E}_n \triangleq \left\{ \omega : X_{\tau(t_{n-1})}^* \leq h(t_n) \right\}.$$

(Compare this with $E_n \triangleq E(t_n)$ defined in (2.3).) By the proof of Lemma (2.4)(i),

$$\mathbb{P}(\tilde{E}_n) = \exp(-t_{n-1}\psi_n/t_n).$$

Applying (2.13), we see that

$$\mathbb{P}(\tilde{E}_n) \leq \exp(-\psi_n) \cdot \exp(-c_7 t_{n-1} \ln n / t_n).$$

By Lemma (2.5)(i), $t_{n-1}/t_n \geq 1 - (c_1/\ln n)$. Therefore, for all $n \geq \exp(c_1 + c_7^{-1})$,

$$\mathbb{P}(\tilde{E}_n) \leq e^{-1} \cdot \exp(-\psi_n) = e^{-1} \mathbb{P}_n,$$

by Lemma (2.4)(i). By (2.14) and the Borel–Cantelli lemma, $\mathbb{P}(\tilde{E}_n, \text{ i.o. }) = 0$. A monotonicity argument à la [E] finishes the proof. \diamond

§3. The Proof of Theorem (1.5). Let $s_n \triangleq 2^n$ and recalling (2.1) and (2.2) define,

$$\begin{aligned} F_n &\triangleq \left\{ \omega : X_{\tau(s_n)}^* \geq h(s_n) \right\} \\ \tilde{F}_n &\triangleq \left\{ \omega : X_{\tau(s_{n+1})}^* \geq h(s_n) \right\}. \end{aligned}$$

(3.1) **Lemma.** *There exists $\varepsilon > 0$ so that for all $n \geq 1$,*

- (i) $\mathbb{P}(\tilde{F}_n) \leq 2s_n m(h(s_n))$;
- (ii) $\varepsilon s_n m(h(s_n)) \leq \mathbb{P}(F_n) \leq s_n m(h(s_n))$.

Proof. In the excursion theory notation of §2, using the fact that for $x \geq 0$, $e^{-x} \geq 1 - x$,

$$\begin{aligned} \mathbb{P}(\tilde{F}_n) &= 1 - \mathbb{P}(N_{h(s_n)}(s_{n+1}) = 0) \\ &= 1 - e^{-s_{n+1}m(h(s_n))} \\ &\leq s_{n+1}m(h(s_n)), \end{aligned}$$

proving (i), since $s_{n+1} = 2s_n$. The proof of the upper bound in (ii) is similar. To prove the lower bound in (ii), we can use the fact that $e^{-x} \leq 1 - x + \frac{1}{2}x^2$ for $x \geq 0$ to see that

$$\begin{aligned} \mathbb{P}(F_n) &= 1 - e^{-s_n m(h(s_n))} \\ &\geq s_n m(h(s_n)) \left(1 - \frac{1}{2} s_n m(h(s_n))\right) \\ &\geq \frac{1}{2} s_n m(h(s_n)), \end{aligned}$$

for all n large, since by assumption $xm(h(x)) \leq 1$ for all x large. The result follows for ε appropriately small. \diamond

We are now ready to prove Theorem (1.5). Suppose $\int_1^\infty m(h(t))dt < \infty$. By Lemma (3.1)(i), $\sum_n \mathbb{P}(\tilde{F}_n) < \infty$. By the Borel–Cantelli lemma and monotonicity, it follows that

$$\mathbb{P}(X_{\tau(s)}^* \leq h(s), \text{ eventually }) = 1,$$

as desired. On the other hand, if $\int_1^\infty m(h(t))dt = \infty$, Lemma (3.1)(ii) implies that

$$(3.2) \quad \sum_n \mathbb{P}(F_n) = \infty.$$

Now for any $n, k \geq 1$, consider,

$$\begin{aligned} \mathbb{P}(F_n \cap F_{n+k}) &= \mathbb{P}(X_{\tau(s_n)}^* \geq h(s_n), X_{\tau(s_{n+k})}^* \geq h(s_{n+k})) \\ &= \mathbb{P}(X_{\tau(s_n)}^* \geq h(s_{n+k})) + \mathbb{P}(h(s_n) \leq X_{\tau(s_n)}^* \leq h(s_{n+k}), \sup_{\tau(s_n) \leq r \leq \tau(s_{n+k})} \geq h(s_{n+k})) \\ &\leq \mathbb{P}(X_{\tau(s_n)}^* \geq h(s_{n+k})) + \mathbb{P}(X_{\tau(s_n)}^* \geq h(s_n))\mathbb{P}(X_{\tau(s_{n+k}-s_n)}^* \geq h(s_{n+k})), \end{aligned}$$

by the strong Markov property. By monotonicity, the second summand is bounded above by $\mathbb{P}(F_n)\mathbb{P}(F_{n+k})$. Furthermore, the first summand equals $1 - \mathbb{P}(N_{h(s_{n+k})}(s_n) = 0) = 1 - \exp(-s_n m(h(s_{n+k}))) \leq 2^{-k} s_n m(h(s_{n+k})) \leq 2^{-k} \varepsilon^{-1} \mathbb{P}(F_{n+k})$, by Lemma (3.1)(ii). Hence,

$$\mathbb{P}(F_n \cap F_{n+k}) \leq \mathbb{P}(F_{n+k})(\varepsilon^{-1} 2^{-k} + \mathbb{P}(F_n)).$$

Together with (3.2) and the lemma of KOCHEN AND STONE [KS], this implies that $\mathbb{P}(F_n, \text{ i.o. }) = 1$ which finishes the proof. \diamond

§4. Examples. Throughout this section, X denotes a symmetric β -stable Lévy process with $\beta \in (1, 2]$. That is, X is an infinitely divisible process whose Fourier transform is given by

$$(4.1) \quad \mathbb{P} \exp(i\xi X_t) = e^{-t\chi|\xi|^\beta},$$

where $\xi \in \mathbb{R}^1$ and $\chi > 0$ is arbitrary. It is well-known that X is recurrent and possesses local times L at 0; cf. GETTOOR AND KESTEN [GK] for this and much more. Let $X_t^* \triangleq \sup_{0 \leq s \leq t} |X_s|$. The main result of this section is the following explicit calculation of the function m of (1.2) in this setting.

(4.2) **Theorem.** For all $\beta \in (1, 2]$ and any $x > 0$, $m(x) = K(\beta)x^{1-\beta}$, where,

$$(4.3) \quad K(\beta) \triangleq \frac{2}{\sqrt{\pi}} \chi \Gamma\left(\frac{\beta}{2}\right) \Gamma\left(\frac{3-\beta}{2}\right) \Gamma(\beta) \left| \cos\left(\frac{\pi\beta}{2}\right) \right|.$$

Together with Theorems (1.4) and (1.5), Theorem (4.2) immediately implies our next two results.

(4.4) **Theorem.** Let $f : (0, \infty) \mapsto (0, \infty)$ be nonincreasing so that $x \mapsto xf(x)$ is nondecreasing. Then for the $K(\beta)$ defined in (4.3),

$$\mathbb{P}\left(X_t^* \geq (K(\beta)/f(L_t))^{\frac{1}{\beta-1}}, \text{ eventually} \right) = \begin{cases} 1, & \text{if } \int_1^\infty f(x) \exp(-xf(x)) dx < \infty \\ 0, & \text{otherwise} \end{cases}.$$

(4.5) **Theorem.** Let $f : (0, \infty) \mapsto (0, \infty)$ be nonincreasing so that for all x large enough, $xf(x) \leq 1$. Then,

$$\mathbb{P}\left(X_t^* \leq f(L_t)^{\frac{1}{1-\beta}}, \text{ eventually} \right) = \begin{cases} 1, & \text{if } \int_1^\infty f(x) dx < \infty \\ 0, & \text{otherwise} \end{cases}.$$

It is possible to show that in the case of β -stable Lévy processes with $\beta \in (1, 2]$, $\ln \ln L_t \sim \ln \ln t$, a.s. . Therefore, Theorem (4.2) implies (1.1) and (1.0).

We will use the rest of this section to prove Theorem (4.2).

For $\alpha > 0$, define for all $x \in \mathbb{R}^1$,

$$(4.6) \quad v^\alpha(x) \triangleq \frac{1}{\pi} \int_0^\infty \frac{\cos(rx)}{\alpha + \chi r^\beta} dr.$$

Note that v^α is a symmetric function and in the language of [GK], the α -resolvent density, u^α , of X is given by

$$(4.7) \quad u^\alpha(a, b) \triangleq v^\alpha(a - b).$$

See BRETAGNOLLE [B] and KESTEN [K]. Since 0 is used as a distinguished point, we will have need for one more piece of notation. Define for all $\alpha > 0$,

$$(4.8) \quad \kappa(\alpha) \triangleq u^\alpha(0, 0) = C_\beta \alpha^{-1+1/\beta},$$

where

$$C_\beta \triangleq \frac{B\left(\frac{1}{\beta}, 1 - \frac{1}{\beta}\right)}{\pi \beta \chi^{1/\beta}}.$$

Next, we need to construct a version of the local times L . In order to do so, let $\gamma(\alpha)$ be an independent exponential random variable with mean α^{-1} . Since $u^\alpha(0, \cdot)$ is excessive, the Doob–Meyer decomposition implies the following which is a special case of the work of [GK]:

$$(4.9) \quad v^\alpha(X_0) - v^\alpha(X_{t \wedge \gamma(\alpha)}) - L_{t \wedge \gamma(\alpha)} = \text{a stopped martingale,}$$

stopped at time $\gamma(\alpha)$. Let us start with some technical lemmas.

(4.10) **Lemma.** *For any $x > 0$, there exists a constant $c_{10} = c_{10}(x, \beta) \in (0, \infty)$ such that for any random time $\zeta > 0$,*

$$\mathbb{P}(v^\alpha(X_\zeta); T(F_x) > \gamma(\alpha)) \leq c_{10} \alpha^{1/\beta}.$$

Proof. It follows from (4.6) and (4.7) that $v^\alpha(x) \leq \kappa(\alpha)$. By (4.8),

$$(4.11) \quad \mathbb{P}(v^\alpha(X_\zeta); T(F_x) > \gamma(\alpha)) \leq C_\beta \alpha^{-1+1/\beta} \mathbb{P}(T(F_x) > \gamma(\alpha)).$$

By an eigenfunction expansion (cf. WIDOM [Wi]; or alternatively use the estimates of FRISTEDT [F, Proposition 10.3]), there exists some $c_{11} = c_{11}(x) \in (0, \infty)$, such that for all $r > 0$, $\mathbb{P}(T(F_x) > r) \leq \exp(-c_{11}r)$. Therefore,

$$\begin{aligned} \mathbb{P}(T(F_x) > \gamma(\alpha)) &\leq \alpha \int_0^\infty e^{-c_{11}r} e^{-\alpha r} dr \\ &\leq c_{11}^{-1} \alpha. \end{aligned}$$

Putting this into (4.11) yields the lemma with $c_{10} \triangleq C_\beta c_{11}^{-1}$. ◇

(4.12) **Lemma.** *Fix some $x > 0$. As $\alpha \rightarrow 0^+$,*

$$\mathbb{P}L_{T(F_x) \wedge \gamma(\alpha)} = \kappa(\alpha) - \mathbb{P}v^\alpha(X_{T(F_x)}) + o(1).$$

Proof. By (4.9), almost surely, $(\kappa(\alpha) - v^\alpha(X_{t \wedge \gamma(\alpha)}) - L_{t \wedge \gamma(\alpha)}; 0 \leq t \leq \gamma(\alpha))$ is a mean zero martingale stopped at time $\gamma(\alpha)$. Hence,

$$\begin{aligned} \mathbb{P}L_{T(F_x) \wedge \gamma(\alpha)} &= \kappa(\alpha) - \mathbb{P}(v^\alpha(X_{T(F_x) \wedge \gamma(\alpha)})) \\ &= \kappa(\alpha) - \mathbb{P}(v^\alpha(X_{T(F_x)}); T(F_x) < \gamma(\alpha)) - \mathbb{P}(v^\alpha(X_{\gamma(\alpha)}); T(F_x) > \gamma(\alpha)) \\ &= \kappa(\alpha) - \mathbb{P}(v^\alpha(X_{T(F_x)})) - \\ &\quad - \left(\mathbb{P}(v^\alpha(X_{\gamma(\alpha)}); T(F_x) > \gamma(\alpha)) - \mathbb{P}(v^\alpha(X_{T(F_x)}); T(F_x) > \gamma(\alpha)) \right) \\ &= \kappa(\alpha) - \mathbb{P}(v^\alpha(X_{T(F_x)})) + O(\alpha^{1/\beta}), \end{aligned}$$

by Lemma (4.10). This proves the result. \diamond

The proof of Theorem (4.2). By Lemma (4.12) and the monotone convergence theorem,

$$(4.13) \quad m(x) = \lim_{\alpha \rightarrow 0^+} \frac{1}{\kappa(\alpha) - \mathbb{P}v^\alpha(X_{T(F_x)})}.$$

By (4.6) and (4.8),

$$\begin{aligned} \kappa(\alpha) - v^\alpha(x) &= \frac{1}{\pi} \int_0^\infty \frac{1 - \cos(rx)}{\alpha + \chi r^\beta} dr \\ &\quad \uparrow \frac{1}{\pi} \int_0^\infty \frac{1 - \cos(rx)}{\chi r^\beta} dr, \end{aligned}$$

as $\alpha \rightarrow 0^+$. By (4.13) and the monotone convergence theorem,

$$(4.14) \quad \frac{1}{m(x)} = \frac{1}{\pi} \int_0^\infty \frac{1 - \mathbb{P} \cos(rX_{T(F_x)})}{\chi r^\beta} dr.$$

By (4.1), when $\beta = 2$, $B_t \triangleq (2\chi)^{-1/2} X_t$ is standard Brownian motion. Using symmetry and path continuity of B we see that,

$$\mathbb{P} \cos(rX_{T(F_x)}) = \mathbb{P} \cos(rX_{T(\{-x, x\})}) = \cos(rx).$$

Hence, (4.14) implies that when $\beta = 2$,

$$\frac{1}{m(x)} = \frac{1}{\pi} \int_0^\infty \frac{1 - \cos(rx)}{\chi r^2} dr = \frac{x}{2\chi}.$$

In light of (4.3), this proves Theorem (4.2) when $\beta = 2$. To finish the proof of (4.2), let $\beta \in (1, 2)$. By WIDOM [Wi], (see also BLUMENTHAL ET AL. [BGR] and its references),

$$(4.15) \quad \mathbb{P}(X_{T(F_1)} \in da) = \frac{1}{\pi} \sin\left(\frac{\pi\beta}{2}\right) |a|^{-1} (a^2 - 1)^{-\beta/2} da, \quad \forall a \in \mathbb{R}^1 \setminus [-1, 1].$$

(Actually, the above references prove (4.15) for the case $\chi = 1$. The general case follows from their work together with scaling considerations.) Moreover, by (4.14), scaling and symmetry,

$$\begin{aligned} \frac{1}{m(x)} &= \frac{2}{\pi\chi} \int_x^\infty \mathbb{P}(X_{T(F_x)} \in da) \int_0^\infty \frac{1 - \cos(ra)}{r^\beta} dr \\ &= \frac{2}{\pi\chi} \int_x^\infty a^{\beta-1} \mathbb{P}(X_{T(F_x)} \in da) \int_0^\infty \frac{1 - \cos r}{r^\beta} dr \\ &= \frac{1}{\pi\chi} \mathbb{P}|X_{T(F_x)}|^{\beta-1} \int_0^\infty \frac{1 - \cos r}{r^\beta} dr \\ (4.16) \quad &= \frac{x^{\beta-1}}{\pi\chi} \mathbb{P}|X_{T(F_1)}|^{\beta-1} \int_0^\infty \frac{1 - \cos r}{r^\beta} dr. \end{aligned}$$

By (4.15) and some calculus,

$$\begin{aligned}
\mathbb{P}|X_{T(F_1)}|^{\beta-1} &= \frac{2}{\pi} \sin\left(\frac{\pi\beta}{2}\right) \int_1^\infty a^{\beta-2} (a^2 - 1)^{-\beta/2} da \\
&= \frac{2}{\pi} \sin\left(\frac{\pi\beta}{2}\right) \int_1^\infty a^{-2} (1 - a^{-2})^{-\beta/2} da \\
&= \frac{1}{\pi} \sin\left(\frac{\pi\beta}{2}\right) \int_0^1 u^{-1/2} (1 - u)^{-\beta/2} du \\
&= \frac{1}{\pi} \sin\left(\frac{\pi\beta}{2}\right) B\left(\frac{1}{2}, 1 - \frac{\beta}{2}\right).
\end{aligned}$$

Contour integration shows that for all $z \in \mathbb{C}$, $\Gamma(z)\Gamma(1-z) = \pi/\sin(\pi z)$. Consequently,

$$B\left(\frac{1}{2}, 1 - \frac{\beta}{2}\right) = \frac{\pi^{3/2}}{\Gamma\left(\frac{\beta}{2}\right)\Gamma\left(\frac{3-\beta}{2}\right)\sin\left(\frac{\pi\beta}{2}\right)}.$$

Therefore,

$$(4.17) \quad \mathbb{P}|X_{T(F_1)}|^{\beta-1} = \frac{\sqrt{\pi}}{\Gamma\left(\frac{\beta}{2}\right)\Gamma\left(\frac{3-\beta}{2}\right)}.$$

Recall the following well-known identity:

$$\int_0^\infty \frac{1 - \cos r}{r^\beta} dr = -\frac{1}{2} \frac{\pi}{\Gamma(\beta) \cos\left(\frac{\pi\beta}{2}\right)}.$$

To evaluate the aforementioned integral, we can write $x^{-\beta}(1 - \cos x)$ as $x^{2-\beta}$ times $x^{-2}(1 - \cos x)$ and perform integration by parts to see that the integral equals $(\beta - 1)^{-1} \int_0^\infty x^{1-\beta} \sin x dx$ and the latter is computable by standard means. Together with (4.16) and (4.17), this finishes the proof. \diamond

§5. Recurrent walks in \mathbb{Z}^d . Let Y_1, Y_2, \dots be i.i.d. random vectors, taking values in \mathbb{Z}^d . The corresponding random walk, X_n is defined by

$$(5.1) \quad X_n \triangleq \sum_{j=1}^n Y_j, \quad n \geq 1.$$

We shall assume that X is a genuinely d -dimensional random walk. By recurrence, this implies that $d \leq 2$. Define the local time of X as

$$(5.2) \quad L_n \triangleq \sum_{j=1}^n 1_{\{0\}}(X_j), \quad n \geq 1.$$

Arguments similar to those presented in Section 1 show that the recurrence of X is equivalent to $\mathbb{P}(\lim_{n \rightarrow \infty} L_n = \infty) = 1$. As in §1, define for any Borel set $A \subset \mathbb{Z}^d$,

$$(5.3) \quad T(A) \triangleq \inf(k \geq 1 : X_k \in A).$$

Interpreting $|x|$ as the ℓ_2 norm of $x \in \mathbb{Z}^2 \setminus \{0\}$, we can define m and F_x as in (1.2) and (1.3), respectively. The main result of this section is the following analogue of Theorem (1.4):

(5.4) **Theorem.** Suppose $h : (0, \infty) \mapsto (0, \infty)$ is nondecreasing so that $x \mapsto x(m \circ h)(x)$ is also nondecreasing. Then,

$$\mathbb{P}\left(\max_{1 \leq k \leq n} |X_k| \geq h(L_n), \text{ eventually}\right) = \begin{cases} 1, & \text{if } \int_1^\infty m(h(x))e^{-xm(h(x))} dx < \infty \\ 0, & \text{otherwise} \end{cases}.$$

A discrete-time analogue of (1.5) is also possible; its statement (and proof) is left to the interested reader.

The proof of (5.4) is much like that of (1.4) except that one need use discrete excursion theory. The latter is not as well-documented as continuous-time excursion theory. Therefore, we will include an outline of the proof of (5.4). Before doing so, however, let us investigate two interesting examples. Namely, the simple walk in \mathbb{Z}^1 and the simple walk in \mathbb{Z}^2 .

(5.5) **Example.** Suppose $d = 1$ and $Y_1 = \pm 1$ with probability $\frac{1}{2}$ each. In order to apply (5.4), we first and foremost need to compute the function m . One can proceed in complete analogy to §4. However, there is a simpler way to compute m . For all $x \in \mathbb{Z}^1 \setminus \{0\}$,

$$m(x) = \frac{1}{\mathbb{P}L_{T(F_x)}} = \mathbb{P}(T(F_x) < T_0),$$

where $T_a \triangleq T(\{a\})$. This is essentially a renewal-theoretic argument; see the proof of (5.4). Let $\langle x \rangle \triangleq \lceil |x| \rceil$. Note that $X_{T(F_x)} = X_{\langle x \rangle}$. Hence, $m(x) = \mathbb{P}(T(\{\langle x \rangle, -\langle x \rangle\}) < T_0)$. By symmetry, $m(x) = 2\mathbb{P}(T_{\langle x \rangle} < T_0)$. Therefore, by the gambler's ruin calculation, $m(x) = 1/\langle x \rangle$. Theorem (5.4) then implies the following:

$$\mathbb{P}\left(\max_{1 \leq k \leq n} |X_k| \geq h(L_n), \text{ eventually}\right) = \begin{cases} 1, & \text{if } \sum_{j=1}^\infty \frac{1}{h(j)} e^{-j/h(j)} < \infty \\ 0, & \text{otherwise} \end{cases}.$$

It is possible to show that $\ln \ln L_n \sim \ln \ln n$. Thus, we obtain the analogue of (1.0): almost surely,

$$\liminf_{n \rightarrow \infty} \frac{\ln \ln n}{L_n} \max_{1 \leq k \leq n} |X_k| = 1.$$

◇

(5.6) **Example.** Suppose $d = 2$ and $Y_1 \in \{(\pm 1, \pm 1), (\mp 1, \pm 1)\}$, with probability $\frac{1}{4}$ each. Analysis similar to that of Example (5.5) shows that for all $x \in \mathbb{Z}^2 \setminus \{0\}$,

$$m(x) = \frac{\pi}{2 \ln |x|} (1 + o(1)),$$

where $o(1)$ is a term which goes to 0 as $|x| \rightarrow \infty$. It can be shown that almost surely, $\ln L_n \sim \ln \ln n$. (Many results of this type can be found in RÉVÉSZ [R], for example). Therefore, we obtain

$$\liminf_{n \rightarrow \infty} \frac{\ln \ln \ln n}{L_n} \max_{1 \leq k \leq n} \ln |X_k| = \frac{\pi}{2},$$

almost surely. An interesting consequence of the above — suggested to us by an anonymous referee — is the following: since $\ln \max_{1 \leq k \leq n} |X_k| \sim \frac{1}{2} \ln n$, almost surely,

$$\limsup_{n \rightarrow \infty} \frac{L_n}{\ln n \cdot \ln \ln \ln n} = \frac{1}{\pi}.$$

Thus, we obtain an alternative proof of the LIL of ERDŐS AND TAYLOR (cf. RÉVÉSZ [R, p.202]):
 \diamond

The proof of Theorem (5.4). As in (2.1) and (2.2) define for all $n \geq 1$,

$$(5.7) \quad \begin{aligned} X_n^* &\triangleq \max_{1 \leq k \leq n} |X_k| \\ \tau(n) &\triangleq \min(k \geq 1 : L_k = n) \end{aligned}.$$

Notice that $\tau(n)$ is none other than the n -th return time to 0. In analogy with (2.3), define for all $n \geq 1$

$$E(n) \triangleq \left\{ \omega : X_{\tau(n)}^* < h(n) \right\}.$$

Since the excursions of X from 0 are i.i.d., $X_{\tau(n)}^* = \max_{1 \leq j \leq n} \max_{\tau(j-1) < k \leq \tau(j)} |X_k|$ is the maximum of n i.i.d. random variables ($\tau(0) \triangleq 0$). Moreover,

$$\mathbb{P}(X_{\tau(1)}^* \geq x) = \mathbb{P}(T(F_x) < T_0),$$

where $T_a \triangleq T(\{a\})$. Hence, by renewal theory, $\mathbb{P}(X_{\tau(1)}^* \geq x) = (\mathbb{P}L_{T(F_x)})^{-1} = m(x)$. To see this, define path-valued processes e_k as follows:

$$e_k(j) \triangleq \left(X_i; \tau(k-1) \leq i \leq \tau(k) \right).$$

The e_k 's are the excursions from 0 and are ordered according to their natural clock, τ , in which things are measured. Let A be any Borel set on the space of paths. (We shall not bother with the topological asides here. They are straight-forward, especially as the state space is discrete.) By the strong Markov property, $N(A)_m \triangleq \sum_{j=1}^m 1_A(e_j)$ is a Bernoulli process in that it is a sum of m i.i.d. random variable and its growth times are geometric random variables with parameter $p(A) \triangleq \mathbb{P}(e_1 \in A)$. Applying the optional sampling theorem, we see that for all stopping times σ (in the filtration of $N(A)$) and all $n \geq 1$, $\mathbb{P}N(A)_{\sigma \wedge n} = p(A)\mathbb{P}(\sigma \wedge n)$. By the monotone convergence theorem, $\mathbb{P}N(A)_\sigma = p(A)\mathbb{P}\sigma$. Now let σ be the first growth time of $N(A)$. In other words, $\sigma = L_{T(A)}$. Clearly, σ is a $N(A)$ -stopping time. Since $N(A)_\sigma = 1$, it follows that $p(A) = (\mathbb{P}L_{T(A)})^{-1}$. The above argument can be pushed further to show that $(m, A) \mapsto N(A)_m$ is a Bernoulli point process in that it is a Bernoulli process for each A and $N(A)$ and $N(B)$ are independent if $A \cap B = \emptyset$. This is the essentials of discrete excursion theory.

Now to see why $\mathbb{P}(X_{\tau(1)}^* \geq x) = (\mathbb{P}L_{T(F_x)})^{-1}$, let $A \triangleq F_x$. Therefore, $\mathbb{P}(E(n)) = (1 - m(h(n)))^n$. Without loss of generality, we can assume that $h(n) \rightarrow \infty$, as $n \rightarrow \infty$ for otherwise, there is nothing to prove. Furthermore, just as in (2.13), we can assume that for some $c_{12} > 1$,

$$c_{12}^{-1} \ln \ln k \leq m(h(k)) \leq c_{12} \ln \ln k.$$

Therefore, for some $c_{13} \in (0, 1)$,

$$c_{13} e^{-nm(h(n))} \leq \mathbb{P}(E(n)) \leq e^{-nm(h(n))}.$$

This is the analogue of Lemma (2.4)(i). Lemma (2.4)(ii)'s analogue is proved along the same lines. Since the combinatorial aspects of the proof of Theorem (5.4) are not at all different from those of Theorem (1.5), the same proof now goes through without any further difficulties. \diamond

Appendix: NBU random variables and local times. Following SHAKED AND SHANTHIKUMAR [SS, p.11], we say that a positive random variable S is *NBU* if for all $a, b > 0$,

$$(A.1) \quad \mathbb{P}(S > a + b) \leq \mathbb{P}(S \geq a)\mathbb{P}(S \geq b).$$

(*NBU* stands for *New Better than Used*.) As is customary, we write $\|S\|_p \triangleq \mathbb{P}^{1/p} S^p$ for the moments of S . Let us begin with a basic $L^p(\mathbb{P})$ estimate for *NBU* random variables.

(A.2) **Lemma.** *If S is NBU, for all $p \geq 1$, $\|S\|_p \leq \Gamma^{1/p}(1+p)\|S\|_1$, where Γ is the standard Gamma function.*

Proof. When $p = 1$, this is clear. Suppose we could prove the result for all integer $p \geq 1$. By the convexity theorem of RIESZ (cf. STEIN [St], Theorem B.1 for a version of this), we are done. To this end, we proceed by induction. We can suppose that for some integer $p \geq 1$, $\|S\|_p \leq \Gamma^{1/p}(1+p)\|S\|_1$. We will strive to prove that it also holds for $p + 1$. To do this, we use the induction hypothesis and integrate by parts as follows:

$$\begin{aligned} \|S\|_1^{p+1} &= \|S\|_1 \cdot \|S\|_1^p \\ &\geq \frac{1}{\Gamma(1+p)} \|S\|_1 \cdot \|S\|_p^p \\ &= \frac{p}{\Gamma(p+1)} \int_0^\infty \mathbb{P}(S \geq x) dx \cdot \int_0^\infty y^{p-1} \mathbb{P}(S \geq y) dy \\ &\geq \frac{p}{\Gamma(1+p)} \int_0^\infty \int_0^\infty y^{p-1} \mathbb{P}(S \geq x+y) dx dy, \end{aligned}$$

by (A.1). Making a change of variables from (x, y) to $(u, v) = (x + y, y)$, we see that

$$\begin{aligned} \|S\|_p^{p+1} &= \frac{p}{\Gamma(1+p)} \int_0^\infty \int_v^\infty v^{p-1} \mathbb{P}(S \geq u) du dv \\ &= \frac{1}{\Gamma(1+p)} \int_0^\infty u^p \mathbb{P}(S \geq u) du \\ &= \frac{1}{(p+1)\Gamma(1+p)} \|S\|_{p+1}^{p+1} \\ &= \frac{1}{\Gamma(p+2)} \|S\|_{p+1}^{p+1}, \end{aligned}$$

as desired. ◇

It is not difficult to see that the above is sharp.

(A.3) **Lemma.** *Let S_n be a sequence of NBU random variables. Suppose further that $\mathbb{P}S_n$ increases to infinity, as $n \rightarrow \infty$. Then with probability one, $\limsup_{n \rightarrow \infty} S_n = \infty$.*

Proof. Take $p > 1$ and $\theta \in (0, 1)$ and let 1_A be the characteristic function of a Borel set $A \subseteq (0, \infty)$. By Lemma (A.2) and Hölder's inequality,

$$\begin{aligned} \|S_n\|_1 &= \|S_n 1_{(-\infty, \theta \|S_n\|_1)}(S_n)\|_1 + \|S_n 1_{[\theta \|S_n\|_1, \infty)}(S_n)\|_1 \\ &\leq \theta \|S_n\|_1 + \|S_n\|_p \left(\mathbb{P}(S_n \geq \theta \|S_n\|_1) \right)^{1/q} \\ &\leq \theta \|S_n\|_1 + \Gamma^{1/p}(p+1) \|S_n\|_1 \left(\mathbb{P}(S_n \geq \theta \|S_n\|_1) \right)^{1/q}, \end{aligned}$$

where $q^{-1} + p^{-1} = 1$. Solving, we obtain,

$$\mathbb{P}(S_n \geq \theta \|S_n\|_1) \geq (1 - \theta)^q \cdot \Gamma^{-q/p}(p + 1),$$

By the monotonicity of $\|S_n\|_1$,

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} S_n = \infty\right) \geq (1 - \theta)^q \cdot \Gamma^{-q/p}(p + 1).$$

First, let $\theta \downarrow 0$ and then $p \downarrow 1$ to obtain the result. \diamond

The relation to local times is the following result which is a consequence of the strong Markov property.

(A.4) **Lemma.** *For each $t \geq 0$, L_t is NBU.*

Since $t \mapsto L_t$ is a.s. increasing, Lemmas (A.3) and (A.4) together imply that $\lim_{t \rightarrow \infty} L_t = \infty$, a.s. if and only if $\mathbb{E}L_\infty = \infty$. This condition is the well-known condition that X is point recurrent. (Indeed, the corresponding potential measure is $U(A) \triangleq \int_0^\infty P(X_s \in A) ds$; cf. BLUMENTHAL AND GETOOR [BG] and SHARPE [Sh].)

Acknowledgements. This paper owes much of its inspiration, content and clarity to Richard Bass, Jean Bertoin, Robert Blumenthal, Bruce Erickson, Patrick Fitzsimmons, Thomas Lewis, Jim Pitman, Zhan Shi and Marc Yor to all of whom many thanks are due. The bulk of this work was done while the author was visiting the Department of Mathematical Sciences at the Johns Hopkins University. I wish to thank them, especially Daniel Naiman, John Wierman and Colin Wu for their hospitality and support. Finally, several hearty thanks are due to an anonymous referee for providing me with corrections as well as connections to other parts of the literature.

References.

- [BN] O. BARNDORFF-NIELSEN (1961). On the rate of growth of partial maxima of a sequence of independent and identically distributed random variables. *Math. Scand.*, **9**, 383–394
- [BG] R. BLUMENTHAL AND R.K. GETOOR (1968). *Markov Processes and Potential Theory*. Academic Press. New York
- [BGR] R. BLUMENTHAL, R.K. GETOOR AND D.B. RAY (1961). On the distribution of first hits for the symmetric stable process. *Trans. Amer. Math. Soc.*, **99**, 540–554
- [B] J. BRETAGNOLLE (1970). Résultat de Kesten sur les processus à accroissements indépendants. *Sém. de Prob.*, Lecture Notes in Mathematics, **191**, 21–36
- [Du] C. DUPUIS (1974). Mesure de Hausdorff de la trajectoire de certains processus à accroissements indépendants et stationnaires, *Sém. de Prob. VIII*, Lecture Notes in Mathematics, **381**, 37–77
- [E] P. ERDŐS (1942). On the law of the iterated logarithm. *Ann. of Math.*, Vol. 43, No. 3, 419–436
- [Er] K.B. ERICKSON (1994). Divergent sums over excursions. *Stoch. Proc. Appl.*, **54** 175–182
- [F] B. FRISTEDT (1974). Sample Functions of Stochastic Processes with Stationary, independent increments. *Advances in Probability III*, Dekker, 241–397
- [GK] R.K. GETOOR AND H. KESTEN (1972). Continuity of local times for Markov processes. *Comp. Math.*, **24**, 277–303
- [GK1] P. GRIFFIN AND J. KUELBS (1991). Some extensions of the LIL via self normalizations. *Ann. Prob.*, **19**, 380–395
- [GK2] P. GRIFFIN AND J. KUELBS (1989). Self-normalized laws of the iterated logarithm. *Ann. Prob.*, **17**, 1571–1601

- [It] K. ITÔ (1970). Poisson processes attached to Markov processes. *Proc. Sixth Berkeley Symp. Math. Stat. Prob.*, Vol. 3. University of California, Berkeley, 225–239
- [K] H. KESTEN (1969). Hitting probabilities of single points for processes with stationary independent increments. *Memoirs A.M.S.*, **93**, American Mathematical Society, Providence, Rhode Island
- [Kl] M.J. KLASS (1985). The Robbins–Siegmund series criterion for partial maxima. *Ann. Prob.*, **4**, 1369–1370
- [Kn] F. KNIGHT (1973). Local Variation of diffusion. *Ann. Prob.*, **1**, 1026–1034
- [KS] S.B. KOCHEN AND C.J. STONE (1964). A note on the Borel–Cantelli lemma. *Ill. J. Math.*, **8**, 248–251
- [Le] T.M. LEWIS (1992). A self normalized law of the iterated logarithm for random walk in random scenery. *J. Th. Prob.*, **5**, 629–659
- [Ma] B. MAISONNEUVE (1975). Exit systems. *Ann. Prob.*, **3**, 395–411
- [MR] M. B. MARCUS AND J. ROSEN (1994). Laws of the iterated logarithm for the local times of symmetric Lévy processes and recurrent random walks. *Ann. Prob.*, **22**, 626–658
- [Mo] M. MOTOO (1959). Proof of the law of the iterated logarithm through diffusion equation. *Ann. Inst. Stat. Math.*, **10**, 21–28
- [R] P. RÉVÉZ (1990). *Random walks on random and nonrandom environments*. World Scientific, Singapore
- [RY] D. REVUZ AND M. YOR (1991). *Continuous Martingales and Brownian Motion*. Springer–Verlag, N.Y.
- [RS] H. ROBBINS AND D. SIEGMUND (1972). On the law of the iterated logarithm for maxima and minima. *Proc. 6th Berkeley Symp. Math. Stat. Prob.*, Vol. III, 51–70
- [SS] M. SHAKED AND J.G. SHANTHIKUMAR (1994). *Stochastic Orders and Their Applications*. Academic Press, San Diego, CA.
- [S] Q.–M. SHAO (1994). Self normalized large deviations. Preprint
- [Sh] M. SHARPE (1988). *General Theory of Markov Processes*. Academic Press, N.Y.
- [St] E. M. STEIN (1986). *Singular Integrals and Differentiability Properties of Functions*. Fifth Edition. Princeton University Press, Princeton, N.J.
- [We1] I.S. WEE (1990). Lower functions for asymmetric Lévy processes, *Prob. Th. Rel. Fields*, **85**, 469–488
- [We2] I.S. WEE (1988). Lower functions for processes with stationary independent increments, *Prob. Th. Rel. Fields*, **77**, 551–566
- [Wi] H. WIDOM (1961). Stable processes and integral equations. *Trans. Amer. Math. Soc.*, **98**, 430–449

Davar Khoshnevisan
 Department of Mathematics
 University of Utah
 Salt Lake City, UT 84112
 davar@math.utah.edu