

Asymptotics of Bernoulli random walks, bridges, excursions and meanders with a given number of peaks

Jean-Maxime Labarbe
Université de Versailles
45 avenue des Etats-Unis
78035 Versailles cedex
labarbe@math.uvsq.fr

Jean-François Marckert
CNRS, LaBRI
Université Bordeaux 1
351 cours de la Libération
33405 Talence cedex
marckert@labri.fr

Abstract

A Bernoulli random walk is a random trajectory starting from 0 and having i.i.d. increments, each of them being +1 or -1 , equally likely. The other families quoted in the title are Bernoulli random walks under various conditions. A peak in a trajectory is a local maximum. In this paper, we condition the families of trajectories to have a given number of peaks. We show that, asymptotically, the main effect of setting the number of peaks is to change the order of magnitude of the trajectories. The counting process of the peaks, that encodes the repartition of the peaks in the trajectories, is also studied. It is shown that suitably normalized, it converges to a Brownian bridge which is independent of the limiting trajectory. Applications in terms of plane trees and parallelogram polyominoes are provided, as well as an application to the “comparison” between runs and Kolmogorov-Smirnov statistics.

Key words: Weak convergence, Bernoulli random walks, Brownian meander, bridge, excursion, peaks.

AMS 2000 Subject Classification: Primary 60J65, 60B10.

Submitted to EJP on January 25 2006, final version accepted February 28 2007.

1 Introduction

Let $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ be the set of non-negative integers. For any $n \in \mathbb{N}$, we denote by \mathcal{W}_n the set of Bernoulli chains with n steps :

$$\mathcal{W}_n = \{\mathbf{S} = (S(i))_{0 \leq i \leq n} : S(0) = 0, S(i+1) = S(i) \pm 1 \text{ for any } i \in \llbracket 0, n-1 \rrbracket\}.$$

The sets of Bernoulli bridges \mathcal{B}_n , Bernoulli excursions \mathcal{E}_n , Bernoulli meanders \mathcal{M}_n with n steps are defined by

$$\begin{aligned} \mathcal{B}_n &= \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, S(n) = 0\}, \\ \mathcal{E}_n &= \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, S(n) = 0, S(i) \geq 0 \text{ for any } i \in \llbracket 0, n \rrbracket\}, \\ \mathcal{M}_n &= \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, S(i) \geq 0 \text{ for any } i \in \llbracket 0, n \rrbracket\}. \end{aligned}$$

The cardinalities of these sets are given by

$$\#\mathcal{W}_n = 2^n, \quad \#\mathcal{B}_{2n} = \binom{2n}{n}, \quad \#\mathcal{E}_{2n} = \frac{1}{n+1} \binom{2n}{n}, \quad \#\mathcal{M}_n = \binom{n}{\lfloor n/2 \rfloor}, \quad (1)$$

and for every odd number n , $\mathcal{B}_n = \mathcal{E}_n = \emptyset$. The two first formulas are obvious, the third can be proved for instance thanks to the cyclical lemma (see also the 66 examples of the appearance of the Catalan numbers $\#\mathcal{E}_{2n}$ in combinatorics in Stanley (23, ex. 6.19 p.219)), and the last one, may be proved iteratively or thanks to a bijection with Bernoulli bridges (see Section 4.4).

Let $n \in \mathbb{N}$. For every $\mathbf{S} \in \mathcal{W}_n$, the set of peaks of \mathbf{S} , denoted by \mathbf{S}_\wedge , is defined by

$$\mathbf{S}_\wedge = \{x : x \in \llbracket 1, n-1 \rrbracket, S(x-1) = S(x+1) = S(x) - 1\}.$$

The set $(-\mathbf{S})_\wedge$ is called the set of valleys of \mathbf{S} : it is easy to check that for any \mathbf{S} , $\#\mathbf{S}_\wedge - \#(-\mathbf{S})_\wedge$ belongs to $\{+1, 0, -1\}$. The value of this difference depends only on the signs of the first and last steps of \mathbf{S} . In this paper, we focus only on the number of peaks and we denote by $\mathcal{W}_n^{(k)}$ (resp. $\mathcal{B}_n^{(k)}$, $\mathcal{E}_n^{(k)}$ and $\mathcal{M}_n^{(k)}$) the subset of \mathcal{W}_n , (resp. \mathcal{B}_n , \mathcal{E}_n , \mathcal{M}_n) of trajectories having exactly k peaks (for any $k > \lfloor n/2 \rfloor$ these sets are empty). We have

Proposition 1. *For any $k \geq 0$ and any $n \geq 0$,*

$$\#\mathcal{W}_n^{(k)} = \binom{n+1}{n-2k}, \quad \#\mathcal{B}_{2n}^{(k)} = \binom{n}{k}^2, \quad \#\mathcal{E}_{2n}^{(k)} = \frac{1}{n} \binom{n}{k} \binom{n}{k-1}, \quad \#\mathcal{M}_n^{(k)} = \binom{\lfloor n/2 \rfloor}{k} \binom{\lceil n/2 \rceil}{k},$$

where, by convention

$$\binom{m}{p} = \begin{cases} \frac{m!}{p!(m-p)!} & \text{if } p \text{ and } m \text{ are non negative integers and } p \in \llbracket 0, m \rrbracket, \\ 0 & \text{in any other cases.} \end{cases}$$




Figure 1: Trajectories from $\mathcal{W}_{12}^{(3)}$, $\mathcal{B}_{12}^{(2)}$, $\mathcal{E}_{12}^{(3)}$, and $\mathcal{M}_{12}^{(4)}$. Black dots correspond to peaks.

The formula giving $\#\mathcal{E}_{2n}^{(k)}$ is due to Narayana (18) computed in relation with pairs of k -compositions of n satisfying some constraints (see also Stanley (23, ex. 6.36 p.237) and Theorem 3.4.3 in Krattenthaler (14), with $\mu = 1$, $e_1 = e_2 = n$). The formula giving $\#\mathcal{B}_{2n}^{(k)}$ can be also found in (14, Formula (3.6)). The survey (14) of Krattenthaler is very related to the present work and we refer the interested reader to this work and references therein. Among other things, he investigates the link between the number of peaks in some Bernoulli trajectories and the trajectories themselves (14, Theorem 3.4.4) (see also Section 1.3). His results concern also the enumerations of trajectories with a prescribed number of peaks staying above a given line, as well as some multidimensional case (the number of non intersecting Bernoulli trajectories with a total number of turns).

Let $\mathbb{P}_n^w, \mathbb{P}_n^b, \mathbb{P}_n^e$ and \mathbb{P}_n^m be the uniform law on $\mathcal{W}_n, \mathcal{B}_n, \mathcal{E}_n$, and \mathcal{M}_n and $\mathbb{P}_n^{w,(k)}, \mathbb{P}_n^{b,(k)}, \mathbb{P}_n^{e,(k)}$ and $\mathbb{P}_n^{m,(k)}$ be the uniform law on $\mathcal{W}_n^{(k)}, \mathcal{B}_n^{(k)}, \mathcal{E}_n^{(k)}$ and $\mathcal{M}_n^{(k)}$. For $x \in \{w, b, e, m\}$, a random variable under $\mathbb{P}_n^{x,(k)}$ is then simply a random variable under \mathbb{P}_n^x conditioned to have k peaks. We are interested in the asymptotic behavior of random chains under the distributions $\mathbb{P}_n^{x,(k)}$, when n and $k = \mathcal{K}_n$ go to infinity.

Let $C[0, 1]$ be the set of continuous functions defined on $[0, 1]$ with real values. For any $\mathbf{S} \in \mathcal{W}_n$, denote u_n the function in $C[0, 1]$ obtained from \mathbf{S} by the following rescaling:

$$u_n(t) = \frac{1}{\sqrt{n}}(S(\lfloor nt \rfloor) + \{nt\}(S(\lceil nt \rceil) - S(\lfloor nt \rfloor))) \text{ for any } t \in [0, 1]. \quad (2)$$

We call Brownian bridge **b**, Brownian excursion **e** and Brownian meander **m** the (normalized) processes characterized as follows : let w be a 1-dimensional standard Brownian motion. Let $d = \inf\{t : t \geq 1, w_t = 0\}$ and $g = \sup\{t : t \leq 1, w_t = 0\}$. Almost surely, we have $d - g > 0$, $g \in (0, 1)$. The processes **b**, **e** and **m** have the following representations :

$$(\mathbf{b}_t)_{t \in [0,1]} \stackrel{(d)}{=} \left(\frac{w_{gt}}{\sqrt{g}} \right)_{t \in [0,1]}, \quad (\mathbf{e}_t)_{t \in [0,1]} \stackrel{(d)}{=} \left(\frac{|w_{d+(g-d)t}|}{\sqrt{d-g}} \right)_{t \in [0,1]}, \quad (\mathbf{m}_t)_{t \in [0,1]} \stackrel{(d)}{=} \left(\frac{|w_{(1-g)t}|}{\sqrt{1-g}} \right)_{t \in [0,1]}.$$

As a consequence of the Donsker (6) theorem (for $x = w$), Kaigh (13) (for $x = e$), Iglehart (11) or Belkin (2) (for $x = m$), and Liggett (15) (for $x = b$),

Theorem 2. For any $x \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$, under \mathbb{P}_n^x , $u_n \xrightarrow{(d)} x$ in $C[0, 1]$ endowed with the topology of the uniform convergence.

In the case $x \in \{\mathbf{b}, \mathbf{e}\}$, even if not specified, it is understood that $n \rightarrow +\infty$ in $2\mathbb{N}$.

In fact, Theorem 2 can be proved directly, thanks to the elementary enumeration of paths passing via some prescribed positions in the model of Bernoulli paths. The method used to show the tightnesses in our Theorem 4 may be used to prove the tightness in Theorem 2; thanks to some probability tricks, this reduces to show the tightness under $\mathbb{P}_n^{\mathbf{w}}$, which is simple.

The finite dimensional distributions of $\mathbf{w}, \mathbf{e}, \mathbf{b}$ and \mathbf{m} are recalled in Section 3.1. Numerous relations exist between these processes, and their trajectories, and a lot of parameters have been computed. We refer to Bertoin & Pitman (4), Biane & Yor (5), Pitman (20) to have an overview of the subject. These convergences have also provided some discrete approaches to the computation of values attached to these Brownian processes, and the literature about that is considerable, see e.g. Csáki & Y. Hu (7), and references therein.

We introduce the counting process of the number of peaks : for any $\mathbf{S} \in \mathcal{W}_n$, denote by $\Lambda(\mathbf{S}) = (\Lambda_l(\mathbf{S}))_{l \in [0, n]}$ the process :

$$\Lambda_l(\mathbf{S}) = \#\mathbf{S}_\wedge \cap [0, l] \text{ for any } l \in [0, n]. \quad (3)$$

For $\mathbf{S} \in \mathcal{W}_n$, $\Lambda_n(\mathbf{S}) = \#\mathbf{S}_\wedge$ is simply the total number of peaks in \mathbf{S} . We have

Proposition 3. For any $x \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$, under \mathbb{P}_n^x ,

$$\frac{\Lambda_n - n/4}{\sqrt{n}} \xrightarrow{(d)} \mathcal{N}(0, 1/16),$$

where $\mathcal{N}(0, 1/16)$ denotes the centered Gaussian distribution with variance 1/16.

We will now describe the main result of this paper. Its aim is to describe the influence of the number of peaks on the shape of the families of trajectories introduced above. We will then condition the different families by $\#\mathbf{S}_\wedge = \mathcal{K}_n$ for a general sequence (\mathcal{K}_n) satisfying the following constraints :

$$(H) = \left(\text{For any } n, \mathcal{K}_n \in \mathbb{N}, \lim_n \mathcal{K}_n = +\infty, \lim_n n/2 - \mathcal{K}_n = +\infty \right).$$

Notice that for every $\mathbf{S} \in \mathcal{W}_n$, $\#\mathbf{S}_\wedge \in [0, \lfloor n/2 \rfloor]$ and then (H) is as large as possible to avoid that the sequences \mathcal{K}_n and $n/2 - \mathcal{K}_n$ have a finite accumulation point.

We set $p_n := 2\mathcal{K}_n/n$ and

$$\beta_n := \sqrt{n(1-p_n)/p_n}, \text{ and } \gamma_n := \sqrt{np_n(1-p_n)} = p_n\beta_n. \quad (4)$$

Each peak can be viewed to be made by two consecutive steps; hence, if you pick at random one step of a trajectory under $\mathbb{P}_n^{\mathbf{x},(\mathcal{K}_n)}$, the probability that this step belongs to a peak is p_n .

We consider \mathbf{S} and $\Lambda(\mathbf{S})$ as two continuous processes on $[0, n]$, the values between integer points being defined by linear interpolation. The normalized versions of \mathbf{S} and $\Lambda(\mathbf{S})$ are respectively denoted by s_n and λ_n :

$$s_n(t) := \frac{S(nt)}{\beta_n} \quad \text{and} \quad \lambda_n(t) := 2 \frac{\Lambda_{nt} - t\mathcal{K}_n}{\gamma_n}. \quad (5)$$

Theorem 4. *If (H) is satisfied, for any $\mathbf{x} \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$, under $\mathbb{P}_n^{\mathbf{x},(\mathcal{K}_n)}$,*

$$(s_n, \lambda_n) \xrightarrow[n]{(d)} (\mathbf{x}, \widehat{\mathbf{b}}) \quad (6)$$

where $\widehat{\mathbf{b}}$ is a Brownian bridge independent of \mathbf{x} and where the weak convergence holds in $C([0, 1])^2$ endowed with the topology of uniform convergence.

Hence, under $\mathbb{P}_n^{\mathbf{x},(\mathcal{K}_n)}$, up to the scaling constant, the process s_n behaves as under $\mathbb{P}_n^{\mathbf{x}}$. The normalizing factor β_n , that will be explained later in the paper, indicates the order of magnitude of the process \mathbf{S} under $\mathbb{P}_n^{\mathbf{x},(\mathcal{K}_n)}$ (β_n is a decreasing function of \mathcal{K}_n). The normalizing constant γ_n is smaller than $\sqrt{n/4}$ whatever is p_n ; γ_n gives the asymptotic order of the ‘‘linearity defect’’ of $t \mapsto \Lambda_{nt}$. The fact that (λ_n) converges to a Brownian bridge independent of the limit trajectory is quite puzzling. For example, under $\mathbb{P}_n^{\mathbf{e}}$, one would expect that only few peaks appear in a neighborhood of 0, this resulting in a negative bias in λ_n near 0. This must be true, but this bias is not important enough to change the asymptotic behavior of λ_n .

A second direct corollary of Theorem 4 is stated below:

Corollary 5. *For any $\mathbf{x} \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$, under $\mathbb{P}_n^{\mathbf{x}}$, we have*

$$\left(\frac{S(nt)}{\sqrt{n}}, 4 \frac{\Lambda_{nt} - tn/4}{\sqrt{n}} \right)_{t \in [0,1]} \xrightarrow[n]{(d)} (\mathbf{x}_t, \widehat{\mathbf{w}}_t)_{t \in [0,1]} \quad (7)$$

and

$$\left(\frac{S(nt)}{\sqrt{n}}, 4 \frac{\Lambda_{nt} - t\Lambda_n}{\sqrt{n}} \right)_{t \in [0,1]} \xrightarrow[n]{(d)} (\mathbf{x}_t, \widehat{\mathbf{b}}_t)_{t \in [0,1]} \quad (8)$$

where $\widehat{\mathbf{w}}$ is a Brownian motion independent of \mathbf{x} , $\widehat{\mathbf{b}}$ is a Brownian bridge independent of \mathbf{x} and where the weak convergences hold in $C([0, 1])^2$ endowed with the topology of uniform convergence.

Theorem 2 is of course a consequence of Corollary 5.

Proof. For any $\mathbf{S} \in \mathcal{W}_n$, set $q_n(\mathbf{S}) = 2\Lambda_n(\mathbf{S})/n$, $\tilde{\beta}_n(\mathbf{S}) = \sqrt{n(1 - q_n(\mathbf{S}))/q_n(\mathbf{S})}$, and $\tilde{\gamma}_n(\mathbf{S}) := \sqrt{nq_n(\mathbf{S})(1 - q_n(\mathbf{S}))}$. We have

$$\left(\frac{S(nt)}{\sqrt{n}}, 4 \frac{\Lambda_{nt} - tn/4}{\sqrt{n}} \right) = \left(\frac{S(nt)}{\tilde{\beta}_n} \frac{\tilde{\beta}_n}{\sqrt{n}}, 2 \frac{\tilde{\gamma}_n}{\sqrt{n}} \left(2 \frac{\Lambda_{nt} - t\Lambda_n}{\tilde{\gamma}_n} \right) + 4t \frac{\Lambda_n - n/4}{\sqrt{n}} \right). \quad (9)$$

By Proposition 3 and Theorem 4, under \mathbb{P}_n^x , the five-tuple

$$\left(4 \frac{\Lambda_n - n/4}{\sqrt{n}}, \left(\frac{S(nt)}{\tilde{\beta}_n} \right)_{t \in [0,1]}, 2 \left(\frac{\Lambda_{nt} - t\Lambda_n}{\tilde{\gamma}_n} \right)_{t \in [0,1]}, 2 \frac{\tilde{\gamma}_n}{\sqrt{n}}, \frac{\tilde{\beta}_n}{\sqrt{n}} \right)$$

converges in distribution to

$$(N, (s_t)_{t \in [0,1]}, (\lambda_t)_{t \in [0,1]}, A, B)$$

where N is a centered Gaussian random variable with variance 1 and where conditionally on N , $(s, \lambda) \stackrel{(d)}{=} (x, \hat{b})$ where x and \hat{b} are independent, \hat{b} is a Brownian bridge, and A and B are two random variables equal to 1 a.s.. By (9), $\left(\frac{S(nt)}{\sqrt{n}}, 4 \frac{\Lambda_{nt} - tn/4}{\sqrt{n}} \right)$ converges to $(x, (\hat{b}_t + tN)_{t \in [0,1]})$ where N is independent of x and \hat{b} , and then the result follows, since $(\hat{b}_t + tN)_{t \in [0,1]}$ is a standard Brownian motion. \square

1.1 Consequences in terms of plane trees

Consider the set T_n of *plane trees* (rooted ordered trees) with n edges (we refer to (1; 16) for more information on these objects). There exists a well known bijection between T_n and \mathcal{E}_{2n} which may be informally described as follows. Consider a plane tree $\tau \in T_n$ (see Figure 2), and a fly walking around the tree τ clockwise, starting from the root, at the speed 1 edge per unit of time. Let $V(t)$ be the distance from the root to the fly at time t . The process $V(t)$ is called in the literature, the contour process or the Harris' walk associated with τ . The contour process




Figure 2: A plane tree and its contour process

is the very important tool for the study of plane trees and their asymptotics and we refer to Aldous (1), Pitman (21, Section 6), Duquesne & Le Gall (10), Marckert & Mokkadem (16) for considerations on the asymptotics of normalized trees. It is straightforward that the set of trees encoded by $\mathcal{E}_{2n}^{(k)}$ is the subset of T_n of trees having exactly k leaves (say $T_n^{(k)}$), a leaf being a node without any child. A corollary of Theorem 4, is that random plane tree with n edges and \mathcal{K}_{2n} leaves, converges, normalized by $\beta_{2n}/2$, to the continuum random tree introduced by Aldous (1), which is encoded by 2e. The variable Λ_{2nt} gives the number of leaves visited at time $2nt$. By Theorem 4, $\sup_{t \in [0,1]} |(\Lambda_{2nt} - t\mathcal{K}_{2n})/n| \xrightarrow{\text{proba.}} 0$. This translates the fact that the leaves are asymptotically uniformly distributed on a random tree chosen equally likely in $T_n^{(\mathcal{K}_{2n})}$.

1.2 Consequences in terms of parallelogram polyominoes

We refer to Delest & Viennot (8) for more information on parallelogram polyominoes. Unit squares having their vertices at integer points in the Cartesian plane are called cells. A polyomino is a finite union of cells with connected interior. The number of cells is the area and the length of the border is called the perimeter (see Figure 3). A polyomino \mathcal{P} is said to be convex if the intersection of \mathcal{P} with any horizontal or vertical line is a convex segment. For any convex polyomino \mathcal{P} there exists a minimal rectangle $R(\mathcal{P})$ (that can be seen as a convex polyomino) containing \mathcal{P} . Then \mathcal{P} touches the border of $R(\mathcal{P})$ along four connected segments. A convex polyomino \mathcal{P} is said to be a parallelogram polyomino if the south-west point and the north-east point of $R(\mathcal{P})$ belong to \mathcal{P} (see Figure 3). Let denote by $H(\mathcal{P})$ and $V(\mathcal{P})$ the horizontal and




Figure 3: The first convex polyomino is not parallelogram, the second is. Their areas are 9 and 11, their perimeters equal that of their minimal rectangles, here 18. For both polyominoes $H(\mathcal{P}) = 4$ and $V(\mathcal{P}) = 5$. The last picture represents the Bernoulli excursion associated by ρ with the parallelogram polyomino.

vertical length of the border of $R(\mathcal{P})$, and let Pol_n be the set of parallelogram polyominoes with perimeter n .

Proposition 6. (Delest & Viennot (8, Section 4)) *For any integer $N \geq 1$, there is a bijection ρ between \mathcal{E}_{2N} and Pol_{2N+2} , such that if $\mathcal{P} = \rho(\mathbf{S})$, the area of \mathcal{P} is equal to the sum of the heights of peaks of \mathbf{S} , moreover $\#\mathbf{S}_\wedge = H(\mathcal{P})$, and $V(\mathcal{P}) = 2N + 2 - 2\#\mathbf{S}_\wedge$ (where $2N - 2\#\mathbf{S}_\wedge$ is the number of steps of \mathbf{S} that do not belong to a peak).*

By symmetry with respect to the first diagonal, the random variables $V(\mathcal{P})$ and $H(\mathcal{P})$ have the same distribution when \mathcal{P} is taken equally likely in Pol_{2N+2} . Hence, the proposition says that under \mathbb{P}_{2N}^e , $2N + 2 - 2\#\mathbf{S}_\wedge$ and $\#\mathbf{S}_\wedge$ have the same distribution.

We describe in a few words Delest & Viennot's bijection: the successive lengths of the columns of the polyomino \mathcal{P} give the successive heights of the peaks of \mathbf{S} . The difference between the heights of the floor-cells of the i th and $i + 1$ th columns of \mathcal{P} **plus one** gives the number of down steps between the i th and $i + 1$ th peaks of \mathbf{S} .

For $i \in \{1, \dots, H(\mathcal{P})\}$, let $v_i(\mathcal{P})$ be the number of cells in the i th column of \mathcal{P} . The values $(v_i(\mathcal{P}))_{i \in [1, H(\mathcal{P})]}$ coincide with the ordered sequence $(S_i)_{i \in \mathbf{S}_\wedge}$. Let $\mathbb{P}_{\text{Pol}(2n+2)}^{(\mathcal{K}_n)}$ be the uniform law

on the set of parallelogram polyominoes with perimeter $2n + 2$ and width \mathcal{K}_n (that is such that $H(\mathcal{P}) = \mathcal{K}_n$). Assume that v is interpolated between integer points, and $v(0) = 0$. We have

Proposition 7. *If (\mathcal{K}_n) satisfies (H), under $\mathbb{P}_{Pol(2n+2)}^{(\mathcal{K}_n)}$*

$$\left(\frac{v(\mathcal{K}_n t)}{\beta_n} \right)_{t \in [0,1]} \xrightarrow{(d)} (\mathbf{e}_t)_{t \in [0,1]}$$

in $C[0,1]$ endowed with the topology of uniform convergence.

Proof. Let $(V_i)_{i \in \{1, \dots, \mathcal{K}_n\}}$ be the successive height of the peaks in \mathbf{S} . Assume also that $V(0) = 0$ and that V is interpolated between integer points. By Delest & Viennot's bijection, $\beta_n^{-1}v(\mathcal{K}_n \cdot)$ under $\mathbb{P}_{Pol(2n+2)}^{(\mathcal{K}_n)}$ has the same distribution as $\beta_n^{-1}V(\mathcal{K}_n \cdot)$ under $\mathbb{P}_n^{\mathbf{e}, (\mathcal{K}_n)}$. Since $(\beta_n^{-1}S(nt))_{t \in [0,1]} \xrightarrow{(d)} (\mathbf{e}_t)_{t \in [0,1]}$, to conclude, it suffices to show that

$$\sup_{t \in [0,1]} \left| \frac{V(\mathcal{K}_n t) - S(nt)}{\beta_n} \right| \xrightarrow{proba.} 0. \quad (10)$$

Let $J(i)$ be (abscissa of) the i th peak in \mathbf{S} . We have, for any $t \in \{0, 1/\mathcal{K}_n, \dots, \mathcal{K}_n/\mathcal{K}_n\}$,

$$V(\mathcal{K}_n t) - S(nt) = S(J(\mathcal{K}_n t)) - S(nt). \quad (11)$$

As one can see using the convergence of λ_n to \widehat{b} ,

$$\sup_t \left| \frac{J(\mathcal{K}_n t) - nt}{n} \right| \xrightarrow{proba.} 0. \quad (12)$$

Indeed, $\sup_t |J(\mathcal{K}_n t) - nt|/n \leq \sup_t |\Lambda_{nt} - t\mathcal{K}_n|/n \xrightarrow{proba.} 0$. Since (s_n) converges in $C[0,1]$ under $\mathbb{P}_n^{\mathbf{e}, (\mathcal{K}_n)}$, by a simple argument about its modulus of continuity, using (11) and (12), formula (10) holds true. \square

We would like to point out the work of de Sainte Catherine & Viennot (9), who exhibit a quite unexpected link between the cardinalities of excursions having their peaks in a given subset of \mathbb{N} and the famous Tchebichev polynomials.

1.3 Consequences in terms of Kolmogorov-Smirnov statistics

We refer again to Krattenthaler (14, Example 3.3.2 and Theorem 3.4.4).

Let $X = (X_1, \dots, X_n)$ and $Y = (Y_1, \dots, Y_n)$ be two independent sets of independent random variables, where the X_i (resp. the Y_i) are identically distributed. Assume that the cumulative distributions F_X and F_Y of the X_i 's and Y_i 's are continuous, and that we want to test the hypothesis H that $F_X = F_Y$ in view of X and Y .

Let $Z = (Z_1, \dots, Z_{2n})$ be the order statistics of $(X_1, \dots, X_n, Y_1, \dots, Y_n)$; in other words Z_i is the i th smallest element in $(X_1, \dots, X_n, Y_1, \dots, Y_n)$. We may associate almost surely to Z a Bernoulli bridge $\mathbf{S} \in \mathcal{B}_{2n}$ saying that $S(i) - S(i-1) = 1$ or $S(i) - S(i-1) = -1$ whether Z_i is in X or in Y .

It is easy to see that \mathbf{S} is uniform in \mathcal{B}_{2n} (for any n) when $F_X = F_Y$ and that it is not the case when $F_X \neq F_Y$: assume that $F_X(x) \neq F_Y(x)$ for some x and denote by $N_X(x) = \#\{i, X_i \leq x\}$ and $N_Y(x) = \#\{i, Y_i \leq x\}$. By the law of large numbers

$$\frac{N_X(x)}{n} \xrightarrow[n]{(a.s.)} F_X(x) \quad \text{and} \quad \frac{N_Y(x)}{n} \xrightarrow[n]{(a.s.)} F_Y(x)$$

which implies that $S(N_X(x) + N_Y(x))$ is at the first order $n(F_X(x) - F_Y(x))$, that is much bigger than the \sqrt{n} order expected by Theorem 2. We examine now, as in (14) the run statistics

$$R_n(\mathbf{S}) = \#\{i \in \llbracket 1, 2n-1 \rrbracket, S(i+1) \neq S(i)\}$$

that counts the number of runs in \mathbf{S} (up to at most 1, twice the number of peaks) and

$$D_{n,n}^+(\mathbf{S}) = \frac{1}{n} \max_i S(i) \quad \text{and} \quad D_{n,n}(\mathbf{S}) = \frac{1}{n} \max_i |S(i)|,$$

the one-sided and two-sided Kolmogorov-Smirnov statistics.

The run statistics R_n has been introduced by Wald and Wolfowitz (24). It is shown to be independent of F_X when $F_X = F_Y$. It is well adapted in many cases since no assumption on F_X (but its continuity) is needed. Theorem 3.4.4 of Krattenthaler (14) (see also the consideration just above Example 13.3.3) provides the exact enumeration of Bernoulli walks \mathbf{S} for which $D_{n,n} \leq t/n$ and $R_n = j$.

The random variable R_n satisfies obviously

$$|R_n - 2\Lambda_{2n}| \leq 1. \tag{13}$$

A simple consequence of assertion (7) and (13) is the following result

Corollary 8. *If $F_X = F_Y$ then*

$$\left(\frac{\sqrt{n}}{\sqrt{2}} D_{n,n}^+, \frac{\sqrt{n}}{\sqrt{2}} D_{n,n}, \frac{R_n - n/2}{\sqrt{n}} \right) \xrightarrow[n]{(d)} (\sup b, \sup |b|, N)$$

where b is a Brownian bridge independent of N , a centered Gaussian random variable with variance 1/2.

This means that asymptotically the run statistics and the Kolmogorov statistics are independent.

Comments 1. *Notice that the process $t \mapsto \Lambda_{nt}$ can also be used to build a test H based on the runs of \mathbf{S} (using Corollary 5). It is clearly much more precise than R_n alone.*

2 Combinatorial facts : decomposition of trajectories

The decomposition of the trajectories is determinant in our approach, since we will prove directly the convergence of finite dimensional distributions under \mathbb{P}_n^x . An important difference with the case where the peaks are not considered is that under $\mathbb{P}_n^{x,(k)}$, \mathbf{S} is not a Markov chain. Indeed, the law of $(S(j))_{l \leq j \leq n}$ depends on $S(l)$, on the number of peaks in $\llbracket 0, l \rrbracket$, and also on the step $(S(l-1), S(l))$. The computation of the distributions of vector $(S(t_1), \dots, S(t_k))$ in these various cases admits the complication that \mathbf{S} may own a peak in some of the positions t_1, \dots, t_k . In order to handle the contribution of these peaks, we have to specify what the types u or d ($+1$ or -1) of the first and last steps of the studied parts $(S(t_i), \dots, S(t_{i+1}))$ are.

We set the following notation : $\Delta S_l = S(l) - S(l-1)$, and write for convenience, $\Delta S_l = u$ when $\Delta S_l = 1$, and $\Delta S_l = d$ when $\Delta S_l = -1$. In this paper, we deal only with discrete trajectories \mathbf{S} such that $\Delta S_k \in \{+1, -1\}$ for any k . We will not recall this condition.

For a and b in $\{d, u\}$, and l, x, y, j in \mathbb{Z} , set

$$\begin{aligned} T_{ab}^j(l, x, y) &= \{\mathbf{S} : \mathbf{S} = (S(i))_{0 \leq i \leq l}, \#\mathbf{S}_\wedge = j, \Delta S_1 = a, \Delta S_l = b, S(0) = x, S(l) = y\} \\ T_{ab}^{j, \geq}(l, x, y) &= \{\mathbf{S} : \mathbf{S} \in T_{ab}^j(l, x, y), S(i) \geq 0 \text{ for any } i \in \llbracket 0, l \rrbracket\}. \end{aligned}$$

For any $l, j_1, j_2, x, y \in \mathbb{Z}$, set

$$[l, j_1, j_2, x, y] = \binom{\frac{l+y-x}{2} - 1}{j_1} \binom{\frac{l-y+x}{2} - 1}{j_2}.$$

We have

Proposition 9. *For any $y \in \mathbb{Z}$, $l \geq 0$, $j \geq 0$,*

$$\#T_{ab}^j(l, 0, y) = [l, j - \mathbf{1}_{b=d}, j - \mathbf{1}_{a=u}, 0, y]. \quad (14)$$

For any $x \geq 0$, $y \geq 1$, $l \geq 0$, $j \geq 0$,

$$\begin{aligned} \#T_{uu}^{j, \geq}(l, x, y) &= [l, j, j-1, x, y] - [l, j-1, j, -x, y], \\ \#T_{ud}^{j, \geq}(l, x, y) &= [l, j-1, j-1, x, y] - [l, j-2, j, -x, y], \end{aligned}$$

For any $x \geq 1$, $y \geq 1$, $l \geq 0$, $j \geq 0$,

$$\begin{aligned} \#T_{du}^{j, \geq}(l, x, y) &= [l, j, j, x, y] - [l, j, j, -x, y], \\ \#T_{dd}^{j, \geq}(l, x, y) &= [l, j-1, j, x, y] - [l, j-1, j, -x, y] \end{aligned}$$

(notice that $\#T_{db}^{j, \geq}(l, 0, y) = 0$ and $\#T_{au}^{j, \geq}(l, x, 0) = 0$). In other words

$$\#T_{ab}^{j, \geq}(l, x, y) = [l, j - \mathbf{1}_{b=d}, j - \mathbf{1}_{a=u}, x, y] - [l, j - \mathbf{1}_{a=u} - \mathbf{1}_{b=d}, j, -x, y]. \quad (15)$$

Proof. Let $n, k \in \mathbb{N}$. A composition of n in k parts is an ordered sequence x_1, \dots, x_k of non negative integers, such that $x_1 + \dots + x_k = n$. The number of compositions of n in k parts (or of compositions of $n + k$ in k positive integer parts) is $\binom{n+k-1}{k-1}$.

We call *run* of the chain $\mathbf{S} = (S(i))_{0 \leq i \leq l}$, a maximal non-empty interval I of $\llbracket 1, l \rrbracket$ such that $(\Delta S_i)_{i \in I}$ is constant. The trajectories \mathbf{S} of $T_{ab}^j(l, y)$ are composed by $j + \mathbb{1}_{b=u}$ runs of u and $j + \mathbb{1}_{a=d}$ runs of d . The u -runs form a composition of $(l + y)/2$ (this is the number of steps u) in positive integer parts, and the d -runs form a composition of $(l - y)/2$ in positive integer parts. Hence,

$$\#T_{ab}^j(l, y) = \binom{(l+y)/2 - 1}{j + \mathbb{1}_{b=u} - 1} \binom{(l-y)/2 - 1}{j + \mathbb{1}_{a=d} - 1},$$

and Formula (14) holds true.

The proofs of the other formulas are more tricky; the main reason for it is that the reflexion principle does not conserve the number of peaks. What still holds is, for any $x \geq 0, y \geq 0, j \geq 0, l \geq 0$

$$\#T_{ab}^{j, \geq}(l, x, y) = \#T_{ab}^j(l, x, y) - \#T_{ab}^{j, \vee}(l, x, y)$$

where $T_{ab}^{j, \vee}(l, x, y)$ is the set of trajectories belonging to $T_{ab}^j(l, x, y)$ that reach the level -1 . Since $\#T_{ab}^j(l, x, y) = \#T_{ab}^j(l, 0, y - x)$ is known, it remains to determine $T_{ab}^{j, \vee}(l, x, y)$.

We define two actions on the set of chains :

■ let $\mathbf{S} = (S(i))_{i \in \llbracket 0, l \rrbracket} \in \mathcal{W}_l$. For any $t \in \llbracket 0, l \rrbracket$ we denote by $\mathbf{S}' = \text{Ref}(\mathbf{S}, t)$ the path $\mathbf{S}' = (S'_i)_{0 \leq i \leq l}$ obtained from \mathbf{S} by a reflexion from the abscissa t ; formally :

$$\begin{cases} S'(i) = S(i) & \text{for any } 0 \leq i \leq t, \\ S'(i+t) = 2S(i) - S(i+t) & \text{for any } 0 \leq i \leq l-t \end{cases}.$$

When g is a function from \mathcal{W}_l taking its values in $\llbracket 0, l \rrbracket$, we write simply $\text{Ref}(\cdot, g)$ for the reflexion at abscissa $g(\mathbf{S})$.

■ let $\mathbf{S} = (S(i))_{i \in \llbracket 0, l \rrbracket} \in \mathcal{W}_l$. For any c and d in \mathbb{N} , $0 \leq c \leq d \leq l$, we denote by $\mathbf{S}' = \text{Cont}(\mathbf{S}, [c, d])$ the path $\mathbf{S}' = (S'(i))_{0 \leq i \leq l-d}$ obtained from \mathbf{S} by a contraction of the interval $[c, d]$:

$$\begin{cases} S'(i) = S(i) & \text{for any } 0 \leq i \leq c, \\ S'(c+i) = S(d+i) - S(d) + S(c) & \text{for any } 0 \leq i \leq l - (d-c) \end{cases}.$$

As before, we write $\text{Cont}(\cdot, [g_1, g_2])$ for the contraction of the interval $[g_1(\mathbf{S}), g_2(\mathbf{S})]$.

We denote by $\mathcal{T}_{-1}(\mathbf{S}) = \inf\{j : S(j) = -1\}$ the hitting time of -1 by \mathbf{S} . We proceed to a classification of the paths \mathbf{S} from $T_{ab}^{j, \vee}(l, x, y)$ according to the two first steps following $\mathcal{T}_{-1}(\mathbf{S})$, that exist since y is taken positive. We encode these two first steps following \mathcal{T}_{-1} above the symbol T : for any $\alpha, \beta \in \{u, d\}$, set

$$T_{ab}^{(\alpha\beta)j, \vee}(l, x, y) = \{\mathbf{S} : \mathbf{S} \in T_{ab}^{j, \vee}(l, x, y), \Delta S_{\mathcal{T}_{-1}(\mathbf{S})+1} = \alpha, \Delta S_{\mathcal{T}_{-1}(\mathbf{S})+2} = \beta\}.$$




Figure 4: On the first column \mathbf{S} and $\text{Ref}(\mathbf{S}, t)$, on the second column \mathbf{S} and $\text{Cont}(\mathbf{S}, [c, d])$

Hence, $T_{ab}^{j,\gamma}(l, x, y)$ is the union of four elements of that type. Let us compute $\# T_{ab}^{(\alpha\beta)}$ for any α, β .

For any $a \in \{u, d\}$, $x \geq 0$, $y \geq 1$, the following bijections (denoted by \leftrightarrow) hold

$$\begin{aligned} T_{ad}^{(du)^j, \gamma}(l, x, y) &\leftrightarrow T_{au}^{(ud)^j, \gamma}(l, x, -2 - y) \leftrightarrow T_{au}^{j-1}(l-2, x, -2 - y) \\ T_{au}^{(du)^j, \gamma}(l, x, y) &\leftrightarrow T_{ad}^{(ud)^{j+1}, \gamma}(l, x, -2 - y) \leftrightarrow T_{ad}^j(l-2, x, -2 - y). \end{aligned}$$

On each line, the first bijection is $\text{Ref}(\cdot, \mathcal{T}_{-1})$, the second one is $\text{Cont}(\cdot, [\mathcal{T}_{-1}, \mathcal{T}_{-1} + 2])$. Notice that this last operation does not create any peak because $\Delta S_{\mathcal{T}_{-1}}(\mathbf{s}) = d$. The cardinalities of the sets in the right hand side are known, hence, $T_{dd}^{(du)}$, $T_{ud}^{(du)}$, $T_{du}^{(du)}$, $T_{uu}^{(du)}$ are determined. Set $T_{ab}^{(\alpha)^j, \gamma}(l, x, y) = T_{ab}^{(\alpha u)^j, \gamma}(l, x, y) \cup T_{ab}^{(\alpha d)^j, \gamma}(l, x, y)$. We have

$$\begin{aligned} T_{ad}^{(u)^j, \gamma}(l, x, y) &\leftrightarrow T_{au}^{(d)^{j-1}, \gamma}(l, x, -y - 2) \leftrightarrow T_{au}^{j-1}(l-1, x, -y - 1) \\ T_{au}^{(u)^j, \gamma}(l, x, y) &\leftrightarrow T_{ad}^{(d)^j, \gamma}(l, x, -y - 2) \leftrightarrow T_{ad}^j(l-1, x, -y - 1) \end{aligned}$$

The first bijection is $\text{Ref}(\cdot, \mathcal{T}_{-1})$, the second one $\text{Cont}(\cdot, [\mathcal{T}_{-1}, \mathcal{T}_{-1} + 1])$. Now,

$$\begin{aligned} \# T_{ab}^{(dd)^j, \gamma} &= \# T_{ab}^{(ud)^{j+1}, \gamma}(l, x, y + 2) \\ &= \# T_{ab}^{(u)^{j+1}, \gamma}(l, x, y + 2) - \# T_{ab}^{(uu)^{j+1}, \gamma}(l, x, y + 2) \\ &= \# T_{ab}^{(u)^{j+1}, \gamma}(l, x, y + 2) - \# T_{ab}^{(du)^{j+1}, \gamma}(l, x, y) \end{aligned}$$

in the first line, we have replaced $\Delta S_{\mathcal{T}_{-1}} = d$ by u , in the third line we have replaced uu by du . \square

2.1 Proof of Proposition 1

(i) To build a path with k peaks, dispose k peaks, that is k pairs ud . Take a composition $x_1, \dots, x_{2(k+1)}$ of $n - 2k$ in $2(k+1)$ non negative parts. Fill in now the $k+1$ intervals between these peaks : in the l th interval dispose x_{2l-1} steps d and x_{2l} steps u .

(ii) Assume $n = 2N$ is even. To build a bridge with k peaks, dispose k pairs ud . Take two compositions $x_1, \dots, x_{(k+1)}$ and $x'_1, \dots, x'_{(k+1)}$ of $N - k$ in $k+1$ parts. Fill now the $k+1$ intervals between these peaks : in the l th interval dispose x_l steps d and x'_l steps u .

$$(iii) \quad \#\mathcal{E}_n^{(k)} = \#T_{ud}^{k, \geq}(n-1, 0, 1) + \#T_{uu}^{k-1, \geq}(n-1, 0, 1)$$

For (iv), one may use the bijections described in Section 4.4, or proceed to a direct computation as follows; first,

$$\#\mathcal{M}_n^{(k)} = \#\mathcal{E}_n^{(k)} + \sum_{y \geq 1} T_{uu}^{k, \geq}(n, 0, y) + T_{ud}^{k, \geq}(n, 0, y). \quad (16)$$

Denote by $W(n, k)$ the sum in (16). The integer $W(n, k)$ is the number of meanders with length n , ending in a positive position. Using that $\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k}$, we have

$$\begin{aligned} W(n, k) &= \sum_{y \geq 1} [n, k, k-1, 0, y] - [n, k-1, k, 0, y] + [n, k-1, k-1, 0, y] - [n, k-2, k, 0, y]. \\ &= \sum_{y \geq 1} \binom{\frac{n+y}{2}}{k} \binom{\frac{n-y}{2}-1}{k-1} - \binom{\frac{n+y}{2}}{k-1} \binom{\frac{n-y}{2}}{k}. \end{aligned}$$

Let a, b, c, k be positive integers, the following formula holds

$$\sum_{y \geq c} \binom{a+y}{k} \binom{b-y}{k-1} - \sum_{y \geq c} \binom{a+y}{k-1} \binom{b-y}{k} = \binom{a+c}{k} \binom{b-c+1}{k}. \quad (17)$$

Indeed: a term in the first sum counts the number of ways to choose $2k$ items among $a+b+1$, choosing, the $a+y+1$ th, and k items among the $a+y$ first ones, when, a term in the second sum counts the number of ways to choose $2k$ items among $a+b+1$, choosing, the $a+y+1$ th, and $k-1$ items among the $a+y$ first ones. The choices counted by the first sum but not by the second one, are those where exactly k items are chosen among the $a+c$ first ones.

We need to consider the two cases n even and n odd :

- If n is even, using that $S(n) = n \bmod 2 = y \bmod 2$, set $N = n/2$, $Y = y/2$ in the sum,

$$W(2N, k) = \sum_{Y \geq 1} \binom{N+Y}{k} \binom{N-Y-1}{k-1} - \binom{N+Y}{k-1} \binom{N-Y-1}{k} = \binom{N+1}{k} \binom{N-1}{k}.$$

If $n = 2N+1$, take $y = 2Y+1$ in the sum

$$W(2N+1, k) = \sum_{Y \geq 0} \binom{N+Y+1}{k} \binom{N-Y-1}{k-1} - \binom{N+Y+1}{k-1} \binom{N-Y-1}{k} = \binom{N+1}{k} \binom{N}{k}. \quad \square$$

3 Asymptotic considerations and proofs

We first recall a classical result of probability theory, simple consequence of Billingsley (6, Theorem 7.8), that allows to prove the weak convergence in \mathbb{R}^d using a local limit theorem.

Proposition 10. *Let k be a positive integer and for each $i \in \llbracket 1, k \rrbracket$, $(\alpha_n^{(i)})$ a sequence of real numbers such that $\alpha_n^{(i)} \xrightarrow{n} +\infty$. For any n , let $X_n = (X_n^{(1)}, \dots, X_n^{(k)})$ be a \mathbb{Z}^k -valued random variable. If $\forall \lambda = (\lambda_1, \dots, \lambda_k) \in \mathbb{R}^k$,*

$$\alpha_n^{(1)} \dots \alpha_n^{(k)} \mathbb{P}((X_n^{(1)}, \dots, X_n^{(k)}) = (\lfloor \lambda_1 \alpha_n^{(1)} \rfloor, \dots, \lfloor \lambda_k \alpha_n^{(k)} \rfloor)) \longrightarrow \phi(\lambda_1, \dots, \lambda_k)$$

where ϕ is a density probability on \mathbb{R}^k , then $(X_n^{(1)}/\alpha_n^{(1)}, \dots, X_n^{(k)}/\alpha_n^{(k)}) \xrightarrow[n]{(d)} X$ where X is a random variable with density ϕ .

Proposition 3 is a simple consequence of this proposition, since by (1) and Proposition 1, the application of the Stirling formula simply yields

$$\sqrt{n} \mathbb{P}_n^x(\lfloor \Lambda_n - n/4 \rfloor = \lfloor t\sqrt{n} \rfloor) \xrightarrow{n} \frac{1}{\sqrt{2\pi/16}} \exp(-8t^2),$$

for any $x \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$ and any $t \in \mathbb{R}$. Note that under $\mathbb{P}_n^{\mathbf{w}}$, one may also compute the limiting distribution using that $\Lambda_n(\mathbf{S}) = \sum_{i=1}^{n-1} \mathbb{1}_{i \in \mathbf{S}^\wedge}$, which is a sum of Bernoulli random variables with an easy to handle dependence.

3.1 Finite dimensional distribution of the Brownian processes

Notation For any sequence $(o_i)_i$ indexed by integers, the sequence (Δo_i) is defined by $\Delta o_i = o_i - o_{i-1}$ and $(\bar{\Delta} o_i)$ by $\bar{\Delta} o_i = o_i + o_{i-1}$.

For any $t > 0$ and $x, y \in \mathbb{R}$, set

$$\mathbf{p}_t(x, y) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(y-x)^2}{2t}\right).$$

Let $\ell \geq 1$ and let $(t_1, \dots, t_\ell) \in [0, 1]^\ell$ satisfying $0 < t_1 < \dots < t_{\ell-1} < t_\ell := 1$. The distributions of the a.s. continuous processes $\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}$ are characterized by their finite dimensional distributions. Let f_{t_1, \dots, t_ℓ}^x be the density of $(x_{t_1}, \dots, x_{t_\ell})$ with respect to the Lebesgue measure on \mathbb{R}^k . We have

$$\begin{aligned} f_{t_1, \dots, t_\ell}^{\mathbf{w}}(x_1, \dots, x_\ell) &= \prod_{i=1}^{\ell} \mathbf{p}_{\Delta t_i}(x_{i-1}, x_i), \quad \text{with } x_0 = 0 \text{ by convention,} \\ f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}}(x_1, \dots, x_{\ell-1}) &= \sqrt{2\pi} f_{t_1, \dots, t_{\ell-1}}^{\mathbf{w}}(x_1, \dots, x_{\ell-1}) \mathbf{p}_{\Delta t_\ell}(x_{\ell-1}, 0), \\ f_{t_1, \dots, t_\ell}^{\mathbf{m}}(x_1, \dots, x_\ell) &= \sqrt{2\pi} \frac{x_1}{t_1} \mathbf{p}_{t_1}(0, x_1) \left(\prod_{i=2}^{\ell} \mathbf{p}_{\Delta t_i}(x_{i-1}, x_i) - \mathbf{p}_{\Delta t_i}(x_{i-1}, -x_i) \right) \mathbb{1}_{x_1, \dots, x_\ell \geq 0}, \\ f_{t_1, \dots, t_{\ell-1}}^{\mathbf{e}}(x_1, \dots, x_{\ell-1}) &= f_{t_1, \dots, t_{\ell-1}}^{\mathbf{m}}(x_1, \dots, x_{\ell-1}) \frac{x_{\ell-1}}{\Delta t_\ell} \mathbf{p}_{\Delta t_\ell}(x_{\ell-1}, 0). \end{aligned}$$

We end this section with two classical facts: first $\mathbf{p}_t(x, y) = \mathbf{p}_t(0, y - x)$ and, for any $\alpha > 0$, $f_{t_1, \dots, t_k}^{\alpha \mathbf{x}}(x_1, \dots, x_k) = \alpha^{-k} f_{t_1, \dots, t_k}^{\mathbf{x}}(x_1/\alpha, \dots, x_k/\alpha)$.

3.2 Finite dimensional convergence

We will show that for any $\mathbf{x} \in \{\mathbf{w}, \mathbf{b}, \mathbf{e}, \mathbf{m}\}$, under $\mathbb{P}_n^{\mathbf{x}}$, for any $\ell \in \{1, 2, 3, \dots\}$ and $0 < t_1 < \dots < t_{\ell-1} < t_\ell := 1$

$$(s_n(t_1), \dots, s_n(t_\ell), \lambda_n(t_1), \dots, \lambda_n(t_\ell)) \xrightarrow[n]{(d)} (\mathbf{x}(t_1), \dots, \mathbf{x}(t_\ell), \widehat{\mathbf{b}}(t_1), \dots, \widehat{\mathbf{b}}(t_\ell)),$$

where $t_\ell := 1$ has been chosen for computation convenience, and $(\mathbf{x}, \widehat{\mathbf{b}})$ has the prescribed distribution (as in Theorem 4). This implies the convergence of the finite dimensional distribution in Theorem 4.

In order to handle easily the binomial coefficients appearing in $[l, j_1, j_2, x, y]$ that involve half-integers, we proceed as follows. Let $N = \lfloor n/2 \rfloor$ and let $\mathcal{E}_n = n - 2N = n \pmod{2}$. For $i \in \llbracket 1, \ell - 1 \rrbracket$, let $t_i^{(n)}$ be defined by

$$2Nt_i^{(n)} := 2\lfloor nt_i/2 \rfloor,$$

$t_0^{(n)} = 0$, and $t_\ell^{(n)}$ by $2Nt_\ell^{(n)} = 2N + \mathcal{E}_n = n$ (notice that $t_\ell^{(n)}$ is in $\{1, 1 + 1/(n-1)\}$).

Using that for any i , $|2Nt_i^{(n)} - nt_i| \leq 2$, we have clearly under $\mathbb{P}_n^{\mathbf{x}, (\mathcal{K}_n)}$,

$$\frac{S(2Nt_i^{(n)}) - S(nt_i)}{\beta_n} \xrightarrow{\text{proba.}} 0 \text{ and } \frac{\Lambda_{2Nt_i^{(n)}} - \Lambda_{nt_i}}{\gamma_n} \xrightarrow{\text{proba.}} 0 \quad (18)$$

since γ_n and β_n goes to $+\infty$. From now on, we focus on the values of the processes on the discretization points $2Nt_i^{(n)}$. For any $i \in \llbracket 1, \ell - 1 \rrbracket$, set

$$\widetilde{\Lambda}_i := \#\mathbf{S}_\wedge \cap \llbracket 2Nt_{i-1}^{(n)} + 1, 2Nt_i^{(n)} - 1 \rrbracket$$

the number of peaks lying strictly between $2Nt_{i-1}^{(n)}$ and $2Nt_i^{(n)}$.

In order to obtain a local limit theorem, we are interested in the number of trajectories passing via some prescribed positions.

3.2.1 Case $\mathbf{x} = \mathbf{w}$

Let $0 = u_0, u_1, \dots, u_\ell, v_1, \dots, v_{\ell-1}$ be fixed real numbers. Set

$$\Theta := (t_1, \dots, t_{\ell-1}, u_1, \dots, u_\ell, v_1, \dots, v_{\ell-1})$$

and for any $i \in \llbracket 1, \ell - 1 \rrbracket$, set

$$\mathcal{K}_n^{(i)} = \lfloor \mathcal{K}_n \Delta t_i \rfloor + \lfloor v_i \gamma_n \rfloor,$$

and

$$A_n^w(\Theta) := \{ \mathbf{S} \in \mathcal{W}_n^{(\mathcal{K}_n)}, S(2Nt_i^{(n)}) = 2\lfloor u_i \beta_n \rfloor, \text{ for } i \in \llbracket 1, \ell - 1 \rrbracket, S(n) = 2\lfloor u_\ell \beta_n \rfloor + \mathcal{E}_n, \\ \tilde{\Lambda}_i = \mathcal{K}_n^{(i)} \text{ for } i \in \llbracket 1, \ell - 1 \rrbracket \}.$$

For any $i \in \llbracket 1, \ell - 1 \rrbracket$, denote by $a_i(\mathbf{S}) = \Delta S_{2Nt_{i-1}^n+1}$ and $b_i(\mathbf{S}) = \Delta S_{2Nt_i^n}$ the first and last increments of the i th part of \mathbf{S} between the discretization points. Some peaks may appear in the positions $2Nt_i^n$, and then, we must take into account the pairs (b_i, a_{i+1}) to compute the cardinality of $A_n^w(\Theta)$. For any $\mathbf{S} \in A_n^w(\Theta)$, the number of peaks in $\llbracket 2Nt_{\ell-1} + 1, n - 1 \rrbracket$ is

$$\mathcal{K}_n^{(\ell)} = \mathcal{K}_n - \sum_{i=1}^{\ell-1} \mathcal{K}_n^{(i)} - \sum_{i=1}^{\ell-1} \mathbb{1}_{(b_i, a_{i+1})=(u, d)}.$$

We have $\#A_n^w(\Theta) =$

$$\sum_c \left(\prod_{i=1}^{\ell-1} \#T_{a_i, b_i}^{\mathcal{K}_n^{(i)}}(2N\Delta t_i^n, 2\lfloor u_{i-1} \beta_n \rfloor, 2\lfloor u_i \beta_n \rfloor) \right) \#T_{a_\ell, b_\ell}^{\mathcal{K}_n^{(\ell)}}(2N\Delta t_\ell^n, 2\lfloor u_{\ell-1} \beta_n \rfloor, 2\lfloor u_\ell \beta_n \rfloor + \mathcal{E}_n) \quad (19)$$

where the sum is taken over every $c := ((a_1, b_1), \dots, (a_\ell, b_\ell)) \in \{u, d\}^\ell$.

In order to evaluate the sum (19), we introduce some binomial random variables $B(l, p_n)$ with parameters l and p_n and we use the following version of the local limit theorem

Lemma 11. *Let $(l(n))$ be a sequence of integers going to $+\infty$ and $\sigma_n^2 = l(n)p_n(1 - p_n)$. We have*

$$\mathbb{P}(B(l(n), p_n) = m) = \binom{l(n)}{m} p_n^m (1 - p_n)^{l(n)-m} = \frac{1}{\sigma_n \sqrt{2\pi}} \exp\left(-\frac{(m - l(n)p_n)^2}{2\sigma_n^2}\right) + o(1/\sigma_n). \quad (20)$$

This may be proved using Stirling formula. As a consequence, if (a_n) , (a'_n) , (a''_n) are sequences of integers such that

$$a_n - nt/2 = O(1) \text{ for } t \in (0, 1), \quad \frac{a'_n}{\beta_n} \longrightarrow u, \quad \frac{a''_n - t\mathcal{K}_n}{\gamma_n} \longrightarrow v,$$

then

$$\mathbb{P}(B(a_n + a'_n, p_n) = a''_n) \sim \frac{1}{\sqrt{\pi t \gamma_n}} \exp\left(-\frac{(v - u)^2}{t}\right).$$

We then get easily that

$$\#T_{a_i, b_i}^{\mathcal{K}_n^{(i)}}(2N\Delta t_i^n, 2\lfloor u_{i-1} \beta_n \rfloor, 2\lfloor u_i \beta_n \rfloor) (1 - p_n)^{2N\Delta t_i^n - 2} \left(\frac{p_n}{1 - p_n}\right)^{2\mathcal{K}_n^{(i)} - \mathbb{1}_{b_i=d} - \mathbb{1}_{a_i=d}}$$

is equivalent to $\frac{1}{\pi \Delta t_i \gamma_n^2} \exp\left(-2\frac{v_i^2 + \Delta u_i^2}{\Delta t_i}\right) = 2\gamma_n^{-2} \mathbf{p}_{\Delta t_i}(0, 2v_i) \mathbf{p}_{\Delta t_i}(0, 2\Delta u_i)$ and

$$\#T_{a_\ell, b_\ell}^{\mathcal{K}_n^{(\ell)}}(2N\Delta t_\ell^n, 2\lfloor u_{\ell-1} \beta_n \rfloor, 2\lfloor u_\ell \beta_n \rfloor + \mathcal{E}_n) (1 - p_n)^{n - 2Nt_{\ell-1}^{(n)} - 2} \left(\frac{p_n}{1 - p_n}\right)^{2\mathcal{K}_n^{(\ell)} - \mathbb{1}_{b_\ell=d} - \mathbb{1}_{a_\ell=d}}$$

is equivalent to $\frac{1}{\pi \Delta t_\ell \gamma_n^2} \exp\left(-2 \frac{(\sum_{i=1}^{\ell-1} v_i)^2 + \Delta u_\ell^2}{\Delta t_\ell}\right) = 2\gamma_n^{-2} \mathbf{p}_{\Delta t_\ell}(2 \sum_{i=1}^{\ell-1} v_i, 0) \mathbf{p}_{\Delta t_\ell}(0, 2\Delta u_\ell)$.

Since $p_n^{2\mathcal{K}_n+1} (1-p_n)^{n-2\mathcal{K}_n} \#\mathcal{W}_n^{(\mathcal{K}_n)} \sim 1/(\gamma_n \sqrt{2\pi})$ we obtain

$$\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}(A_n^{\mathbf{w}}(\Theta)) = \frac{\#A_n^{\mathbf{w}}(\Theta)}{\#\mathcal{W}_n^{(\mathcal{K}_n)}} \sim c_n^{\mathbf{w}} f_{t_1, \dots, t_\ell}^{\mathbf{w}/2}(u_1, \dots, u_\ell) f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(v_1, v_1 + v_2, \dots, v_1 + \dots + v_{\ell-1})$$

where

$$\begin{aligned} c_n^{\mathbf{w}} &:= 2^{1-\ell} \gamma_n^{1-2\ell} p_n (1-p_n)^{2\ell} \sum_c \left(\frac{p_n}{1-p_n} \right)^{\mathbb{1}_{a_1=u} + \mathbb{1}_{b_\ell=d} + \sum_{i=1}^{\ell-1} 2\mathbb{1}_{(b_i, a_{i+1})=(u,d)} + \mathbb{1}_{b_i=d} + \mathbb{1}_{a_{i+1}=u}} \\ &= \gamma_n^{1-2\ell} p_n^\ell = \gamma_n^{1-\ell} \beta_n^{-\ell} \end{aligned}$$

The contribution of the sum over c has been computed as follows:

$$\begin{aligned} \sum_c x^{f(c)} &= \left(\sum_{(a_1, b_\ell) \in \{+1, -1\}^2} x^{\mathbb{1}_{a_1=u} + \mathbb{1}_{b_\ell=d}} \right) \prod_{i=1}^{\ell-1} \sum_{(b_i, a_{i+1}) \in \{+1, -1\}^2} x^{\mathbb{1}_{(b_i, a_{i+1})=(u,d)} + \mathbb{1}_{b_i=d} + \mathbb{1}_{a_{i+1}=u}} \\ &= (1 + 2x + x^2)(2x + 2x^2)^{\ell-1} = 2^{\ell-1} x^{\ell-1} (1+x)^{\ell+1}. \end{aligned}$$

Finally, this says that

$$\beta_n^\ell \gamma_n^{\ell-1} \mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)} \left(\frac{S(2Nt_i)}{2} = \lfloor u_i \beta_n \rfloor, i \in \llbracket 1, \ell \rrbracket, \tilde{\Lambda}_i - \lfloor \mathcal{K}_n \Delta t_i \rfloor = \lfloor v_i \gamma_n \rfloor, i \in \llbracket 1, \ell-1 \rrbracket \right) \rightarrow f_{t_1, \dots, t_\ell}^{\mathbf{w}/2}(u_1, \dots, u_\ell) f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(v_1, v_1 + v_2, \dots, v_1 + \dots + v_{\ell-1}).$$

Hence by Proposition 10 and (5), and taking into account that for any i , $\lambda_n(t_i) - 2 \sum_{j=1}^i \frac{\tilde{\Lambda}_i - \lfloor \mathcal{K}_n \Delta t_i \rfloor}{\gamma_n} \xrightarrow{\text{proba}} 0$, this allows to conclude to the finite dimensional convergence in Theorem 4 in the case $\mathbf{x} = \mathbf{w}$.

Comments 2. To compute a local limit theorem under the other distributions the numbers $u_1, \dots, u_\ell, v_1, \dots, v_\ell$ and the set $A_n^{\mathbf{w}}$ have to be suitably changed. First, in each case, the set $\mathcal{W}_n^{(\mathcal{K}_n)}$ has to be replaced by the right set.

- In the case of excursions and bridges, n is an even number and u_ℓ is taken equal to 0.
- In the case of excursions $a_1 = u, b_\ell = d$
- In the case of excursions and meanders all the reals u_i are chosen positive. Moreover, T^\geq must replace T in the summation (19).

Up to these changes, the computations are very similar to the case of Bernoulli chains.

3.2.2 Case $\mathbf{x} = \mathbf{b}$

The computation is very similar to the previous case; the only differences are : here $n = 2N$ is even, $\#\mathcal{B}_n^{(\mathcal{K}_n)} = \binom{N}{\mathcal{K}_n}^2$, we set u_ℓ to 0 and we take $\Theta' := (t_1, \dots, t_{\ell-1}, u_1, \dots, u_{\ell-1}, 1, v_1, \dots, v_{\ell-1})$. We get

$$\mathbb{P}_n^{\mathbf{b}, (\mathcal{K}_n)}(A_n^{\mathbf{b}}(\Theta')) \sim c_n^{\mathbf{b}} f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(u_1, \dots, u_{\ell-1}) f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(v_1, v_1 + v_2, \dots, v_1 + \dots + v_{\ell-1}),$$

where $c_n^b := c_n^w \gamma_n p_n^{-1} = \gamma_n^{1-\ell} \beta_n^{1-\ell}$.

3.2.3 Case $x = e$

In this case $n = 2N$ is even, $u_\ell = 0$, $a_1 = u$, $b_\ell = d$. In order to avoid some problems with the formulas provided in Proposition 9 that have to be handled with precautions when x or y are 0, we will compute the local limit theorem “far” from 0. This will however suffice to conclude. For $i \in \llbracket 1, \ell - 1 \rrbracket$ we take $u_i > 0$, and β_n is assumed large enough so that $\lfloor u_i \beta_n \rfloor > 0$. For the calculus in the case of $x = e$, in formula (19), we replace T by T^\geq . Finally, $\#\mathcal{E}_n^{(\mathcal{K}_n)} = \frac{1}{N} \binom{N}{\mathcal{K}_n} \binom{N}{\mathcal{K}_n - 1}$.

We first treat the contribution of the non extreme parts of the trajectories, namely, $i \in \llbracket 2, \ell - 1 \rrbracket$,

$$\#T_{a_i, b_i}^{\mathcal{K}_n^{(i)}, \geq} (2N \Delta t_i^n, 2\lfloor u_{i-1} \beta_n \rfloor, 2\lfloor u_i \beta_n \rfloor) (1 - p_n)^{2N \Delta t_i^n - 2} \left(\frac{p_n}{1 - p_n} \right)^{2\mathcal{K}_n^{(i)} - \mathbb{1}_{b_i=d} - \mathbb{1}_{a_i=u}}$$

is equivalent to $2\gamma_n^{-2} \mathbf{p}_{\Delta t_i}(0, 2v_i) (\mathbf{p}_{\Delta t_i}(0, 2\Delta u_i) - \mathbf{p}_{\Delta t_i}(0, 2\bar{\Delta} u_i))$.

Let us consider $i = 1$. Notice that $\#T_{uu}^{j, \geq}(l, 0, y)$ and $\#T_{ud}^{j, \geq}(l, 0, y)$ may be very different :

$$T_{uu}^{j, \geq}(l, 0, y) = \binom{\frac{l+y}{2} - 1}{j-1} \binom{\frac{l-y}{2} - 1}{j-1} \frac{y}{j} \quad \text{and} \quad T_{ud}^{j, \geq}(l, 0, y) = \binom{\frac{l+y}{2} - 1}{j-1} \binom{\frac{l-y}{2} - 1}{j-1} \frac{2jy + l - y}{j(l+y-2j+2)}$$

$$A_{b_1}(n) := \#T_{u, b_1}^{\mathcal{K}_n^{(1)}, \geq} (2N \Delta t_1^n, 0, 2\lfloor u_1 \beta_n \rfloor) (1 - p_n)^{2N \Delta t_1^n - 2} \left(\frac{p_n}{1 - p_n} \right)^{2\mathcal{K}_n^{(1)} - \mathbb{1}_{b_1=d}}$$

We notice that under (H)

$$n = o(\mathcal{K}_n \beta_n), \quad \gamma_n = o(n - 2\mathcal{K}_n), \quad \gamma_n \leq \beta_n, \quad \gamma_n = o(\mathcal{K}_n).$$

We then get,

$$A_{b_1}(n) \sim \frac{1}{n \gamma_n (1 - p_n)^2} \frac{8u_1}{t_1} \mathbf{p}_{\Delta t_1}(0, 2u_1) \mathbf{p}_{\Delta t_1}(0, 2v_1).$$

The case $i = \ell$ is treated with the same method. We obtain

$$\mathbb{P}_n^{\mathbf{e}, (\mathcal{K}_n)}(A_n^{\mathbf{e}}(\Theta')) = \frac{\#A_n^{\mathbf{e}}(\Theta')}{\#\mathcal{E}_n^{(\mathcal{K}_n)}} \sim c_n^{\mathbf{e}} f_{t_1, \dots, t_{\ell-1}}^{\mathbf{e}/2}(u_1, \dots, u_{\ell-1}) f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(v_1, v_1 + v_2, \dots, v_1 + \dots + v_{\ell-1})$$

where $c_n^{\mathbf{e}} = \gamma_n^{-\ell+1} \beta_n^{-\ell+1}$.

3.2.4 Case $x = m$

The computation is the same as in the case $x = e$, except that u_ℓ is taken > 0 (and n large enough such that $\lfloor u_\ell \beta_n \rfloor \geq 1$). The last piece in the decomposition of meanders is of the same type as a standard excursion piece. We obtain

$$\mathbb{P}_n^{\mathbf{m}, (\mathcal{K}_n)}(A_n^{\mathbf{m}}(\Theta)) = \frac{\#A_n^{\mathbf{m}}(\Theta)}{\#\mathcal{M}_n^{(\mathcal{K}_n)}} \sim c_n^{\mathbf{m}} f_{t_1, \dots, t_{\ell-1}}^{\mathbf{m}/2}(u_1, \dots, u_{\ell-1}) f_{t_1, \dots, t_{\ell-1}}^{\mathbf{b}/2}(v_1, v_1 + v_2, \dots, v_1 + \dots + v_{\ell-1})$$

where $c_n^{\mathbf{m}} = \gamma_n^{-\ell+1} \beta_n^{-\ell}$.

4 Tightness

We begin with some recalls of some classical facts regarding tightness in $C[0, 1]$. First, tightness and relative compactness are equivalent in $C[0, 1]$ (and in any Polish space, by Prohorov). Consider the function modulus of continuity,

$$\begin{aligned} \omega : [0, 1] \times C[0, 1] &\longrightarrow \mathbb{R}^+ \\ (\delta, f) &\longmapsto \omega_\delta(f) \end{aligned}$$

defined by

$$\omega_\delta(f) := \sup_{s, t \in [0, 1], |s-t| \leq \delta} |f(t) - f(s)|.$$

A sequence of processes (x_n) such that $x_n(0) = 0$ is tight in $C[0, 1]$, if for any $\varepsilon > 0, \eta > 0$, there exists $\delta > 0$, such that for n large enough

$$\mathbb{P}(\omega_\delta(x_n) \geq \varepsilon) \leq \eta.$$

If the sequences (x_n) and (y_n) are tight in $C[0, 1]$ (and if for each n , x_n and y_n are defined on the same probability space Ω_n), then the sequence (x_n, y_n) is tight in $C([0, 1])^2$. We will use this result here, and prove the tightness separately for (s_n) and (λ_n) for every model \mathbb{P}_n^x .

We say that a sequence (x_n) in $C[0, 1]$ is tight on $[0, 1/2]$ if the sequence of restrictions $(x_n|_{[0, 1/2]})$ is tight in $C[0, 1/2]$. We would like to stress on the fact that we deal only with processes piecewise interpolated (on intervals $[k/n, (k+1)/n]$); for these processes, for n large enough such that $1/n < \delta$,

$$\left(\sup\{|x_n(t) - x_n(s)|, s, t \in \{k/n, k \in \llbracket 0, n \rrbracket\}, |s-t| \leq \delta\} \leq \varepsilon/3 \right) \Rightarrow (\omega_\delta(x_n) \leq \varepsilon).$$

In other words, one may assume that s and t are discretization points, in our proofs of tightness.

We recall a result by Petrov (19, Exercise 2.6.11) :

Lemma 12. *Let $(X_i)_i$ be i.i.d. centered random variables, such that $\mathbb{E}(e^{tX_1}) \leq e^{gt^2/2}$ for $|t| \in [0, T]$ and $g > 0$. Let $Z_k = X_1 + \dots + X_k$. Then*

$$\mathbb{P}\left(\max_{1 \leq k \leq N} |Z_k| \geq x\right) \leq 2 \begin{cases} \exp(-x^2/2Ng) & \text{for any } x \in [0, NgT] \\ \exp(-Tx) & \text{for any } x \geq NgT. \end{cases}$$

The tightness of (s_n, λ_n) is proved as follows: first, under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$, the passage via an alternative model of “simple random walk” allows to remove the conditioning by $\Lambda_n = \mathcal{K}_n$. Then, the tightness under $\mathbb{P}_n^{\mathbf{b}, (\mathcal{K}_n)}$ is deduced from that under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$, thanks to the fact that the conditioning by $S(n) = 0$ does not really change the distribution the first half of the trajectories. The tightness under $\mathbb{P}_n^{\mathbf{e}, (\mathcal{K}_n)}$ and $\mathbb{P}_n^{\mathbf{m}, (\mathcal{K}_n)}$ are then obtained from that under $\mathbb{P}_n^{\mathbf{b}, (\mathcal{K}_n)}$ by some usual trajectory transformations that preserve the main properties of the variations and peak distributions of the trajectories.

4.1 A correspondence between simple chains and Bernoulli chains

We denote by \mathcal{H}_n the set of “simple chains”, starting from 0 and having $n + 1$ steps :

$$\mathcal{H}_n = \{\mathbf{H} = (H_i)_{0 \leq i \leq n+1} : H_0 = 0, H_{i+1} = H_i \text{ or } H_{i+1} = H_i + 1 \text{ for any } i \in \llbracket 0, n \rrbracket\}.$$

We consider the map

$$\begin{aligned} \Phi_n : \mathcal{W}_n &\longrightarrow \mathcal{H}_n \\ \mathbf{S} &\longmapsto \mathbf{H} = \Phi_n(\mathbf{S}) \end{aligned}$$

where \mathbf{H} is the simple chain with increments: for any $i \in \llbracket 1, n + 1 \rrbracket$,

$$\begin{cases} \text{if } \Delta S_i \neq \Delta S_{i-1} & \text{then } \Delta H_i = 1 \\ \text{if } \Delta S_i = \Delta S_{i-1} & \text{then } \Delta H_i = 0 \end{cases}$$

where by convention $\Delta S_0 = -1$ and $\Delta S_{n+1} = 1$ (see illustration on Figure 5).




Figure 5: Correspondence between simple chains and Bernoulli chains

The mapping Φ_n is a combinatorial trick. Obviously, the map $\mathbf{S} \mapsto \mathbf{H}$ where \mathbf{H} is defined by $\Delta H_i = (\Delta S_i + 1)/2$ is a bijection from \mathcal{W}_n onto \mathcal{H}_{n-1} . The map Φ_n is then certainly not a bijection (it is an injection). But, Φ_n owns some interesting properties that will really simplify our task.

Each increasing step in \mathbf{H} corresponds to a changing of direction in \mathbf{S} . Since $\Delta S_0 = -1$, the first one corresponds then to a valley, and the last one to a valley (which can not be in position n , since $\Delta S_{n+1} = 1$). Hence, for any $j \in \llbracket 0, n \rrbracket$,

$$\Lambda_j(\mathbf{S}) = \#\mathbf{S}_\wedge \cap \llbracket 0, j \rrbracket = \left\lfloor \frac{H_{j+1}}{2} \right\rfloor.$$

Hence,

$$\mathbf{S}_\wedge = \{\mathcal{T}_{2l}(\mathbf{H}) - 1, l \in \mathbb{N}\} \quad \text{and} \quad (-\mathbf{S})_\wedge = \{\mathcal{T}_{2l+1}(\mathbf{H}) - 1, l \in \mathbb{N}\},$$

where $(-\mathbf{S})_\wedge$ is the set of valleys of \mathbf{S} and where $\mathcal{T}_l(\mathbf{H}) = \inf\{j, H_j = l\}$ is the hitting time by \mathbf{H} of the level l . The process \mathbf{S} may then be described with \mathbf{H} :

$$S(k) = \sum_{i=0}^{H_k-1} (-1)^{i+1} (\mathcal{T}_{i+1}(\mathbf{H}) - \mathcal{T}_i(\mathbf{H})) + (-1)^{H_k+1} (k - \mathcal{T}_{H_k}(\mathbf{H})). \quad (21)$$

To end these considerations, consider now the subset of simple chains with k increasing steps,

$$\mathcal{H}_n^k = \{\mathbf{H} \in \mathcal{H}_n, H_{n+1} = k\},$$

and focus on \mathcal{H}_n^{2k+1} . Each element $\mathbf{H} \in \mathcal{H}_n^{2k+1}$ is image by Φ_n of a unique trajectory \mathbf{S} that has k peaks in $\llbracket 1, n-1 \rrbracket$ and $k+1$ valleys in $\llbracket 0, n \rrbracket$ (that may be in position 0 and n by construction), in other words, to a trajectory of $\mathcal{W}_n^{(k)}$.

This may alternatively be viewed as follows: to build a trajectory of $\mathcal{W}_n^{(k)}$ choose $2k+1$ integers $i_1 < i_2 < \dots < i_{2k+1}$ in the set $\llbracket 0, n \rrbracket$. Then construct a trajectory from \mathcal{W}_n in placing a valley in $i_1, i_3, \dots, i_{2k+1}$, a peak in i_2, i_4, \dots, i_{2k} and fill in the gaps between these points by straight lines. Hence

Lemma 13. *For any $k \in \llbracket 0, \lfloor n/2 \rfloor \rrbracket$, the restriction of Φ_n on $\mathcal{W}_n^{(k)}$ is a bijection onto \mathcal{H}_n^{2k+1} .*

For any $p \in [0, 1]$, let \mathbb{Q}_p^n be the distribution on \mathcal{H}_n of the Bernoulli random walks with $n+1$ i.i.d. increments, Bernoulli $B(p)$ distributed (that is $\mathbb{Q}_p^n(\Delta H_i = 1) = 1 - \mathbb{Q}_p^n(\Delta H_i = 0) = p$). For any \mathbf{H} in \mathcal{H}_n ,

$$\mathbb{Q}_p^n(\{\mathbf{H}\}) = p^{H_{n+1}}(1-p)^{n+1-H_{n+1}},$$

and then, \mathbb{Q}_p^n gives the same weight to the trajectories ending at the same level. Hence the conditional distribution $\mathbb{Q}_p^n(\cdot | \mathcal{H}_n^{2k+1})$ is the uniform law on \mathcal{H}_n^{2k+1} . On the other hand, since $\mathbb{P}_n^{w,(k)}$ is the uniform distribution on $\mathcal{W}_n^{(k)}$, by Lemma 13, $\mathbb{P}_n^{w,(k)} \circ \Phi_n^{-1}$ is also the uniform law on \mathcal{H}_n^{2k+1} . Hence

Lemma 14. *For any $p \in (0, 1)$, $n \in \mathbb{N}$, $k \in \llbracket 0, \lfloor n/2 \rfloor \rrbracket$, $\mathbb{Q}_p^n(\cdot | \mathcal{H}_n^{2k+1}) = \mathbb{P}_n^{w,(k)} \circ \Phi_n^{-1}$.*

Using simple properties of binomial distribution, the value of p that maximizes $\mathbb{Q}_p^n(\mathcal{H}_n^{2\mathcal{K}_n+1})$ is $\tilde{p}_n = (2\mathcal{K}_n + 1)/(n + 1)$. This morally explains why in Section 3, p_n appears as a suitable parameter. For sake of simplicity, we will work again with $p_n = 2\mathcal{K}_n/n$ instead of \tilde{p}_n . We will see that under $\mathbb{Q}_{p_n}^n$, the conditioning by \mathcal{H}_n^{2k+1} is a “weak conditioning”, and to bound certain quantities, this conditioning may be suppressed, leading to easy computations. The archetype of this remark is the following property

Lemma 15. *Assume (H). There exists $c > 0$ such that for n large enough, for any set A_n on \mathcal{H}_n depending only on the first half part of the trajectories, (that is $\sigma(H_0, H_1, \dots, H_{\lfloor n/2 \rfloor})$ -measurable),*

$$\tilde{\mathbb{Q}}_n(A_n) := \mathbb{Q}_{p_n}^n(A_n | \mathcal{H}_n^{2\mathcal{K}_n+1}) \leq c \mathbb{Q}_{p_n}^n(A_n). \quad (22)$$

Proof. The idea is taken from the proof of Lemma 1 in (12) :

$$\begin{aligned}
\tilde{Q}_n(A_n) &= \sum_j \tilde{Q}_n(A_n, H_{\lfloor n/2 \rfloor} = j) \\
&= \sum_j \mathbb{Q}_{p_n}^n(A_n \mid H_{\lfloor n/2 \rfloor} = j, H_{n+1} = 2\mathcal{K}_n + 1) \tilde{Q}_n(H_{\lfloor n/2 \rfloor} = j) \\
&= \sum_j \mathbb{Q}_{p_n}^n(A_n \mid H_{\lfloor n/2 \rfloor} = j) \tilde{Q}_n(H_{\lfloor n/2 \rfloor} = j).
\end{aligned}$$

The latter equality comes from the Markov property of \mathbf{H} under $\mathbb{Q}_{p_n}^n$ that implies that $\mathbb{Q}_{p_n}^n(H_{n+1} = 2\mathcal{K}_n + 1 \mid A_n, H_{\lfloor n/2 \rfloor} = j) = \mathbb{Q}_{p_n}^n(H_{n+1} = 2\mathcal{K}_n + 1 \mid H_{\lfloor n/2 \rfloor} = j)$. It suffices to establish that there exists $c \geq 0$ such that for n large enough, for any j

$$\tilde{Q}_n(H_{\lfloor n/2 \rfloor} = j) \leq c \mathbb{Q}_{p_n}^n(H_{\lfloor n/2 \rfloor} = j).$$

Write

$$\tilde{Q}_n(H_{\lfloor n/2 \rfloor} = j) = \mathbb{Q}_{p_n}^n(H_{\lfloor n/2 \rfloor} = j) \frac{\mathbb{Q}_{p_n}^n(H_{\lfloor n/2 \rfloor} = 2\mathcal{K}_n + 1 - j)}{\mathbb{Q}_{p_n}^n(H_{n+1} = 2\mathcal{K}_n + 1)};$$

using Lemma 11, the last quotient is bounded, uniformly on j and $n \geq 1$. \square

A simple consequence of Lemma 15 is the following : let X_n be a positive random variable that depends only on the first half part of the trajectories, then the expectation of X_n under \tilde{Q}_n is bounded by the expectation of cX_n under $\mathbb{Q}_{p_n}^n$.

4.2 Tightness under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$

Assume that (\mathcal{K}_n) satisfies (H), that $\mathbf{S} \in \mathcal{W}_n$, and let $\mathbf{H} = \Phi_n(\mathbf{S})$. Set

$$h_n(t) = \frac{H_{(n+1)t}}{2\gamma_n} - \frac{t\mathcal{K}_n}{\gamma_n} \text{ for } t \in [0, 1], \quad (23)$$

where \mathbf{H} is assumed to be interpolated between integer points. Thanks to formulas (4), (5), (21),

$$|\lambda_n(t) - \lambda_n(s)| \leq |h_n(t) - h_n(s)| + 2/\gamma_n. \quad (24)$$

Hence, the tightness of (h_n) under \tilde{Q}_n implies the tightness of (λ_n) under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$.

By symmetry of the random walk under these distributions, we may prove the tightness only on $[0, 1/2]$. By Lemma 15, the tightness of (h_n) on $[0, 1/2]$ under $\mathbb{Q}_{p_n}^n$ implies the tightness of (h_n) on $[0, 1/2]$ under \tilde{Q}_n , and then that of (λ_n) on $[0, 1/2]$ under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$. Hence, it suffices to prove the tightness of (h_n) on $[0, 1/2]$ under $\mathbb{Q}_{p_n}^n$ to prove that of (λ_n) on $[0, 1]$ under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$.

Comments 3. *The conditioning by the number of peaks is a strong conditioning on \mathcal{W}_n . Indeed, $\mathcal{W}_n^{(\mathcal{K}_n)}$ may have a very small (even exponentially small) probability under $\mathbb{P}_n^{\mathbf{w}}$ when \mathcal{K}_n is far away from $n/4$: no tight bound can be derived using comparison between $\mathbb{P}_n^{\mathbf{w}}$ and $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$ by just removing the conditioning by $\Lambda_n = \mathcal{K}_n$. The passage by $\mathbb{Q}_{p_n}^n$ allows to remove this conditioning.*

Tightness of the sequence (λ_n) under $\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)}$

At first sight, under $\mathbb{Q}_{p_n}^n$, (h_n) is a random walk with the right normalization, and it should converge to the Brownian motion (and then the tightness should follow). However, we were unable to find a reference for this result under the present setting. We will then prove it.

In the sub-case where there exists $\delta > 0$ such that, for n large enough, p_n satisfies $n^{\delta-1} \leq p_n \leq 1 - n^{\delta-1}$ then under $\mathbb{Q}_{p_n}^n$, $h_n \xrightarrow[n]{(d)} \mathbf{w}$ in $C[0, 1]$: it is a consequence of Rackauskas, & Suquet (22, Theorem 2). In this case the tightness holds in a space of Hölder functions, with exponent smaller than $1/2$. When $p_n = o(n^{\delta-1})$ or $1 - p_n = o(n^{\delta-1})$, for any δ , (h_n) is not tight in any Hölder space; this may be checked by considering a single normalized step.

Let $\varepsilon > 0$ and $\eta > 0$ be fixed, and let us prove that for any n large enough, $\mathbb{Q}_{p_n}^n(\omega_\delta(h_n) \geq \varepsilon) \leq \eta$ for δ sufficiently small. So take a parameter $\delta \in (0, 1)$. We have

$$\omega_\delta(h_n) \leq 2 \max_{0 \leq j \leq \lfloor 1/(2\delta) \rfloor} \left(\max_{I_j^\delta(n)} h_n - \min_{I_j^\delta(n)} h_n \right) \quad (25)$$

where

$$I_j^\delta(n) = \left[\frac{2j \lfloor \delta(n+1) \rfloor}{n+1} \wedge 1, \frac{2(j+1) \lfloor \delta(n+1) \rfloor}{n+1} \wedge 1 \right]$$

(notice that the length of $I_j^\delta(n)$ is larger than δ for n large enough, and smaller than 3δ). The factor 2 in (25) simply comes from the splitting up $[0, 1]$ into parts. Since the extremities of the $I_j^\delta(n)$'s coincide with the discretization points, by the Markov property of h_n ,

$$\mathbb{Q}_{p_n}^n(\omega_\delta(h_n) \geq \varepsilon) \leq (1/(2\delta) + 1) \mathbb{Q}_{p_n}^n \left(\sup_{I_0^\delta(n)} |h_n| \geq \varepsilon/2 \right). \quad (26)$$

We need to control the supremum of a random walk, and we then use Lemma 12.

Lemma 16. *Let $B(p_n)$ be a Bernoulli random variable with parameter p_n . There exists $K > 0$, such that for any $n \geq 1$, any $|t| \leq \gamma_n$,*

$$\mathbb{E}(e^{t(B(p_n)-p_n)/\gamma_n}) \leq \exp \left(\frac{2K t^2}{n} \right). \quad (27)$$

Proof. There exists $K > 0$ such that, for any $|x| \leq 1$, $e^x \leq 1 + x + Kx^2$. Hence, for any $|t| \leq 1$, $\mathbb{E}(e^{t(B(p_n)-p_n)}) = p_n e^{t(1-p_n)} + (1-p_n) e^{-tp_n} \leq 1 + Kt^2(p_n(1-p_n)) \leq e^{2Kt^2 p_n(1-p_n)}$. Hence, for any $|t| \leq \gamma_n$, (27) holds (recall that $(1-p_n)p_n/\gamma_n^2 = 1/n$). \square

Let us end the proof of tightness of (h_n) . Since, for $N \in \llbracket 0, n \rrbracket$, $h_n(N/(n+1)) = \frac{Np_n}{2(n+1)\gamma_n} + D_N/2$ where D_N is a sum of N i.i.d. r.v. with law of $(B(p_n)-p_n)/\gamma_n$. Notice also that $\frac{Np_n}{2(n+1)\gamma_n} = o(1)$. By Lemmas 12 and 16, we get

$$\mathbb{Q}_{p_n}^n \left(\sup_{j \leq N} |D_j| \geq \varepsilon/2 \right) \leq 2 \exp(-\varepsilon^2 n / (16NK)) \text{ for any } \varepsilon \leq 4N\gamma_n K/n.$$

Let $\varepsilon > 0$ and $\delta > 0$ be fixed. For n large enough,

$$\begin{aligned} \mathbb{Q}_{p_n}^n(\omega_\delta(h_n) \geq \varepsilon) &\leq (1/(2\delta) + 1) \mathbb{Q}_{p_n}^n \left(\sup_{0 \leq t \leq 3\delta} |h_n(\lfloor (n+1)t \rfloor)| \geq \varepsilon/2 \right) \\ &\leq 2(1/(2\delta) + 1) \exp \left(-\frac{\varepsilon^2}{16K(3\delta)} \right) \end{aligned}$$

and this is smaller than any $\eta > 0$ for δ small enough, and n large enough. \square

Tightness of (s_n) under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$

Let $\delta \in (0, 1)$. First, it suffices to prove the tightness of (s_n) under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$ on $[0, 1/2]$. Consider formula (21). Denote by $G_{i+1}(\mathbf{H}) = \mathcal{T}_{i+1}(\mathbf{H}) - \mathcal{T}_i(\mathbf{H})$. For any s and t such that ns and nt are integer, and $s < t$, we have

$$s_n(t) - s_n(s) = \beta_n^{-1} \sum_{i=H_{ns}}^{H_{nt}-1} (-1)^{i+1} G_{i+1}(\mathbf{H}) + (-1)^{H_{ns}} ((ns - \mathcal{T}_{H_{ns}}(\mathbf{H})) - (nt - \mathcal{T}_{H_{nt}}(\mathbf{H}))). \quad (28)$$

The range of s_n in a subinterval $I \subset [0, 1/2]$ is then a function of the values of h_n in the same interval. Denote by $y_n(s, t)$ the right hand side of (28). We may control the range of s_n under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$ by making some computations on $y_n(s, t)$ under $\tilde{\mathbb{Q}}_n$, and then, by Lemma 15, we may work with $y_n(s, t)$ under $\mathbb{Q}_{p_n}^n$. Under $\mathbb{Q}_{p_n}^n$, the variables $G_i(\mathbf{H})$ are geometrically $\mathcal{G}(p_n)$ distributed ($\mathbb{P}(G_1 = k) = p_n(1 - p_n)^{k-1} \mathbf{1}_{k \geq 1}$), and then the computations are simplified. By Formula (28), we have

$$|y_n(s, t)| \leq \left| \sum_{i=H_{ns}}^{H_{nt}-1} \beta_n^{-1} (G_{i+2}(\mathbf{H}) - G_{i+1}(\mathbf{H})) \mathbf{1}_{i \equiv H_{ns} \pmod{2}} \right| + 3\beta_n^{-1} \max_{j \leq H_n} G_j(\mathbf{H})$$

where, in the sum, we have packed the variables $G_i(\mathbf{H})$ per 2.

Denote by $\tilde{y}_n(s, t)$ the sum. Using that $\max_{k \leq m} G_k$ is a non decreasing function of m , we have

$$\mathbb{Q}_{p_n}^n \left(\frac{\sup_{j \leq H_n} G_j(\mathbf{H})}{\beta_n} \geq \varepsilon \right) \leq \mathbb{Q}_{p_n}^n (H_n > 4\mathcal{K}_n) + \mathbb{Q}_{p_n}^n \left(\frac{\sup_{j \leq 4\mathcal{K}_n} G_j(\mathbf{H})}{\beta_n} \geq \varepsilon \right). \quad (29)$$

Since H_n is a binomial random variable with parameters n and $p_n = 2\mathcal{K}_n/n$, by the Bienaymé-Tchebichev's inequality, the first term in the right hand side is $O(\mathcal{K}_n^{-1})$. For the second term, write $\mathbb{Q}_{p_n}^n (\sup_{j \leq 4\mathcal{K}_n} G_j(\mathbf{H}) \geq \varepsilon \beta_n) = 1 - (1 - \mathbb{P}(G_1 \geq \varepsilon \beta_n))^{4\mathcal{K}_n}$. Since $\mathbb{P}(G_1 \geq \varepsilon \beta_n) = (1 - p_n)^{\lceil \varepsilon \beta_n \rceil - 1}$, we find that the second term goes to 0 when $n \rightarrow +\infty$.

It remains to control the variables $\tilde{y}_n(s, t)$. Using the Markov property of the random walk $z_k = \sum_{i=1}^k \beta_n^{-1} (G_{2i}(\mathbf{H}) - G_{2i-1}(\mathbf{H}))$, we get that under $\mathbb{Q}_{p_n}^n$,

$$\sup_{s, t \in I_j^\delta(n)} |\tilde{y}_n(s, t)| \stackrel{(d)}{=} \sup_{s, t \in I_1^\delta(n)} |\tilde{y}_n(s, t)| \leq 2 \sup_{t \in [0, 3\delta]} |\tilde{y}_n(0, t)|.$$

Writing \tilde{G}_i instead of $G_{2i} - G_{2i-1}$, we have

$$\left\{ \sup_{t \leq 3\delta} |\tilde{y}_n(0, t)| \geq \frac{\varepsilon}{2} \right\} \subset \left(\{H_{3n\delta} > 12\mathcal{K}_n\delta\} \cup \left\{ \sup_{j \leq 6\mathcal{K}_n\delta} \left| \beta_n^{-1} \sum_{i=1}^j \tilde{G}_i(\mathbf{H}) \right| \geq \varepsilon \right\} \right).$$

Once again, by Bienaymé-Tchebichev, $\mathbb{Q}_{p_n}^n(\{H_{3n\delta} > 12\mathcal{K}_n\delta\}) = O(1/(\delta\mathcal{K}_n))$. For the second set in the union, we have to control the maximum of a random walk with increments the variables \tilde{G} .

Lemma 17. *Let G_1 and G_2 be two independent geometrical random variables with parameter p_n . There exists $c > 0, c' > 0$, such that for any $|t| \leq c'\gamma_n$,*

$$\mathbb{E} \left(\exp \left(t \frac{G_1 - G_2}{\beta_n} \right) \right) \leq \exp \left(\frac{ct^2}{np_n} \right) \quad (30)$$

Proof. Write $\mathbb{E}(\exp(u(G_1 - G_2))) = \frac{p_n^2}{(e^{-u+p_n-1})(e^{u+p_n-1})}$. The denominator D is equal to $1 + (1 - p_n)^2 - (1 - p_n)(e^u + e^{-u})$. There exists a constant $c > 0$, such that for any $|u| \leq 1$, $(e^u + e^{-u}) \leq 2 + cu^2$. And then $D \geq p_n^2 - c(1 - p_n)u^2 \geq p_n^2(1 - \frac{c(1-p_n)u^2}{p_n^2}) \geq p_n^2 \exp(-\frac{c}{2} \frac{(1-p_n)u^2}{p_n^2})$ this last equality holds when $\frac{c}{2} \frac{(1-p_n)u^2}{p_n^2} \leq 1$. Hence $\mathbb{E}(\exp(u(G_1 - G_2))) \leq \exp(\frac{c}{2} \frac{(1-p_n)u^2}{p_n^2})$ for $|u| \leq \sqrt{\frac{2}{c(1-p_n)}} p_n$. Hence, (30) holds for $|t| \leq c'\beta_n p_n / \sqrt{1-p_n}$, and then for $|t| \leq c'\gamma_n$. \square

We end now the proof of tightness for the family (s_n) . According to Lemmas 12 and 17, for $\varepsilon > 0, \delta > 0$ fixed, for a constant $c'' > 0$ and n large enough

$$\mathbb{Q}_{p_n}^n \left(\beta_n^{-1} \sup_{j \leq 6\delta\mathcal{K}_n} \tilde{G}_j(\mathbf{H}) \geq \varepsilon/2 \right) \leq \exp(-\varepsilon^2/(c''\delta)).$$

This allows to conclude as for (h_n) . \square

4.3 Tightness under $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}$

In this section, n is an even number. Since $\mathbb{P}_n^{\mathbf{b},(k)}$ is the uniform distribution on $\mathcal{B}_n^{(k)}$, it coincides with the conditional law $\mathbb{P}_n^{\mathbf{w},(k)}(\cdot | S(n) = 0)$. We first establish a lemma that allows to control the probability of a set under $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}$, by the probability of the same set under $\mathbb{P}_n^{\mathbf{w},(k)}$.

Lemma 18. *Assume (H). There exists a constant $c > 0$ such that for n large enough, any $A_n \subset \mathcal{W}_n$,*

$$\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}(A_n) \leq c \beta_n \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n).$$

Proof. Write

$$\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}(A_n) = \frac{\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n, S(n) = 0)}{\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(S(n) = 0)} \leq \frac{\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n)}{\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(S(n) = 0)}.$$

Now $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(S(n) = 0) = \#\mathcal{B}_n^{(\mathcal{K}_n)} / \#\mathcal{W}_n^{(\mathcal{K}_n)} \sim \sqrt{2/\pi} \beta_n^{-1}$, by the local limit theorem. \square

Since $\gamma_n \rightarrow +\infty$, this lemma is interesting only for sets with probability $o(\gamma_n^{-1})$, e.g. :

Lemma 19. Assume (H). Let (α_n) be a sequence such that $\alpha_n \rightarrow +\infty$. There exist $c > 0$, $c' > 0$, such that for n large enough

$$\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)} (|\Lambda_{n/2} - \mathcal{K}_n/2| \geq \alpha_n \gamma_n) \leq \exp(-c' \alpha_n^2),$$

and then

$$\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)} (|\Lambda_{n/2} - \mathcal{K}_n/2| \geq \alpha_n \gamma_n) \leq c \beta_n \exp(-c' \alpha_n^2).$$

Proof. Assume again that $\mathbf{H} = \Phi_n(\mathbf{S})$. The variable $\Lambda_{n/2-1}(\mathbf{S}) = \lfloor H_{n/2}/2 \rfloor$ depends only on the first half of the trajectories. By Lemma 15, for any set I_n ,

$$\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)} (\Lambda_{n/2-1}(\mathbf{S}) \in I_n) = \tilde{Q}_n (\lfloor H_{n/2}/2 \rfloor \in I_n) \leq c'' \mathbb{Q}_{p_n}^n (\lfloor H_{n/2}/2 \rfloor \in I_n).$$

Under $\mathbb{Q}_{p_n}^n$, $H_{n/2}$ is a binomial random variable with parameters $n/2$ and p_n . Now, using Lemmas 12 and 16 with $I_n = \mathbb{C}[\mathcal{K}_n/2 - \alpha_n \gamma_n, \mathcal{K}_n/2 + \alpha_n \gamma_n]$, we get the first assertion. The second assertion is a consequence of Lemma 18. \square

Consider $\overline{\mathcal{W}}_n$ the set of simple walks satisfying

$$\overline{\mathcal{W}}_n = \{\mathbf{S} : \mathbf{S} \in \mathcal{W}_n, |\Lambda_{n/2} - \mathcal{K}_n/2| \leq \gamma_n^{5/4}\}.$$

Since by Lemma 19, $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}(\overline{\mathcal{W}}_n) \rightarrow 1$, we will from now on concentrate on these trajectories. We stress on the fact that $\gamma_n^{5/4} = o(\mathcal{K}_n)$. Assume that the following lemma is proved.

Lemma 20. Assume (H). There exists a constant $c > 0$ such that for n large enough, for any subset $A_n \subset \overline{\mathcal{W}}_n$ depending only on the first half of the trajectories,

$$\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}(A_n) = \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n | S(n) = 0) \leq c \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n). \quad (31)$$

This lemma, very similar to Lemma 15, allows to obtain the tightness of (s_n, λ_n) under $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}$ from that under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$; proceed as follows. By symmetry of bridges under $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}$, it suffices to prove the tightness on $[0, 1/2]$. Since $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}(\overline{\mathcal{W}}_n) \rightarrow 1$, we restrict our study to the trajectories of $\overline{\mathcal{W}}_n$. By (31), the tightness of (s_n, λ_n) under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$ on $[0, 1]$ (and then on $[0, 1/2]$) implies that under $\mathbb{P}_n^{\mathbf{b},(\mathcal{K}_n)}$ on $[0, 1/2]$. It only remains to prove Lemma 20.

Proof of Lemma 20 First, for any A_n , depending only on the first half of the trajectories,

$$\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n | S(n) = 0) = \sum_{(l,x,a)} \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n | (\Lambda_{n/2}, S(n/2), \Delta S_{n/2}) = (l, x, a)) \quad (32)$$

$$\times \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}((\Lambda_{n/2}, S(n/2), \Delta S_{n/2}) = (l, x, a) | S(n) = 0) \quad (33)$$

where the summation is taken on all possible triples (l, x, a) . Indeed, under $\mathbb{P}_n^{\mathbf{w}}$, the sequence $Y_k = (\Lambda_k, S(k), \Delta S_k)$ is a Markov chain, and also under $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}$. Then write $\mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(A_n | S(n) = 0) = \sum_{(l,x,a)} \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(S(n) = 0 | A_n, Y_{n/2} = (l, x, a)) \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(Y_{n/2} = (l, x, a)) / \mathbb{P}_n^{\mathbf{w},(\mathcal{K}_n)}(S(n) = 0)$,

then the first conditioning by A_n may be deleted, by Markov. A trite computation leads to the result.

Now, assume that $A_n \subset \overline{\mathcal{W}_n}$. The summation in (32) can be done on the triples (l, x, a) , such that $l \in J_n := \llbracket \mathcal{K}_n/2 - \gamma_n^{5/4}, \mathcal{K}_n/2 + \gamma_n^{5/4} \rrbracket$, $x \in \llbracket -n/2, n/2 \rrbracket$, $a \in \{u, d\}$. To end the proof, we check that there exists $c > 0$, valid for any $(l, x, a) \in J_n \times \llbracket -n/2, n/2 \rrbracket \times \{u, d\}$, and n large enough, such that

$$\mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)} (Y_{n/2} = (l, x, a) \mid S(n) = 0) \leq c \mathbb{P}_n^{\mathbf{w}, (\mathcal{K}_n)} (Y_{n/2} = (l, x, a)). \quad (34)$$

We choose to condition by the last increment of \mathbf{S} in $\llbracket 0, n/2 \rrbracket$ for computation reasons.

For any $a \in \{u, d\}$, denote $T_{-a} = T_{ua} \cup T_{da}$ and similar notation for T_{a-} and T_{--} .

- Case $a = d$. In this case, the left hand side of (34) equals

$$\frac{\#T_{-d}^l(n/2, x, 0) \#T_{--}^{\mathcal{K}_n-l}(n/2, x, 0)}{\#\mathcal{B}_n^{(\mathcal{K}_n)}}$$

and the right hand side

$$\frac{\#T_{-d}^l(n/2, x, 0) \#\mathcal{W}_{n/2}^{(\mathcal{K}_n-l)}}{\#\mathcal{W}_n^{(\mathcal{K}_n)}}.$$

Since for any $c > 0$, the map $x \mapsto \Gamma(x)/\Gamma(x-c)$ is log-concave,

$$\#T_{--}^{\mathcal{K}_n-l}(n/2, x, 0) = \binom{\frac{n/2-x}{2}}{\mathcal{K}_n-l} \binom{\frac{n/2+x}{2}}{\mathcal{K}_n-l} \leq \binom{\lfloor n/4 \rfloor}{\mathcal{K}_n-l} \binom{\lceil n/4 \rceil}{\mathcal{K}_n-l} := g_{n,l}.$$

Using that

$$\frac{\#\mathcal{B}_n^{(\mathcal{K}_n)}}{\#\mathcal{W}_n^{(\mathcal{K}_n)}} = \frac{p_n \#\mathcal{B}_n^{(\mathcal{K}_n)} p_n^{2\mathcal{K}_n} (1-p_n)^{n-2\mathcal{K}_n}}{\#\mathcal{W}_n^{(\mathcal{K}_n)} p_n^{2\mathcal{K}_n+1} (1-p_n)^{n-2\mathcal{K}_n}} \sim p_n \frac{\sqrt{2\pi}\gamma_n}{2\pi\gamma_n^2/2}$$

To prove (34) when $a = d$ it suffices to prove that

$$\limsup_n \max_{l \in J_n} \frac{\gamma_n}{p_n} \frac{g_{n,l}}{\#\mathcal{W}_{n/2}^{(\mathcal{K}_n-l)}} < +\infty$$

(notice that $\lfloor n/4 \rfloor + \lceil n/4 \rceil = n/2$ for any even n .) We have

$$\frac{\gamma_n}{p_n} \frac{g_{n,l}}{\#\mathcal{W}_{n/2}^{\mathcal{K}_n-l}} = \gamma_n \frac{\binom{\lfloor n/4 \rfloor}{\mathcal{K}_n-l} \binom{\lceil n/4 \rceil}{\mathcal{K}_n-l}}{\binom{n/2}{2(\mathcal{K}_n-l)} \left(\frac{n/2+1}{2\mathcal{K}_n-2l+1} p_n \right)}$$

The (second) parenthesis in the denominator is bounded. It remains to prove that $\limsup_n \max_{l \in J_n} G_n(l)$ is bounded, for

$$G_n(l) := \frac{\gamma_n \binom{\lfloor n/4 \rfloor}{\mathcal{K}_n-l} \binom{\lceil n/4 \rceil}{\mathcal{K}_n-l}}{\binom{n/2}{2(\mathcal{K}_n-l)}}.$$

We will prove this assertion by showing that for any sequence (l_n) of integers, that satisfies $l_n \in J_n$ for any n , $(G_n(l_n))$ converges to a constant that does not depend on (l_n) . This allows to conclude, since one may take the sequence (l_n) s.t. $G_n(l_n)$ maximizes $G_n(l)$ on J_n for any n . Set $\rho_n = 4(\mathcal{K}_n - l_n)/n$. Since $l_n \in J_n$, for any n , $\rho_n \in (0, 1)$. Now, one checks easily that

$$G_n(l_n) = \frac{\gamma_n \mathbb{P}(B(\lfloor n/4 \rfloor, \rho_n) = \mathcal{K}_n - l_n) \mathbb{P}(B(\lceil n/4 \rceil, \rho_n) = \mathcal{K}_n - l_n)}{\mathbb{P}(B(n/2, \rho_n) = 2(\mathcal{K}_n - l_n))}.$$

Since $\mathcal{K}_n - l_n \sim \mathcal{K}_n/2$, by the central local limit theorem, it converge to $2/\sqrt{\pi}$.

• Case $a = u$. In this case, the left hand side of (34) equals

$$\frac{\#T_{-u}^l(n/2, x, 0)(\#T_{-u}^{\mathcal{K}_n-l}(n/2, x, 0) + \#T_{-d}^{\mathcal{K}_n-l-1}(n/2, x, 0))}{\#\mathcal{B}_n^{(\mathcal{K}_n)}}$$

and the right hand side

$$\frac{\#T_{-d}^l(n/2, x, 0)(\#\mathcal{W}_{n/2,u}^{(\mathcal{K}_n-l)} + \#\mathcal{W}_{n/2,d}^{(\mathcal{K}_n-l-1)})}{\#\mathcal{W}_n^{(\mathcal{K}_n)}},$$

where $\mathcal{W}_{n,a}^k$ is the set of trajectories \mathbf{S} with k peaks with $\Delta S_1 = a$.

Once again, it suffices to check that the quotient

$$\#T_{-u}^{\mathcal{K}_n-l}(n/2, x, 0) + \#T_{-d}^{\mathcal{K}_n-l-1}(n/2, x, 0) = \binom{1 + \frac{n/2-x}{2}}{k-l} \binom{\frac{n/2+x}{2} - 1}{\mathcal{K}_n - l - 1}$$

divided by

$$\#\mathcal{W}_{n/2,u}^{\mathcal{K}_n-l} + \#\mathcal{W}_{n/2,d}^{(\mathcal{K}_n-l-1)} = \binom{n/2+1}{2(\mathcal{K}_n-l)}$$

is bounded by $c \#\mathcal{B}_n^{(\mathcal{K}_n)} / \#\mathcal{W}_n^{(\mathcal{K}_n)}$. The same arguments lead to the same conclusion. \square

4.4 Tightness under $\mathbb{P}_n^{m,(\mathcal{K}_n)}$

The case n even

Assume first that $n = 2N$ is even. We recall a bijection $\Psi_{2N} : \mathcal{B}_{2N} \rightarrow \mathcal{M}_{2N}$, illustrated on Figure 6, that maps $\mathcal{B}_{2N}^{(k)}$ on $\mathcal{M}_{2N}^{(k)}$, and that moreover preserves sufficiently the trajectories, to prove that the tightness of (s_{2N}, λ_{2N}) under $\mathbb{P}_{2N}^{b,(\mathcal{K}_{2N})}$ yields that under $\mathbb{P}_{2N}^{m,(\mathcal{K}_{2N})}$.

The map $\Psi_{2N} : \mathcal{B}_{2N} \rightarrow \mathcal{W}_{2N}$ (we will see later that $\Psi_{2N}(\mathcal{B}_{2N}) = \mathcal{M}_{2N}$) is defined as follows. Let $\mathbf{S} \in \mathcal{B}_{2N}$ and $m = \min \mathbf{S} \leq 0$ its minimum. For $j \in \{1, \dots, -m\}$, let $t_j = \tau_{-j}(\mathbf{S})$ the reaching time of $-j$ by \mathbf{S} . Write $I_{\mathbf{S}} = \{t_j, j \geq 1\}$. Notice that when $m = 0$, $I_{\mathbf{S}} = \emptyset$.

The trajectory $\Psi_{2N}(\mathbf{S}) = \mathbf{Z} = (Z_i)_{i=0, \dots, n}$ is defined by $Z_0 = 0$ and :

$$\Delta Z_i = \begin{cases} \Delta S_i & \text{if } i \notin I_{\mathbf{S}}, \\ -\Delta S_i + 1 & \text{if } i \in I_{\mathbf{S}}. \end{cases}$$

Proposition 21. *For any even $2N$, Ψ_{2N} is a bijection from \mathcal{B}_{2N} onto \mathcal{M}_{2N} ; moreover, for any k , its restriction to $\mathcal{B}_{2N}^{(k)}$ is a bijection onto $\mathcal{M}_{2N}^{(k)}$ that preserves the peak positions.*

Proof. First, it is easy to see that if $\mathbf{Z} = \Psi_{2N}(\mathbf{S})$, for any $i \leq 2N$,

$$Z_i = S(i) + 2 \min_{j \leq i} S(j). \quad (35)$$

Hence, $\Psi_{2N}(\mathcal{B}_{2N}) \subset \mathcal{M}_{2N}$. Since Ψ_{2N} is clearly an injection, the first assertion of the Proposition follows $\#\mathcal{B}_{2N} = \#\mathcal{M}_{2N}$. Since Ψ_{2N} neither creates nor destroys any peaks, nor even changes the position of the peaks, the restriction of Ψ_{2N} onto $\mathcal{B}_{2N}^{(k)}$ is a bijection onto $\Psi_{2N}(\mathcal{B}_{2N}^{(k)}) \subset \mathcal{M}_{2N}^{(k)}$. The equality $\#\mathcal{B}_{2N}^{(k)} = \#\mathcal{M}_{2N}^{(k)}$ suffices then to conclude. \square




Figure 6: Synthetic description of Ψ_{2N} . The map Ψ_{2N} turns over each increment corresponding to a reaching time of a negative position. The map Ψ_{2N}^{-1} turns over the last increments reaching each position $x \in \llbracket 1, Z_{2N}/2 \rrbracket$ (Z_{2N} is even).

By Proposition 21, the tightness of (λ_{2N}) under $\mathbb{P}_{2N}^{m,(\mathcal{K}_{2N})}$ is a consequence of that under $\mathbb{P}_{2N}^{b,(\mathcal{K}_{2N})}$. For (s_{2N}) , (35) implies that the modulus of continuity of the non normalized trajectories are related by

$$\omega_{\mathbf{Z}}(\delta) \leq 3\omega_{\mathbf{S}}(\delta) \text{ for any } \delta \in [0, n]$$

and then, the tightness of (s_{2N}) under $\mathbb{P}_{2N}^{m,(\mathcal{K}_{2N})}$ follows that under $\mathbb{P}_{2N}^{b,(\mathcal{K}_{2N})}$.

The case n odd

The case $n = 2N + 1$ odd is very similar. There exists a bijection Ψ_{2N+1} between $\tilde{\mathcal{B}}_{2N+1}$ and \mathcal{M}_{2N+1} where $\tilde{\mathcal{B}}_{2N+1}$ is the subset of \mathcal{W}_{2N+1} of trajectories ending at position $+1$. The map Ψ_{2N+1} has the same properties as Ψ_{2N} to conserve the peak positions, and the set $\tilde{\mathcal{B}}_{2N+1}^{(\mathcal{K}_{2N+1})} := \tilde{\mathcal{B}}_{2N+1} \cap \mathcal{W}_{2N+1}^{(\mathcal{K}_{2N+1})}$ is sent on $\mathcal{M}_{2N+1}^{(\mathcal{K}_{2N+1})}$. To conclude, we need a tightness result for the uniform distribution on $\tilde{\mathcal{B}}_{2N+1}^{(\mathcal{K}_{2N+1})}$. But the result of Section 4.3 regarding $\mathcal{B}_{2N}^{(\mathcal{K}_{2N})}$ may be generalized to $\mathcal{B}_{2N+1}^{(\mathcal{K}_{2N+1})}$.




Figure 7: Synthetic description of Ψ_{2N+1} . The map Ψ_{2N+1} turns over each increment corresponding to a reaching time of a negative position. The map Ψ_{2N+1}^{-1} turns over the last increments reaching each position $x \in \llbracket 1, \lfloor Z_{2N+1}/2 \rfloor \rrbracket$ (Z_{2N+1} is odd).

Hence, $(s_{2N+1}, \lambda_{2N+1})$ is tight under $\mathbb{P}_{2N+1}^{m, (\mathcal{K}_{2N+1})}$, and then we may conclude that (s_n, λ_n) is tight under $\mathbb{P}_n^{m, (\mathcal{K}_n)}$.

4.5 Tightness under $\mathbb{P}_n^{e, (\mathcal{K}_n)}$

Here $n = 2N$ is an even number. Consider

$$\begin{aligned} \check{B}_{2N+1} &= \{\mathbf{S}, \mathbf{S} \in \mathcal{W}_{2N+1}, S(2N+1) = -1\}, \\ \check{E}_{2N+1} &= \{\mathbf{S}, \mathbf{S} \in \mathcal{W}_{2N+1}, S(j) \geq 0 \text{ for any } j \in \llbracket 0, 2N \rrbracket, S(2N+1) = -1\} \end{aligned}$$

and $\check{B}_{2N+1}^{(K)} = \check{B}_{2N+1} \cap \mathcal{W}_{2N+1}^{(K)}$, $\check{E}_{2N+1}^{(K)} = \check{E}_{2N+1} \cap \mathcal{W}_{2N+1}^{(K)}$. Informally, \check{E}_{2N+1} (resp. $\check{E}_{2N+1}^{(K)}$) are Bernoulli excursion from \mathcal{E}_{2N} (resp. with K peaks) with an additional ending d -step, and \check{B}_{2N+1} and $\check{B}_{2N+1}^{(K)}$ are trajectories ending at -1 (resp. with K peaks).

Consider the map

$$\begin{aligned} \mathbf{R} : \check{E}_{2N+1} \times \llbracket 0, 2N \rrbracket &\longrightarrow \check{B}_{2N+1} \\ (\mathbf{S}, \theta) &\longmapsto \mathbf{R}(\mathbf{S}, \theta) = \mathbf{S}^{(\theta)} = (S^{(\theta)})_{i=0, \dots, 2N+1} \end{aligned}$$

defined by

$$\Delta S_k^{(\theta)} = \Delta S_{k+\theta \bmod 2N+1}$$

or equivalently $S_k^{(\theta)} = S(k + \theta \bmod 2N+1) - S(\theta) - \mathbf{1}_{k+\theta > 2N+1}$. Informally, $\mathbf{S} \mapsto \mathbf{S}^{(\theta)}$ exchanges the θ first steps of \mathbf{S} with the last $2N+1 - \theta$'s ones. The map \mathbf{R} is a bijection between $\check{E}_{2N+1} \times \llbracket 0, 2N \rrbracket$ and \check{B}_{2N+1} : this is the so-called cyclical lemma attributed to Dvoretzky-Motzkin, or Kemperman, or Otter see Pitman (21, Chapter 5) and Bertoin & al. (3). The peaks positions of $\mathbf{S}^{(\theta)}$ are obtained from that of \mathbf{S} by a shift of $-\theta \bmod 2N+1$, and $\#\mathbf{S}_\wedge^{(\theta)} = \#\mathbf{S}_\wedge - 1$

iff $\theta \in \mathbf{S}_\wedge$ (if $\theta \notin \mathbf{S}_\wedge$ then $\#\mathbf{S}_\wedge^{(\theta)} = \#\mathbf{S}_\wedge$).

For any $\mathbf{S} \in \mathcal{W}_{2N+1}$, set $\Xi_0(\mathbf{S}) = 0$, and for any $k \leq 2N - \#\mathbf{S}_\wedge$,

$$\Xi_m(\mathbf{S}) := \min\{j, j \geq \Xi_{m-1}(\mathbf{S}), j \leq 2N, j \notin \mathbf{S}_\wedge\},$$

the successive non-peak positions of \mathbf{S} in $\llbracket 0, 2N \rrbracket$. Consider the map

$$\begin{aligned} \widehat{\mathbf{R}} : \check{E}_{2N+1}^{(K)} \times \llbracket 0, 2N - K \rrbracket &\longrightarrow \check{B}_{2N+1}^{(K)} \\ (\mathbf{S}, \ell) &\longmapsto \widehat{\mathbf{R}}(\mathbf{S}, \ell) = \mathbf{S}^{(\Xi_\ell(\mathbf{S}))} \end{aligned} \cdot$$

Proposition 22. *For any $K \leq N$, any $N \geq 0$, the map $\widehat{\mathbf{R}}$ is a bijection from $\check{E}_{2N+1}^{(K)} \times \llbracket 0, 2N - K \rrbracket$ onto $\check{B}_{2N+1}^{(K)}$.*

Proof. It is a consequence of the two following points: for any \mathbf{S} , $m \mapsto \Xi_m(\mathbf{S})$ is a bijection from $\llbracket 0, \#\llbracket 0, 2N \rrbracket \setminus \mathbf{S}_\wedge \rrbracket$ onto $\llbracket 0, 2N \rrbracket \setminus \mathbf{S}_\wedge$, and \mathbf{R} is a bijection. \square

Consider (\mathbf{S}, θ) in $\check{E}_{2N+1}^{(K)} \times \llbracket 0, 2N - K \rrbracket$; for any u ,

$$\begin{aligned} \sup_{|m_1 - m_2| \leq u} |S(m_1) - S(m_2)| &\leq 2 \sup_{|m_1 - m_2| \leq u} |S^{(\theta)}(m_1) - S^{(\theta)}(m_2)| \quad (36) \\ \sup_{|m_1 - m_2| \leq u} \left| (\Lambda_{m_1} - \Lambda_{m_2})(\mathbf{S}) - \frac{m_1 - m_2}{2N} \mathcal{K}_{2N} \right| &\leq 2 \sup_{|m_1 - m_2| \leq u} \left| (\Lambda_{m_1} - \Lambda_{m_2})(\mathbf{S}^{(\theta)}) - \frac{m_1 - m_2}{2N} \mathcal{K}_{2N} \right|. \end{aligned}$$

Endow $\check{E}_{2N+1}^{(K)} \times \llbracket 0, 2N - K \rrbracket$ with the uniform distribution and consider a random element (\mathbf{S}, θ) under this law (\mathbf{S} is then uniform on $\check{E}_{2N+1}^{(K)}$). By the last proposition, $\mathbf{S}^{(\theta)}$ is uniform on $\check{B}_{2N+1}^{(K)}$. By (36), we have

$$\mathbb{P}_{2N}^{\varepsilon, (\mathcal{K}_{2N})}(\omega_\delta(s_{2N}) \geq \varepsilon) \leq \mathbb{P}_{2N+1}^{\mathbf{w}, (\mathcal{K}_{2N})}(2\omega_\delta(s_{2N}) \geq \varepsilon | S(2N+1) = -1)$$

and the same result holds for λ_{2N} . Once again the result of Section 4.3 concerning $\mathbb{P}_{2N}^{\mathbf{b}, (\mathcal{K}_{2N})} = \mathbb{P}_{2N}^{\mathbf{w}, (\mathcal{K}_{2N})}(\cdot | S(2N) = 0)$ can be generalized to $\mathbb{P}_{2N}^{\mathbf{w}, (\mathcal{K}_{2N})}(\cdot | S(2N+1) = -1)$. \square

Acknowledgments

We would like to thank Mireille Bousquet-Mélou who pointed many references, and for helpful discussions. We thank the referee for pointing out some important references.

References

- [1] D. Aldous, (1991) *The continuum random tree. II: An overview.*, Stochastic analysis, Proc. Symp., Durham/UK 1990, Lond. Math. Soc. Lect. Note Ser. 167, 23-70. MR1166406
- [2] B. Belkin, (1972) *An invariance principle for conditioned recurrent random walk attracted to a stable law.*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 21, 45-64. MR0309200

- [3] J. Bertoin, L. Chaumont & J. Pitman, (2003) *Path transformations of first passage bridges*, Elec. Comm. in Probab., **8**, Paper 17. MR2042754
- [4] J. Bertoin, & J. Pitman, (1994) *Path transformations connecting Brownian bridge, excursion and meander*, Bull. Sci. Math., II. Sér. 118, No.2, 147-166.
- [5] Ph. Biane, & M. Yor (1987) *Valeurs principales associées aux temps locaux browniens.*, Bull. Sci. Math. (2) 111 (1987), no. 1, 23–101.
- [6] P. Billingsley, (1968) *Convergence of probability measures*, New York-London-Sydney-Toronto: Wiley and Sons. MR0233396
- [7] E. Csáki & Y. Hu, (2004) *Invariance principles for ranked excursion lengths and heights.*, Electron. Commun. Probab. 9, 14-21.
- [8] M.P. Delest, & G. Viennot, (1984) *Algebraic languages and polyominoes enumeration.*, Theoret. Comput. Sci. 34, No. 1-2, 169–206. MR0774044
- [9] M. de Sainte Catherine, G. Viennot, (1985) *Combinatorial interpretation of integrals of products of Hermite, Laguerre and Tchebycheff polynomials.* in “Polynômes orthogonaux et applications,” Proc. Laguerre Symp., Bar-le- Duc/France 1984, Lect. Notes Math. 1171, 120-128.
- [10] T. Duquesne, & J.F. Le Gall (2002) *Random trees, Lévy processes and spatial branching processes*, Astérisque, 281. MR1954248
- [11] D.L. Iglehart (1974) *Functional central limit theorems for random walks conditioned to stay positive.*, Ann. Probability 2, 608–619.
- [12] S. Janson & J.F. Marckert, (2005) *Convergence of discrete snake.*, J. Theor. Probab. 18, No.3, 615-645. MR2167644
- [13] W.D.Kaigh, (1976), *An invariance principle for random walk conditioned by a late return to zero.*, Ann. Probability 4, No. 1, 115–121. MR0415706
- [14] C. Krattenthaler, (1997), *The enumeration of lattice paths with respect to their number of turns*, Advances in combinatorial methods and applications to probability and statistics, Stat. Ind. Technol., Birkhäuser Boston, 29–58. MR1456725
- [15] T.M. Liggett, (1968), *An invariance principle for conditioned sums of independent random variables*, J. Math. Mech. 18, 559–570. MR0238373
- [16] J.F. Marckert & A. Mokkadem, (2003) *The depth first processes of Galton-Watson trees converge to the same Brownian excursion.*, Ann. of Probab., Vol. 31, No. 3. MR1989446
- [17] P. Marchal, (2003) *Constructing a sequence of random walks strongly converging to Brownian motion*, 181–190 (electronic), Discrete Math. Theor. Comput. Sci. Proc., AC, Assoc. Discrete Math. Theor. Comput. Sci. MR2042386
- [18] T.V. Narayana, (1959) *A partial order and its applications to probability theory*, Sankhyā, 21, 91–98. MR0106498
- [19] V.V. Petrov, (1995) *Limit theorems of probability theory. Sequences of independent random variables.*, Oxford Studies in Probab. MR1353441
- [20] J. Pitman (1999) *Brownian motion, bridge excursion, and meander characterized by sampling at independent uniform times.*, Electron. J. Probab. 4, No.11. MR1690315
- [21] J. Pitman (2005) *Combinatorial Stochastic Processes*, Lectures from St. Flour Course, July 2002. (<http://stat.berkeley.edu/users/pitman/621.ps>), To appear in *Springer Lecture Notes in Mathematics*. MR2245368
- [22] A. Rackauskas, & Ch. Suquet, (2003) *Hölderian invariance principle for triangular arrays of random variables. (Principe d'invariance Hölderien pour des tableau triangulaires de variables aléatoires)*, Lith. Math. J. 43, No.4, 423-438 (translation from Liet. Mat. Rink 43, No.4, 513-532 (2003)).
- [23] R.P. Stanley, Richard (1999) *Enumerative combinatorics. Vol. 2.* Cambridge University Press. MR1676282
- [24] A. Wald, & J. Wolfowitz, (1940) *On a test whether two samples are from the same population*, Ann. Math. Statistics, 11, 147–162. MR0002083