

## PRINCIPAL EIGENVECTORS OF IRREGULAR GRAPHS\*

SEBASTIAN M. CIOABĂ<sup>†</sup> AND DAVID A. GREGORY<sup>‡</sup>

**Abstract.** Let  $G$  be a connected graph. This paper studies the extreme entries of the principal eigenvector  $x$  of  $G$ , the unique positive unit eigenvector corresponding to the greatest eigenvalue  $\lambda_1$  of the adjacency matrix of  $G$ . If  $G$  has maximum degree  $\Delta$ , the greatest entry  $x_{\max}$  of  $x$  is at most  $1/\sqrt{1 + \lambda_1^2/\Delta}$ . This improves a result of Papendieck and Recht. The least entry  $x_{\min}$  of  $x$  as well as the principal ratio  $x_{\max}/x_{\min}$  are studied. It is conjectured that for connected graphs of order  $n \geq 3$ , the principal ratio is always attained by one of the lollipop graphs obtained by attaching a path graph to a vertex of a complete graph.

**Key words.** Spectral radius, Irregular graph, Eigenvectors.

**AMS subject classifications.** 05C50, 15A18.

**1. Introduction.** The study of the eigenvectors of the adjacency matrix of a graph has led to many applications. The principal eigenvectors of graphs form the basis of the PageRank algorithm used by Google (see [2]). The eigenvector corresponding to the second largest eigenvalue of a connected graph has been used in spectral partitioning algorithms (see [14]). The ratio of a non-negative irreducible matrix has been studied by Ostrowski [9, 10], Minc [5], De Oliveira [8], Latham [4] and Zhang [17] among others.

In this paper, we study principal eigenvectors of connected irregular graphs. Our graph theoretic notation is standard, see West [15]. The *eigenvalues* of a graph  $G$  are the eigenvalues  $\lambda_i$  of its adjacency matrix  $A$ , indexed so that  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . The greatest eigenvalue,  $\lambda_1$ , is also called the *spectral radius*. If  $G$  is connected, then the multiplicity of  $\lambda_1$  is 1. The positive eigenvector of length 1 corresponding to  $\lambda_1$  will be called the *principal* eigenvector of  $G$ . Note that if  $G$  is  $k$ -regular and connected, then  $k = \lambda_1 > \lambda_2$  and the principal eigenvector of  $G$  is  $\frac{1}{\sqrt{n}}\mathbf{1}$ , where  $\mathbf{1}$  is the all one vector.

If  $x \in \mathbb{R}^n$ , we denote by  $x_{\max}$  the greatest entry of  $x$  and by  $x_{\min}$  the least entry of  $x$ . The *ratio* of  $x$  is defined as  $x_{\max}/x_{\min}$ . If  $A$  is a non-negative irreducible matrix and  $\rho$  is the positive eigenvalue of  $A$  of maximum modulus, then  $\sigma(A)$  denotes the ratio of  $x$ , where  $x$  is a positive eigenvector corresponding to  $\rho$ . The *principal ratio*,  $\gamma(G)$ , of a connected graph  $G$  is the ratio of the principal eigenvector of  $G$ . It is well known that  $G$  is regular if and only if  $\gamma(G) = 1$ . Thus, one can regard  $\gamma(G)$  as a measure of the irregularity of the graph  $G$ .

Schneider [12] has proved that if  $G$  is a connected graph, then  $\gamma(G) \leq \lambda_1^{n-1}$ . In Section 2, we improve Schneider's result as well as some recent results of Nikiforov

---

\*Received by the editors 2 April 2007. Accepted for publication 8 October 2007. Handling Editor: Hans Schneider.

<sup>†</sup>Department of Mathematics, University of California at San Diego, La Jolla, CA 92093-0112, USA (scioaba@math.ucsd.edu). Research supported by an NSERC Postdoctoral Fellowship

<sup>‡</sup>Department of Mathematics and Statistics, Queen's University, Kingston, Ontario K7L 3N6, Canada (gregoryd@mast.queensu.ca). Research supported by NSERC Canada.

[7]. We also present some lower bounds on  $\gamma(G)$  which improve previous results of Ostrowski [9] and Zhang [17].

If  $G$  is a connected graph and  $x$  is the principal eigenvector of  $G$ , then Papendieck and Recht [11] proved that  $x_{\max} \leq \frac{1}{\sqrt{2}}$  with equality if and only if  $G = K_{1,n-1}$ . In Section 3, we generalize and extend this result and we also prove some lower and upper bounds for the extreme entries of the principal eigenvector of a connected graph.

**2. The principal ratio of a graph.** In this section, we study the principal ratio  $\gamma(G)$  of a connected graph. As stated before, this parameter can be regarded as a measure of the irregularity of a graph.

**2.1. Upper bounds for  $\gamma(G)$ .** Let  $G$  be a connected graph with vertex set  $\{1, 2, \dots, n\}$ . Suppose that the path  $P_r$  on the first  $r$  vertices  $1, 2, \dots, r$  is a subgraph of  $G$ . Subgraphs found by taking the shortest path between two vertices in  $G$  and renumbering the vertices will have this property and will be induced paths. Such examples will have length at most  $D$ , the diameter of  $G$ .

Assume that  $\lambda_1 = \lambda_1(G) > 2$ . Let  $x$  be the principal eigenvector of  $G$ . Because  $Ax = \lambda_1 x$ , it follows that, for each vertex  $i$ ,

$$\lambda_1 x_i = \sum_{j \sim i} x_j.$$

Since the entries of  $x$  are all positive, if we let  $x_0 = 0$ , it follows that

$$(2.1) \quad x_2 \leq \lambda_1 x_1 \quad \text{and} \quad x_k \leq \lambda_1 x_{k-1} - x_{k-2} \quad \text{for} \quad 2 \leq k \leq r,$$

where  $x_2 = \lambda_1 x_1$  if and only if vertex 1 has degree 1 while  $x_k = \lambda_1 x_{k-1} - x_{k-2}$  if and only if vertex  $k-1$  has degree 2. Thus

$$\begin{bmatrix} x_k \\ x_{k-1} \end{bmatrix} \leq \begin{bmatrix} \lambda_1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x_{k-1} \\ x_{k-2} \end{bmatrix} \quad \text{for} \quad 2 \leq k \leq r.$$

Thus

$$\begin{bmatrix} x_k \\ x_{k-1} \end{bmatrix} \leq \begin{bmatrix} \lambda_1 & -1 \\ 1 & 0 \end{bmatrix}^{k-1} \begin{bmatrix} x_1 \\ x_0 \end{bmatrix} \quad \text{for} \quad 1 \leq k \leq r,$$

with equality for  $2 \leq k \leq r$  if and only if vertices  $2, 3, \dots, r-1$  all have degree 2.

Let  $\sigma, \tau = \frac{1}{2}(\lambda_1 \pm \sqrt{\lambda_1^2 - 4})$ , the eigenvalues of the  $2 \times 2$  matrix above. Because  $\lambda_1 > 2$ , then  $\sigma \neq \tau$  and there is an invertible matrix  $P$  such that

$$\begin{bmatrix} \lambda_1 & -1 \\ 1 & 0 \end{bmatrix} = P \begin{bmatrix} \sigma & 0 \\ 0 & \tau \end{bmatrix} P^{-1}.$$

Then

$$x_k \leq \begin{bmatrix} 1 & 0 \end{bmatrix} P \begin{bmatrix} \sigma^{k-1} & 0 \\ 0 & \tau^{k-1} \end{bmatrix} P^{-1} \begin{bmatrix} x_1 \\ x_0 \end{bmatrix} \quad \text{for} \quad 1 \leq k \leq r.$$

Thus, there are constants  $a, b$  such that

$$x_k \leq a\sigma^{k-1} + b\tau^{k-1} \quad \text{for } 1 \leq k \leq r.$$

Solving the recursion in the case of equality gives  $a+b = 1$  when  $k = 1$  and  $\sigma a + \tau b = \lambda_1$  when  $k = 2$ . Thus we may take  $a = (\lambda_1 - \tau)/(\sigma - \tau)$  and  $b = (\sigma - \lambda_1)/(\sigma - \tau)$ . Substituting these values and noting that  $\sigma\tau = 1$  and  $\sigma + \tau = \lambda_1$ , we get

$$(2.2) \quad x_k \leq \frac{\sigma^k - \tau^k}{\sigma - \tau} x_1 \quad \text{for } 1 \leq k \leq r,$$

with equality at a particular value of  $k \geq 2$  if and only if vertex 1 has degree 1 and vertices  $2, 3, \dots, k - 1$  all have degree 2.

**THEOREM 2.1.** *Let  $G$  be a connected graph of order  $n$  with spectral radius  $\lambda_1 > 2$  and principal eigenvector  $x$ . Let  $d$  be the shortest distance from a vertex on which  $x$  is maximum to a vertex on which it is minimum. Then*

$$(2.3) \quad \gamma(G) \leq \frac{\sigma^{d+1} - \tau^{d+1}}{\sigma - \tau},$$

where  $\sigma = \frac{1}{2}(\lambda_1 + \sqrt{\lambda_1^2 - 4})$  and  $\tau = \sigma^{-1}$ .

*Equality is attained if and only if either  $G$  is regular or there is an induced path of length  $d > 0$  whose endpoints index  $x_{\min}$  and  $x_{\max}$  and the degrees of the endpoints are 1 and 3 or more, respectively, while all other vertices of the path have degree 2 in  $G$ .*

*Proof.* Let  $x$  denote the principal eigenvector of  $G$ . If  $G$  is regular, then  $d = 0$  and the bound equals 1. If  $G$  is irregular, relabel the vertices so that  $x_1$  is a smallest entry of  $x$  and  $x_{d+1}$  a largest and let  $P_{d+1}$  be a path of length  $d$  connecting vertex 1 and vertex  $d + 1$ . Because  $\lambda_1 > 2$ , the upper bound (2.3) follows from (2.2) with  $r = d + 1$ .

If  $d = 0$ , then equality is attained if and only if  $\gamma(G) = 1$ , that is, if and only if  $G$  is regular. If  $d > 0$  and equality occurs then, by (2.2), vertex 1 has degree 1 and vertices  $2, 3, \dots, r - 1$  all have degree 2. Vertex  $r$  cannot have degree 1, otherwise  $x_{r-1} = \lambda_1 x_r > x_{r-1}$ , a contradiction. If vertex  $r$  has just one more neighbour,  $x_{r+1}$  say, then  $\lambda_1 x_r = x_{r+1} + x_r < 2x_r$  and so  $\lambda_1 < 2$ , a contradiction. Thus,  $x_r$  has degree at least 3. Conversely, if such a path exists and the endpoints index extreme entries, then all the inequalities in (2.2) are equalities and we have equality for the ratio.  $\square$

In a recent paper [7], Nikiforov proved that if  $H$  is a proper subgraph of a connected graph  $G$  with  $n$  vertices and diameter  $D$ , then

$$(2.4) \quad \lambda_1(G) - \lambda_1(H) > \frac{1}{n\lambda_1^{2D}(G)}.$$

In proving (2.4), Nikiforov uses the following inequality. If  $r$  is the distance between two vertices  $i$  and  $j$  of a connected graph  $G$ , then

$$(2.5) \quad \frac{x_i}{x_j} < \lambda_1(G)^r.$$

If  $\lambda_1 = \lambda_1(G) > 2$ , then (2.2) gives the following slight improvement of (2.5)

$$(2.6) \quad \frac{x_i}{x_j} \leq \frac{\sigma^{r+1} - \tau^{r+1}}{\sigma - \tau},$$

where  $\sigma = \frac{\lambda_1 + \sqrt{\lambda_1^2 - 4}}{2}$  and  $\tau = \frac{\lambda_1 - \sqrt{\lambda_1^2 - 4}}{2} = \frac{1}{\sigma}$ . Following the arguments from [7], inequality (2.6) can be used to show that if  $H$  is a proper subgraph of a connected graph  $G$  with  $n$  vertices and diameter  $D$ , then

$$\lambda_1(G) - \lambda_1(H) > \frac{(\sigma - \tau)^2}{n(\sigma^{D+1} - \tau^{D+1})^2} = \frac{1}{n(\sigma^D + \sigma^{D-2} + \dots + \sigma^{-(D-2)} + \sigma^{-D})^2}.$$

This is a slight improvement of inequality (2.4) since

$$\frac{1}{n\lambda_1^{2D}} = \frac{1}{n(\sigma + \tau)^{2D}} = \frac{1}{n\left(\sigma^D + \binom{D}{1}\sigma^{D-2} + \dots + \binom{D}{D-1}\sigma^{-(D-2)} + \sigma^{-D}\right)^2}.$$

When  $\lambda_1(G) \leq 2$ , inequality (2.4) can be further improved by using Smith's classification [13] of the connected graphs with spectral radius at most 2.

**2.2. Maximizing  $\gamma(G)$  over connected graphs on  $n$  vertices.** Let  $\gamma(n)$  be the maximum value of  $\gamma(G)$  taken over all connected graphs of order  $n$ . Let  $P_r \cdot K_s$  denote the graph of order  $n = r + s - 1$  formed by identifying an end vertex of a path  $P_r$  on  $r \geq 2$  vertices with a vertex of the complete graph  $K_s$  on  $s \geq 2$  vertices. In [1], Brightwell and Winkler call such graphs *lollipop* graphs and show that, for a graph on  $n$  vertices, the maximum expected time for a random walk between two vertices is attained on a lollipop graph  $P_r \cdot K_s$  of order  $n$  with  $s = \lceil (2n - 2)/3 \rceil$ . A computer search reveals that for  $3 \leq n \leq 9$ ,  $\gamma(n)$  is always attained by one of the two lollipop graphs  $P_r \cdot K_s$  of order  $n$  with  $s = \lceil (n + 1)/4 \rceil + \epsilon$  where  $\epsilon = 1$  or  $2$ . We conjecture that  $\gamma(n)$  is always attained by a lollipop graph of order  $n$ . As a first step in studying this conjecture, we examine the entries of the principal eigenvectors of the graphs  $P_r \cdot K_s$ .

**2.3. Principal eigenvectors of lollipop graphs.** Let the vertices of  $P_r$  in  $P_r \cdot K_s$  be  $\{1, 2, \dots, r\}$  and the vertices of  $K_s$  be  $\{r, r + 1, \dots, r + s - 1\}$ . We continue to assume that  $n = r + s - 1$  denotes the order of  $P_r \cdot K_s$  and that  $r \geq 2$ . However, we assume now that  $s \geq 3$ . This ensures that  $\lambda_1 = \lambda_1(P_r \cdot K_s) > 2$  since  $K_s$  is a proper induced subgraph. As before, we let  $\sigma = \frac{1}{2}(\lambda_1 + \sqrt{\lambda_1^2 - 4})$  and  $\tau = \sigma^{-1}$ .

LEMMA 2.2. *Let  $\lambda_1$  be the greatest eigenvalue of  $P_r \cdot K_s$  and let  $x$  be the principal eigenvector. Then*

$$x_k = \frac{\sigma^k - \tau^k}{\sigma - \tau} x_1 \quad \text{for } 1 \leq k \leq r,$$

while

$$x_k = \frac{1}{s-1} \frac{\sigma^{r+1} - \tau^{r+1}}{\sigma - \tau} x_1 \quad \text{for } r+1 \leq k \leq n.$$

*Proof.* Because the vertices  $\{1, 2, \dots, r\}$  are the vertices of a path, the inequalities (2.1) and (2.2) are equalities for  $k = 1, 2, \dots, r$ . This gives the first expression.

Because  $x$  is unique, by symmetry we have  $x_{r+1} = x_{r+2} = \dots = x_n$ . Thus,  $\lambda_1 x_r = (Ax)_r = x_{r-1} + (s-1)x_n$ . So, for  $k = r+1, \dots, n$ ,

$$\begin{aligned} x_k &= \frac{1}{s-1}(\lambda_1 x_r - x_{r-1}) \\ &= \frac{1}{s-1} \frac{(\sigma + \tau)(\sigma^r - \tau^r) - (\sigma^{r-1} - \tau^{r-1})}{\sigma - \tau} x_1 \\ &= \frac{1}{s-1} \frac{\sigma^{r+1} - \tau^{r+1}}{\sigma - \tau} x_1. \quad \square \end{aligned}$$

LEMMA 2.3. *Let  $\lambda_1$  be the greatest eigenvalue of  $P_r \cdot K_s$  and let  $x$  be the principal eigenvector. Then  $x_{r+1} = \dots = x_n$  and  $x_1 < x_2 < \dots < x_{r-1} < x_n < x_r$ . Thus*

$$\gamma(P_r \cdot K_s) = \frac{x_r}{x_1} = \frac{\sigma^r - \tau^r}{\sigma - \tau}.$$

Also,

$$\lambda_1^{r-1} - \lambda_1^{-r-1} < \gamma(P_r \cdot K_s) < \lambda_1^{r-1}.$$

*em Proof.* These results follow from Lemma 2.2, but it is perhaps clearer and shorter to give direct proofs based on the recursion (2.1). Because the graph has maximum degree  $s$  and  $K_s$  is an induced subgraph, it follows that

$$2 \leq s-1 < \lambda_1 < s.$$

Thus,  $\lambda_1 - s + 2 > 1$ ,  $\lambda_1 > 2$  and  $\sigma$  is real. Because  $1, 2, \dots, r$  are the vertices of a path, the inequalities (2.1) are equalities for  $k = 1, 2, \dots, r$ . Thus

$$x_k = \lambda_1 x_{k-1} - x_{k-2} = (\lambda_1 - 1)x_{k-1} + (x_{k-1} - x_{k-2}),$$

and it follows by induction that  $x_k < \lambda_1 x_{k-1}$  for  $k = 2, \dots, r$ . Thus,  $x_1 < x_2 < \dots < x_r$  and  $x_r < \lambda_1^{r-1} x_1$ . Because  $x$  is unique, by symmetry we have  $x_{r+1} = x_{r+2} = \dots = x_n$ . Thus  $\lambda_1 x_n = (Ax)_n = x_r + (s-2)x_n$  and so

$$(2.7) \quad x_r = (\lambda_1 - s + 2)x_n > x_n.$$

Also,  $(\lambda_1 - s + 2)x_n = x_r > (\lambda_1 - 1)x_{r-1} \geq (\lambda_1 - s + 2)x_{r-1}$  so  $x_{r-1} < x_n$ . Thus  $\gamma = x_r/x_1$ . Finally, because  $\sigma > \lambda_1$  and  $\sigma\tau = 1$ , we have  $\tau < 1/\lambda_1$  and so

$$\gamma = \frac{x_r}{x_1} = \frac{\sigma^r - \tau^r}{\sigma - \tau} > \frac{\lambda_1^r - \lambda_1^{-r}}{\sqrt{\lambda_1^2 - 4}} > \lambda_1^{r-1} - \lambda_1^{-r-1}. \quad \square$$

Note that substituting the expression (2.7) into the formulae in Lemma 2.2 gives

$$(2.8) \quad \frac{\sigma^{r+1} - \tau^{r+1}}{\sigma^r - \tau^r} = \frac{s-1}{\lambda_1 - s + 2}$$

a relation that determines  $\lambda_1$  in terms of  $r$  and  $s$ .

In the next lemma, we see that the parameter  $r$  has only a slight effect on the spectral radius of the graphs  $P_r \cdot K_s$ .

LEMMA 2.4. For  $r \geq 2$  and  $s \geq 3$ ,

$$s - 1 + \frac{1}{s(s-1)} < \lambda_1(P_r \cdot K_s) < s - 1 + \frac{1}{(s-1)^2}.$$

*Proof.* Because  $P_2 \cdot K_s$  is an induced subgraph of  $P_r \cdot K_s$ , we have  $\lambda_1(P_r \cdot K_s) \geq \lambda_1(P_2 \cdot K_s)$ . Thus, it is sufficient to establish the lower bound when  $r = 2$ .

If  $r = 2$ , then in (2.8),

$$\frac{\sigma^3 - \tau^3}{\sigma^2 - \tau^2} = \frac{\sigma^2 + \sigma\tau + \tau^2}{\sigma + \tau} = \frac{\lambda_1^2 - 1}{\lambda_1} = \frac{s - 1}{\lambda_1 - s + 2}.$$

Thus,  $\lambda_1$  is a root of the polynomial

$$p(x) = x^3 - (s - 2)x^2 - sx + s - 2 = (x - s + 1)(x^2 + x - 1) - 1$$

and so

$$(2.9) \quad \lambda_1 = s - 1 + \frac{1}{\lambda_1^2 + \lambda_1 - 1}.$$

Since  $\lambda_1 > s - 1$ , we deduce that

$$\lambda_1(P_2 \cdot K_s) < s - 1 + \frac{1}{s^2 - s - 1}.$$

Substituting this upper bound for  $\lambda_1$  in the denominator in (2.9), a lengthy calculation shows that, for  $s \geq 4$ ,

$$(2.10) \quad \lambda_1(P_2 \cdot K_s) > s - 1 + \frac{1}{s(s-1)}.$$

This also holds for  $s = 3$  because  $p(x) < 0$  when  $x = s - 1 + \frac{1}{s(s-1)} = 2 + \frac{1}{6}$ .

Suppose now that  $r \geq 3$ . Then from (2.8), we have

$$\lambda_1(P_r \cdot K_s) = s - 2 + (s - 1) \cdot \frac{\sigma^r - \tau^r}{\sigma^{r+1} - \tau^{r+1}} < s - 2 + (s - 1)\tau.$$

Thus, to obtain the upper bound, it is sufficient to show that

$$s - 2 + (s - 1)\tau < s - 1 + \frac{1}{(s-1)^2},$$

or, equivalently, that

$$2\tau = \lambda_1 - \sqrt{\lambda_1^2 - 4} < \frac{2}{s-1} + \frac{2}{(s-1)^3}.$$

Isolating and squaring the radical, we find that the above inequality is equivalent to

$$\lambda_1 \left( \frac{1}{s-1} + \frac{1}{(s-1)^3} \right) > 1 + \frac{1}{(s-1)^2} + \frac{2}{(s-1)^4} + \frac{1}{(s-1)^6}.$$

Substituting the lower bound (2.10) for  $\lambda_1$  shows that the inequality holds.  $\square$

Given  $n$ , it would be interesting to see if there are bounds that can be used to determine the values of  $r$  and  $s$  with  $r + s = n + 1$ , for which  $\gamma(P_r \cdot K_s)$  is maximum. A computer run indicates that for  $n \leq 50$ , the maximum is always attained by one of the two graphs  $P_r \cdot K_s$  of order  $n$  with  $s = \lceil (n + 1)/4 \rceil + \epsilon$  where  $\epsilon = 1$  or  $2$ .

**2.4. The principal ratio of an irreducible matrix.** In this subsection, we review some of the previous results regarding the ratio of the principal eigenvector of an irreducible matrix. This parameter has been studied by many researchers. Ostrowski [10] proved the following result.

LEMMA 2.5 (Ostrowski [10]). *If  $A$  is an  $n$  by  $n$  positive, irreducible matrix, then*

$$(2.11) \quad \gamma(A) \leq \frac{\max_{i,j \in [n]} a_{ij}}{\min_{i,j \in [n]} a_{ij}}.$$

Inequality 2.11 was improved by Minc [5].

THEOREM 2.6 (Minc [5]). *If  $A$  is an  $n$  by  $n$  positive irreducible matrix, then*

$$\gamma(A) \leq \max_{j,s,t \in [n]} \frac{a_{sj}}{a_{tj}}.$$

*Equality holds if and only if the  $p$ -th row of  $A$  is a multiple of the  $q$ -th row for some pairs of indices  $p$  and  $q$  satisfying*

$$\frac{a_{ph}}{a_{qh}} = \max_{j,s,t} \frac{a_{sj}}{a_{tj}}.$$

From Minc's proof [5], one can deduce the stronger inequality

$$\gamma(A) \leq \max_j \frac{a_{1j}}{a_{nj}},$$

where  $x_1 = x_{\max}$  and  $x_n = x_{\min}$ . This result along with other refinements of Minc's inequality were obtained by De Oliveira [8] and Latham [4].

Let  $k^{(i)}$  be the smallest positive entry of the  $i$ -th row of  $A$ . Ostrowski [10] also proved the following theorem.

THEOREM 2.7 (Ostrowski [10]). *If  $A$  is an irreducible non-negative matrix and assume that  $1, \dots, r$  is a path ( $a_{i,i+1} \neq 0$  for  $i \in [r-1]$ ) from 1 to  $r$ , where  $x_1 = x_{\max}$  and  $x_{\min} = x_r$ . Then*

$$(2.12) \quad \gamma(A) \leq \prod_{i=1}^{r-1} \frac{\rho - a_{ii}}{k^{(i)}}.$$

The proof of this result is done by using the inequality

$$(2.13) \quad \frac{x_{i+1}}{x_i} \leq \frac{\rho - a_{ii}}{k^{(i)}}$$

for each  $i \in [r - 1]$ . This follows from

$$(\rho - a_{ii})x_i = \sum_{j \neq i} a_{ij}x_j \geq k^{(i)}x_{i+1}.$$

The previous result implies a result of Schneider [12] who proved that

$$\gamma(A) \leq \left( \frac{\rho - k_1}{k} \right)^{n-1},$$

where  $k$  is the smallest positive entry of  $A$  and  $k_1 = \min a_{ii}$ . When  $A$  is the adjacency matrix of a connected graph  $G$  on  $n$  vertices, the previous inequality implies

$$\gamma(G) \leq \lambda_1^{n-1}(G)$$

which is weaker than Theorem 2.1.

Using the stronger inequality

$$(\rho - a_{ii})x_i = \sum_{j \neq i} a_{ij}x_j \geq k^{(i)}(x_{i+1} + x_{i-1})$$

and the argument from the proof of Theorem 2.1, one can improve Theorem 2.7.

**2.5. Lower bounds for  $\gamma$ .** For a matrix  $A$ , let  $k_1 = \min_{i \in [n]} a_{ii}$  and for  $i \in [n]$ , let  $d_i = \sum_{j=1}^n a_{ij}$ . Also, let  $\Delta = \max_{i \in [n]} d_i$  and  $\delta = \min_{i \in [n]} d_i$ .

Ostrowski [9] proved the following result.

**THEOREM 2.8** (Ostrowski [9]). *If  $A$  is a non-negative irreducible matrix, then*

$$(2.14) \quad \gamma \geq \max \left( \frac{\Delta - k_1}{\rho - k_1}, \frac{\rho - k_1}{\delta - k_1} \right) \geq \sqrt{\frac{\Delta - k_1}{\delta - k_1}}.$$

If we apply this inequality to the adjacency matrix of an irregular graph  $G$ , we obtain

$$\gamma(G) \geq \max \left( \frac{\Delta}{\lambda_1}, \frac{\lambda_1}{\delta} \right) \geq \sqrt{\frac{\Delta}{\delta}}.$$

This inequality has been also observed by Zhang [17].

We improve Theorem 2.8 as follows.

**THEOREM 2.9.** *If  $A$  is a non-negative irreducible matrix. Let  $T = \{t : d_t > \rho\}$  and  $S = \{s : d_s < \rho\}$ . Also, let  $i \in T$  such that  $d_i = \Delta$  and  $j \in S$  such that  $d_j = \delta$ . Then*

$$(2.15) \quad \gamma \geq \max \left( \frac{\Delta - a_{ii} + \sum_{r \in T \setminus \{i\}} a_{ir} \frac{d_r - \rho}{\rho - a_{rr}}}{\rho - a_{ii}}, \frac{\rho - a_{jj}}{\delta - a_{jj} - \sum_{r \in S \setminus \{j\}} \frac{a_{jr}(\rho - d_r)}{\rho - a_{rr}}} \right).$$



*Proof.* For each  $r$ , we have

$$(\rho - a_{rr})x_r = \sum_{l \neq r} a_{rl}x_l \geq (d_r - a_{rr})x_{\min} > 0,$$

where the final expression is positive since  $A$  is nonnegative and irreducible. Thus, for each  $r$ ,

$$\frac{x_r}{x_{\min}} \geq \frac{d_r - a_{rr}}{\rho - a_{rr}}.$$

Now

$$(\rho - a_{ii})x_i = \sum_{r \in T \setminus \{i\}} a_{ir}x_r + \sum_{s \notin T} a_{is}x_s.$$

Since  $d_i = \Delta$ , this implies

$$(\rho - a_{ii})x_i \geq x_{\min} \sum_{r \in T \setminus \{i\}} \frac{a_{ir}(d_r - a_{rr})}{\rho - a_{rr}} + (\Delta - \sum_{s \in T} a_{is})x_{\min}.$$

Putting  $d_r - a_{rr} = (d_r - \rho) + (\rho - a_{rr})$  and simplifying, we obtain the first lower bound

$$\gamma \geq \frac{x_i}{x_{\min}} \geq \frac{\Delta - a_{ii} + \sum_{r \in T \setminus \{i\}} a_{ir} \frac{d_r - \rho}{\rho - a_{rr}}}{\rho - a_{ii}}.$$

To obtain the second lower bound on  $\gamma$ , we note first that for each  $r$ ,

$$0 < (\rho - a_{rr})x_r = \sum_{l \neq r} a_{rl}x_l \leq (d_r - a_{rr})x_{\max}$$

and so

$$\frac{x_{\max}}{x_r} \geq \frac{\rho - a_{rr}}{d_r - a_{rr}} > 0.$$

Now

$$(\rho - a_{jj})x_j = \sum_{r \in S \setminus \{j\}} a_{jr}x_r + \sum_{s \notin S} a_{js}x_s.$$

Since  $d_j = \delta$ , this implies

$$(\rho - a_{jj})x_j \leq x_{\max} \sum_{r \in S \setminus \{j\}} \frac{a_{jr}(d_r - a_{rr})}{\rho - a_{rr}} + (\delta - \sum_{s \in S} a_{js})x_{\max}.$$

Putting  $d_r - a_{rr} = (d_r - \rho) + (\rho - a_{rr})$  and simplifying, we obtain

$$(\rho - a_{jj})x_{\min} \leq (\rho - a_{jj})x_j \leq x_{\max} \sum_{r \in S \setminus \{j\}} \frac{a_{jr}(d_r - \rho)}{\rho - a_{rr}} + (\delta - a_{jj})x_{\max}.$$

The previous inequality implies

$$\gamma = \frac{x_{\max}}{x_{\min}} \geq \frac{\rho - a_{jj}}{\delta - a_{jj} - \sum_{r \in S \setminus \{j\}} \frac{a_{jr}(\rho - d_r)}{\rho - a_{rr}}}. \quad \square$$

Note that if  $T = \{i\}$  and  $S = \{j\}$ , then the previous result is the same as Ostrowski's bound.

If  $G$  is a graph with  $\delta = \Delta - 1$  having exactly one vertex of degree  $\Delta - 1$ , then the previous result implies that

$$\gamma \geq \frac{\Delta + \Delta \cdot \frac{\Delta - \lambda_1(G)}{\lambda_1(G)}}{\lambda_1(G)} = \frac{\Delta^2}{\lambda_1^2(G)}$$

which improves Ostrowski's inequality

$$\gamma \geq \frac{\Delta}{\lambda_1(G)}.$$

**3. Extreme entries of principal eigenvectors.** In this section, we determine some upper bounds for the entries of  $x$  when  $G$  is an irregular graph.

Papendieck and Recht [11] obtained the following upper bound on  $x_{\max}$ . Zhao and Hong [16] generalized Papendieck and Recht's result to symmetric nonnegative matrices with zero trace.

**THEOREM 3.1.** *If  $G$  is a connected graph on  $n$  vertices, then*

$$(3.1) \quad x_{\max} \leq \frac{1}{\sqrt{2}}$$

with equality if and only if  $G = K_{1,n-1}$ .

The next result improves Theorem 3.1.

**THEOREM 3.2.** *Let  $G$  be a connected graph on  $n$  vertices whose principal eigenvector is  $x$ . For  $i \in [n]$ , if  $d_i$  is the degree of vertex  $i$ , then*

$$(3.2) \quad x_i \leq \frac{1}{\sqrt{1 + \frac{\lambda_1^2}{d_i}}}.$$

Equality is attained if and only if  $x_i = x_{\max}$  and  $G$  is the join of vertex  $i$  and a regular graph on  $n - 1$  vertices.

*Proof.* Using the Cauchy-Schwarz inequality, we have

$$\sum_{j \sim i} x_j^2 \geq \frac{(\sum_{j \sim i} x_j)^2}{d_i} = \frac{\lambda_1^2 x_i^2}{d_i}.$$

It follows that

$$1 = \sum_{l=1}^n x_l^2 \geq x_i^2 + \sum_{j \sim i} x_j^2 \geq x_i^2 \left( 1 + \frac{\lambda_1^2}{d_i} \right).$$

This proves the inequality.

Equality is attained if and only if  $x_j = a$  for  $j \sim i$  and  $x_l = 0$  for  $l$  not adjacent to  $i$ . Since  $G$  is connected, it follows that every vertex different from  $i$  is adjacent to  $i$ . For  $j \neq i$ , we have  $\lambda_1 x_j = x_i + (d_j - 1)x_j$ . This implies that the graph  $G \setminus \{i\}$  is regular.  $\square$

Theorem 3.1 follows now easily since  $\lambda_1 \geq \sqrt{\Delta} \geq \sqrt{d_1}$ .

An easy lower bound on  $x_{\max}$  can be obtained from the fact that

$$(n-1)x_{\max}^2 + x_{\min}^2 \geq \sum_{i=1}^n x_i^2 = 1.$$

This implies the following result.

LEMMA 3.3. *If  $G$  is a graph on  $n$  vertices with maximum degree  $\Delta$  and minimum degree  $\delta$ , then*

$$x_{\max} \geq \frac{1}{\sqrt{n-1 + \frac{1}{\gamma^2(G)}}} \geq \frac{1}{\sqrt{n - \frac{\Delta-\delta}{\Delta}}}$$

with equality if and only if  $G$  is regular. If  $G$  is irregular, then

$$x_{\max} > \frac{1}{\sqrt{n - \frac{1}{\Delta}}}.$$

*Proof.* From the previous inequality, we have

$$x_{\max}^2 \left( (n-1) + \frac{1}{\gamma^2(G)} \right) \geq 1.$$

Because  $\gamma(G) \geq \sqrt{\frac{\Delta}{\delta}}$ , we obtain

$$x_{\max} \geq \frac{1}{\sqrt{n - \frac{\Delta-\delta}{\Delta}}}$$

with equality if and only if  $G$  is regular.

If  $G$  is irregular, we have  $\Delta - \delta \geq 1$  which implies  $x_{\max} \geq \frac{1}{\sqrt{n - \frac{1}{\Delta}}}$ . The previous inequality is strict because otherwise, it would imply that  $n-1$  vertices of the graph have their eigenvector entry equal to  $x_{\max}$  and one vertex has its eigenvector entry equal to  $x_{\min}$ . It can be shown easily that this situation cannot happen when  $G$  is irregular.  $\square$

Note that the graph on  $n$  vertices obtained from  $K_n$  by deleting an edge has  $x_{\max} < \frac{1}{\sqrt{n-\frac{4}{n-1}}} = \frac{1}{\sqrt{n-\frac{4}{\Delta}}}$  so the second inequality from Lemma 3.3 comes very close to approximating  $x_{\max}$  in this case.

Next, we present another lower bound for  $x_1$  in terms of the spectral radius and the degree sequence of the graph.

**THEOREM 3.4.** *Let  $G$  be a connected graph with degrees  $d_1, \dots, d_n$ . Then*

$$x_{\max} \geq \frac{\lambda_1}{\sqrt{\sum_{i=1}^n d_i^2}} \geq \frac{1}{\sqrt{n}}$$

with equality iff  $G$  is regular.

*Proof.* For each  $i$ , we have

$$\lambda_1 x_i = \sum_{j \sim i} x_j \leq d_i x_{\max}.$$

Squaring and summing over all  $i$ , we obtain

$$\lambda_1^2 \leq x_{\max}^2 \sum_{i=1}^n d_i^2$$

which implies the inequality stated in the theorem.

Equality happens iff  $x_{\max} = x_j$  for each  $j$  which is equivalent to  $G$  being regular.  $\square$

One can also apply this inequality to powers of  $A$  since the eigenvectors of  $A^k$  are the same as the eigenvectors of  $A$ . Let  $w_l(G)$  denote the number of walks of length  $l$  in  $G$ .

**COROLLARY 3.5.** *If  $G$  is a graph on  $n$  vertices, then*

$$x_{\max} \geq \sqrt{\frac{\lambda_1^{2k}}{w_{2k}(G)}}$$

for each  $k \geq 1$ .

It follows from the results in [6] that the right hand-side of the previous inequality is monotonically increasing with  $k$ . If, in addition,  $G$  is connected and not bipartite, it actually tends to  $\frac{1}{\sum_{i=1}^n x_i}$  as  $k$  goes to infinity.

We have provided upper and lower bounds for the maximum entry of the principal eigenvector of a connected graph that are sharp in some cases. We conclude this section with the following result which gives an upper bound for the minimum entry of the principal eigenvector of a connected graph.

**THEOREM 3.6.** *If  $G$  is a graph on  $n$  vertices with maximum degree  $\Delta$  and  $e$  edges, then*

$$\left(\Delta - \frac{2e}{n}\right) x_{\min} \leq \frac{\Delta - \lambda_1}{\sqrt{n}} \leq \frac{\Delta - \frac{2e}{n}}{\sqrt{n}}.$$

Equality happens in the first inequality if and only if  $d_i = \Delta$  for each vertex  $i$  with  $x_i > x_{\min}$ . The second inequality is equality if and only if  $G$  is regular.

*Proof.* Since  $\lambda_1 \sum_{i=1}^n x_i = \sum_{i=1}^n d_i x_i$ , it follows that

$$(3.3) \quad (\Delta - \lambda_1) \sum_{i=1}^n x_i = \sum_{i=1}^n (\Delta - d_i) x_i \geq x_{\min} \sum_{i=1}^n (\Delta - d_i) = nx_{\min} \left( \Delta - \frac{2e}{n} \right).$$

Equality happens if and only if  $d_i = \Delta$  for each  $i$  with  $x_i > x_{\min}$ . Since  $\lambda_1 \geq \frac{2e}{n}$  with equality if and only if  $G$  is regular, this finishes the proof.  $\square$

Actually, using the inequality  $\sum_{i=1}^n x_i \leq x_{\min} + \sqrt{(n-1)(1-x_{\min}^2)}$  in (3.3), one can obtain the following better although more complicated inequality

$$x_{\min} \cdot \sqrt{(\Delta - \lambda_1)^2 + \frac{((n-1)\Delta + \lambda_1 - 2e)^2}{n-1}} \leq \Delta - \lambda_1.$$

Applying Theorem 3.6 and results from [3], we can show that if  $G$  is an irregular graph, then

$$x_{\min} \leq \frac{\Delta - \lambda_1}{\Delta - \frac{2e}{n}} \cdot \frac{1}{\sqrt{n}} < \left( 1 - \frac{1}{(\Delta + 2)(n\Delta - 2e)} \right) \frac{1}{\sqrt{n}}.$$

Note that while the maximum entry of the principal eigenvector can be as large as  $\frac{1}{\sqrt{2}}$ , the principal eigenvector can contain entries which are exponentially smaller than  $\frac{1}{\sqrt{n}}$  as shown by the graphs of the form  $P_r \cdot K_s$ .

**Acknowledgment.** We thank Vladimir Nikiforov for comments on an earlier version of this paper.

#### REFERENCES

- [1] G. Brightwell and P. Winkler. Maximum hitting time for random walks on graphs. *Random Structures and Algorithms*, 1:263-276, 1993.
- [2] S. Brin and L. Page. The anatomy of a large-scale hypertextual search engine. *Computer Networks and ISDN Systems*, 30:107-117, 1998.
- [3] S. M. Cioabă and D. A. Gregory. Large matchings from eigenvalues. *Linear Algebra and its Applications*, 422:308-317, 2007.
- [4] G. A. Latham. pp. 403-406 in: A remark on Minc's maximal eigenvector bound for positive matrices. *SIAM Journal on Matrix Analysis and Applications*, 16:307-311, 1995.
- [5] H. Minc. On the maximal eigenvector of a positive matrix. *SIAM Journal on Numerical Analysis*, 7:424-427, 1970.
- [6] V. Nikiforov. Walks and the spectral radius of graphs. *Linear Algebra and its Applications*, 418:257-268, 2006.
- [7] V. Nikiforov. Revisiting two classical results on graph spectra. *Electronic Journal of Combinatorics*, 14: Research Paper 14, 7 pp., 2007.
- [8] G. N. De Oliveira. On the characteristic vectors of a matrix. *Linear Algebra and its Applications*, 5:189-196, 1972.
- [9] A. M. Ostrowski. Bounds for the greatest latent root of a positive matrix. *The Journal of the London Mathematical Society* 27:253-256, 1952.
- [10] A. M. Ostrowski. On the eigenvector belonging to the maximal root of a non-negative matrix. *Proceedings of the Edinburgh Mathematical Society. Series II*, 12:107-112, 1960/1961.
- [11] B. Papendieck and P. Reicht. On maximal entries in the principal eigenvector of graphs. *Linear Algebra and its Applications*, 310:129-138, 2000.
- [12] H. Schneider. Note on the fundamental theorem on irreducible non-negative matrices. *Proceedings of the Edinburgh Mathematical Society*, 11:127-130, 1958/1959.

- [13] J. H. Smith. Some properties of the spectrum of a graph. *Combinatorial Structures and their applications (Proc. Conf. Calgary, 1969)*, pp. 403-406, Gordon and Breach, New York, 1970.
- [14] D. Spielman and S. H. Teng. Spectral partitioning works: Planar graphs and finite element meshes. *Linear Algebra and its Applications*, 421:284-305, 2007.
- [15] D. B. West. *Introduction to graph theory*, Edition 2. Prentice-Hall, Upper Saddle River, NJ, 2001.
- [16] S. Zhao and Y. Hong. On the bounds of maximal entries in the principal eigenvector of symmetric nonnegative matrix. *Linear Algebra and its Applications*, 340:245-252, 2002.
- [17] X.-D. Zhang. Eigenvectors and eigenvalues of non-regular graphs. *Linear Algebra and its Applications*, 49:79-86, 2005.