Electronic Transactions on Numerical Analysis.
Volume 9, 1999, pp. 128-136.

ISSN 1068-9613.

ETNA
Kent State University
etna@mcs.kent.edu

AN ITERATIVE METHOD FOR COMPUTING THE EIGENVALUES OF SECOND KIND FREDHOLM OPERATORS AND APPLICATIONS*

P. NATALINI†, S. NOSCHESE‡, AND P.E. RICCI§

Abstract. An inverse iteration method is used to approximate the eigenvalues of second kind Fredholm operators. Numerical examples are given.

Key words. Fredholm Integral equation, eigenvalues, inverse iteration method, Rayleigh-Ritz method.

AMS subject classifications. 45C05, 45B05, 65R20.

*Received November 1, 1998. Accepted for publicaton December 1, 1999. Recommended by F. Marcellán.

†Università degli Studi Roma III, Facoltà di Ingegneria, Largo San Leonardo Murialdo, 1, 00146 – Roma. (natalini@matrm3.mat.uniroma3.it).

‡Dipartimento di Matematica "Guido CASTELNUOVO", Università degli Studi di Roma "La Sapienza", P.le A. Moro, 2, 00185 – Roma. (noschese@axrma.uniroma1.it).

§Dipartimento di Matematica "Guido CASTELNUOVO", Università degli Studi di Roma "La Sapienza", P.le A. Moro, 2, 00185 – Roma. (riccip@axrma.uniroma1.it), Fax: (396) - 44701007.