

**ON THE CORRECT FORMULATION OF A
MULTIDIMENSIONAL PROBLEM FOR STRICTLY
HYPERBOLIC EQUATIONS OF HIGHER ORDER**

S. KHARIBEGASHVILI

ABSTRACT. A theorem of the unique solvability of the first boundary value problem in the Sobolev weighted spaces is proved for higher-order strictly hyperbolic systems in the conic domain with special orientation.

In the space R^n , $n > 2$, let us consider a strictly hyperbolic equation of the form

$$p(x, \partial)u(x) = f(x), \quad (1)$$

where $\partial = (\partial_1, \dots, \partial_n)$, $\partial_j = \partial/\partial x_j$, $p(x, \xi)$ is a real polynomial of order $2m$, $m > 1$, with respect to $\xi = (\xi_1, \dots, \xi_n)$, f is the known function and u is the unknown function. It is assumed that in equation (1) the coefficients at higher derivatives are constant and the other coefficients are finite and infinitely differentiable in R^n .

Let D be a conic domain in R^n , i.e., D together with a point $x \in D$ contains the entire beam tx , $0 < t < \infty$. Denote by Γ the cone ∂D . It is assumed that D is homeomorphic onto the conic domain $x_1^2 + \dots + x_{n-1}^2 - x_n^2 < 0$, $x_n > 0$ and $\Gamma' = \Gamma \setminus O$ is a connected $(n-1)$ -dimensional manifold of the class C^∞ , where O is the vertex of the cone Γ .

Consider the problem: Find in the domain D the solution $u(x)$ of equation (1) by the boundary conditions

$$\frac{\partial^i u}{\partial \nu^i} \Big|_{\Gamma'} = g_i, \quad i = 0, \dots, m-1, \quad (2)$$

where $\nu = \nu(x)$ is the outward normal to Γ' at a point $x \in \Gamma'$, and g_i , $i = 0, \dots, m-1$, are the known real functions.

Note that the problem (1), (2) is considered in [1-6] for a hyperbolic-type equation of second order when Γ is a characteristic conoid. In [7] this

1991 *Mathematics Subject Classification.* 35L35.

problem is considered for a wave equation when the conic surface Γ is not characteristic at any point and has a time-type orientation. A multidimensional analogue of the problem is treated in [8–10] for the case when one part of the cone Γ is characteristic and the other part is a time-type hyperplane. Other multidimensional analogues of the Goursat problem for hyperbolic systems of first and second order are investigated in [11–15].

In this paper we consider the question whether the problem (1), (2) can be correctly formulated in special weighted spaces $W_\alpha^k(D)$ when the cone Γ is assumed not to be characteristic but having a quite definite orientation.

Denote by $p_0(\xi)$ the characteristic polynomial of the equation (1), i.e., the higher homogeneous part of the polynomial $p(x, \xi)$. The strict hyperbolicity of the equation (1) implies the existence of a vector $\zeta \in R^n$ such that the straight line $\xi = \lambda\zeta + \eta$, where $\eta \in R^n$ is an arbitrarily chosen vector not parallel to ζ and λ is the real parameter, intersects the cone of normals $K : p_0(\xi) = 0$ of the equation (1) at $2m$ different real points. In other words, the equation $p_0(\lambda\zeta + \eta) = 0$ with respect to λ has $2m$ different real roots. The vector ζ is called a spatial-type normal. As is well-known, a set of all spatial-type normals form two connected centrally-symmetric convex conic domains whose boundaries K_1 and K_{2m} give the internal cavity of the cone of normals K [3]. The surface $S \subset R^n$ is called characteristic at a point $x \in S$ if the normal to S at the point x belongs to the cone K .

Let the vector ζ be a spatial-type normal and the vector $\eta \neq 0$ change in the plane orthogonal to ζ . Then for λ the roots of the characteristic polynomial $p_0(\lambda\zeta + \eta)$ can be reenumerated so that $\lambda_{2m}(\eta) < \lambda_{2m-1}(\eta) < \dots < \lambda_1(\eta)$. It is obvious that the vectors $\lambda_i(\eta)\zeta + \eta$ cover the cavities K_i of K when the η changes on the plane orthogonal to ζ . Since $\lambda_{m-j}(\eta) = -\lambda_{m+j+1}(-\eta)$, $0 \leq j \leq m-1$, the cones K_{m-j} and K_{m+j+1} are centrally symmetric with respect to the point $(0, \dots, 0)$. As is well-known, by the bicharacteristics of the equation (1) we understand straight beams whose orthogonal planes are tangential planes to one of the cavities K_i at the point different from the vertex.

Assume that there exists a plane π_0 such that $\pi_0 \cap K_m = \{(0, \dots, 0)\}$. This means that the cones K_1, \dots, K_m are located on one side of π_0 and the cones K_{m+1}, \dots, K_{2m} on the other. Set $K_i^* = \cap_{\eta \in K_i} \{\xi \in R^n : \xi \cdot \eta < 0\}$, where $\xi \cdot \eta$ is the scalar product of ξ and η . Since $\pi_0 \cap K_m = \{(0, \dots, 0)\}$, K_i^* is a conic domain and $K_m^* \subset K_{m-1}^* \subset \dots \subset K_1^*$, $K_{m+1}^* \subset K_{m+2}^* \subset \dots \subset K_{2m}^*$. It is easy to verify that $\partial(K_i^*)$ is a convex cone whose generatrices are bicharacteristics; note that in this case none of the bicharacteristics of the equation (1) comes from the point $(0, \dots, 0)$ into the cone $\partial(K_m^*)$ or $\partial(K_{m+1}^*)$ [3].

Let us consider

Condition 1. The surface Γ' is characteristic at none of its points and

each generatrix of the cone Γ has the direction of a spatial-type normal; moreover, $\Gamma \subset K_m^* \cup 0$ or $\Gamma \subset K_{m+1}^* \cup 0$.

Denote by $W_\alpha^k(D)$, $k \geq 2m$, $-\infty < \alpha < \infty$, the functional space with the norm [16]

$$\|u\|_{W_\alpha^k(D)}^2 = \sum_{i=0}^k \int_D r^{-2\alpha-2(k-i)} \left\| \frac{\partial^i u}{\partial x^i} \right\|^2 dx,$$

where

$$r = (x_1^2 + \dots + x_n^2)^{\frac{1}{2}}, \quad \frac{\partial^i u}{\partial x^i} = \frac{\partial^i u}{\partial x_1^{i_1} \dots \partial x_n^{i_n}}, \quad i = i_1 + \dots + i_n.$$

The space $W_\alpha^k(\Gamma)$ is defined in a similar manner. Consider the space

$$V = W_{\alpha-1}^{k+1-2m}(D) \times \prod_{i=0}^{m-1} W_{\alpha-\frac{1}{2}}^{k-i}(\Gamma).$$

Assume that to the problem (1), (2) there corresponds the unbounded operator

$$T : W_\alpha^k(D) \rightarrow V$$

with the domain of definition $\Omega_T = W_{\alpha-1}^{k+1}(D) \subset W_\alpha^k(D)$, acting by the formula

$$Tu = \left(p(x, \partial)u, u \Big|_{\Gamma'}, \dots, \frac{\partial^i u}{\partial \nu^i} \Big|_{\Gamma'}, \dots, \frac{\partial^{m-1} u}{\partial \nu^{m-1}} \Big|_{\Gamma'} \right), \quad u \in \Omega_T.$$

It is obvious that the operator T admits the closure \bar{T} .

The function u is called a strong solution of the problem (1), (2) of the class $W_\alpha^k(D)$ if $u \in \Omega_{\bar{T}}$, $\bar{T}u = (f, g_0, \dots, g_{m-1}) \in V$, which is equivalent to the existence of a sequence $u_i \in \Omega_T = W_{\alpha-1}^{k+1}(D)$ such that $u_i \rightarrow u$ in $W_\alpha^k(D)$ and $(p(x, \partial)u_i, u_i \Big|_{\Gamma'}, \dots, \frac{\partial^{m-1} u_i}{\partial \nu^{m-1}} \Big|_{\Gamma'}) \rightarrow (f, g_0, \dots, g_{m-1})$ in V .

Below, by a solution of the problem (1), (2) of the class $W_\alpha^k(D)$ we will mean a strong solution of this problem in the sense as indicated above.

We will prove

Theorem. *Let condition 1 be fulfilled. Then there exists a real number $\alpha_0 = \alpha_0(k) > 0$ such that for $\alpha \geq \alpha_0$ the problem (1), (2) is uniquely solvable in the class $W_\alpha^k(D)$ for any $f \in W_{\alpha-1}^{k+1-2m}(D)$, $g_i \in W_{\alpha-\frac{1}{2}}^{k-i}(\Gamma)$, $i = 0, \dots, m-1$, and to obtain the solution u we have the estimate*

$$\|u\|_{W_\alpha^k(D)} \leq c \left(\sum_{i=1}^{m-1} \|g_i\|_{W_{\alpha-\frac{1}{2}}^{k-i}(\Gamma)} + \|f\|_{W_{\alpha-1}^{k+1-2m}(D)} \right), \quad (3)$$

where c is a positive constant not depending on f , g_i , $i = 0, \dots, m - 1$.

Proof. First we will show that the corollaries of condition 1 are the following conditions: Take any point $P \in \Gamma'$ and choose a Cartesian system x_1^0, \dots, x_n^0 connected with this point and having vertex at P such that the x_n^0 -axis is directed along the generatrix of Γ passing through P and the x_{n-1}^0 -axis is directed along the inward normal to Γ at this point.

Condition 2. The surface Γ' is characteristic at none of its points. Each generatrix of the cone Γ has the direction of a spatial-type normal, and exactly m characteristic planes of equation (1) pass through the $(n - 2)$ -dimensional plane $x_n^0 = x_{n-1}^0 = 0$ connected with an arbitrary point $P \in \Gamma'$ into the angle $x_n^0 > 0$, $x_{n-1}^0 > 0$.

Denote by $\tilde{p}_0(\xi)$ the characteristic polynomial of the equation (1) written in terms of the coordinate system x_1^0, \dots, x_n^0 , connected with an arbitrarily chosen point $P \in \Gamma'$.

Condition 3. The surface Γ' is characteristic at none of its point. Each generatrix of the cone Γ has the direction of a spatial-type normal and for $\text{Re } s > 0$ the number of roots $\lambda_j(\xi_1, \dots, \xi_{n-2}, s)$, if we take into account the multiplicity of the polynomial $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s)$ with $\text{Re } \lambda_j < 0$, is equal to m , $i = \sqrt{-1}$.

When condition 3 is fulfilled, the polynomial $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s)$ can be written as the product $\Delta_-(\lambda)\Delta_+(\lambda)$, where for $\text{Re } s > 0$ the roots of the polynomials $\Delta_-(\lambda)$ and $\Delta_+(\lambda)$ lie, respectively, to the left and to the right of the imaginary axis, while the coefficients are continuous for s , $\text{Re } s \geq 0$, $(\xi_1, \dots, \xi_{n-2}) \in R^{n-2}$, $\xi_1^2 + \dots + \xi_{n-2}^2 + |s|^2 = 1$ [17]. On the left side of the boundary conditions (2) to the differential operator $b_j(x, \partial)$, $0 \leq j \leq m - 1$, written in terms of the coordinate system x_1^0, \dots, x_n^0 connected with the point $P \in \Gamma'$, there corresponds the characteristic polynomial $b_j(\xi) = \xi_{n-1}^j$. Therefore, since the degree of the polynomial $\Delta_-(\lambda)$ is equal to m , the following condition will be fulfilled:

Condition 4. For any point $P \in \Gamma'$ and any s , $\text{Re } s \geq 0$ and $(\xi_1, \dots, \xi_{n-2}) \in R^{n-2}$ such that $\xi_1^2 + \dots + \xi_{n-2}^2 + |s|^2 = 1$, the polynomials $b_j(i\xi_1, \dots, i\xi_{n-2}, \lambda, s) = \lambda^j$, $j = 0, \dots, m - 1$, are linearly independent, like the polynomials of λ modulo $\Delta_-(\lambda)$.

We will now show that condition 1 implies condition 2, while the latter implies condition 3. Let us consider the case $\Gamma \subset K_{m+1}^* \cup O$. The second case $\Gamma \subset K_m^* \cup O$ is treated similarly.

Let $P \in \Gamma'$ and x_1^0, \dots, x_n^0 be the coordinate system connected with this point. Since the generatrix γ of the cone Γ passing through this point is a spatial-type normal, the plane $x_n^0 = 0$ passing through the point P is

a spatial-type plane. Denote by K_j^\wedge the boundary of the convex shell of the set K_j and by K_j^\perp the set which is the union of all bicharacteristics corresponding to the cone K_j and coming out of the point O along the outward normal to K_j , $1 \leq j \leq 2m$. It is obvious that $(K_j^\wedge)^* = K_j^*$, $\partial(K_j^*) = (K_j^\wedge)^\perp$. We will show that the plane π_1 , parallel to the plane $x_n^0 = 0$ and passing through the point $(0, \dots, 0)$, is the plane of support to the cone K_m^\wedge at the point $(0, \dots, 0)$. Indeed, it is obvious that the plane $N \cdot \xi = 0$, $N \in R^n \setminus (0, \dots, 0)$, $\xi \in R^n$ is the plane of support to K_m^\wedge at the point $(0, \dots, 0)$ iff the normal vector N to this plane taken with the sign $+$ or $-$ belongs to the conic domain closure $(K_m^\wedge)^* = K_m^*$. Now it remains for us to note that the conic domains K_m^* and K_{m+1}^* are centrally symmetric with respect to the point $(0, \dots, 0)$, and the generatrix Γ passing through the point P is perpendicular to the plane π_1 and, by the condition, belongs to the set $K_{m+1}^* \cup O$. Since $x_n^0 = 0$ is a spatial-type plane, the two-dimensional plane $\sigma : x_1^0 = \dots = x_{n-2}^0 = 0$ passing through the generatrix γ which is directed along the spatial-type normal intersects the cone of normals K_p of equation (1) with vertex at the point P by $2m$ different real straight lines [3]. The planes orthogonal to these straight lines and passing through the $(n-2)$ -dimensional plane $x_n^0 = x_{n-1}^0 = 0$ give all $2m$ characteric planes passing through the $(n-2)$ -dimensional plane $x_n^0 = x_{n-1}^0 = 0$. The straight lines $x_n^0 = 0$ and $x_{n-1}^0 = 0$ divide the two-dimensional plane σ into four right angles

$$\begin{aligned} \sigma_1 : x_{n-1}^0 > 0, x_n^0 > 0; \quad \sigma_2 : x_{n-1}^0 < 0, x_n^0 > 0; \\ \sigma_3 : x_{n-1}^0 < 0, x_n^0 < 0; \quad \sigma_4 : x_{n-1}^0 > 0, x_n^0 < 0. \end{aligned}$$

One can readily see that exactly m characteristic planes of equation (1) pass through the $(n-2)$ -dimensional plane $x_n^0 = x_{n-1}^0 = 0$ into the angle $x_n^0 > 0$, x_{n-1}^0 iff exactly m straight lines from the intersection of σ_4 with the two-dimensional plane σ pass into the angle K_P . The latter fact really occurs, since: 1) the plane $x_n^0 = 0$ is the plane of support to K_m^\wedge and therefore to all K_1, \dots, K_{2m} ; 2) the planes $x_n^0 = 0$, $x_{n-1}^0 = 0$ are not characteristic because the generatrices of Γ have a spatial-type direction and Γ is not characteristic at the point P .

Now it will be shown that condition 2 implies condition 3. By virtue of condition 2 the plane $x_{n-1}^0 = 0$ is not characteristic and therefore for λ the polynomial $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s)$ has exactly $2m$ roots. In this case, if $\text{Re } s > 0$, the number of roots $\lambda_j(\xi_1, \dots, \xi_{n-2}, s)$, with the multiplicity of the polynomial $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s)$ taken into account, will be equal to m provided that $\text{Re } \lambda_j < 0$. Indeed, recalling that equation (1) is hyperbolic, the equation $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s) = 0$ has no purely imaginary roots with respect to λ . Since the roots λ_j are continuous functions of s , we can determine the number of roots λ_j with $\text{Re } \lambda_j < 0$ by passing to the limits

as $\operatorname{Re} s \rightarrow +\infty$. Since the equality

$$\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}\lambda, s) = s^{2m}\tilde{p}_0\left(i\frac{\xi_1}{s}, \dots, i\frac{\xi_{n-2}}{s}, \frac{\lambda}{s}, 1\right)$$

holds, it is clear that the ratios λ_j/s , where λ_j are the roots of the equation $\tilde{p}_0(i\xi_1, \dots, i\xi_{n-2}, \lambda, s) = 0$, tend to the roots μ_j of the equation $\tilde{p}_0(0, \dots, 0, \mu, 1) = 0$ as $\operatorname{Re} s \rightarrow +\infty$. The latter roots are real and different because equation (1) is hyperbolic. If s is taken positive and sufficiently large, then for $\mu_j \neq 0$ we have $\lambda_j = s\mu_j + o(s)$. But $\mu_j \neq 0$, since the plane $x_n^0 = 0$ is not characteristic. Therefore the number of roots λ_j with $\operatorname{Re} \lambda_j < 0$ coincides with the number of roots μ_j with $\mu_j < 0$. Since the characteristic planes of equation (1), passing through the $(n-2)$ -dimensional plane $x_n^0 = x_{n-1}^0 = 0$, are determined by the equalities $\mu_j x_{n-1}^0 + x_n^0 = 0$, $j = 1, \dots, 2m$, condition 2 implies that for $\operatorname{Re} \lambda_j < 0$ the number of roots λ_j is equal to m .

We give another equivalent description of the space $W_\alpha^k(D)$. On the unit sphere $S^{n-1} : x_1^2 + \dots + x_n^2 = 1$ choose a coordinate system $(\omega_1, \dots, \omega_{n-1})$ such that in the domain D the transformation

$$I : \tau = \log r, \quad \omega_j = \omega_j(x_1, \dots, x_n), \quad j = 1, \dots, n-1,$$

is one-to-one, nondegenerate, and infinitely differentiable. Since the cone $\Gamma = \partial D$ is strictly convex at the point $O(0, \dots, 0)$, such coordinates evidently exist. As a result of the above transformation, the domain D will become the infinite cylinder G bounded by the infinitely differentiable surface $\partial G = I(\Gamma')$.

Introduce the functional space $H_\gamma^k(G)$, $-\infty < \gamma < \infty$, with the norm

$$\|v\|_{H_\gamma^k(G)}^2 = \sum_{i_1+j=0}^k \int_G e^{-2\gamma\tau} \left\| \frac{\partial^{i_1+j} v}{\partial \tau^{i_1} \partial \omega^j} \right\|^2 d\omega d\tau$$

where

$$\frac{\partial^{i_1+j} v}{\partial \tau^{i_1} \partial \omega^j} = \frac{\partial^{i_1+j} v}{\partial \tau^{i_1} \partial \omega_1^{j_1} \dots \partial \omega_{n-1}^{j_{n-1}}}, \quad j = j_1 + \dots + j_{n-1}.$$

As shown in [16], a function $u(x) \in W_\alpha^k(D)$ iff $\tilde{u} = u(I^{-1}(\tau, \omega)) \in H_{(\alpha+k)-\frac{n}{2}}^k(G)$, and the estimates

$$c_1 \|\tilde{u}\|_{H_{(\alpha+k)-\frac{n}{2}}^k(G)} \leq \|u\|_{W_\alpha^k(D)} \leq c_2 \|\tilde{u}\|_{H_{(\alpha+k)-\frac{n}{2}}^k(G)}$$

hold, where I^{-1} is the inverse transformation of I and the positive constants c_1 and c_2 do not depend on u .

It can be easily verified that the condition $v \in H_\gamma^k(G)$ is equivalent to the condition $e^{-\gamma\tau} v \in W^k(G)$, where $W^k(G)$ is the Sobolev space. Denote

by $H_\gamma^k(\partial G)$ a set of ψ such that $e^{-\gamma\tau}\psi \in W^k(\partial G)$, and by $W_{\alpha-\frac{1}{2}}^k(\Gamma)$ a set of all φ for which $\tilde{\varphi} = \varphi(I^{-1}(\tau, \omega)) \in H_{(\alpha+k)-\frac{n}{2}}^k(\partial G)$. Assume that

$$\|\varphi\|_{W_{\alpha-\frac{1}{2}}^k(\Gamma)} = \|\tilde{\varphi}\|_{H_{(\alpha+k)-\frac{n}{2}}^k(\partial G)}.$$

Spaces $W_\alpha^k(D)$ possess the following simple properties:

- 1) if $u \in W_\alpha^k(D)$, then $\frac{\partial^i u}{\partial x^i} \in W_\alpha^{k-i}(D)$, $0 \leq i \leq k$;
- 2) $W_{\alpha-1}^{k+1}(D) \subset W_\alpha^k(D)$;
- 3) if $u \in W_{\alpha-1}^{k+1}(D)$, then by the well-known embedding theorems we have $u|_\Gamma \in W_{\alpha-\frac{1}{2}}^k(\Gamma)$, $\frac{\partial^i u}{\partial \nu^i}|_{\Gamma'} \in W_{\alpha-\frac{1}{2}}^{k-i}(\Gamma)$, $i = 1, \dots, m-1$;
- 4) if $u \in W_{\alpha-1}^{k+1}(D)$, then $f = p(x, \partial)u \in W_{\alpha-1}^{k+1-2m}(D)$.

In what follows we will need, in spaces $W_\alpha^k(D)$, $W_{\alpha-\frac{1}{2}}^k(\Gamma)$, other norms depending on the parameter $\gamma = (\alpha + k) - \frac{n}{2}$ and equivalent to the original norms.

Set

$$\begin{aligned} R_{\omega, \tau}^n &= \{-\infty < \tau < \infty, -\infty < \omega_i < \infty, i = 1, \dots, n-1\}, \\ R_{\omega, \tau, +}^n &= \{(\omega, \tau) \in R_{\omega, \tau}^n : \omega_{n-1} > 0\}, \quad \omega' = (\omega_1, \dots, \omega_{n-2}), \\ R_{\omega', \tau}^{n-1} &= \{-\infty < \tau < \infty, -\infty < \omega_i < \infty, i = 1, \dots, n-2\}. \end{aligned}$$

Denote by $\tilde{v}(\xi_1, \dots, \xi_{n-2}, \xi_{n-1}, \xi_n - i\gamma)$ the Fourier transform of the function $e^{-\gamma\tau}v(\omega, \tau)$, i.e.,

$$\begin{aligned} \tilde{v}(\xi_1, \dots, \xi_{n-1}, \xi_n - i\gamma) &= (2\pi)^{-\frac{n}{2}} \int v(\omega, \tau) e^{-i\omega\xi' - i\tau\xi_n - \gamma\tau} d\omega d\tau, \\ i &= \sqrt{-1}, \quad \xi' = (\xi_1, \dots, \xi_{n-1}), \end{aligned}$$

and by $\hat{v}(\xi, \dots, \xi_{n-2}, \omega_{n-1}, \xi_n - i\gamma)$ the partial Fourier transform of the function $e^{-\gamma\tau}v(\omega, \tau)$ with respect to ω', τ .

We can introduce the following equivalent norms:

$$\begin{aligned} \|v\|_{R^n, k, \gamma}^2 &= \int_{R^n} (\gamma^2 + |\xi|^2)^k \|\tilde{v}(\xi_1, \dots, \xi_{n-1}, \xi_n - i\gamma)\|^2 d\xi, \\ \|v\|_{R_+^n, k, \gamma}^2 &= \int_0^\infty \int_{R^{n-1}} \sum_{j=0}^k (\gamma^2 + |\xi'|^2)^{k-j} \times \\ &\times \left\| \frac{\partial^j}{\partial \omega_{n-1}^j} \hat{v}(\xi_1, \dots, \xi_{n-2}, \omega_{n-1}, \xi_n - i\gamma) \right\|^2 d\xi' d\omega_{n-1}, \end{aligned}$$

in the above-considered spaces $H_\gamma^k(R_{\omega, \tau}^n)$ and $H_\gamma^k(R_{\omega, \tau, +}^n)$.

Let $\varphi_1, \dots, \varphi_N$ be the partitioning of unity into $G' = G \cap \{\tau = 0\}$, where $G = I(D)$, i.e., $\sum_{j=1}^N \varphi_j(\omega) \equiv 1$ in G' , $\varphi_j \in C^\infty(\overline{G}')$, the supports of

functions $\varphi_1, \dots, \varphi_{N-1}$ lie in the boundary half-neighborhoods, while the support of function φ_N lies inside G' . Then for $\gamma = (\alpha + k) - \frac{n}{2}$ the equalities

$$\begin{aligned} \|u\|_{G,k,\gamma}^2 &= \sum_{j=1}^{N-1} \|\varphi_j u\|_{R_{+,k,\gamma}^n}^2 + \|\varphi_N u\|_{R^{n,k,\gamma}}^2, \\ \|u\|_{\partial G,k,\gamma}^2 &= \sum_{j=1}^{N-1} \|\varphi_j u\|_{R_{\omega',\tau,k,\gamma}^{n-1}}^2 \end{aligned} \quad (4)$$

define equivalent norms in the spaces $W_\alpha^k(D)$ and $W_{\alpha-\frac{1}{2}}^k(\Gamma)$, where the norms on the right sides of these equalities are taken in the terms of local coordinates [17].

First we assume that equation (1) contains only higher terms, i.e., $p(x, \xi) \equiv p_0(\xi)$. Equation (1) and the boundary conditions (2) written in terms of the coordinates ω, τ have the form

$$\begin{aligned} e^{-2m\tau} A(\omega, \partial)u &= f, \\ e^{-i\tau} B_i(\omega, \partial)u \Big|_{\partial G} &= g_i, \quad i = 0, \dots, m-1, \end{aligned}$$

or

$$\begin{aligned} A(\omega, \partial)u &= \tilde{f}, \\ B_i(\omega, \partial)u \Big|_{\partial G} &= \tilde{g}_i, \quad i = 0, \dots, m-1, \end{aligned} \quad (5)$$

where $A(\omega, \partial)$ and $B_i(\omega, \partial)$ are, respectively, the differential operators of orders $2m$ and i , with infinitely differentiable coefficients depending only on ω , while $\tilde{f} = e^{2m\tau} f$ and $\tilde{g}_i = e^{i\tau} g_i$, $i = 0, 1, \dots, m-1$.

Thus, for the transformation $I : D \rightarrow G$, the unbounded operator T of the problem (1), (2) transforms to the unbounded operator

$$\tilde{T} : H_\gamma^k(G) \rightarrow H_\gamma^{k+1-2m}(G) \times \prod_{i=0}^{m-1} H_\gamma^{k-i}(\partial G)$$

with the domain of definition $H_\gamma^{k+1}(G)$, acting by the formula

$$\tilde{T}u = (A(\omega, \partial)u, B_0(\omega, \partial)u \Big|_{\partial G}, \dots, B_{m-1}(\omega, \partial)u \Big|_{\partial G})$$

where $\gamma = (\alpha + k) - \frac{n}{2}$. Note that written in terms of the coordinates ω, τ the functions $f = (\omega, \tau) \in H_{\gamma-2m}^{k+1-2m}(G)$, $g_i(\omega, \tau) \in H_{\gamma-i}^{k-i}(\partial G)$, $i = 0, \dots, m-1$, if $f(x) \in W_{\alpha-1}^{k+1-2m}(D)$, $g_i(x) \in W_{\alpha-\frac{1}{2}}^{k-i}(\Gamma)$, $i = 0, \dots, m-1$.

Therefore the functions $\tilde{f} = e^{2m\tau} f \in H_\gamma^{k+1-2m}(G)$, $\tilde{g}_i = e^{i\tau} g_i \in H_\gamma^{k-i}(\partial G)$, $i = 0, \dots, m-1$.

Since by condition 1 each generatrix of the cone Γ has the direction of a spatial-type normal, due to the convexity of K_m each beam coming from the vertex O into the conic domain D also has the direction of a spatial-type normal. Therefore equation (4) is strictly hyperbolic with respect the τ -axis. It was shown above that the fulfillment of condition 1 implies the fulfillment of condition 4. Therefore, according to the results of [17], for $\gamma \geq \gamma_0$, where γ_0 is a sufficiently large number, the operator \widetilde{T} has the bounded right inverse operator \widetilde{T}^{-1} . Thus for any $\widetilde{f} \in H_\gamma^{k+1-2m}(G)$, $\widetilde{g}_i \in H_\gamma^{k-i}(\partial G)$, $i = 0, \dots, m-1$, when $\gamma \geq \gamma_0$, the problem (5), (6) is uniquely solvable in the space $H_\gamma^k(G)$, and for the solution u we have the estimate

$$\| \| u \| \|_{G,k,\gamma}^2 \leq C \left(\sum_{i=0}^{m-1} \| \| \widetilde{g}_i \| \|_{\partial G,k-i,\gamma} + \frac{1}{\gamma} \| \| \widetilde{f} \| \|_{G,k+1-2m,\gamma} \right) \quad (7)$$

with the positive constant C not depending on γ , f and \widetilde{g}_i , $i = 0, \dots, m-1$.

Hence it immediately follows that the theorem and the estimate (3) are valid in the case $p(x, \xi) \equiv p_0(\xi)$. \square

Remark. The estimate (7) with the coefficient $\frac{1}{\gamma}$ at $\| \| \widetilde{f} \| \|_{G,k+1-2m,\gamma}$, obtained in the appropriately chosen norms (4), enables one to prove the theorem also when equation (1) contains lower terms, since the latter give arbitrarily small perturbations for sufficiently large γ .

REFERENCES

1. A.V.Bitsadze, Some classes of partial differential equations. (Russian) *Nauka, Moscow*, 1981.
2. S.L.Sobolev, Some applications of functional analysis in mathematical physics. (Russian) *Publ. Sib. Otd. Akad. Nauk SSSR, Novosibirsk*, 1962.
3. R.Courant, Partial differential equations. *New York-London*, 1962.
4. M.Riesz, L'integrale de Riemann-Liouville et le problem de Cauchy. *Acta Math.* **81**(1949), 107-125.
5. L.Lundberg, The Klein-Gordon equation with light-cone data. *Commun. Math. Phys.* **62**(1978), No. 2, 107-118.
6. A.A.Borgardt and D.A.Karnenko, The characteristic problem for the wave equation with mass. (Russian) *Differentsial'nye Uravneniya* **20**(1984), No. 2, 302-308.
7. S.L.Sobolev, Some new problems of the theory of partial differential equations of hyperbolic type. (Russian) *Mat. Sb.* **11**(53)(1942), No. 3, 155-200.

8. A.V.Bitsadze, On mixed type equations on three-dimensional domains. (Russian) *Dokl. Akad. Nauk SSSR* **143**(1962), No. 5, 1017-1019.
9. A.M.Nakhushev, A multidimensional analogy of the Darboux problem for hyperbolic equations. (Russian) *Dokl. Akad. Nauk SSSR* **194**(1970), No. 1, 31-34.
10. T.Sh.Kalmenov, On multidimensional regular boundary value problems for the wave equation. (Russian) *Izv. Akad. Nauk Kazakh. SSR. Ser. Fiz.-Mat.* (1982), No. 3, 18-25.
11. A.A.Dezin, Invariant hyperbolic systems and the Goursat problem. (Russian) *Dokl. Akad. Nauk SSSR* **135**(1960), No. 5, 1042-1045.
12. F.Cagnac, Probleme de Cauchy sur la conoide caracteristique. *Ann. Mat. Pure Appl.* **104**(1975), 355-393.
13. J.Tolen, Problème de Cauchy sur la deux hypersurfaces caracteristiques sécantes. *C.R. Acad. Sci. Paris Sér. A-B* **291**(1980), No. 1, A49-A52.
14. S.S.Kharibegashvili, The Goursat problems for some class of hyperbolic systems. (Russian) *Differentsial'nye Uravneniya* **17**(1981), No. 1. 157-164.
15. —, On a multidimensional problem of Goursat type for second order strictly hyperbolic systems. (Russian) *Bull. Acad. Sci. Georgian SSR* **117**(1985), No. 1, 37-40.
16. V.A.Kondratyev, Boundary value problems for elliptic equations in domains with conic or corner points. (Russian) *Trudy Moskov. Mat. Obshch.* **16**(1967), 209-292.
17. M.S.Agranovich, Boundary value problems for systems with a parameter. (Russian) *Mat. Sb.* **84**(126)(1971), No. 1, 27-65.

(Received 25.12.1992)

Author's address:
I.Vekua Institute of Applied Mathematics
of Tbilisi State University
2 University St., 380043 Tbilisi
Republic of Georgia