

**LITTLEWOOD–PALEY OPERATORS ON THE
GENERALIZED LIPSCHITZ SPACES**

SHANZHEN LU, CHANGMEI TAN, AND KOZO YABUTA

ABSTRACT. Littlewood–Paley operators defined on a new kind of generalized Lipschitz spaces $\mathcal{E}_0^{\alpha,p}$ are studied. It is proved that the image of a function under the action of these operators is either equal to infinity almost everywhere or is in $\mathcal{E}_0^{\alpha,p}$, where $-n < \alpha < 1$ and $1 < p < \infty$.

1. INTRODUCTION

For $x \in \mathbb{R}^n$, $y > 0$, the Poisson kernel is $P(x, y) = c_n y(y^2 + |x|^2)^{-(n+1)/2}$. Denote the Poisson integral of f by

$$f(x, y) = \int_{\mathbb{R}^n} f(z)P(x - z, y) dz.$$

We have (see [1])

$$|\nabla f(x, y)| \leq c_n \int_{\mathbb{R}^n} |f(z)| (y + |x - z|)^{-(n+1)} dz. \tag{1}$$

Let us now consider the following two kinds of Littlewood–Paley functions:

$$S(f)(x) = \left(\iint_{\Gamma(x)} y^{1-n} |\nabla f(z, y)|^2 dz dy \right)^{1/2}$$

and

$$g_\lambda^*(f)(x) = \left\{ \iint_{\mathbb{R}_+^{n+1}} \left(\frac{y}{y + |x - z|} \right)^{\lambda n} y^{1-n} |\nabla f(z, y)|^2 dz dy \right\}^{1/2}.$$

1991 *Mathematics Subject Classification.* 42B25.

Key words and phrases. Littlewood–Paley functions, generalized Lipschitz spaces, Poisson integral, Poisson kernel.

The generalized Lipschitz space $\mathcal{E}^{\alpha,p}$ consists of functions f which are locally integrable and satisfy the following condition: there exists a constant C such that for any cube Q

$$\int_Q |f(x) - f_Q|^p dx \leq C|Q|^{1+\frac{\alpha p}{n}}, \quad (2)$$

where $f_Q = \frac{1}{|Q|} \int_Q f(x) dx$. Denote the norm of f in $\mathcal{E}^{\alpha,p}$ by

$$\|f\|_{\alpha,p} = \inf \{C^{1/p} : C \text{ satisfies (2)}\}.$$

Recently, Qiu [2] has obtained the following result.

Theorem A. *Let $1 < p < \infty$, $-n/p \leq \alpha < 1/2$, $\alpha \neq 0$, and $\lambda > \max(1, 2/p)$. If $f \in \mathcal{E}^{\alpha,p}$ and Tf is $S(f)$ or $g_\lambda^*(f)$, then either $Tf(x) = \infty$ a.e. or $Tf(x) < \infty$ a.e., and there exists a constant C independent of f such that*

$$\|Tf\|_{\alpha,p} \leq C\|f\|_{\alpha,p}.$$

We notice that the range of α in Theorem A seems somewhat rough. It is natural to consider whether the conclusion of the above theorem holds for $-n < \alpha < 1$. The last named author of this paper proved that the conclusion of Theorem A holds for $-n/p < \alpha < 1$ (see [3]). In this paper, with the aid of the idea in [4], we shall introduce a variant of $\mathcal{E}^{\alpha,p}$, $\mathcal{E}_0^{\alpha,p}$, and prove that the conclusion of Theorem A holds for $\mathcal{E}_0^{\alpha,p}$ with $-n < \alpha < 1$. Let us first define $\mathcal{E}_0^{\alpha,p}$.

Definition. A locally integrable function f is called a generalized Lipschitz function of central type if there exists a constant C such that (2) holds for any cube Q centered at the origin. Moreover, the space consisting of all generalized Lipschitz functions of central type is denoted by $\mathcal{E}_0^{\alpha,p}$. We call $\mathcal{E}_0^{\alpha,p}$ the generalized Lipschitz space of central type.

It is easy to see that $\mathcal{E}^{\alpha,p} \subset \mathcal{E}_0^{\alpha,p}$ and $\mathcal{E}_0^{\alpha,p}$ is just the bounded mean oscillation space of central type, BMO_0 in [4]. Let us now formulate our results.

Theorem 1. *Let $1 < p < \infty$ and $-n < \alpha < 1$. If $f \in \mathcal{E}_0^{\alpha,p}$, then either $S(f)(x) = \infty$ a.e. or $S(f)(x) < \infty$ a.e., and there exists a constant C independent of f such that*

$$\|S(f)\|_{\alpha,p} \leq C\|f\|_{\alpha,p}.$$

Theorem 2. *Let $1 < p < \infty$, $-n < \alpha < 1$, and $\lambda > \max(1, 2/p) + 2/n$. If $f \in \mathcal{E}_0^{\alpha,p}$, then either $g_\lambda^*(f)(x) = \infty$ a.e. or $g_\lambda^*(f)(x) < \infty$ a.e., and there exists a constant $C = C(n, \alpha, p, \lambda)$ such that*

$$\|g_\lambda^*(f)\|_{\alpha,p} \leq C\|f\|_{\alpha,p}.$$

2. SOME LEMMAS

Lemma 1. *Let $1 < p < \infty$, $-n < \alpha < 1$, $\alpha \neq 0$, and $0 < d, \alpha < d$. If $f \in \mathcal{E}_0^{\alpha,p}$ and Q is a cube centered at the origin with the edge length r , then there exists a constant $C = C(n, p, \alpha, d)$ such that for any $y > 0$*

$$\int_{\mathbb{R}^n} \frac{|f(x) - f_Q|}{y^{n+d} + |x|^{n+d}} dx \leq Cy^{-d}(y^\alpha + r^\alpha)\|f\|_{\alpha,p}. \quad (3)$$

See [1] and [2] for its proof.

Lemma 1'. *Let $1 < p < \infty$ and $d > 0$. If $f \in BMO_0 = \mathcal{E}_0^{0,p}$ and Q is a cube centered at the origin with the edge length r , then there exists a constant $C = C(n, p, d)$ such that for any $y > 0$,*

$$\int_{\mathbb{R}^n} \frac{|f(x) - f_Q|}{y^{n+d} + |x|^{n+d}} dx \leq Cy^{-d}\left(1 + \left|\log_2 \frac{y}{r}\right|\right)\|f\|_{0,p}.$$

Proof. By the known result in [5] we have

$$\int_{\mathbb{R}^n} \frac{|f(x) - f_Q|}{r^{n+d} + |x|^{n+d}} dx \leq Cr^{-d}\|f\|_{0,p}.$$

Let R be the cube centered at the origin with the edge length y . Then

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{|f(x) - f_Q|}{y^{n+d} + |x|^{n+d}} dx &\leq \int_{\mathbb{R}^n} \frac{|f(x) - f_R|}{y^{n+d} + |x|^{n+d}} dx \\ + |f_R - f_Q| \int_{\mathbb{R}^n} \frac{dx}{y^{n+d} + |x|^{n+d}} &\leq Cy^{-d}\|f\|_{0,p} + Cy^{-d}|f_R - f_Q|. \end{aligned}$$

Thus it remains to prove

$$|f_R - f_Q| \leq C\left(1 + \left|\log_2 \frac{y}{r}\right|\right)\|f\|_{0,p}.$$

Let $y > r$, and let k satisfy $2^k \leq y < 2^{k+1}r$. Then $k \leq \log_2 \frac{y}{r}$ and

$$\begin{aligned} |f_R - f_Q| &\leq |f_R - f_{Q_k}| + \sum_{j=1}^k |f_{Q_j} - f_{Q_{j-1}}| \\ &\leq 2^n \left(\frac{1}{|R|} \int_R |f(x) - f_R|^p dx \right)^{1/p} + \sum_{j=1}^k 2^n \|f\|_{0,p} \\ &\leq 2^n (1+k) \|f\|_{0,p} \\ &\leq C \left(1 + \log_2 \frac{y}{r} \right) \|f\|_{0,p}, \end{aligned}$$

where Q_k is the concentric extension of Q by 2^k times.

When $y < r$, by exchanging y and r , we shall get the same estimate as above with $\log_2 \frac{r}{y} = |\log_2 \frac{y}{r}|$. \square

Let χ_E be the characteristic function of E . For a cube Q in \mathbb{R}^n and $d > 0$ let dQ be the concentric extension of Q by d times.

Lemma 2. *Suppose that $1 < p < \infty$, $-n < \alpha < 1$, and $f \in \mathcal{E}_0^{\alpha,p}$. Let Q be a cube centered at the origin with the edge length r , and $h_Q(x) = [f(x) - f_Q] \chi_{Q^c}(x)$. If there is $x' \in dQ$ such that $S(h_Q)(x') < \infty$, where $d = (8\sqrt{n})^{-1}$, then there exists a constant $C = C(n, \alpha, p)$ such that*

$$S(h_Q)(x) < \infty, \quad \forall x \in dQ$$

and

$$|S(h_Q)(x) - S(h_Q)(x')| < Cr^\alpha \|f\|_{\alpha,p}, \quad \forall x \in dQ.$$

Proof. Let us first consider the case of $\alpha \neq 0$. Fix $x \in dQ$. Set

$$\Gamma^-(x) = \{(z, y) \in \Gamma(x) : y \leq dr\}$$

and

$$\Gamma^+(x) = \{(z, y) \in \Gamma(x) : y > dr\}.$$

Then

$$S(h_Q)(x) \leq S^- + S^+, \quad x \in dQ,$$

where

$$S^- = \left(\iint_{\Gamma^-(x)} y^{1-n} |\nabla h_Q(z, y)|^2 dz dy \right)^{1/2}$$

and

$$S^+ = \left(\iint_{\Gamma^+(x)} y^{1-n} |\nabla h_Q(z, y)|^2 dz dy \right)^{1/2}.$$

Estimating S^- as in [2], we have

$$S^- \leq Cr^\alpha \|f\|_{\alpha,p}. \quad (4)$$

For S^+ we have

$$\begin{aligned} S^+ &= \left(\iint_{\Gamma^+(0)} y^{1-n} |\nabla h_Q(x+z, y)|^2 dz dy \right)^{1/2} \\ &\leq \left(\iint_{\Gamma^+(0)} y^{1-n} |\nabla h_Q(x'+z, y)|^2 dz dy \right)^{1/2} \\ &\quad + \left(\iint_{\Gamma^+(0)} y^{1-n} |\nabla h_Q(x+z, y) - \nabla h_Q(x'+z, y)|^2 dz dy \right)^{1/2} \\ &\leq S(h_Q)(x') + \left\{ \iint_{\Gamma^+(0)} y^{1-n} \right. \\ &\quad \left. \times \left(\int_{\tilde{Q}^c} |\nabla P(x+z-t, y) \nabla P(x'+z+t, y)| |f(t) - f_Q| dt \right)^2 dz dy \right\}^{1/2}. \end{aligned}$$

Note that

$$|\nabla P(x, y) - \nabla P(x', y)| = \left(\sum_{j=1}^{n+1} \left| \frac{\partial}{\partial x_j} p(x, y) - \frac{\partial}{\partial x_j} P(x', y) \right| \right)^{1/2},$$

where $\frac{\partial}{\partial x_{n+1}} = \frac{\partial}{\partial y}$. By the mean value theorem we have

$$\begin{aligned} &\left| \frac{\partial}{\partial x_j} p(x, y) - \frac{\partial}{\partial x_j} P(x', y) \right| \\ &= \left| \nabla \frac{\partial}{\partial x_j} P(x, y) \right|_{x+\theta_j(x-x')} |x-x'|, \quad 0 < \theta_j < 1, \end{aligned}$$

where

$$\left| \nabla \frac{\partial}{\partial x_j} P(x, y) \right| \leq \frac{C}{(y+|x|)^{n+2}}.$$

Thus

$$\begin{aligned} &|\nabla P(x+z-t, y) - \nabla P(x'+z-t, y)| \\ &\leq C|x-x'| \left\{ \sum_{j=1}^{n+1} (y+|x+z-t+\theta_j(x-x')|)^{-2(n+2)} \right\}^{1/2}. \quad (5) \end{aligned}$$

Since $(x, y) \in \Gamma^+(0)$, $x, x' \in dQ$, and $t \notin Q$, we have $|t| > r/2$, $|z| < y$, $|x| < r/16 < |t|/8$, and $|x - x'| < r/8 < |t|/4$. Thus,

$$\begin{aligned} |t| &\leq |x + z - t + \theta_j(x - x')| + |x| + |z| + |x - x'| \\ &\leq |x + z - t + \theta_j(x - x')| + |t|/8 + y + |t|/4 \end{aligned}$$

and

$$\frac{5}{16} (y + |t|) \leq |x + z - t + \theta_j(x - x')| + y,$$

where $1 \leq j \leq n| + 1$. Therefore

$$|\nabla P(x + z - t, y) - \nabla P(x' + z - t, y)| \leq \frac{Cr}{(y + |t|)^{n+2}}. \quad (6)$$

Using (6) and (3), we obtain

$$\begin{aligned} S^+ &\leq S(h_Q)(x') + C \left\{ \iint_{\Gamma^+(0)} y^{1-n} \left[\int_{Q^c} \frac{r|f(t) - f_Q|}{(y + |t|)^{n+2}} dt \right]^2 dz dy \right\}^{1/2} \\ &\leq S(h_Q)(x') + C \left\{ \iint_{\Gamma^+(0)} y^{1-n} r^2 [y^{-2}(y^\alpha + r^\alpha) \|f\|_{\alpha,p}]^2 dz dy \right\}^{1/2} \\ &\leq S(h_Q)(x') + Cr \|f\|_{\alpha,p} \left\{ \int_{dr}^\infty \int_{|z|<y} y^{1-n} y^{-4} (y^{2\alpha} + r^{2\alpha}) dz dy \right\}^{1/2} \\ &\leq S(h_Q)(x') + Cr^\alpha \|f\|_{\alpha,p}. \end{aligned} \quad (7)$$

Combining (4) with (7) we have

$$S(h_Q)(x) \leq S(h_Q)(x') + Cr^\alpha \|f\|_{\alpha,p}.$$

Thus $S(h_Q)(x) < \infty$. Exchanging x and x' , we obtain

$$|S(h_Q)(x) - S(h_Q)(x')| \leq Cr^\alpha \|f\|_{\alpha,p}.$$

Hence the proof of Lemma 2 is complete for the case of $\alpha \neq 0$.

When $\alpha = 0$, by using Lemma 1' instead of Lemma 1 we obtain

$$\begin{aligned} S^+ &\leq S(h_Q)(x') + C \left\{ \iint_{\Gamma^+(0)} y^{1-n} \left[\int_{Q^c} \frac{r|f(t) - f_Q|}{(y + |t|)^{n+2}} dt \right]^2 dz dy \right\}^{1/2} \\ &\leq S(h_Q)(x') + C \left\{ \iint_{\Gamma^+(0)} y^{1-n} r^2 [y^{-2} (1 + |\log_2 \frac{y}{r}|) \|f\|_{0,p}]^2 dz dy \right\}^{1/2} \\ &\leq S(h_Q)(x') + Cr \|f\|_{0,p} \left\{ \int_{dr}^\infty \int_{|z|<y} y^{-3-n} (1 + |\log_2 \frac{y}{r}|)^2 dz dy \right\}^{1/2} \end{aligned}$$

$$\begin{aligned}
&\leq S(h_Q)(x') + Cr^\alpha \|f\|_{0,p} \left\{ \int_1^\infty u^{-3} (1 + |\log_2 u|)^2 du \right\}^{1/2} \\
&\leq S(h_Q)(x') + Cr^\alpha \|f\|_{0,p}. \tag{8}
\end{aligned}$$

Now, it is easy to see that the conclusion of the lemma for $\alpha = 0$ follows from (8) and (4) with $\alpha = 0$. \square

Lemma 3. *Under the hypothesis of Lemma 2, if there is $x' \in dQ$ such that $g_\lambda^*(h_Q)(x') < \infty$, where $\lambda > \max(1, 2/p) + 2/n$, then there exists a constant $C = C(n, \alpha, \lambda, p)$ such that $g_\lambda^*(h_Q)(x) < \infty$ and*

$$|g_\lambda^*(h_Q)(x) - g_\lambda^*(x')| \leq Cr^\alpha \|f\|_{\alpha,p}, \quad \forall x \in dQ.$$

Proof. We only consider the case of $\alpha \neq 0$. As in Lemma 2, the proof in the case $\alpha = 0$ is similar. Let

$$J_k = \{(z, y) \in \mathbb{R}_+^{n+1} : |z| < 2^{k-2}r, 0 < y < 2^{k-2}r\}, \quad k \geq 0.$$

For fixed $x \in dQ$ we have

$$g_\lambda^*(h_Q)(x) \leq G^- + G^+,$$

where

$$G^- = \left(\iint_{J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} |\nabla h_Q(x+z, y)|^2 dz dy \right)^{1/2}$$

and

$$G^+ = \left(\iint_{\mathbb{R}_+^{n+1} \setminus J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} |\nabla h_Q(x+z, y)|^2 dz dy \right)^{1/2}.$$

Note that if $(z, y) \in J_0$, $x \in dQ$, and $t \neq Q$, then $|z| < r/4$, $|x| < r/16$, and $|t| > r/2$. Thus

$$|t| \leq |t - x - z| + |x| + |z| \leq |x + z - t| + \frac{5}{8}|t|$$

and

$$\frac{1}{16}(r + |t|) \leq |x + z - t| + y.$$

By (1) and Lemma 1 we get

$$\begin{aligned}
G^- &\leq C \left\{ \iint_{J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} \left[\int_{Q^c} \frac{r|f(t) - f_Q|}{(y+|x+z-t|)^{n+1}} dt \right]^2 dz dy \right\}^{1/2} \\
&\leq C \left\{ \iint_{J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} \left[\int_{Q^c} \frac{r|f(t) - f_Q|}{(r+|t|)^{n+1}} dt \right]^2 dz dy \right\}^{1/2}
\end{aligned}$$

$$\begin{aligned}
&\leq C \left\{ \int_{dr}^{\infty} \int_{|z|<y} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} [r^{\alpha-1} \|f\|_{\alpha,p}]^2 dz dy \right\}^{1/2} \\
&\leq Cr^{\alpha-1} \|f\|_{\alpha,p} \left(\int_0^r y^{1-n} r^n dy \right)^{1/2} \\
&\leq Cr^{\alpha} \|f\|_{\alpha,p}.
\end{aligned}$$

To estimate C^+ we observe that

$$\begin{aligned}
G^+ &\leq \left\{ \iint_{\mathbb{R}_+^{n+1} \setminus J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} |\nabla h_Q(x' + z, y)|^2 dz dy \right\}^{1/2} \\
&\quad + \left\{ \iint_{\mathbb{R}_+^{n+1} \setminus J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} |\nabla h_Q(x + z, y) - \right. \\
&\quad \quad \left. - \nabla h_Q(x' + z, y)|^2 dz dy \right\}^{1/2} \\
&\leq g_{\lambda}^*(h_Q)(x') + D,
\end{aligned}$$

where

$$\begin{aligned}
D &= \left\{ \iint_{\mathbb{R}_+^{n+1} \setminus J_0} \left(\frac{y}{y+|z|} \right)^{\lambda n} y^{1-n} |\nabla h_Q(x + z, y) - \right. \\
&\quad \left. - \nabla h_Q(x' + z, y)|^2 dz dy \right\}^{1/2} \\
&\leq C \left\{ \sum_{k=1}^{\infty} (2^k r)^{-\lambda n} \iint_{J_k \setminus J_{k-1}} y^{\lambda n+1-n} \right. \\
&\quad \left. \times \left[\int_{Q^c} |\nabla P(x + z - t, y) - \nabla P(x' + z - t, y)| |f(t) - f_Q| dt \right]^2 dz dy \right\}^{1/2} \\
&\leq C \left\{ \sum_{k=1}^{\infty} (2^k r)^{-\lambda n} (A_k + B_k) \right\}^{1/2}.
\end{aligned}$$

Here

$$\begin{aligned}
A_k &= \iint_{J_k \setminus J_{k-1}} y^{\lambda n+1-n} \\
&\quad \times \left[\int_{Q_{k+1}^c} |\nabla P(x + z - t, y) - \nabla P(x' + z - t, y)| |f(t) - f_Q| dt \right]^2 dz dy,
\end{aligned}$$

$$B_k = \iint_{J_k \setminus J_{k-1}} y^{\lambda_{n+1}-n} \times \left[\int_{Q_{k+1} \setminus Q} |\nabla P(x+z-t, y) - \nabla P(x'+z-t, y)| |f(t) - f_Q| dt \right]^2 dz dy,$$

and $Q_{k+1} = 2^{k+1}Q$. Without loss of generality we may assume that $\max(1, 2/p) + 2/n < \lambda < 3 + 2/n$. By the easy inequality (see [3])

$$|\nabla P(x, y) - \nabla P(x', y)| \leq C|x - x'| \left(\frac{1}{(y + |x|)^{n+2}} + \frac{1}{(y + |x'|)^{n+2}} \right), \quad \forall x, x' \in \mathbb{R}^n, \quad y > 0,$$

together with the Minkowski inequality for integrals, we have

$$\begin{aligned} B_k &\leq Cr^2 \iint_{J_k \setminus J_{k-1}} y^{\lambda_{n+1}-n} \left\{ \int_{Q_{k+1} \setminus Q} |f(t) - f_Q| \left(\frac{1}{(y + |x + z - t|)^{n+2}} + \frac{1}{(y + |x' + z - t|)^{n+2}} \right) dt \right\}^2 dz dy \\ &\leq Cr^2 \int_0^\infty \int_{\mathbb{R}^n} y^{\lambda_{n+1}-n} \left\{ \int_{Q_{k+1}} |f(t) - f_Q| \left(\frac{1}{(y + |x + z - t|)^{n+2}} + \frac{1}{(y + |x' + z - t|)^{n+2}} \right) dt \right\}^2 dz dy \\ &\leq Cr^2 \int_{\mathbb{R}^n} \left[\int_{Q_{k+1}} |f(t) - f_Q| \left(\int_0^\infty \frac{y^{\lambda_{n+1}-n}}{(y + |x + z - t|)^{2(n+2)}} dy \right)^{1/2} dt \right]^2 dz \\ &\quad + Cr^2 \int_{\mathbb{R}^n} \left[\int_{Q_{k+1}} |f(t) - f_Q| \left(\int_0^\infty \frac{y^{\lambda_{n+1}-n}}{(y + |x' + z - t|)^{2(n+2)}} dy \right)^{1/2} dt \right]^2 dz \\ &= Cr^2 \int_{\mathbb{R}^n} \left[\int_{Q_{k+1}} \frac{|f(t) - f_Q|}{|z + x - t|^{(3n-\lambda n+2)/2}} \left(\int_0^\infty \frac{y^{\lambda_{n+1}-n}}{(1+y)^{2(n+2)}} dy \right)^{1/2} dt \right]^2 dz \\ &\quad + Cr^2 \int_{\mathbb{R}^n} \left[\int_{Q_{k+1}} \frac{|f(t) - f_Q|}{|z + x' - t|^{(3n-\lambda n+2)/2}} \left(\int_0^\infty \frac{y^{\lambda_{n+1}-n}}{(1+y)^{2(n+2)}} dy \right)^{1/2} dt \right]^2 dz \\ &= Cr^2 \int_{\mathbb{R}^n} \left(\int_{Q_{k+1}} \frac{|f(t) - f_Q|}{|u - t|^{n - [(\lambda n - 1) - 2]/2}} dt \right)^2 du. \end{aligned}$$

Using the Hardy–Littlewood–Sobolev theorem on fractional integration with $\gamma = [(\lambda - 1)n - 2]/2$, $q = 2$, and $1/s = 1/q + \gamma/n = \lambda/2 - 1/n$ (see [6]), we obtain

$$B_k \leq Cr^2 \left(\int_{Q_{k+1}} |f(z) - f_Q|^s dz \right)^{2/s}.$$

Since $\lambda \geq 2/p + 2/n$, then $p \geq s$. Thus

$$\begin{aligned} B_k &\leq Cr^2 \left(\int_{Q_{k+1}} |f(z) - f_Q|^p dz \right)^{2/p} |Q_{k+1}|^{2(1/s-1/p)} \\ &\leq Cr^2 \left\{ \left(\int_{Q_{k+1}} |f(z) - f_Q|^p dz \right)^{1/p} \right. \\ &\quad \left. + |Q_{k+1}|^{1/p} |f_{Q_{k+1}} - f_Q| \right\}^2 |Q_{k+1}|^{2(1/s-1/p)} \\ &\leq Cr^2 \{ |Q_{k+1}|^{1/p+\alpha/n} \|f\|_{\alpha,p} \\ &\quad + |Q_{k+1}|^{1/p} (2^k r)^\alpha \|f\|_{\alpha,p} \}^2 |Q_{k+1}|^{2(1/s+1/p)} \\ &\leq Cr^2 (2^k r)^{2\alpha} (2^k r)^{\lambda n - 2} \|f\|_{\alpha,p} \\ &\leq C (2^k r)^{\lambda n} (2^{2k(\alpha-1)} r^{2\alpha}) \|f\|_{\alpha,p}. \end{aligned}$$

To estimate A_k we observe that if $(z, y) \in J_k \setminus J_{k-1}$ and $t \notin Q_{k+1}$, then $|t| > 2^k r$, $k \geq 1$, and $|z| < 2^{k-2} r < |t|/4$. Thus,

$$\begin{aligned} |t| &\leq |x + z - t + \theta_j(x - x')| + |x| + |z| + |x - x'| \\ &\leq |x + z - t + \theta(x - x')| + \frac{5}{16}|t|. \end{aligned}$$

By Lemma 1 we have

$$\begin{aligned} A_k &\leq C \iint_{J_k \setminus J_{k+1}} y^{\lambda n + 1 - n} \left[\int_{Q_{k+1}^c} \frac{r|f(t) - f_Q|}{(2^k r + |t|)^{n+2}} dt \right]^2 dz dy \\ &\leq Cr^2 \iint_{J_k \setminus J_{k+1}} y^{\lambda n + 1 - n} \{ (2^k r)^{-2} [(2^k r)^\alpha + r^\alpha] \|f\|_{\alpha,p} \}^2 dz dy \\ &\leq Cr^2 (2^k r)^{-4+2\alpha} \|f\|_{\alpha,p} \int_0^{2^k r} \int_{|z| < 2^k r} y^{\lambda n + 1 - n} dz dy \\ &\leq Cr^{2\alpha} (2^k r)^{\lambda n} 2^{2k(\alpha-1)} \|f\|_{\alpha,p}. \end{aligned}$$

Combining the estimate of A_k with that of B_k , we obtain

$$D \leq C \left\{ \sum_{k=1}^{\infty} (2^k r)^{-\lambda n} (2^k r)^{\lambda n} r^{2\alpha} 2^{2k(\alpha-1)} \|f\|_{\alpha,p} \right\}^{1/2} \leq Cr^\alpha \|f\|_{\alpha,p}.$$

Therefore

$$g_\lambda^*(h_Q)(x) \leq g_\lambda^*(h_Q)(x') + Cr^\alpha \|f\|_{\alpha,p}.$$

As in the last part of the proof of Lemma 2, we have

$$|g_\lambda^*(h_Q)(x) - g_\lambda^*(h_Q)(x')| \leq Cr^\alpha \|f\|_{\alpha,p}. \quad \square$$

3. THE PROOFS OF THE THEOREMS

Let T be one of the Littlewood–Paley functions as in Section 1. Suppose that $Tf(x) \neq \infty$ a.e. Then $|E| \triangleq |\{x : Tf(x) < \infty\}| > 0$. Thus there is a cube Q' centered at the origin such that $|Q' \cap E| > 0$. Set $Q = (1/d)Q'$ (then $Q' = dQ$). We write f as

$$\begin{aligned} f(x) &= f_Q + [f(x) - f_Q]\chi_Q(x) + [f(x) - f_Q]\chi_{Q^c}(x) \\ &\triangleq f_Q + g_Q(x) + h_Q(x). \end{aligned}$$

Since

$$Tf(x) \leq Tg_Q(x) + Th_Q(x) \quad (9)$$

and

$$Th_Q(x) \leq Tf(x) + Tg_Q(x), \quad (10)$$

it is easy to see that the inequality

$$\|g_Q\|_p = \left(\int_Q |f(t) - f_Q|^p dt \right)^{1/p} \leq C|Q|^{1/p+\alpha/n} \|f\|_{\alpha,p} \quad (11)$$

implies that $g_Q \in L^p$. Then it follows from the L^p -boundedness of the Littlewood–Paley operator that $Tg_Q(x) < \infty$ a.e.. Since $|Q' \cap E| > 0$, there is $x' \in Q' \cap E \subset dQ$ such that $Tf(x') < \infty$ and $Tg_Q(x') < \infty$. By (10) and Lemmas 2 and 3, we have $Th_Q(x') < \infty$ and

$$Th_Q(x) < \infty, \quad \forall x \in dQ = Q'.$$

By (9) we obtain

$$Tf(x) < \infty \quad \text{a.e. } x \in Q'.$$

Finally, let the edge length of Q' tend to ∞ ; we have $Tf(x) < \infty$ a.e., $x \in \mathbb{R}^n$.

Let Q' be a cube centered at the origin, and $Q = (1/d)Q'$. Choose $x' \in dQ$ so that $Th_Q(x') < \infty$. Then it follows from (11) and Lemmas 2 and 3 that

$$\begin{aligned} & \left(\int_{Q'} |Tf(x) - Th_Q(x')|^p dx \right)^{1/p} \\ & \leq \left(\int_{Q'} |Tg_Q(x)|^p dx \right)^{1/p} + \left(\int_{Q'} |Th_Q(x) - Th_Q(x')|^p dx \right)^{1/p} \\ & \leq C \|g_Q\|_p + C |Q'|^{1/p} r^\alpha \|f\|_{\alpha,p} \\ & \leq C |Q'|^{1/p+\alpha/n} \|f\|_{\alpha,p}. \end{aligned}$$

This completes the proof of the theorems. \square

ACKNOWLEDGEMENT

The work is supported by the NSFC (Tian Yuan). Shanzhen Lu would like to thank Kozo Yabuta for inviting him to visit Nara during his sabbatical in Japan.

REFERENCES

1. D. S. Kurtz, Littlewood–Paley operators on BMO. *Proc. Amer. Math. Soc.* **99**(1987), 657–666.
2. S. G. Qiu, Boundedness of Littlewood–Paley operators and Marcinkiewicz integral on $\mathcal{E}^{\alpha,p}$. *J. Math. Res. Exposition* **12**(1992), 41–50.
3. K. Yabuta, Boundedness of Littlewood–Paley operators. *Preprint*.
4. S. Z. Lu and D. C. Yang, The Littlewood–Paley function and φ -transform characterizations of a new Hardy space HK_2 associated with the Herz space. *Studia Math.* **101**(1992), 285–298.
5. F. B. Fabes, R. L. Janson, and U. Neri, Spaces of harmonic functions representable by Poisson integrals of functions in BMO and $\mathcal{L}_{p,\lambda}$. *Indiana Univ. Math. J.* **25**(1976), 159–170.
6. E. M. Stein, Singular integrals and differentiability properties of functions. *Princeton University Press, Princeton*, 1970.

(Received 04.05.1994)

Authors' addresses:

Shanzhen Lu and Changmei Tan
Department of Mathematics
Beijing Normal University
Beijing 100875, P. R. China

Kozo Yabuta
Department of Mathematics
Nara Women's University
Nara 630, Japan