

MANN ITERATES OF DIRECTIONALLY NONEXPANSIVE MAPPINGS IN HYPERBOLIC SPACES

ULRICH KOHLENBACH AND LAURENȚIU LEUȘTEAN

Received 2 July 2002

In a previous paper, the first author derived an explicit quantitative version of a theorem due to Borwein, Reich, and Shafrir on the asymptotic behaviour of Mann iterations of nonexpansive mappings of convex sets C in normed linear spaces. This quantitative version, which was obtained by a logical analysis of the ineffective proof given by Borwein, Reich, and Shafrir, could be used to obtain strong uniform bounds on the asymptotic regularity of such iterations in the case of bounded C and even weaker conditions. In this paper, we extend these results to hyperbolic spaces and directionally nonexpansive mappings. In particular, we obtain significantly stronger and more general forms of the main results of a recent paper by W. A. Kirk with explicit bounds. As a special feature of our approach, which is based on logical analysis instead of functional analysis, no functional analytic embeddings are needed to obtain our uniformity results which contain all previously known results of this kind as special cases.

1. Introduction

This paper continues the approach of applying methods from mathematical logic to proofs in the metric fixed-point theory started by the first author in [9, 13, 14]. In particular, [9, 14] were concerned with explicit bounds on the asymptotic behaviour of the so-called Mann iterations of nonexpansive mappings in the following setting.

Let $(X, \|\cdot\|)$ be a normed linear space, $C \subseteq X$ convex, and $f : C \rightarrow C$ nonexpansive, that is,

$$\forall x, y \in C, \quad \|f(x) - f(y)\| \leq \|x - y\|. \quad (1.1)$$

Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence of real numbers in $[0, 1)$. Then, a Mann iteration starting from $x_0 := x \in C$ is defined as (the special case of $\lambda_n := 1/2$ was already

considered by Krasnosel'skii in [16])

$$x_{n+1} := (1 - \lambda_n)x_n + \lambda_n f(x_n). \quad (1.2)$$

In [2], the following important result is proved. If $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum and is bounded away from 1, then

$$\forall x \in C, \quad \|x_n - f(x_n)\| \rightarrow r_C(f), \quad (1.3)$$

where $r_C(f) := \inf\{\|x - f(x)\| \mid x \in C\}$.

In many cases, for example, for bounded C , $r_C(f)$ can be shown to be 0, that is, $\|x_n - f(x_n)\| \rightarrow 0$, which (for bounded C) was first proved by Ishikawa in the classical paper [6]. The special case of constant $\lambda_k = \lambda$ also follows from [3], which even proves uniform (in x) convergence. Later, [4] extended this to uniformity in both x and f .

Using specially designed techniques from mathematical logic, the first author established in a series of papers general theorems on the extractability of explicit bounds from large classes of *prima facie* ineffective existence proofs, together with procedures, to transform such proofs into new ones from which these bounds can be read off (see [10, 11, 12] and, for a general survey, [15]). The proof given by Borwein et al. in [2] of the result just cited happens to be of the required form. In [14], as a result of the logical transformation of the proof, a new quantitative version of the Borwein-Reich-Shafir theorem was obtained. From this version, explicit uniform bounds for the case of bounded C could simply be read off. These bounds only depend on the error ε , an upper bound for the diameter of C , a distance by which (λ_n) stays away from 1, and a rate of divergence of the sum of that sequence towards infinity. Subsequently (see [9]), this could be extended to the case where not C as a whole is required to be bounded but only some Mann iteration sequence.

The logical approach does not use any tools from functional analysis to establish these uniformity results which suggests that it should be possible to generalize the results to other settings in which the basic proof idea of the Borwein-Reich-Shafir theorem applies.

In this paper, we show that, indeed, all results from [14] (as well as the one from [9] just mentioned) extend to the more general class of hyperbolic spaces (in the sense of [18]) and (with minor changes in the assumptions) to the more general class of directionally nonexpansive mappings (in the sense of [8]).

In particular, we prove significantly stronger forms of the main results in [8]. Although some of the proofs follow closely those in [14], we include them in this paper for completeness.

2. Hyperbolic spaces basic results

In this section, we present hyperbolic spaces, defined by Reich and Shafrir [18] as an appropriate context for the study of operator theory in general, and of iterative processes for nonexpansive mappings in particular. This class of metric spaces includes all normed linear spaces and Hadamard manifolds, as well as the Hilbert ball equipped with the hyperbolic metric [7] and the Cartesian products of Hilbert balls. Extensive information on hyperbolic spaces and a detailed treatment of examples like the Hilbert ball can be found in [5] (see also [4, 7, 17, 19]).

A still more general class of metric spaces is the class of spaces of hyperbolic type (see [4, 7]), which are contained in the class of convex metric spaces [20]. In particular, every hyperbolic space is a space of hyperbolic type.

In the following, we collect some basic facts on hyperbolic spaces which we will need later. To make the paper self-contained we include the (short) proofs.

Let (X, ρ) be a metric space and let \mathbb{R} denote the real line. We say that a mapping $c : \mathbb{R} \rightarrow X$ is a *metric embedding* of \mathbb{R} into X if

$$\rho(c(s), c(t)) = |s - t| \tag{2.1}$$

for all real s and t . The image of \mathbb{R} under a metric embedding will be called a *metric line*.

Any isometry $c : \mathbb{R} \rightarrow X$ is a metric embedding and the metric line associated with it is X . In fact, a metric embedding is an isometry if and only if it is surjective.

The image $c([a, b]) \subseteq X$ of a real interval under a metric embedding $c : \mathbb{R} \rightarrow X$ will be called a *metric segment*.

Let $x, y \in X$ and let $c : \mathbb{R} \rightarrow X$ be a metric embedding. We say that the metric line $c(\mathbb{R})$ *passes through x and y* if $x, y \in c(\mathbb{R})$ and that the metric segment $c([a, b])$ *joins x and y* if $(c(a) = x \text{ and } c(b) = y)$ or $(c(a) = y \text{ and } c(b) = x)$.

In the sequel, we assume that (X, ρ) contains a nonempty family M of metric lines such that for each pair of distinct points x and y in X , there is a unique metric line which passes through x and y . Hence, there is a nonempty family $\{c_i\}_{i \in I}$ of metric embeddings such that for all $x \neq y \in X$ there is a unique $i \in I$ such that $x, y \in c_i(\mathbb{R})$.

Remark 2.1. Since $M \neq \emptyset$, there is at least one metric embedding $c : \mathbb{R} \rightarrow X$. Since c is injective, it follows that $\text{card}(X) \geq \text{card}(\mathbb{R}) = \aleph_1$.

The following lemmas collect some simple facts. For the sake of completeness, we prove them.

LEMMA 2.2. *For any $x \in X$, there is at least one metric line from M that passes through x .*

Proof. By the above remark, X is infinite, so there is $y \in X, y \neq x$. Take now the unique metric line that passes through x and y . □

LEMMA 2.3. *For any distinct points x and y in X , there is a unique metric segment joining them.*

Proof. There is a unique $i \in I$ such that $x, y \in c_i(\mathbb{R})$. Since c_i is injective, there are unique $a, b \in \mathbb{R}$, $a \neq b$, such that $c_i(a) = x$ and $c_i(b) = y$. Hence, the unique metric segment joining x and y is $c_i([a, b])$ if $a < b$ or $c_i([b, a])$ if $b < a$. \square

We denote by $[x, y]$ or $[y, x]$ the unique metric segment joining two distinct points x and y from X .

For any $x \in X$, by $[x, x]$, we understand the singleton $\{x\}$. By Lemma 2.2, there is $c : \mathbb{R} \rightarrow X$ and $a \in \mathbb{R}$ such that $c(a) = x$, hence $\{x\} = c([a, a])$. Thus, $[x, x]$ is a degenerate metric segment.

LEMMA 2.4. *Let $x, y \in X$, $x \neq y$, and $z, w \in [x, y]$. Then,*

- (i) $0 \leq \rho(x, z) \leq \rho(x, y)$;
- (ii) if $\rho(x, z) = \rho(x, w)$, then $z = w$.

Proof. Let $[x, y] = c([a, b])$.

(i) Let $s \in [a, b]$ such that $c(s) = z$. If $c(a) = x$ and $c(b) = y$, then $\rho(x, z) = \rho(c(a), c(s)) = |s - a| = s - a \leq b - a = \rho(x, y)$. If $c(a) = y$ and $c(b) = x$, then $\rho(x, z) = \rho(c(b), c(s)) = |b - s| = b - s \leq b - a = \rho(x, y)$.

(ii) Since $z, w \in [x, y]$, there are $s_1, s_2 \in [a, b]$ such that $c(s_1) = z$ and $c(s_2) = w$. Suppose that $c(a) = x$ and $c(b) = y$. It follows that $\rho(x, z) = \rho(c(a), c(s_1)) = |a - s_1| = s_1 - a$, and, similarly, $\rho(x, w) = s_2 - a$. Thus, $\rho(x, z) = \rho(x, w)$ if and only if $s_1 - a = s_2 - a$ if and only if $s_1 = s_2$ if and only if $z = w$. \square

LEMMA 2.5. *Let $c : \mathbb{R} \rightarrow X$ be a metric embedding, $a \leq b \in \mathbb{R}$, and $t \in [0, 1]$. Then,*

$$\begin{aligned} \rho(c(a), c((1-t)a + tb)) &= t\rho(c(a), c(b)), \\ \rho(c(b), c((1-t)a + tb)) &= (1-t)\rho(c(a), c(b)). \end{aligned} \tag{2.2}$$

Proof. This follows from $\rho(c(a), c((1-t)a + tb)) = |a - ((1-t)a + tb)| = t|a - b| = t\rho(c(a), c(b))$, and, similarly, $\rho(c(b), c((1-t)a + tb)) = |b - ((1-t)a + tb)| = (1-t)|a - b| = (1-t)\rho(c(a), c(b))$. \square

PROPOSITION 2.6. *Let $x, y \in X$. For each $t \in [0, 1]$, there is a unique point $z \in [x, y]$ such that*

$$\rho(x, z) = t\rho(x, y), \quad \rho(y, z) = (1-t)\rho(x, y). \tag{2.3}$$

Proof. If $x = y$, then, obviously, $z = x = y$. Suppose that $x \neq y$. Let $[x, y] = c([a, b])$. If $c(a) = x$ and $c(b) = y$, then take $z = c((1-t)a + tb)$. If $c(b) = x$ and $c(a) = y$, then take $z = c((1-t)b + ta)$. Then, $z \in [x, y]$ and z satisfies (2.3) by Lemma 2.5. Unicity of z follows from Lemma 2.4(ii). \square

The unique point satisfying (2.3) will be denoted as $(1 - t)x \oplus ty$. Then, for any $x \in X$ and $t \in [0, 1]$, $(1 - t)x \oplus tx = x$.

If $z \in [x, y]$ satisfies only one of conditions (2.3), then it is necessary that $z = (1 - t)x \oplus ty$. Hence, any point of the segment $[x, y]$ satisfying one of conditions (2.3) satisfies also the other.

Remark 2.7. Let $x, y \in X$, $x \neq y$, and $s, t \in [0, 1]$. Then,

- (i) $(1 - t)x \oplus ty = (1 - s)x \oplus sy$ if and only if $s = t$;
- (ii) $(1 - t)x \oplus ty = ty \oplus (1 - t)x$.

LEMMA 2.8. Let $x, y \in X$, $x \neq y$. Then,

- (i) $[x, y] = \{(1 - t)x \oplus ty \mid t \in [0, 1]\}$;
- (ii) the mapping $f : [0, 1] \rightarrow [x, y]$, $f(t) = (1 - t)x \oplus ty$ is continuous and bijective;
- (iii) $\rho(x, z) + \rho(z, y) = \rho(x, y)$ for all $z \in [x, y]$;
- (iv) if $z \neq w \in X$ are such that $\rho(x, y) \leq \rho(z, w)$, then there is a unique $v \in [z, w]$ such that $\rho(z, v) = \rho(x, y)$.

Proof. (i) The inclusion \supseteq follows by definition.

For the converse inclusion \subseteq , let $z \in [x, y]$ and $t = \rho(x, z)/\rho(x, y)$. Then, by Lemma 2.4(i), $t \in [0, 1]$ and $\rho(x, z) = t\rho(x, y)$. It follows that $z = (1 - t)x \oplus ty$.

(ii) Applying (i) and Remark 2.7(i), we immediately get that f is well defined and bijective. Let $c([a, b]) = [x, y]$. Then for all $t \in [0, 1]$, $f(t) = c((1 - t)a + tb)$. Since c is continuous and the map $[0, 1] \rightarrow [a, b]$, $t \mapsto (1 - t)a + tb$ is also continuous, it follows that f is continuous.

(iii) Let $z \in [x, y]$. By (i), there is $t \in [0, 1]$ such that $z = (1 - t)x \oplus ty$, hence $\rho(x, z) + \rho(z, y) = t\rho(x, y) + (1 - t)\rho(x, y) = \rho(x, y)$.

(iv) Let $t = \rho(x, y)/\rho(z, w)$, so $t \in [0, 1]$. Let $v = (1 - t)z \oplus tw$. Then $v \in [z, w]$ and $\rho(z, v) = t\rho(z, w) = \rho(x, y)$. □

Definition 2.9 [18]. A triple (X, ρ, M) is a hyperbolic space if

$$\rho\left(\frac{1}{2}x \oplus \frac{1}{2}y, \frac{1}{2}x \oplus \frac{1}{2}z\right) \leq \frac{1}{2}\rho(y, z) \tag{2.4}$$

for all $x, y, z \in X$.

Remark 2.10 [18]. Equation (2.4) is equivalent to

$$\rho\left(\frac{1}{2}x \oplus \frac{1}{2}y, \frac{1}{2}w \oplus \frac{1}{2}z\right) \leq \frac{1}{2}(\rho(x, w) + \rho(y, z)) \tag{2.5}$$

for all $x, y, z, w \in X$.

Proof. (2.5) \Rightarrow (2.4) is obvious, just take $w = x$. It remains to prove (2.4) \Rightarrow (2.5). For any $x, y, z, w \in X$,

$$\begin{aligned} \rho\left(\frac{1}{2}x \oplus \frac{1}{2}y, \frac{1}{2}w \oplus \frac{1}{2}z\right) &\leq \rho\left(\frac{1}{2}x \oplus \frac{1}{2}y, \frac{1}{2}x \oplus \frac{1}{2}z\right) + \rho\left(\frac{1}{2}x \oplus \frac{1}{2}z, \frac{1}{2}w \oplus \frac{1}{2}z\right) \\ &\leq \frac{1}{2}(\rho(y, z) + \rho(x, w)). \end{aligned} \tag{2.6}$$

□

Let (X, ρ, M) be a hyperbolic space. A nonempty subset $C \subseteq X$ is *convex* if $[x, y] \in C$ for all $x, y \in C$. We denote by $d(C)$ the diameter of C . Hence,

$$d(C) = \sup \{\rho(x, y) \mid x, y \in C\}. \tag{2.7}$$

The set C is bounded if $d(C) < \infty$. A sequence $(x_n)_{n \in \mathbb{N}} \subseteq X$ is *bounded* if the set $\{x_n \mid n \in \mathbb{N}\}$ is bounded.

At a few places, we use the following fact.

PROPOSITION 2.11 [5, 18]. *Let (X, ρ, M) be a hyperbolic space. Then,*

$$\rho((1-t)x \oplus tz, (1-t)y \oplus tw) \leq (1-t)\rho(x, y) + t\rho(z, w) \tag{2.8}$$

for all $t \in [0, 1]$ and $x, y, z, w \in X$.

Proof. The idea of the proof is presented in [5, pages 77, 104]. We first prove the result for $t = k/2^n$, where $k, n \in \mathbb{N}$ are such that $k \leq 2^n$. We use induction on n . If $n = 0$, then $k/2^n = k$ and $k \in \{0, 1\}$. If $k = 0$, then (2.8) $\Leftrightarrow (\rho(1x \oplus 0z, 1y \oplus 0w) \leq \rho(x, y)) \Leftrightarrow (\rho(x, y) \leq \rho(x, y))$. If $k = 1$, then (2.8) $\Leftrightarrow (\rho(0x \oplus 1z, 0y \oplus 1w) \leq \rho(z, w)) \Leftrightarrow (\rho(z, w) \leq \rho(z, w))$. Hence, (2.8) is true even with equality.

Now, suppose that (2.8) is true for $t = k/2^n$. Hence,

$$\rho\left(\left(1 - \frac{k}{2^n}\right)x \oplus \frac{k}{2^n}z, \left(1 - \frac{k}{2^n}\right)y \oplus \frac{k}{2^n}w\right) \leq \left(1 - \frac{k}{2^n}\right)\rho(x, y) + \frac{k}{2^n}\rho(z, w) \tag{2.9}$$

for all $k \in \mathbb{N}, k \leq 2^n$ and for all $x, y, z, w \in X$.

We have to prove (2.8) for $t = k/2^{n+1}$, where $k \in \mathbb{N}, k \leq 2^{n+1}$. If we denote $u := (1 - k/2^{n+1})x \oplus (k/2^{n+1})z$ and $v := (1 - k/2^{n+1})y \oplus (k/2^{n+1})w$, then we have to prove

$$\rho(u, v) \leq \left(1 - \frac{k}{2^{n+1}}\right)\rho(x, y) + \frac{k}{2^{n+1}}\rho(z, w). \tag{2.10}$$

First, we show (2.10) for $k \leq 2^n$, that is, $k/2^n \in [0, 1]$. Let $\alpha := (1 - k/2^n)x \oplus (k/2^n)z$, $\beta := (1 - k/2^n)y \oplus (k/2^n)w$, $\alpha_1 := (1/2)x \oplus (1/2)\alpha$, and $\beta_1 := (1/2)y \oplus (1/2)\beta$. Then, $\rho(x, \alpha_1) = (1/2)\rho(x, \alpha) = (k/2^{n+1})\rho(x, z) = \rho(x, u)$ and $\alpha_1, u \in [x, z]$

since $u, \alpha \in [x, z]$ and $\alpha_1 \in [x, \alpha]$. Applying Lemma 2.4(ii), it follows that $u = \alpha_1$. We similarly get that $v = \beta_1$. Now, applying (2.5) and the induction hypothesis, it follows that

$$\begin{aligned} \rho(u, v) &= \rho(\alpha_1, \beta_1) = \rho\left(\frac{1}{2}x \oplus \frac{1}{2}\alpha, \frac{1}{2}y \oplus \frac{1}{2}\beta\right) \leq \frac{1}{2}(\rho(x, y) + \rho(\alpha, \beta)) \\ &\leq \frac{1}{2}\rho(x, y) + \frac{1}{2}\left(\left(1 - \frac{k}{2^n}\right)\rho(x, y) + \frac{k}{2^n}\rho(z, w)\right) \\ &= \left(1 - \frac{k}{2^{n+1}}\right)\rho(x, y) + \frac{k}{2^{n+1}}\rho(z, w). \end{aligned} \tag{2.11}$$

Now, suppose that $2^n < k \leq 2^{n+1}$ and let $p := 2^{n+1} - k$. Then $p \leq 2^n$, so we can apply (2.10) for p . We obtain

$$\begin{aligned} \rho(u, v) &= \rho\left(\frac{p}{2^{n+1}}x \oplus \left(1 - \frac{p}{2^{n+1}}\right)z, \frac{p}{2^{n+1}}y \oplus \left(1 - \frac{p}{2^{n+1}}\right)w\right) \\ &= \rho\left(\left(1 - \frac{p}{2^{n+1}}\right)z \oplus \frac{p}{2^{n+1}}x, \left(1 - \frac{p}{2^{n+1}}\right)w \oplus \frac{p}{2^{n+1}}y\right) \\ &\leq \left(1 - \frac{p}{2^{n+1}}\right)\rho(z, w) + \frac{p}{2^{n+1}}\rho(x, y) \\ &= \left(1 - \frac{k}{2^{n+1}}\right)\rho(x, y) + \frac{k}{2^{n+1}}\rho(z, w). \end{aligned} \tag{2.12}$$

In the sequel, we use the fact that the set $D := \{k/2^n \mid k, n \in \mathbb{N}, k \leq 2^n\}$ is dense in $[0, 1]$. Let $t \in [0, 1]$. Then, there is $(t_p)_{p \in \mathbb{N}} \subseteq D$ such that $\lim_{p \rightarrow \infty} t_p = t$. For all $p \in \mathbb{N}$,

$$\rho((1 - t_p)x \oplus t_p z, (1 - t_p)y \oplus t_p w) \leq (1 - t_p)\rho(x, y) + t_p \rho(z, w). \tag{2.13}$$

Letting $p \rightarrow \infty$ and using Lemma 2.8(ii) and the fact that ρ is continuous, we get (2.8). □

COROLLARY 2.12. *Let (X, ρ, M) be a hyperbolic space. Then, for all $t \in [0, 1]$ and $x, y, z \in X$,*

$$\rho((1 - t)x \oplus tz, y) \leq (1 - t)\rho(x, y) + t\rho(z, y). \tag{2.14}$$

Proof. Apply (2.8) with $w = y$. □

We now present the related concept of metric space of hyperbolic type [4, 7] (see also [20]).

Let (X, ρ) be a metric space and S a family of metric segments. We say that (X, ρ, S) is of hyperbolic type if the following are satisfied:

- (i) for each two points $x, y \in X$, there is a unique metric segment from S that joins them, denoted as $[x, y]$;

(ii) if $p, x, y, m \in X$ and if $m \in [x, y]$ satisfies $\rho(x, m) = t\rho(x, y)$ for $t \in [0, 1]$, then

$$\rho(p, m) \leq (1 - t)\rho(p, x) + t\rho(p, y). \tag{2.15}$$

PROPOSITION 2.13. *Any hyperbolic space is of hyperbolic type.*

Proof. Let (X, ρ, M) be a hyperbolic space. Let S be the family of all metric segments determined by the metric embeddings associated with M . Let $x, y \in M$. If $x \neq y$, then (i) is satisfied, by Lemma 2.3. If $x = y$, then the unique metric segment from S that joins x and x is $[x, x] = \{x\}$. We verify (ii). Since $m \in [x, y]$ and $\rho(x, m) = t\rho(x, y) = \rho(x, (1 - t)x \oplus ty)$, by Lemma 2.4(ii), we must have $m = (1 - t)x \oplus ty$. Now apply (2.14). \square

In the sequel, let $(\lambda_n)_{n \in \mathbb{N}} \subseteq [0, 1]$.

Denote, for all $i, n \in \mathbb{N}$,

$$S_{i,n} := \sum_{s=i}^{i+n-1} \lambda_s, \quad P_{i,n} := \prod_{s=i}^{i+n-1} \frac{1}{1 - \lambda_s}. \tag{2.16}$$

Let $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ be two sequences in X such that for all $n \in \mathbb{N}$,

$$x_{n+1} = (1 - \lambda_n)x_n \oplus \lambda_n y_n. \tag{2.17}$$

The following very important result was proved in [4] for spaces of hyperbolic type. Hence, by Proposition 2.13, it is true also for hyperbolic spaces.

PROPOSITION 2.14 [4]. *Let (X, ρ, M) be a hyperbolic space. Suppose that $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ satisfy for all $n \in \mathbb{N}$,*

$$\rho(y_n, y_{n+1}) \leq \rho(x_n, x_{n+1}). \tag{2.18}$$

Then, the sequence $(\rho(x_n, y_n))_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is nonincreasing and, for all $i, n \in \mathbb{N}$,

$$(1 + S_{i,n})\rho(x_i, y_i) \leq \rho(x_i, y_{i+n}) + P_{i,n}[\rho(x_i, y_i) - \rho(x_{i+n}, y_{i+n})]. \tag{2.19}$$

We use in the sequel the following consequence of the above inequality.

PROPOSITION 2.15 [2]. *In the assumptions of Proposition 2.14,*

$$S_{i,n}\rho(x_i, y_i) \leq \rho(x_i, x_{i+n}) + P_{i,n}[\rho(x_i, y_i) - \rho(x_{i+n}, y_{i+n})]. \tag{2.20}$$

Proof. Apply Proposition 2.14 and the fact that $\rho(x_i, y_{i+n}) - \rho(x_i, y_i) \leq \rho(x_i, x_{i+n}) + \rho(x_{i+n}, y_{i+n}) - \rho(x_i, y_i) \leq \rho(x_i, x_{i+n})$; since $(\rho(x_n, y_n))_{n \in \mathbb{N}}$ is nonincreasing, hence $\rho(x_{i+n}, y_{i+n}) - \rho(x_i, y_i) \leq 0$. \square

PROPOSITION 2.16 [4]. *In addition to the assumptions of Proposition 2.14, assume that*

- (i) *the set $\{\rho(x_n, y_{n+i}) \mid n, i \in \mathbb{N}\}$ is bounded;*
- (ii) *$(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum;*
- (iii) *there is $b \in (0, 1)$ such that $\lambda_n \leq b$ for all $n \in \mathbb{N}$.*

Then $\lim_{n \rightarrow \infty} \rho(x_n, y_n) = 0$.

Proof. This result is proved in [4] for any space of hyperbolic type. Applying again Proposition 2.13, it follows that it is true for any hyperbolic space, too. \square

LEMMA 2.17. *In the hypotheses of Proposition 2.14, the following are equivalent:*

- (i) *$(x_n)_{n \in \mathbb{N}}$ is bounded;*
- (ii) *$(y_n)_{n \in \mathbb{N}}$ is bounded;*
- (iii) *the set $\{\rho(x_n, y_{n+i}) \mid n, i \in \mathbb{N}\}$ is bounded.*

Proof. Let $n, i \in \mathbb{N}$.

(i) \Rightarrow (ii). $\rho(y_n, y_{n+i}) \leq \rho(y_n, x_n) + \rho(x_n, x_{n+i}) + \rho(x_{n+i}, y_{n+i}) \leq 2\rho(x_0, y_0) + \rho(x_n, x_{n+i})$ since $(\rho(x_n, y_n))_{n \in \mathbb{N}}$ is nonincreasing, by Proposition 2.14.

(ii) \Rightarrow (iii). $\rho(x_n, y_{n+i}) \leq \rho(x_n, y_n) + \rho(y_n, y_{n+i}) \leq \rho(x_0, y_0) + \rho(y_n, y_{n+i})$.

(iii) \Rightarrow (i). $\rho(x_n, x_{n+i}) \leq \rho(x_n, y_{n+i}) + \rho(y_{n+i}, x_{n+i}) \leq \rho(x_n, y_{n+i}) + \rho(x_0, y_0)$. \square

LEMMA 2.18. *The following are equivalent:*

- (i) $\limsup_{n \rightarrow \infty} \lambda_n < 1$;
- (ii) *there is $b \in (0, 1)$ such that $\lambda_n \leq b < 1$ for all $n \in \mathbb{N}$.*

Proof. Obviously, since $\lambda_n < 1$ for all $n \in \mathbb{N}$. \square

Using these lemmas, we obtain the following reformulation of Proposition 2.16.

THEOREM 2.19. *Let (X, ρ, M) be a hyperbolic space and $(\lambda_n)_{n \in \mathbb{N}} \subseteq [0, 1)$. Suppose that $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum and $\limsup_{n \rightarrow \infty} \lambda_n < 1$.*

Let $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ be two sequences in X which satisfy for all $n \in \mathbb{N}$,

$$x_{n+1} = (1 - \lambda_n)x_n \oplus \lambda_n y_n, \quad \rho(y_n, y_{n+1}) \leq \rho(x_n, x_{n+1}). \tag{2.21}$$

If $(x_n)_{n \in \mathbb{N}}$ is bounded, then $\lim_{n \rightarrow \infty} \rho(x_n, y_n) = 0$.

3. Uniform asymptotic regularity for directionally nonexpansive mappings

The main purpose of the present paper is to generalize the core results from [9, 14] not only to hyperbolic spaces (which is largely straightforward) but at the same time to directionally nonexpansive mappings, which requires quite some care. Directionally nonexpansive mappings were considered in [8]. In this section, we, in particular, strengthen the main results from [8].

Definition 3.1 [8]. Let (X, ρ, M) be a hyperbolic space and $C \subseteq X$ a nonempty convex subset. A mapping $f : C \rightarrow C$ is called *directionally nonexpansive* if

$$\rho(f(x), f(y)) \leq \rho(x, y), \quad (3.1)$$

for all $x \in C$ and $y \in [x, f(x)]$.

Recall that $f : C \rightarrow C$ is called *nonexpansive* if for all $x, y \in C$,

$$\rho(f(x), f(y)) \leq \rho(x, y). \quad (3.2)$$

Obviously, any nonexpansive mapping is directionally nonexpansive, but the converse fails as directionally nonexpansive mappings do not even need to be continuous on the whole space.

Example 3.2 (simplified by Paulo Oliva). Consider the normed space $(\mathbb{R}^2, \|\cdot\|_{\max})$ and the function

$$f : [0, 1]^2 \rightarrow [0, 1]^2, \quad f(x, y) := \begin{cases} (1, y), & \text{if } y > 0, \\ (0, y), & \text{if } y = 0. \end{cases} \quad (3.3)$$

Clearly, f is directionally nonexpansive (even directionally constant) but discontinuous at $(0, 0)$.

In the following, (X, ρ, M) will be an arbitrary hyperbolic space, $C \subseteq X$ a nonempty convex subset of X , and $f : C \rightarrow C$ a directionally nonexpansive mapping. Define [2]

$$r_C(f) := \inf \{\rho(x, f(x)) \mid x \in C\}. \quad (3.4)$$

We consider the so-called *Krasnosel'skiĭ-Mann iteration* starting from $x \in C$

$$x_0 := x, \quad x_{n+1} := (1 - \lambda_n)x_n \oplus \lambda_n f(x_n), \quad (3.5)$$

where $(\lambda_n)_{n \in \mathbb{N}}$ is a sequence of real numbers in $[0, 1)$.

LEMMA 3.3. *For all $n \in \mathbb{N}$,*

$$\rho(f(x_n), f(x_{n+1})) \leq \rho(x_n, x_{n+1}). \quad (3.6)$$

Proof. Since $x_{n+1} \in [x_n, f(x_n)]$, we can apply the fact that f is directionally nonexpansive to obtain that $\rho(f(x_n), f(x_{n+1})) \leq \rho(x_n, x_{n+1})$. \square

Thus, the sequences $(x_n)_{n \in \mathbb{N}}$ and $(f(x_n))_{n \in \mathbb{N}}$ satisfy the hypotheses of [Proposition 2.14](#) with $y_n := f(x_n)$. We get immediately the following results.

PROPOSITION 3.4. *The sequence $(\rho(x_n, f(x_n)))_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is nonincreasing and for all $i, n \in \mathbb{N}$,*

$$S_{i,n}\rho(x_i, f(x_i)) \leq \rho(x_i, x_{i+n}) + P_{i,n}[\rho(x_i, f(x_i)) - \rho(x_{i+n}, f(x_{i+n}))]. \quad (3.7)$$

Proof. Apply [Lemma 3.3](#), [Proposition 2.14](#), and [Proposition 2.15](#). □

For nonexpansive mappings, the following proposition is due to [6] (normed spaces) and [4] for hyperbolic spaces. Using [Lemma 3.3](#), the proof from [4] extends to directionally nonexpansive mappings.

PROPOSITION 3.5. *Suppose that $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum and $\limsup_{n \rightarrow \infty} \lambda_n < 1$. If C is bounded, then $\lim_{n \rightarrow \infty} \rho(x_n, f(x_n)) = 0$.*

Proof. By [Theorem 2.19](#) and [Lemma 3.3](#). □

COROLLARY 3.6. *Suppose that $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum and $\limsup_{n \rightarrow \infty} \lambda_n < 1$. If C is bounded, then for every $x \in X$, $\lim_{n \rightarrow \infty} \rho(x_n, f(x_n)) = 0$.*

COROLLARY 3.7. *Suppose that $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum and $\limsup_{n \rightarrow \infty} \lambda_n < 1$. If C is bounded or—even weaker—there is $x \in C$ such that $(x_n)_{n \in \mathbb{N}}$ is bounded, then $r_C(f) = 0$.*

Let $x^* \in C$ and $(x_n^*)_{n \in \mathbb{N}}$ be the Krasnosel’skiĭ-Mann iteration that starts from x^* .

The next inequality is due to [2].

LEMMA 3.8. *If f is nonexpansive, then for all $n \in \mathbb{N}$,*

$$\rho(x_{n+1}, x_{n+1}^*) \leq \rho(x_n, x_n^*). \tag{3.8}$$

Proof. Applying inequality (2.8) and the definition of a nonexpansive mapping, we get that

$$\begin{aligned} \rho(x_{n+1}, x_{n+1}^*) &= \rho((1 - \lambda_n)x_n \oplus \lambda_n f(x_n), (1 - \lambda_n)x_n^* \oplus \lambda_n f(x_n^*)) \\ &\leq (1 - \lambda_n)\rho(x_n, x_n^*) + \lambda_n\rho(f(x_n), f(x_n^*)) \\ &\leq (1 - \lambda_n)\rho(x_n, x_n^*) + \lambda_n\rho(x_n, x_n^*) \\ &= \rho(x_n, x_n^*). \end{aligned} \tag{3.9}$$

□

Since, in general, $x_n^* \notin [x_n, f(x_n)]$, we cannot prove the inequality

$$\rho(f(x_n), f(x_n^*)) \leq \rho(x_n, x_n^*), \tag{3.10}$$

on which the proof of [Lemma 3.8](#) is based for directionally nonexpansive mappings f . The absence of [Lemma 3.8](#) causes some changes in the generalizations of the main results from [9, 14] to directionally nonexpansive mappings carried out below.

In [2], the following theorem is proved.

THEOREM 3.9 [2]. *Let $(X, \| \cdot \|)$ be a normed linear space, $C \subseteq X$ convex, and $f : C \rightarrow C$ nonexpansive. Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence of real numbers in $[0, 1)$, which is divergent in sum and satisfies $\limsup_{n \rightarrow \infty} \lambda_n < 1$. Then,*

$$\|x_n - f(x_n)\| \xrightarrow{n \rightarrow \infty} r_C(f), \tag{3.11}$$

where $(x_n)_{n \in \mathbb{N}}$ is the Krasnosel'skiĭ-Mann iteration starting from $x \in C$.

In [14], applying proofs transformations developed in the context of mathematical logic (see [15]) to the proof of Theorem 3.8 from [2], the first author obtained an effective quantitative version of that theorem (see also Remark 3.11). From this quantitative version, various strong (effective) uniformity results for the case of bounded C were derived (improving previous results in this direction from [3, 4]), as well as (for the first time) for the more general case of bounded $(x_n)_{n \in \mathbb{N}}$ (see [9]). Since these uniformity results were obtained by logical analysis and, in particular, did not use any functional analytic embedding techniques (in contrast to [3, 4]), this suggests that it should be possible to extend these results to the more general setting of hyperbolic spaces and directionally nonexpansive mappings. The main content of this paper is to show that this is indeed true to a large extent. Whereas the extension to hyperbolic spaces does not cause any problems at all, the absence of Lemma 3.8 for directionally nonexpansive mappings results in an additional hypothesis which, however, is trivially satisfied, for example, in the bounded case.

THEOREM 3.10. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex subset, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $[0, 1)$ which is divergent in sum and satisfies*

$$\forall n \in \mathbb{N}, \quad \lambda_n \leq 1 - \frac{1}{K}, \tag{3.12}$$

for some $K \in \mathbb{N}$.

Let $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be such that

$$\forall i, n \in \mathbb{N}, \quad (\alpha(i, n) \leq \alpha(i + 1, n)) \wedge \left(n \leq \sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s \right). \tag{3.13}$$

Let $x, x^* \in C$ and $d > 0$ be such that

$$\forall n \in \mathbb{N}, \quad \rho(x_n, x_n^*) \leq d, \tag{3.14}$$

where $(x_n)_{n \in \mathbb{N}}$ and $(x_n^*)_{n \in \mathbb{N}}$ are the Krasnosel'skiĭ-Mann iterations starting from x and x^* .

Then,

$$\forall \varepsilon > 0, \quad \forall n \geq h(\varepsilon, x, d, f, K, \alpha), \quad \rho(x_n, f(x_n)) < \rho(x^*, f(x^*)) + \varepsilon \tag{3.15}$$

holds, where $(n \div 1 = \max(0, n - 1))$

$$h(\varepsilon, x, d, f, K, \alpha) := \hat{\alpha}(\lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil + 1, M), \tag{3.16}$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$, $c(f, x) \in \mathbb{R}$ is such that $c(f, x) \geq \rho(x, f(x))$, and $\hat{\alpha}(0, n) := \tilde{\alpha}(0, n)$, $\hat{\alpha}(i + 1, n) := \tilde{\alpha}(\hat{\alpha}(i, n), n)$ with $\tilde{\alpha}(i, n) := i + \alpha(i, n)$ ($i, n \in \mathbb{N}$).

Proof. Most parts of the proof follow closely the one given in [14] for the non-expansive case (and normed spaces). For completeness we present, nevertheless, all details.

Let

$$\gamma := \rho(x^*, f(x^*)). \tag{3.17}$$

Let $\varepsilon > 0$ be arbitrary and $M \in \mathbb{N}$ be such that

$$M \geq \frac{1 + 2d}{\varepsilon}. \tag{3.18}$$

For example, $M := \lceil (1 + 2d)/\varepsilon \rceil$.

Let $\delta > 0$ be so small that

$$\delta \exp(K(M + 1)) < 1. \tag{3.19}$$

For example, $\delta := 1/(2 \exp(K(M + 1)))$.

Let $i, n \in \mathbb{N}$. Then, reasoning as in [6],

$$\begin{aligned} P_{i,n} &= \prod_{s=i}^{i+n-1} \left(1 + \frac{\lambda_s}{1 - \lambda_s} \right) \\ &= \exp \left(\ln \prod_{s=i}^{i+n-1} \left(1 + \frac{\lambda_s}{1 - \lambda_s} \right) \right) = \exp \left(\sum_{s=i}^{i+n-1} \ln \left(1 + \frac{\lambda_s}{1 - \lambda_s} \right) \right) \\ &\leq \exp \left(\sum_{s=i}^{i+n-1} \frac{\lambda_s}{1 - \lambda_s} \right) \quad \text{since } \ln(1 + x) \leq x \text{ for } x \geq 0 \\ &\leq \exp \left(K \sum_{s=i}^{i+n-1} \lambda_s \right) = \exp(K \cdot S_{i,n}) \end{aligned} \tag{3.20}$$

since $\lambda_s \leq 1 - 1/K$ implies $1 - \lambda_s \geq 1/K$, so $1/(1 - \lambda_s) \leq K$ for all $s \in \mathbb{N}$.

Hence, we have proved that for all $i, n \in \mathbb{N}$,

$$P_{i,n} \leq \exp(K \cdot S_{i,n}). \tag{3.21}$$

Define $\alpha^* : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ by

$$\alpha^*(i, n) := \min \{m \in \mathbb{N} \mid n \leq S_{i,m}\}. \tag{3.22}$$

Since $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum, it follows that for all $i \in \mathbb{N}$, the sequence $(S_{i,m})_{m \in \mathbb{N}}$ is not bounded above; so for all $n \in \mathbb{N}$, the set $A_{i,n} := \{m \in \mathbb{N} \mid n \leq S_{i,m}\}$ is nonempty; hence it has a least element. Thus, α^* is well defined. We also get that $\alpha^*(i, n) - 1 \notin A_{i,n}$, which means that $S_{i,\alpha^*(i,n)-1} < n$, that is, $S_{i,\alpha^*(i,n)} - \lambda_{i+\alpha^*(i,n)-1} < n$, so $S_{i,\alpha^*(i,n)} < n + \lambda_{i+\alpha^*(i,n)-1} < n + 1$. Hence, for all $i, n \in \mathbb{N}$,

$$n \leq S_{i,\alpha^*(i,n)} < n + 1. \tag{3.23}$$

Consider the Krasnosel'skiĭ-Mann iteration $(x_n^*)_{n \in \mathbb{N}}$ starting from x^* . Then,

$$\begin{aligned} \rho(x_i^*, x_{i+n}^*) &\leq \sum_{s=i}^{i+n-1} \rho(x_s^*, x_{s+1}^*) = \sum_{s=i}^{i+n-1} \lambda_s \rho(x_s^*, f(x_s^*)) \\ &\leq \left(\sum_{s=i}^{i+n-1} \lambda_s \right) \rho(x_i^*, f(x_i^*)) = S_{i,n} \cdot \rho(x_i^*, f(x_i^*)) \\ &\leq S_{i,n} \cdot \rho(x^*, f(x^*)) \end{aligned} \tag{3.24}$$

since, by [Proposition 3.4](#), $(\rho(x_n^*, f(x_n^*)))_{n \in \mathbb{N}}$ is nonincreasing. Hence, for all $i, n \in \mathbb{N}$,

$$\rho(x_i^*, x_{i+n}^*) \leq S_{i,n} \cdot \rho(x^*, f(x^*)). \tag{3.25}$$

Consider now the Krasnosel'skiĭ-Mann iteration $(x_n)_{n \in \mathbb{N}}$ starting from x . Applying again [Proposition 3.4](#), we get that the sequence $(\rho(x_n, f(x_n)))_{n \in \mathbb{N}}$ is nonincreasing and, since it is bounded below by 0, it is convergent and hence Cauchy. Thus, for δ satisfying [\(3.19\)](#), there exists an i such that

$$\rho(x_i, f(x_i)) - \rho(x_{i+\alpha^*(i,M)}, f(x_{i+\alpha^*(i,M)})) \leq \delta. \tag{3.26}$$

In the sequel, we consider an i satisfying [\(3.26\)](#).

Applying [Proposition 3.4](#) and [\(3.26\)](#), we get that

$$\begin{aligned} S_{i,\alpha^*(i,M)} \cdot \rho(x_i, f(x_i)) &\leq \rho(x_i, x_{i+\alpha^*(i,M)}) + \delta \cdot P_{i,\alpha^*(i,M)} \\ &\leq \rho(x_i, x_i^*) + \rho(x_i^*, x_{i+\alpha^*(i,M)}^*) \\ &\quad + \rho(x_{i+\alpha^*(i,M)}^*, x_{i+\alpha^*(i,M)}) + \delta \cdot P_{i,\alpha^*(i,M)} \\ &\leq 2d + S_{i,\alpha^*(i,M)} \cdot \rho(x^*, f(x^*)) \\ &\quad + \delta \cdot P_{i,\alpha^*(i,M)} \quad \text{by the hypothesis and (3.25)} \\ &= 2d + S_{i,\alpha^*(i,M)} \cdot \gamma + \delta \cdot P_{i,\alpha^*(i,M)} \quad \text{by (3.17)}. \end{aligned} \tag{3.27}$$

That is, we have got

$$S_{i,\alpha^*(i,M)} \cdot \rho(x_i, f(x_i)) \leq 2d + S_{i,\alpha^*(i,M)} \cdot \gamma + \delta \cdot P_{i,\alpha^*(i,M)}. \tag{3.28}$$

We now prove

$$\rho(x_i, f(x_i)) < \gamma + \varepsilon. \tag{3.29}$$

Suppose that $\rho(x_i, f(x_i)) \geq \gamma + \varepsilon$. It follows that

$$S_{i,\alpha^*(i,M)}(\gamma + \varepsilon) \leq S_{i,\alpha^*(i,M)} \cdot \rho(x_i, f(x_i)); \tag{3.30}$$

so applying (3.28), we get that

$$S_{i,\alpha^*(i,M)}(\gamma + \varepsilon) \leq 2d + S_{i,\alpha^*(i,M)} \cdot \gamma + \delta \cdot P_{i,\alpha^*(i,M)}. \tag{3.31}$$

Hence,

$$S_{i,\alpha^*(i,M)} \cdot \varepsilon \leq 2d + \delta \cdot P_{i,\alpha^*(i,M)}. \tag{3.32}$$

It follows that

$$\begin{aligned} 1 + 2d &\leq M \cdot \varepsilon && \text{by (3.18)} \\ &\leq S_{i,\alpha^*(i,M)} \cdot \varepsilon && \text{by (3.23)} \\ &\leq 2d + \delta \cdot P_{i,\alpha^*(i,M)} && \text{by (3.32)} \\ &\leq 2d + \delta \cdot \exp(K \cdot S_{i,\alpha^*(i,M)}) && \text{by (3.21)} \\ &< 2d + \delta \cdot \exp(K(M + 1)) && \text{by (3.23)} \\ &< 2d + 1 && \text{by (3.19)}. \end{aligned} \tag{3.33}$$

That is, we have got a contradiction.

Hence, we have proved that if $i \in \mathbb{N}$ is such that

$$\rho(x_i, f(x_i)) - \rho(x_{i+\alpha^*(i,M)}, f(x_{i+\alpha^*(i,M)})) \leq \delta, \tag{3.34}$$

then

$$\rho(x_i, f(x_i)) < \gamma + \varepsilon. \tag{3.35}$$

Define $\widetilde{\alpha}^*, \widehat{\alpha}^* : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ by

$$\begin{aligned} \widetilde{\alpha}^*(k, n) &:= k + \alpha^*(k, n), \\ \widehat{\alpha}^*(0, n) &:= \widetilde{\alpha}^*(0, n), \\ \widehat{\alpha}^*(k + 1, n) &:= \widetilde{\alpha}^*(\widehat{\alpha}^*(k, n), n). \end{aligned} \tag{3.36}$$

Since $\widehat{\alpha}^*(k+1, n) = \widehat{\alpha}^*(\widehat{\alpha}^*(k, n), n) = \widehat{\alpha}^*(k, n) + \alpha^*(\widehat{\alpha}^*(k, n), n) \geq \widehat{\alpha}^*(k, n)$, it follows that for all $k, n \in \mathbb{N}$,

$$\widehat{\alpha}^*(k, n) \leq \widehat{\alpha}^*(k+1, n). \quad (3.37)$$

Claim 1. Let $j := \lceil \rho(x, f(x))/\delta \rceil + 1$. For all $n \in \mathbb{N}$,

$$\exists k \leq j, \quad \rho(x_{\widehat{\alpha}^*(k, n)}, f(x_{\widehat{\alpha}^*(k, n)})) - \rho(x_{\widehat{\alpha}^*(k+1, n)}, f(x_{\widehat{\alpha}^*(k+1, n)})) \leq \delta. \quad (3.38)$$

Proof of Claim 1. Let $n \in \mathbb{N}$ and for every $k \in \mathbb{N}$ let

$$T_k := \rho(x_{\widehat{\alpha}^*(k, n)}, f(x_{\widehat{\alpha}^*(k, n)})) - \rho(x_{\widehat{\alpha}^*(k+1, n)}, f(x_{\widehat{\alpha}^*(k+1, n)})). \quad (3.39)$$

Suppose that the claim is false. Then, $T_k > \delta$ for all $k \leq j$, so $\sum_{k=0}^j T_k > \delta \cdot (j+1)$, that is,

$$\begin{aligned} & \rho(x_{\widehat{\alpha}^*(0, n)}, f(x_{\widehat{\alpha}^*(0, n)})) - \rho(x_{\widehat{\alpha}^*(j+1, n)}, f(x_{\widehat{\alpha}^*(j+1, n)})) \\ & > \delta \cdot (j+1) = \delta \cdot \left\lceil \frac{\rho(x, f(x))}{\delta} \right\rceil \geq \rho(x, f(x)). \end{aligned} \quad (3.40)$$

From this we get that

$$\rho(x_{\widehat{\alpha}^*(0, n)}, f(x_{\widehat{\alpha}^*(0, n)})) > \rho(x, f(x)), \quad (3.41)$$

which is a contradiction to the fact that the sequence $(\rho(x_n, f(x_n)))_{n \in \mathbb{N}}$ is non-increasing and finishes the proof of the claim.

Let k satisfy (3.38) with $n := M$ and let $i := \widehat{\alpha}^*(k, M)$. Applying (3.38) and the definition of $\widehat{\alpha}^*$, it follows immediately that i satisfies (3.34). Hence, i also satisfies (3.35).

Let $c(f, x) \in \mathbb{R}$ be such that $c(f, x) \geq \rho(x, f(x))$. Let

$$h(\varepsilon, x, d, f, K, \alpha^*) := \widehat{\alpha}^*(\lceil 2c(f, x) \cdot \exp(K(M+1)) \rceil + 1, M). \quad (3.42)$$

Since we can put $\delta := 1/(2 \exp(K(M+1)))$, we get that

$$\begin{aligned} \frac{\rho(x, f(x))}{\delta} &= 2\rho(x, f(x)) \cdot \exp(K(M+1)) \\ &\leq 2c(f, x) \cdot \exp(K(M+1)). \end{aligned} \quad (3.43)$$

Hence,

$$k \leq \left\lceil \frac{\rho(x, f(x))}{\delta} \right\rceil + 1 \leq \lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil + 1. \tag{3.44}$$

Applying (3.37), it follows that $i \leq h(\varepsilon, x, d, f, K, \alpha^*)$. Now, using the fact that i satisfies (3.35), we immediately get that

$$\forall n \geq h(\varepsilon, x, f, d, K, \alpha^*), \quad (\rho(x_n, f(x_n)) < \rho(x^*, f(x^*)) + \varepsilon). \tag{3.45}$$

Hence, we have obtained the conclusion of the theorem with α^* instead of α . We now show that we can replace α^* with α satisfying the more flexible requirement from the hypothesis

$$\forall i, n \in \mathbb{N}, \quad (\alpha(i, n) \leq \alpha(i + 1, n)) \wedge \left(n \leq \sum_{s=i}^{i+\alpha(i, n)-1} \lambda_s \right). \tag{3.46}$$

Since $n \leq S_{i, \alpha(i, n)}$, by the definition of α^* it follows that for all $i, n \in \mathbb{N}$,

$$\alpha^*(i, n) \leq \alpha(i, n). \tag{3.47}$$

We now prove that for all $i, n \in \mathbb{N}$,

$$\widehat{\alpha}^*(i, n) \leq \widehat{\alpha}(i, n). \tag{3.48}$$

We use induction on i . For $i = 0$, we get that

$$\widehat{\alpha}^*(0, n) = \widetilde{\alpha}^*(0, n) = \alpha^*(0, n) \leq \alpha(0, n) = \widetilde{\alpha}(0, n) = \widehat{\alpha}(0, n). \tag{3.49}$$

Suppose that $\widehat{\alpha}^*(i, n) \leq \widehat{\alpha}(i, n)$. Using (3.47) and the fact that, by the hypothesis, α is nondecreasing in the first argument, we get that $\widehat{\alpha}^*(i + 1, n) = \widetilde{\alpha}^*(\widehat{\alpha}^*(i, n), n) = \alpha^*(i, n) + \alpha^*(\widehat{\alpha}^*(i, n), n) \leq \widehat{\alpha}(i, n) + \alpha(\widehat{\alpha}^*(i, n), n) \leq \widehat{\alpha}(i, n) + \alpha(\widehat{\alpha}(i, n), n) = \widetilde{\alpha}(\widehat{\alpha}(i, n), n) = \widehat{\alpha}(i + 1, n)$. It follows that

$$\begin{aligned} h(\varepsilon, x, d, f, K, \alpha^*) &= \widehat{\alpha}^*(\lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil + 1, M) \\ &\leq \widehat{\alpha}(\lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil + 1, M) \\ &= h(\varepsilon, x, d, f, K, \alpha). \end{aligned} \tag{3.50}$$

Finally, applying (3.45), we obtain

$$\forall n \geq h(\varepsilon, x, d, f, K, \alpha), \quad \rho(x_n, f(x_n)) < \rho(x^*, f(x^*)) + \varepsilon. \quad (3.51)$$

□

Remark 3.11. If f is nonexpansive, then applying Lemma 3.8, it follows that the sequence $(\rho(x_n, x_n^*))_{n \in \mathbb{N}}$ is nonincreasing; so letting $d := \rho(x, x^*)$, we get that

$$\forall n \in \mathbb{N}, \quad \rho(x_n, x_n^*) \leq d. \quad (3.52)$$

Hence, Theorem 3.10 holds with

$$h(\varepsilon, x, x^*, f, K, \alpha) = \hat{\alpha}(\lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil \div 1, M), \quad (3.53)$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2\rho(x, x^*))/\varepsilon$ and $c(f, x)$, $\tilde{\alpha}$, and $\hat{\alpha}$ are as above.

It is this restricted form (for normed spaces) of Theorem 3.10 which is proved in [14].

The following remarks from [14] apply in our context as well.

Remark 3.12. Let $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be such that

$$\forall i, n \in \mathbb{N}, \quad n \leq \sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s. \quad (3.54)$$

Define $\alpha^+ : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ by

$$\alpha^+(i, n) := \max_{j \leq i} \alpha(j, n). \quad (3.55)$$

Then, α^+ is nondecreasing in the first argument and also satisfies (3.54) so that Theorem 3.10 holds with $h(\varepsilon, x, d, f, K, \alpha^+)$.

Remark 3.13. A function α satisfying the conditions of Theorem 3.10 can be easily computed from a function $\beta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying the weaker requirement

$$\forall n, \quad n \leq \sum_{s=0}^{\beta(n)} \lambda_s. \quad (3.56)$$

Just define $\beta'(i, n) := \beta(n + i) - i + 1$ and $\beta^+(i, n) := \max_{j \leq i} \beta'(j, n)$.

Then, β^+ satisfies the conditions imposed on α so that Theorem 3.10 holds with $h(\varepsilon, x, d, f, K, \beta^+)$, where β satisfies (3.56).

Proof. We only have to verify that β' satisfies condition (3.54) from Remark 3.12. Let $i, n \in \mathbb{N}$. Then,

$$\sum_{s=i}^{i+\beta'(i,n)-1} \lambda_s = \sum_{s=i}^{\beta(n+i)} \lambda_s = \sum_{s=0}^{\beta(n+i)} \lambda_s - \sum_{s=0}^{i-1} \lambda_s \geq n+i - \sum_{s=0}^{i-1} \lambda_s > n+i-i = n, \tag{3.57}$$

since $\lambda_s < 1$ for all $s \in \mathbb{N}$. □

Just note that as a corollary to Theorem 3.10, we get the following (nonquantitative) strengthened version of the original Borwein-Reich-Shafir theorem.

COROLLARY 3.14. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex subset, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $[0, 1)$ which is divergent in sum and satisfies that $\limsup_{n \rightarrow \infty} \lambda_n < 1$. Then, for all $x \in C$ if*

$$\forall \varepsilon > 0, \exists x^* \in C(\rho(x_n, x_n^*) \text{ bounded} \wedge \rho(x^*, f(x^*)) \leq r_C(f) + \varepsilon), \tag{3.58}$$

then

$$\rho(x_n, f(x_n)) \xrightarrow{n \rightarrow \infty} r_C(f). \tag{3.59}$$

The main application of the quantitative version of the Borwein-Reich-Shafir theorem given in [14] was a fully uniform bound on the asymptotic regularity $\|x_n - f(x_n)\| \rightarrow 0$ in the case of bounded C . ‘‘Fully uniform’’ here means that the rate of convergence depends only on the error ε , an upper bound d for the diameter of C , and the quantities K, α on λ_k but not on x, f , or any special features of C . Uniformity in x (for constant $\lambda_k := \lambda$) was first established in [3]. In [4], uniformity in x and f has been proved for general λ_k , but no uniformity in C or λ_k . Moreover, no effective bounds were obtained. Recently (see [8, Theorem 1]), Kirk has established uniformity in x, f for directionally nonexpansive mappings in the case of constant $\lambda_k := \lambda$. All these results are based on functional analytic embeddings. We now show that the results obtained in [14] by logical analysis of the proof of Theorem 3.9 extend even with the same numerical bounds to the case of hyperbolic spaces and directionally nonexpansive mappings (containing Theorem 1 from [8] just mentioned as a special case). This is due to the fact that the only additional assumption that $\forall n \in \mathbb{N}(\rho(x_n, x_n^*) \leq d)$, which we had to impose in the directionally nonexpansive case, holds trivially for sets C whose diameter is bounded by d . The proofs of Corollaries 3.15, 3.17, 3.18, and 3.20 follow the ones in [14] for the corresponding results in the case of nonexpansive mappings in normed spaces except that we now have to use our more general Theorem 3.10.

COROLLARY 3.15. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex bounded subset with diameter $d(C) < \infty$, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $[0, 1)$, which is divergent in sum and*

satisfies

$$\forall n \in \mathbb{N}, \quad \lambda_n \leq 1 - \frac{1}{K}, \tag{3.60}$$

for some $K \in \mathbb{N}$.

Let $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be such that

$$\forall i, n \in \mathbb{N}, \quad (\alpha(i, n) \leq \alpha(i + 1, n)) \wedge \left(n \leq \sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s \right). \tag{3.61}$$

Then,

$$\forall x \in C, \quad \forall \varepsilon > 0, \quad \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon \tag{3.62}$$

holds, where

$$h(\varepsilon, d, K, \alpha) := \hat{\alpha}(\lceil 2d \cdot \exp(K(M + 1)) \rceil - 1, M), \tag{3.63}$$

where $d \in \mathbb{R}$ is such that $d \geq d(C)$, $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$, and $\hat{\alpha}(0, n) := \tilde{\alpha}(0, n)$, $\hat{\alpha}(i + 1, n) := \tilde{\alpha}(\hat{\alpha}(i, n), n)$ with $\tilde{\alpha}(i, n) := i + \alpha(i, n)$.

Proof. Let $x \in C$ and $\varepsilon > 0$. Let $d \geq d(C)$. Then, for every $x^* \in C$, we have that $\rho(x_n, x_n^*) \leq d(C) \leq d$ for all $n \in \mathbb{N}$. Hence, for every $x^* \in C$, we can apply [Theorem 3.10](#) to get

$$\forall n \geq h(\varepsilon, x, d, f, K, \alpha), \quad \rho(x_n, f(x_n)) < \rho(x^*, f(x^*)) + \varepsilon, \tag{3.64}$$

where

$$h(\varepsilon, x, d, f, K, \alpha) := \hat{\alpha}(\lceil 2c(f, x) \cdot \exp(K(M + 1)) \rceil - 1, M), \tag{3.65}$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$, $c(f, x) \in \mathbb{R}$ is such that $c(f, x) \geq \rho(x, f(x))$, and $\tilde{\alpha}, \hat{\alpha}$ are defined as above.

Since $d \geq d(C) \geq \rho(x, f(x))$, we can take $c(x, f) := d$.

Thus, we get that

$$h(\varepsilon, x, d, f, K, \alpha) = \hat{\alpha}(\lceil 2d \cdot \exp(K(M + 1)) \rceil - 1, M) = h(\varepsilon, d, K, \alpha). \tag{3.66}$$

Let $n \geq h(\varepsilon, d, K, \alpha)$. It follows that

$$\forall x^* \in C, \quad \rho(x_n, f(x_n)) < \rho(x^*, f(x^*)) + \varepsilon, \tag{3.67}$$

hence

$$\rho(x_n, f(x_n)) \leq \inf \{ \rho(x^*, f(x^*)) \mid x^* \in C \} + \varepsilon, \tag{3.68}$$

that is,

$$\rho(x_n, f(x_n)) \leq r_C(f) + \varepsilon. \tag{3.69}$$

Now, apply the fact that $r_C(f) = 0$, by [Corollary 3.7](#). □

Remark 3.16. In [Corollary 3.15](#), the bound $h(\varepsilon, d, K, \alpha)$ can be replaced by $h(\varepsilon/d, 1, K, \alpha)$ just by applying the old bound to the modified hyperbolic space, where $\rho_d(x, y) := (1/d)\rho(x, y)$ and $c_d(s) := c(d \cdot s)$.

COROLLARY 3.17. *Let $d, \varepsilon > 0$, $K \in \mathbb{N}$ and $\beta : \mathbb{N} \rightarrow \mathbb{N}$ be an arbitrary mapping. Then, there exists an $N \in \mathbb{N}$ such that for any hyperbolic space (X, ρ, M) , any nonempty bounded convex set $C \subseteq X$ such that $d(C) \leq d$, any directionally nonexpansive mapping $f : C \rightarrow C$, any sequence $\lambda_n \in [0, 1 - 1/K]$ satisfying $n \leq \sum_{s=0}^{\beta(n)} \lambda_s$ (for all $n \in \mathbb{N}$), and any $x \in C$, the following holds:*

$$\forall n \geq N, \quad \rho(x_n, f(x_n)) \leq \varepsilon. \tag{3.70}$$

Proof. From $n \leq \sum_{s=0}^{\beta(n)} \lambda_s$ for all $n \in \mathbb{N}$, it follows that $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum. Apply [Remark 3.13](#) and [Corollary 3.15](#). □

COROLLARY 3.18. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex bounded subset with diameter $d(C) < \infty$, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $K \in \mathbb{N}$, $K \geq 2$ and $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $[1/K, 1 - 1/K]$. Then, the following holds:*

$$\forall x \in C, \forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K), \quad \rho(x_n, f(x_n)) \leq \varepsilon, \tag{3.71}$$

where

$$h(\varepsilon, d, K) := K \cdot M \cdot \lceil 2d \cdot \exp(K(M+1)) \rceil \tag{3.72}$$

with $d \in \mathbb{R}$, $d \geq d(C)$ and $M \in \mathbb{N}$, $M \geq (1 + 2d)/\varepsilon$.

Proof. Define $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ by

$$\alpha(i, n) = Kn. \tag{3.73}$$

Then, $\sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s \geq \sum_{s=i}^{i+\alpha(i,n)-1} 1/K = (1/K)\alpha(i, n) = n$ and $\alpha(i, n) = \alpha(i + 1, n) = Kn$, so α satisfies the conditions of [Corollary 3.15](#).

We also get immediately that

$$\begin{aligned} \tilde{\alpha}(i, n) &= i + \alpha(i, n) = i + Kn, \\ \hat{\alpha}(i, n) &= K(i + 1)n. \end{aligned} \tag{3.74}$$

Applying [Corollary 3.15](#), it follows that

$$\forall x \in C, \forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon, \quad (3.75)$$

where

$$\begin{aligned} h(\varepsilon, d, K, \alpha) &= \hat{\alpha}([2d \cdot \exp(K(M+1))] - 1, M) \\ &= K \cdot M \cdot [2d \cdot \exp(K(M+1))] \\ &= h(\varepsilon, d, K). \end{aligned} \quad (3.76)$$

□

Remark 3.19. (1) We could have also used in the proof of the above corollary [Remark 3.13](#) for the function $\beta : \mathbb{N} \rightarrow \mathbb{N}$ defined by

$$\beta(n) = Kn \div 1 \quad (3.77)$$

instead of [Corollary 3.15](#). However, this would have resulted in a much less good bound

$$h(\varepsilon, d, K) = \frac{K}{K-1} \cdot M \cdot (K^{\lceil 2d \cdot \exp(K(M+1)) \rceil - 1} - 1), \quad (3.78)$$

where $d \geq d(C)$ and $M \in \mathbb{N}, M \geq (1 + 2d)/\varepsilon$.

(2) For the special case of constant $\lambda_n = \lambda \in (0, 1)$, normed spaces, and nonexpansive functions, the exponential bound in [Corollary 3.18](#) is not optimal. In fact, [1] establishes—using an extremely complicated proof involving computer-aided calculations—an optimal quadratic bound in this special case. However, even for normed spaces and nonexpansive mappings, the bounds in the present paper and [14] are the only effective bounds known at all for the nonconstant sequences λ_n .

The next corollary strengthens Theorem 1 in [8].

COROLLARY 3.20. *Let $d, \varepsilon > 0$ and $K \in \mathbb{N}, K \geq 2$. Then, there exists an $N \in \mathbb{N}$ such that for any hyperbolic space (X, ρ, M) , any nonempty bounded convex set $C \subseteq X$ such that $d(C) \leq d$, any directionally nonexpansive mapping $f : C \rightarrow C$, any sequence $(\lambda_n)_{n \in \mathbb{N}}$ in $[1/K, 1 - 1/K]$, and any $x \in C$, the following holds:*

$$\forall n \geq N, \quad \rho(x_n, f(x_n)) \leq \varepsilon. \quad (3.79)$$

Proof. Apply [Corollary 3.18](#). □

In [9, Theorem 2.5], the first author extended (for normed spaces and nonexpansive mappings) [Corollary 3.15](#) to the situation where C is no longer required to be bounded but only the existence of a point $x^* \in C$, whose iteration sequence $(x_n^*)_{n \in \mathbb{N}}$ is bounded, is assumed. We obtained a fully uniform bound which only

depends on an upper bound d on $\|x - x^*\|$ and $\|x_n^*\|$ (and ε, K, α). This is of interest since the functional analytic embedding techniques from [4, 8] seem to require that C be bounded. Using the results above, it is easy to see that Theorem 2.5 from [9] extends to hyperbolic spaces.

THEOREM 3.21. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex subset, and $f : C \rightarrow C$ a nonexpansive mapping, and let $(\lambda_n)_{n \in \mathbb{N}}$, α , and K be as before. Let $d > 0$, $x, x^* \in C$ be such that*

$$\rho(x, x^*) \leq d \wedge \forall n, m \in \mathbb{N} (\rho(x_n^*, x_m^*) \leq d). \tag{3.80}$$

Then,

$$\forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon, \tag{3.81}$$

holds, where

$$h(\varepsilon, d, K, \alpha) := \hat{\alpha}([\!|12d \cdot \exp(K(M + 1))|\!] - 1, M), \tag{3.82}$$

$M \in \mathbb{N}$ is such that $M \geq (1 + 6d)/\varepsilon$, and $\hat{\alpha}$ as before.

Proof. It is the same as in the proof of Theorem 2.5 in [9] using Remark 3.11 and Proposition 3.5. □

For the case of directionally nonexpansive mappings, however, the additional assumption in our Theorem 3.10 causes various problems and changes in the proofs. In the following, we only consider the case where $(x_n)_{n \in \mathbb{N}}$ itself is bounded (i.e., $x = x^*$). We need an additional assumption which for the case of the constant $\lambda_k := \lambda$, though is redundant. The proof differs significantly from that given in [9] since the argument which was used there to derive the bound $\rho(x, f(x)) \leq 6d$ in the nonexpansive case does not seem to hold for directionally nonexpansive mappings. However, a different bound can be obtained depending on α .

For any $k \in \mathbb{N}$, we define the sequence $((x_k)_m)_{m \in \mathbb{N}}$ by

$$(x_k)_0 = x_k, \quad (x_k)_{m+1} = (1 - \lambda_m)(x_k)_m \oplus \lambda_m f((x_k)_m). \tag{3.83}$$

Hence, for any $k \in \mathbb{N}$, $((x_k)_m)_{m \in \mathbb{N}}$ is the Krasnosel’skiĭ-Mann iteration starting with x_k .

Remark 3.22. The sequence $((x_k)_m)_{m \in \mathbb{N}}$ is not in general a subsequence of $(x_n)_{n \in \mathbb{N}}$. But if $(\lambda_n)_{n \in \mathbb{N}}$ is a constant sequence, $\lambda_n = \lambda$, then $(x_k)_m = x_{k+m}$ for all $m, k \in \mathbb{N}$, hence $((x_k)_m)_{m \in \mathbb{N}}$ is a subsequence of $(x_n)_{n \in \mathbb{N}}$.

THEOREM 3.23. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex subset, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $[0, 1)$ which is divergent in sum and satisfies*

$$\forall n \in \mathbb{N}, \quad \lambda_n \leq 1 - \frac{1}{K}, \tag{3.84}$$

for some $K \in \mathbb{N}$.

Let $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be such that

$$\forall i, n \in \mathbb{N}, \quad (\alpha(i, n) \leq \alpha(i + 1, n)) \wedge \left(n \leq \sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s \right). \tag{3.85}$$

Let $d > 0$ and $x \in C$ such that

$$\forall n, k, m \in \mathbb{N}, \quad \rho(x_n, (x_k)_m) \leq d. \tag{3.86}$$

Then,

$$\forall \varepsilon > 0, \quad \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon \tag{3.87}$$

holds, where

$$h(\varepsilon, d, K, \alpha) := \alpha(0, 1) + \widehat{\alpha}^*([2d \cdot \alpha(0, 1) \cdot \exp(K(M + 1))] - 1, M), \tag{3.88}$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$, $\widehat{\alpha}^*(0, n) := \widetilde{\alpha}^*(0, n)$, $\widehat{\alpha}^*(i + 1, n) := \widetilde{\alpha}^*(\widehat{\alpha}^*(i, n), n)$ with $\widetilde{\alpha}^*(i, n) := i + \alpha^*(i, n)$, $\alpha^*(i, n) := \alpha(i + \alpha(0, 1), n)$ ($i, n \in \mathbb{N}$).

Proof. The sequence $(x_n)_{n \in \mathbb{N}}$ is bounded since

$$\forall m, n \in \mathbb{N}, \quad \rho(x_n, x_m) = \rho(x_n, (x_0)_m) \leq d. \tag{3.89}$$

By the hypothesis on α , we have that $\sum_{s=0}^{\alpha(0,1)-1} \lambda_s \geq 1$. From this, it is easy to see that there is $N \in \mathbb{N}$, $N \leq \alpha(0, 1) - 1$ such that

$$\lambda_N \geq \frac{1}{\alpha(0, 1)}. \tag{3.90}$$

It follows that

$$\rho(x_{\alpha(0,1)}, f(x_{\alpha(0,1)})) \leq \rho(x_N, f(x_N)) = \frac{1}{\lambda_N} \rho(x_N, x_{N+1}) \leq d \cdot \alpha(0, 1). \tag{3.91}$$

Let $\mu_n = \lambda_{\alpha(0,1)+n}$ for all $n \in \mathbb{N}$. It is obvious that $(\mu_n)_{n \in \mathbb{N}}$ is divergent in sum and $\mu_n \leq 1 - 1/K$ for all $n \in \mathbb{N}$.

Consider the sequence $(y_n)_{n \in \mathbb{N}}$ defined by

$$y_0 := y := x_{\alpha(0,1)}, \quad y_{n+1} := (1 - \mu_n)y_n \oplus \mu_n f(y_n). \quad (3.92)$$

Hence, $(y_n)_{n \in \mathbb{N}}$ is the Krasnosel'skii-Mann iteration associated with $(\mu_n)_{n \in \mathbb{N}}$, starting with $x_{\alpha(0,1)}$. It follows by an easy induction on n that

$$y_n = x_{\alpha(0,1)+n}, \quad (3.93)$$

so

$$\forall m, n \in \mathbb{N}, \quad \rho(y_n, y_m) = \rho(x_{\alpha(0,1)+n}, x_{\alpha(0,1)+m}) \leq d. \quad (3.94)$$

Thus, we can apply [Proposition 3.5](#) to get that $\lim_{n \rightarrow \infty} \rho(y_n, f(y_n)) = 0$. It follows that

$$\forall \delta > 0, \exists N_\delta, \forall n \geq N_\delta, \quad \rho(y_n, f(y_n)) < \delta. \quad (3.95)$$

Let $y^* := y_{N_\delta}$. Then, by the hypothesis,

$$\forall n \in \mathbb{N}, \quad \rho(y_n, y_n^*) = \rho(x_{\alpha(0,1)+n}, (x_{N_\delta+\alpha(0,1)})_n) \leq d. \quad (3.96)$$

Define for all $i, n \in \mathbb{N}$

$$\alpha^*(i, n) := \alpha(i + \alpha(0, 1), n). \quad (3.97)$$

It follows immediately that $\alpha^*(i, n) \leq \alpha^*(i + 1, n)$ and that

$$\sum_{s=i}^{i+\alpha^*(i,n)-1} \mu_s = \sum_{s=i}^{i+\alpha(i+\alpha(0,1),n)-1} \lambda_{\alpha(0,1)+s} = \sum_{s=i+\alpha(0,1)}^{i+\alpha(0,1)+\alpha(i+\alpha(0,1),n)-1} \lambda_s \geq n. \quad (3.98)$$

This satisfies the hypotheses of [Theorem 3.10](#) with μ_n, α^*, y, y^* instead of $\lambda_n, \alpha, x, x^*$, so we can apply it to get

$$\forall \varepsilon > 0, \forall n \geq h^*(\varepsilon, y, d, f, K, \alpha^*), \quad \rho(y_n, f(y_n)) < \rho(y^*, f(y^*)) + \varepsilon, \quad (3.99)$$

where

$$h^*(\varepsilon, y, d, f, K, \alpha^*) := \widehat{\alpha^*}([\!|2c(f, y) \cdot \exp(K(M + 1))|\!] \div 1, M), \quad (3.100)$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$ and $c(f, y) \in \mathbb{R}$ is such that $c(f, y) \geq \rho(y, f(y))$.

By (3.91), we have that

$$\rho(y, f(y)) = \rho(x_{\alpha(0,1)}, f(x_{\alpha(0,1)})) \leq d \cdot \alpha(0, 1), \quad (3.101)$$

so we can take $c(f, y) := d \cdot \alpha(0, 1)$.

We get that

$$\begin{aligned} h^*(\varepsilon, y, d, f, K, \alpha^*) &= \widehat{\alpha^*}([2d \cdot \alpha(0, 1) \cdot \exp(K(M+1))] - 1, M) \\ &= h^*(\varepsilon, d, K, \alpha) \end{aligned} \quad (3.102)$$

since α^* is defined in terms of α .

Now applying (3.95), it follows that

$$\forall \varepsilon > 0, \forall n \geq h^*(\varepsilon, d, K, \alpha), \quad \rho(y_n, f(y_n)) < \delta + \varepsilon. \quad (3.103)$$

Since (3.103) is true for every $\delta > 0$, we obtain

$$\forall \varepsilon > 0, \forall n \geq h^*(\varepsilon, d, K, \alpha), \quad \rho(y_n, f(y_n)) \leq \varepsilon, \quad (3.104)$$

that is,

$$\forall \varepsilon > 0, \forall n \geq h^*(\varepsilon, d, K, \alpha), \quad \rho(x_{\alpha(0,1)+n}, f(x_{\alpha(0,1)+n})) \leq \varepsilon. \quad (3.105)$$

Finally, letting $h(\varepsilon, d, K, \alpha) := \alpha(0, 1) + h^*(\varepsilon, d, K, \alpha)$, we get

$$\forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon. \quad (3.106)$$

□

As already mentioned, the condition

$$\forall n, k, m \in \mathbb{N}, \quad \rho(x_n, (x_k)_m) \leq d \quad (3.107)$$

is equivalent to the boundedness of (x_n) by d

$$\forall n, m \in \mathbb{N}, \quad \rho(x_n, x_m) \leq d \quad (3.108)$$

in the case of constant $\lambda_n = \lambda$. Hence, we obtain the following strong uniform version of [8, Theorem 2] (note that [8, Theorem 2] does not state any uniformity of the convergence at all).

COROLLARY 3.24. *Let (X, ρ, M) be a hyperbolic space, $C \subseteq X$ a nonempty convex subset, and $f : C \rightarrow C$ a directionally nonexpansive mapping. Let $d > 0$, $K \in \mathbb{N}$, $K \geq 2$, and $\lambda \in [1/K, 1 - 1/K]$. Let $\lambda_n := \lambda$ for all $n \in \mathbb{N}$. Let $x \in C$ such that $\rho(x_n, x_m) \leq d$ for all $m, n \in \mathbb{N}$. Then,*

$$\forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K), \quad \rho(x_n, f(x_n)) \leq \varepsilon \tag{3.109}$$

holds, where

$$h(\varepsilon, d, K) := K + K \cdot M \cdot \lceil 2d \cdot K \cdot \exp(K(M + 1)) \rceil, \quad M \in \mathbb{N}, M \geq \frac{1 + 2d}{\varepsilon}. \tag{3.110}$$

Proof. Obviously, $(\lambda_n)_{n \in \mathbb{N}}$ is divergent in sum.

Define $\alpha : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ by

$$\alpha(i, n) = Kn. \tag{3.111}$$

Then, $\alpha(i, n) = \alpha(i + 1, n) = Kn$ and

$$\sum_{s=i}^{i+\alpha(i,n)-1} \lambda_s \geq \sum_{s=i}^{i+\alpha(i,n)-1} \frac{1}{K} = \frac{1}{K} \cdot \alpha(i, n) = n. \tag{3.112}$$

It is an easy exercise to see that

$$\begin{aligned} \alpha^*(i, n) &= \alpha(i + \alpha(0, 1), n) = Kn = \alpha(i, n), \\ \widetilde{\alpha}^*(i, n) &= \widetilde{\alpha}(i, n) = i + \alpha(i, n) = i + Kn, \\ \widehat{\alpha}^*(i, n) &= \widehat{\alpha}(i, n) = K(i + 1)n. \end{aligned} \tag{3.113}$$

Since $\lambda_n = \lambda$ for all $n \in \mathbb{N}$, it follows that $(x_k)_m = x_{k+m}$, hence for all $m, n, k \in \mathbb{N}$,

$$\rho(x_n, (x_k)_m) = \rho(x_n, x_{k+m}) \leq d. \tag{3.114}$$

Hence, we can apply [Theorem 3.23](#) to obtain

$$\forall \varepsilon > 0, \forall n \geq h(\varepsilon, d, K, \alpha), \quad \rho(x_n, f(x_n)) \leq \varepsilon, \tag{3.115}$$

where $M \in \mathbb{N}$ is such that $M \geq (1 + 2d)/\varepsilon$ and

$$\begin{aligned} h(\varepsilon, d, K, \alpha) &:= \alpha(0, 1) + \widehat{\alpha}^*([2d \cdot \alpha(0, 1) \cdot \exp(K(M+1))] - 1, M) \\ &= K + K \cdot M \cdot [2d \cdot K \cdot \exp(K(M+1))] \\ &= h(\varepsilon, d, K). \quad \square \end{aligned} \tag{3.116}$$

Remark 3.25. Inspection of the proofs in this paper shows that the only places where we used the requirement (2.4) from the definition of hyperbolic spaces was in Lemma 3.8, which, in turn, was only used in Remark 3.11, as well as the proof of Theorem 3.21. Thus, all other results in this paper even hold for spaces of hyperbolic type in the sense of [4].

Acknowledgment

The research of the second author has been supported by a Marie Curie Fellowship of the European Community Programme “Improving the Human Research Potential and the Socio-economic Knowledge Base” under contract number HPTM-CT-2000-00093 at Basic Research in Computer Science (BRICS), funded by the Danish National Research Foundation.

References

- [1] J. Baillon and R. E. Bruck, *The rate of asymptotic regularity is $O(1/n)$ [$O(1/\sqrt{n})$]*, Theory and Applications of Nonlinear Operators of Accretive and Monotone Type, Lecture Notes in Pure and Appl. Math., vol. 178, Dekker, New York, 1996, pp. 51–81.
- [2] J. Borwein, S. Reich, and I. Shafrir, *Krasnosel’skiĭ-Mann iterations in normed spaces*, Canad. Math. Bull. **35** (1992), no. 1, 21–28.
- [3] M. Edelstein and R. C. O’Brien, *Nonexpansive mappings, asymptotic regularity and successive approximations*, J. London Math. Soc. (2) **17** (1978), no. 3, 547–554.
- [4] K. Goebel and W. A. Kirk, *Iteration processes for nonexpansive mappings*, Topological Methods in Nonlinear Functional Analysis (Toronto, Ont., 1982), Contemp. Math., vol. 21, American Mathematical Society, Rhode Island, 1983, pp. 115–123.
- [5] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 83, Dekker, New York, 1984.
- [6] S. Ishikawa, *Fixed points and iteration of a nonexpansive mapping in a Banach space*, Proc. Amer. Math. Soc. **59** (1976), no. 1, 65–71.
- [7] W. A. Kirk, *Krasnosel’skiĭ’s iteration process in hyperbolic space*, Numer. Funct. Anal. Optim. **4** (1981/1982), no. 4, 371–381.
- [8] ———, *Nonexpansive mappings and asymptotic regularity*, Nonlinear Anal. **40** (2000), no. 1–8, Ser. A: Theory Methods, 323–332.
- [9] U. Kohlenbach, *Uniform asymptotic regularity for Mann iterates*, to appear in J. Math. Anal. Appl.
- [10] ———, *Analysing proofs in analysis*, Logic: from Foundations to Applications (Staffordshire, 1993), Oxford Sci. Publ., Oxford University Press, New York, 1996, pp. 225–260.

- [11] ———, *Mathematically strong subsystems of analysis with low rate of growth of provably recursive functionals*, Arch. Math. Logic **36** (1996), no. 1, 31–71.
- [12] ———, *Arithmetizing proofs in analysis*, Logic Colloquium '96 (San Sebastián), Lecture Notes Logic, vol. 12, Springer, Berlin, 1998, pp. 115–158.
- [13] ———, *On the computational content of the Krasnosel'skiĭ and Ishikawa fixed point theorems*, Computability and Complexity in Analysis (Swansea, 2000), Lecture Notes in Comput. Sci., vol. 2064, Springer, Berlin, 2001, pp. 119–145.
- [14] ———, *A quantitative version of a theorem due to Borwein-Reich-Shafrir*, Numer. Funct. Anal. Optim. **22** (2001), no. 5-6, 641–656.
- [15] U. Kohlenbach and P. Oliva, *Proof mining: a systematic way of analysing proofs in mathematics*, to appear in Proc. Steklov Inst. Math.
- [16] M. A. Krasnosel'skiĭ, *Two remarks on the method of successive approximations*, Uspehi Mat. Nauk (N.S.) **10** (1955), no. 1(63), 123–127 (Russian).
- [17] S. Reich, *The alternating algorithm of von Neumann in the Hilbert ball*, Dynam. Systems Appl. **2** (1993), no. 1, 21–25.
- [18] S. Reich and I. Shafrir, *Nonexpansive iterations in hyperbolic spaces*, Nonlinear Anal. **15** (1990), no. 6, 537–558.
- [19] S. Reich and A. J. Zaslavski, *Generic aspects of metric fixed point theory*, Handbook of Metric Fixed Point Theory, Kluwer Academic Publishers, Dordrecht, 2001, pp. 557–575.
- [20] W. Takahashi, *A convexity in metric space and nonexpansive mappings. I*, Kōdai Math. Sem. Rep. **22** (1970), 142–149.

Ulrich Kohlenbach: Department of Computer Science, University of Aarhus, Ny Munkegade, DK-8000 Aarhus C, Denmark
E-mail address: kohlenb@brics.dk

Laurențiu Leuştean: National Institute for Research and Development in Informatics, 8-10 Maresal Al. Averescu Avenue, Sector 1, 71316 Bucharest, Romania
E-mail address: leo@u3.ici.ro