

A NOTE ON THE VARIATIONAL STRUCTURE OF AN ELLIPTIC SYSTEM INVOLVING CRITICAL SOBOLEV EXPONENT

MARIO ZULUAGA

Received 22 August 2002 and in revised form 10 January 2003

We consider an elliptic system involving critical growth conditions. We develop a technique of variational methods for elliptic systems. Using the well-known results of maximum principle for systems developed by Fleckinger et al. (1995), we can find positive solutions. Also, we generalize the systems results obtained (for the scalar case) by Brézis and Nirenberg (1983). Also, we give applications to biharmonic equations.

1. Introduction

In this paper, we are concerned with the existence of solutions of the elliptic system

$$\begin{aligned} -\Delta u &= \lambda u + \delta v + g_1(u, v), \\ -\Delta v &= \theta u + \gamma v + g_2(u, v) \end{aligned} \tag{1.1}$$

on Ω and $u = v = 0$ on $\partial\Omega$, where $\Omega \subset \mathbb{R}^n$, $n > 2$, is a bounded domain with the smooth boundary $\partial\Omega$, λ , δ , θ , and γ are real numbers, and g_1 , g_2 are real-valued functions with critical growth.

The purpose of this paper is to extend the results, obtained in [4], of elliptic equations for the case of only one equation (the scalar case) to the case of elliptic systems as (1.1). Our main tools are a variational approach developed for functionals with values on \mathbb{R}^2 (we want to remark that it is an important innovation in this paper), a maximum principle for systems developed in [6], and a minimax approach as in [1].

Letting

$$\begin{aligned} U &= (u, v), & -\vec{\Delta}U &= (-\Delta u, -\Delta v), \\ \mathbf{A} &= \begin{pmatrix} \lambda & \delta \\ \theta & \gamma \end{pmatrix}, & G(U) &= (g_1(u, v), g_2(u, v)), \end{aligned} \quad (1.2)$$

we can write (1.1) as

$$-\vec{\Delta}U = \mathbf{A}(U) + G(U), \quad U = \Theta = (0, 0) \quad (1.3)$$

on Ω and $U = \Theta$ on $\partial\Omega$.

It is not common to find in the literature a variational approach of problems like (1.1). Our starting point is [5] where resonance cases were considered. In that paper, the authors considered the functional

$$J_{\pm}(U) = \frac{1}{2} \int_{\Omega} [|\nabla u|^2 \pm |\nabla v|^2 - (\lambda u^2 \pm 2\delta uv + \gamma v^2)] - \int_{\Omega} F(U), \quad (1.4)$$

where $\nabla F = (g_1, g_2)$, for the study problem (1.1) in the cases

$$\mathbf{A} = \begin{pmatrix} \lambda & \delta \\ \delta & \gamma \end{pmatrix}, \quad \mathbf{A} = \begin{pmatrix} \lambda & -\delta \\ \delta & \gamma \end{pmatrix}. \quad (1.5)$$

The first case is known as cooperative problem and the second one as noncooperative problem. It is important to remark that J_{\pm} are real-valued functionals and thus it is no clear how critical points of J_{\pm} , called weak solutions in that paper, became classical solutions of (1.3). So, it is necessary to maintain the classical concept of weak solutions extended now to systems like (1.1) and then to develop a critical point theory for functionals with values on \mathbb{R}^2 .

Weak solutions of (1.1)

It is natural to define weak solutions of (1.1) as follows: $u, v \in H_0^1(\Omega)$ are weak solutions of (1.1) if, for all $\phi \in H_0^1(\Omega)$,

$$\begin{aligned} \int_{\Omega} \nabla u \nabla \phi - \lambda u \phi - \delta v \phi - g_1(u, v) \phi &= 0, \\ \int_{\Omega} \nabla v \nabla \phi - \theta u \phi - \gamma v \phi - g_2(u, v) \phi &= 0. \end{aligned} \quad (1.6)$$

The novelty here is that we can choose a functional whose critical points are weak solutions of (1.1) in the sense of (1.6). In Section 3, we

present such a *functional*. Also, we can use the regularity theory to show that weak solutions of (1.1) are classical solutions as well.

Our paper mainly focuses on the cases $g_1(u, v) = |u|^{(n+2)/(n-2)}$ and $g_2(u, v) = |v|^{(n+2)/(n-2)}$, $n > 2$, and as in [4] it is necessary to distinguish between $n = 3$ and $n \geq 4$ cases.

A superlinear case was considered in [9] where $g_1(u, v) = |u|^r$ and $g_2(u, v) = |v|^r$, $r < (n + 2)/(n - 2)$; sufficient conditions were given in that paper for the existence of positive solutions. The techniques used there was Leray-Schauder degree theory and measure theory.

2. Preliminaries and notation

We define $L^r(\Omega) = L^r(\Omega) \times L^r(\Omega)$, $r \geq 1$, and the following operations: for all $U = (u, v)$ and $\Phi = (\phi, \psi)$, we have

- (1) $U * \Phi = (u\phi, v\psi)$;
- (2) $-\vec{\Delta}U = (-\Delta u, -\Delta v)$;
- (3) $U^p = (u^p, v^p)$;
- (4) $D_i U = (D_i u, D_i v)$;
- (5) $|\nabla U|^2 = (|\nabla u|^2, |\nabla v|^2)$;
- (6) $|U| = (|u|, |v|)$;
- (7) $U \geq \Theta = (0, 0)$ if and only if $u \geq 0$ and $v \geq 0$. Also, $U > \Theta$ if and only if $u > 0$ or $v > 0$;
- (8) $\alpha U = (\alpha u, \alpha v)$ for $\alpha \in \mathbb{R}$;
- (9) $\vec{\alpha} * U = (au, bv)$ for all $\vec{\alpha} = (a, b) \in \mathbb{R}^2$;
- (10) $\|U\|_r = (\|u\|_r, \|v\|_r) \in \mathbb{R}^2$ for all $U \in L^r(\Omega)$, $r \geq 1$;
- (11) $\int_{\Omega} U = (\int_{\Omega} u, \int_{\Omega} v)$.

Pohozaev’s identity

Consider the general elliptic system

$$-\vec{\Delta}U = G(U), \quad U = \Theta \tag{2.1}$$

on Ω and $U = \Theta$ on $\partial\Omega$, where $G(U) = (g_1(u, v), g_2(u, v))$. Then we define

$$F(U) = \int_{\Theta}^U G(t) dt = \left(\int_0^u g_1(t, v) dt, \int_0^v g_2(u, t) dt \right). \tag{2.2}$$

As in the scalar case, if U is a smooth function satisfying (2.1), then it is easy to check that

$$n \int_{\Omega} F(U) + \frac{2-n}{2} \int_{\Omega} U * G(U) = \frac{1}{2} \int_{\partial\Omega} \langle X, \vec{\eta} \rangle \left(\frac{\partial U}{\partial \vec{\eta}} \right)^2, \tag{2.3}$$

where $\partial\vec{\eta}$ denotes the normal outer vector on $\partial\Omega$.

Particular cases

(a) If, for example, $G(U) = |U|^{(n+2)/(n-2)}$, $n > 2$, we see that

$$\begin{aligned} n \int_{\Omega} F(U) &= \frac{n}{2^*} \int_{\Omega} |U|^{2^*}, \\ \frac{2-n}{2} \int_{\Omega} U * G(U) &= \frac{2-n}{2} \int_{\Omega} |U|^{2^*}, \end{aligned} \tag{2.4}$$

where $2^* = 2n/(n-2)$.

Then, if Ω is *starshaped* ($\langle X, \vec{\eta} \rangle > 0$) by (2.3), we conclude that there is no positive solution of (2.1). This result is well known from 1965, see [8].

(b) If, for example, $G(U) = \mathbf{A}(U) + |U|^{(n+2)/(n-2)}$, $n > 2$, Pohozaev's identity tells us that

$$\int_{\Omega} (\lambda, \gamma) * U^2 + \frac{n+2}{2} \int_{\Omega} (\delta, \theta) * U * \tilde{U} = \frac{1}{2} \int_{\partial\Omega} \langle X, \vec{\eta} \rangle \left(\frac{\partial U}{\partial \vec{\eta}} \right)^2, \tag{2.5}$$

where $\tilde{U} = (v, u) \in \mathbf{H}_0^1(\Omega)$ and $\mathbf{H}_0^1(\Omega) = H_0^1(\Omega) \times H_0^1(\Omega)$. Identity (2.5) gives us the following negative result.

THEOREM 2.1. *Suppose that Ω is starshaped and that $\lambda, \delta, \theta, \gamma \leq 0$ or also $(\delta, \theta) \geq \Theta$ and $(\lambda, \gamma) \geq (\lambda_1, \lambda_1)$, where λ_1 is the first eigenvalue of $-\Delta$, then there is no positive solution for problem (1.1) for the case $g_1(u, v) = |u|^{(n+2)/(n-2)}$ and $g_2(u, v) = |v|^{(n+2)/(n-2)}$.*

Proof. For the first case, it is only sufficient to observe that if U is a solution of (1.1), then $U * \tilde{U} > \Theta$, which is contradictory with (2.5). In the second case, if $U > \Theta$ is a solution of

$$\begin{aligned} -\vec{\Delta}U &= \mathbf{A}(U) + |U|^{(n+2)/(n-2)} \quad \text{on } \Omega, \\ U &= \Theta \quad \text{on } \partial\Omega, \end{aligned} \tag{2.6}$$

then

$$\int_{\Omega} (-\vec{\Delta}U) * \Phi_1 = \int_{\Omega} \mathbf{A}(U) * \Phi_1 + \int_{\Omega} |U|^{(n+2)/(n-2)} * \Phi_1, \tag{2.7}$$

where $\Phi_1 = (\phi_1, \phi_1)$ and ϕ_1 is the first eigenfunction of $-\Delta$. Then we have

$$\begin{aligned} &\int_{\Omega} (\lambda_1, \lambda_1) * U * \Phi_1 \\ &= \int_{\Omega} (\lambda, \gamma) * U * \Phi_1 + \int_{\Omega} (\delta, \theta) * \tilde{U} * \Phi_1 + \int_{\Omega} |U|^{(n+2)/(n-2)} * \Phi_1. \end{aligned} \tag{2.8}$$

Since $\int_{\Omega} (\delta, \theta) * \tilde{U} * \Phi_1 \geq \Theta$, the foregoing identity tells us that $U = \Theta$, contrary to our hypothesis $U > \Theta$. This theorem extends to elliptic systems, a well-known result of [4]. \square

Now, we will borrow the ideas of a maximum principle developed in [6] to prove the following theorem.

THEOREM 2.2. *If $0 < \lambda, \gamma < \lambda_1$, $(\delta, \theta) \geq \Theta$, and $\det(\lambda_1 \mathbf{I} - \mathbf{A}) > 0$, then all nonzero solutions of*

$$\begin{aligned} -\Delta u &= \lambda u + \delta v + |u|^{(n+2)/(n-2)}, \\ -\Delta v &= \theta u + \gamma v + |v|^{(n+2)/(n-2)} \end{aligned} \tag{2.9}$$

on Ω and $u = v = 0$ on $\partial\Omega$ are no negative solutions.

Proof. In this proof, we use the arguments of maximum principle for systems developed in [6], which plays a crucial role in the proof of Theorem 3.1. Suppose that $(u, v) = U \in \mathbf{H}_0^1(\Omega)$ is a nonzero solution of foregoing system. Let $\Phi = (\phi, \psi) \in \mathbf{H}_0^1(\Omega)$ such that

$$\Phi = \max\{\Theta, -U\}. \tag{2.10}$$

If we multiply $-\vec{\Delta}U = \mathbf{A}(U) + |U|^{(n+2)/(n-2)}$ by Φ , we get

$$\begin{aligned} -\int_{\Omega} \vec{\Delta}U * \Phi &= \int_{\Omega} \nabla U * \nabla \Phi = -\int_{\Omega} |\nabla \Phi|^2 \\ &= -\int_{\Omega} (\lambda, \gamma) * |\Phi|^2 + \int_{\Omega} (\delta, \theta) * \tilde{U} * \Phi + \int_{\Omega} |U|^{(n+2)/(n-2)} * \Phi, \end{aligned} \tag{2.11}$$

which produces

$$\int_{\Omega} (\lambda_1, \lambda_1) * |\Phi|^2 \leq \int_{\Omega} |\nabla \Phi|^2 \leq \int_{\Omega} (\lambda, \gamma) * |\Phi|^2 + \int_{\Omega} (\delta, \theta) * \tilde{\Phi} * \Phi \tag{2.12}$$

and then

$$\int_{\Omega} (\lambda_1 - \lambda, \lambda_1 - \gamma) * |\Phi|^2 \leq \int_{\Omega} (\delta, \theta) * \tilde{\Phi} * \Phi. \tag{2.13}$$

From (2.13) and Cauchy-Schwarz inequality, we conclude that

$$\begin{aligned}(\lambda_1 - \lambda) \|\phi\|_{L^2}^2 &\leq \delta \|\phi\|_{L^2} \|\psi\|_{L^2}, \\(\lambda_1 - \gamma) \|\psi\|_{L^2}^2 &\leq \theta \|\phi\|_{L^2} \|\psi\|_{L^2},\end{aligned}\tag{2.14}$$

then

$$\det(\lambda_1 \mathbf{I} - \mathbf{A}) \|\phi\|_{L^2}^2 \|\psi\|_{L^2}^2 \leq 0.\tag{2.15}$$

Inequality (2.15) implies that $\phi = 0$ or $\psi = 0$ and then $\phi = 0$ and $\psi = 0$. By regularity, we conclude that $U \geq \Theta$. \square

Also, for weakly coupled cooperative elliptic systems, a maximum and strong maximum principle and its characterization have been developed by López-Gómez and Molina-Meyer in [7, Theorems 2.1 and 2.6] which can be used in Theorem 2.2 in the cooperative case.

3. Main results

For all $U = (u, v)$, $\Phi = (\varphi, \psi) \in L^2(\Omega)$. We define an \mathbb{R}^2 inner product with the following bracket:

$$[U, \Phi] = \left(\int_{\Omega} u\varphi, \int_{\Omega} v\psi \right) \in \mathbb{R}^2.\tag{3.1}$$

It is easy to check that

- (1) $[U, \Phi] = [\Phi, U]$ for any $U, \Phi \in L^2(\Omega)$;
- (2) $[U, \Phi + \Lambda] = [U, \Phi] + [U, \Lambda]$ for any $U, \Phi, \Lambda \in L^2(\Omega)$;
- (3) $[\lambda U, \Phi] = \lambda [U, \Phi] = (\int_{\Omega} \lambda u\varphi, \int_{\Omega} \lambda v\psi)$ for any $\lambda \in \mathbb{R}$ and $U, \Phi \in L^2(\Omega)$;
- (4) $[(\vec{\Delta})^{-1}U, \Phi] = [U, (\vec{\Delta})^{-1}\Phi]$ for any $U, \Phi \in L^2(\Omega)$.

The weak solutions of our problem (1.1), like we have defined in (1.6), can be represented as critical points of functional $\mathbf{J} : \mathbf{H}_0^1(\Omega) \rightarrow \mathbb{R}^2$, defined as

$$\mathbf{J}(U) = \frac{1}{2}[\nabla U, \nabla U] - \frac{1}{2}[\mathbf{C}U, U] - \int_{\Omega} F(U),\tag{3.2}$$

where $F(U) = \int_{\Theta}^U G(t)dt = (\int_0^u g_1(t, v)dt, \int_0^v g_2(u, t)dt)$, $U = (u, v) \in \mathbf{H}_0^1(\Omega)$, and

$$\mathbf{C} = \begin{pmatrix} \lambda - \delta & 2\delta \\ 2\theta & \gamma - \theta \end{pmatrix}.\tag{3.3}$$

In fact, a calculation shows that for $U = (u, v)$ and $\Phi = (\phi, \psi)$,

$$\begin{aligned} J'(U)(\Phi) &= \frac{d}{dt} J(U + t * \Phi)|_{t=0} \\ &= [\nabla U, \nabla \Phi] - \frac{1}{2} \{ [CU, \Phi] + [C\Phi, U] \} - [G(U), \Phi]. \end{aligned} \tag{3.4}$$

Now, in the case $\Phi = (\phi, \phi)$, the previous equality is transformed in

$$J'(U)(\Phi) = [\nabla U, \nabla \Phi] - [AU, \Phi] - [G(U), \Phi], \tag{3.5}$$

then critical points of J become weak solutions of (1.1).

For our main theorem, we use a Lagrange multiplier method which has been adapted to our purpose. Let

$$\begin{aligned} \vec{S}_C &= \inf_{\|U\|_{p+1} = \vec{1}} \{ [\nabla U, \nabla U] - [CU, U] \}, \\ \vec{S} &= \inf_{\|U\|_{p+1} = \vec{1}} \{ [\nabla U, \nabla U] \}, \end{aligned} \tag{3.6}$$

where $\|U\|_{p+1} = \vec{1}$ means that $\|u\|_{p+1} = 1$ and $\|v\|_{p+1} = 1$, and whether $u = 0$ then $\|v\|_{p+1} = 1$, also, or $v = 0$ then $\|u\|_{p+1} = 1$, $U = (u, v) \in L^{p+1}(\Omega)$ and $p = (n + 2)/(n - 2)$.

It is important to note that $U > \Theta$ in (3.6) because, in other case, we replace U by $|U|$.

Now we have our main theorem.

THEOREM 3.1. *Suppose that*

- (a) $0 < \lambda$ and $\gamma < \lambda_1$;
- (b) $\delta, \theta \geq 0$;
- (c) $\det(\lambda_1 I - A) > 0$;
- (d) $\lambda + \delta$ and $\gamma + \theta$ less than λ_1 .

Then the problem

$$-\vec{\Delta}U = AU + U^p, \quad U > \Theta \tag{3.7}$$

on Ω and $U = \Theta$ on $\partial\Omega$, where $p = (n + 2)/(n - 2)$, $n \geq 4$, has a weak solution.

Proof. Here we follow similar arguments to the one used in [4] for the scalar case. Let $\{U_n\} \subset H_0^1(\Omega)$ be a minimizing sequence of \vec{S}_C . Then

$$[\nabla U_n, \nabla U_n] - [CU_n, U_n] = \vec{S}_C + \overline{o(1)} \tag{3.8}$$

as $n \rightarrow \infty$.

Since $\| |U_n| \|_{p+1} = \vec{\mathbf{I}}$ for all n , then $\{U_n\}$ is bounded in $L^2(\Omega)$ and from (3.8), we conclude that $\{U_n\}$ is bounded in $\mathbf{H}_0^1(\Omega)$. So, we see that there exists $U \in \mathbf{H}_0^1(\Omega)$ and $\| |U| \|_{p+1} \leq \vec{\mathbf{I}}$ such that

- (a) $U_n \rightharpoonup U$ in $\mathbf{H}_0^1(\Omega)$;
- (b) $U_n \rightarrow U$ in $L^2(\Omega)$;
- (c) $U_n \rightarrow U$ almost everywhere.

Let $V_n = U_n - U$, then $V_n \rightharpoonup \Theta$ in $\mathbf{H}_0^1(\Omega)$, $V_n \rightarrow \Theta$ in $L^2(\Omega)$, and $V_n \rightarrow \Theta$ a.e.

Since $\mathbf{H}_0^1(\Omega) \hookrightarrow L^{p+1}(\Omega)$, $\| |U_n| \|_{p+1} = \vec{\mathbf{I}}$, and $\vec{\mathbf{S}} * \| |X| \|_{p+1}^2 \leq [\nabla X, \nabla X]$, for all $X \in \mathbf{H}_0^1(\Omega)$, we conclude, using (3.8), that

$$[CU_n, U_n] > \vec{\mathbf{S}} - \vec{\mathbf{S}}_C \tag{3.9}$$

as $n \rightarrow \infty$.

A direct calculation shows that $\vec{\mathbf{S}}_C \leq \vec{\mathbf{S}}_D$, where

$$\mathbf{D} = \begin{pmatrix} \lambda + \delta & 0 \\ 0 & \gamma + \theta \end{pmatrix}. \tag{3.10}$$

Now, by hypothesis (a) and (b), we see that $\lambda + \delta$ and $\gamma + \theta$ are greater than zero, then $\vec{\mathbf{S}}_C \leq \vec{\mathbf{S}}_D$. Also, as in [4, Lemma 1.1], since $n \geq 4$, then $\vec{\mathbf{S}}_D < \vec{\mathbf{S}}$. Then, by (3.9), we affirm that $U \neq \Theta$.

Using (3.8), we obtain

$$[\nabla U, \nabla U] + [\nabla V_n, \nabla V_n] - [CU, U] = \vec{\mathbf{S}}_C + \overrightarrow{o(1)}. \tag{3.11}$$

A well-known result, due to Brézis and Lieb [3], tells us that

$$\| |U + V_n| \|_{p+1}^{p+1} = \| |U| \|_{p+1}^{p+1} + \| |V_n| \|_{p+1}^{p+1} + \overrightarrow{o(1)}, \tag{3.12}$$

and therefore

$$\vec{\mathbf{I}} = \| |U| \|_{p+1}^{p+1} + \| |V_n| \|_{p+1}^{p+1} + \overrightarrow{o(1)}. \tag{3.13}$$

So we get

$$\vec{\mathbf{I}} \leq \| |U| \|_{p+1}^2 + \| |V_n| \|_{p+1}^2 + \overrightarrow{o(1)}. \tag{3.14}$$

By (3.14) and since $\vec{\mathbf{S}} * \| |X| \|_{p+1}^2 \leq [\nabla X, \nabla X]$ for all $X \in \mathbf{H}_0^1(\Omega)$, we have

$$\vec{\mathbf{S}} \leq \vec{\mathbf{S}} * \| |U| \|_{p+1}^2 + [\nabla V_n, \nabla V_n] + \overrightarrow{o(1)}. \tag{3.15}$$

Now, since $\vec{S} - \vec{S}_C > \Theta$ and $\|U\|_{p+1}^2 \leq \vec{1}$, we use (3.11) and (3.15) and we get

$$\begin{aligned} [\nabla U, \nabla U] - [CU, U] &= \vec{S}_C - [\nabla V_n, \nabla V_n] + \overline{o(\vec{1})} \\ &\leq \vec{S}_C + \vec{S} * \|U\|_{p+1}^2 - \vec{S} + \overline{o(\vec{1})} \\ &\leq \vec{S}_C * \|U\|_{p+1}^2. \end{aligned} \tag{3.16}$$

Inequality (3.16) states that \vec{S}_C is really achieved. Now we will use a Lagrange multiplier argument. Let

$$\begin{aligned} \mathbf{f}(V) &= \frac{1}{2}[\nabla V, \nabla V] - \frac{1}{2}[CV, V], \\ \mathbf{h}(V) &= \|V\|_{p+1}^{p+1}. \end{aligned} \tag{3.17}$$

A direct calculation shows that for all $V, \Phi \in H_0^1(\Omega)$,

$$\begin{aligned} \mathbf{f}'(V)(\Phi) &= [\nabla V, \nabla \Phi] - \frac{1}{2}\{[CV, \Phi] + [V, C\Phi]\}, \\ \mathbf{h}'(V)(\Phi) &= (p+1) \int_{\Omega} |V|^{4/(n-2)} * V * \Phi. \end{aligned} \tag{3.18}$$

Let $U \in H_0^1(\Omega)$ be the function for which \vec{S}_C is achieved, then there exists a Lagrange multiplier $\mu \in \mathbb{R}^2$ such that, for all $\Phi \in H_0^1(\Omega)$,

$$\mathbf{f}'(U)(\Phi) = \mu * \mathbf{h}'(U)(\Phi). \tag{3.19}$$

In particular, for $\Phi = U$, we get, from (3.18) and (3.19), that

$$\mu = \frac{1}{p+1} \{[\nabla U, \nabla U] - [CU, U]\} = \frac{1}{p+1} \vec{S}_C. \tag{3.20}$$

By (3.18), (3.19), and (3.20) and since we can take $U > \Theta$, we have

$$[\nabla U, \nabla \Phi] - \frac{1}{2}\{[CU, \Phi] + [U, C\Phi]\} = \vec{S}_C * \int_{\Omega} |U|^{(n+2)/(n-2)} * \Phi \tag{3.21}$$

for all $\Phi \in H_0^1(\Omega)$. In particular, if in (3.20) we take Φ as $\Phi = (\varphi, \varphi)$, (3.21) turns out

$$[\nabla U, \nabla \Phi] - [AU, \Phi] = \vec{S}_C * \int_{\Omega} |U|^{(n+2)/(n-2)} * \Phi. \tag{3.22}$$

Equality (3.21) tells us that

$$-\vec{\Delta}U = AU + \vec{S}_C * |U|^p \tag{3.23}$$

on Ω , $U = \Theta$ on $\partial\Omega$, $n \geq 4$, has a nonzero weak solution. Now, we conclude from hypothesis (d), after a direct calculation, that $\vec{S}_C > \Theta$.

Finally, from (3.22) and Theorem 2.2, it follows that $(\vec{S}_C)^{1/(p-1)} * U$ is a nonnegative weak solution of problem (3.7). \square

Remark 3.2. It is important to note that, for the case $\delta = \theta = 0$, hypothesis (d) of Theorem 3.1 is superfluous. In this case, our system is uncoupled and each equation can be handled separately, so we are in the context of [4, Theorem (1.1)].

Regularity of solutions

As in [2], if $\vec{\Delta}U + \mathbf{a}(x) * U = \Theta$ for $\mathbf{a}(x) \in L^{n/2}(\Omega)$, $n \geq 3$, then $U \in L^r(\Omega)$ for all $r > 0$. Thereafter $U \in C^\infty(\Omega)$. In our case, we are considering $\mathbf{a}(x) * U = A(U) + |U|^p$. Since $U \in L^{p+1}(\Omega)$, then $\mathbf{a}(x) \in L^{n/2}(\Omega)$.

4. A problem related to (3.7)

In this section, we deal with the problem

$$-\vec{\Delta}U = AU + |U|^{2^*-2} * U \tag{4.1}$$

on Ω , $U = \Theta$ on $\partial\Omega$, and $2^* = 2n/(n - 2)$, $n \geq 4$.

It is clear that weak solutions of (4.1) are the critical points of the functional

$$J(U) = \frac{1}{2}[\nabla U, \nabla U] - \frac{1}{2}[CU, U] - \frac{1}{2^*} \int_{\Omega} |U|^{2^*}. \tag{4.2}$$

Our main tool is the following theorem.

THEOREM 4.1. *For all $\vec{c} \in \mathbb{R}^2$ such that $\Theta < \vec{c} < (1/n)\vec{S}^{n/2}$, $n \geq 4$, the functional J satisfies the Palais-Smale condition on \vec{c} .*

Proof. Let $\{U_i\} \in H_0^1(\Omega)$ such that $J(U_i) \rightarrow \vec{c}$ and $J'(U_i) \rightarrow \Theta$, as $i \rightarrow \infty$. This is

$$\frac{1}{2}[\nabla U_i, \nabla U_i] - \frac{1}{2}[CU_i, U_i] - \frac{1}{2^*} \int_{\Omega} |U_i|^{2^*} - \vec{c} = o(\vec{1}), \tag{4.3}$$

$$[\nabla U_i, \nabla \Phi] - \frac{1}{2} \{ [CU_i, \Phi] + [U_i, C\Phi] \} - [|U_i|^{2^*-2} * U_i, \Phi] = \overrightarrow{o(1)}, \tag{4.4}$$

as $i \rightarrow \infty$, for all $U, \Phi \in H_0^1(\Omega)$. In (4.4), we put $U = \Phi = U_i$ and we get

$$[\nabla U_i, \nabla U_i] - [CU_i, U_i] - \int_{\Omega} |U_i|^{2^*} = \overrightarrow{o(1)} \tag{4.5}$$

as $i \rightarrow \infty$. Then, the left-hand side of (4.3) minus the left-hand side of (4.5) produces

$$\int_{\Omega} |U_i|^{2^*} = n\vec{c} + \overrightarrow{o(1)} \tag{4.6}$$

as $i \rightarrow \infty$.

From (4.6), we conclude that $\|U_i\|_{L^{2^*}(\Omega)}$ is bounded. Now, since $L^{2^*}(\Omega) \hookrightarrow L^2(\Omega)$ is continuous, then $\{U_i\}$ is bounded on $L^2(\Omega)$; by (4.5) we see that $\{U_i\}$ is bounded on $H_0^1(\Omega)$ as well. Therefore, let $U_0 \in H_0^1(\Omega)$ such that

$$U_i \rightharpoonup U_0 \text{ in } H_0^1(\Omega), \tag{4.7a}$$

$$U_i \rightarrow U_0 \text{ in } L^r(\Omega), 1 \leq r < 2^*, \tag{4.7b}$$

also, $U_i \rightarrow U_0$ a.e. Now, for all $\Phi \in H_0^1(\Omega)$,

$$\begin{aligned} & (J'(U_0) - J'(U_i))(\Phi) \\ &= [\nabla(U_0 - U_i), \nabla \Phi] \\ & \quad - \frac{1}{2} \{ [C(U_0 - U_i), \Phi] + [U_0 - U_i, C\Phi] \} \\ & \quad + \int_{\Omega} (|U_i|^{4/(n-2)} * U_i - |U_0|^{4/(n-2)} * U_0) * \Phi. \end{aligned} \tag{4.8}$$

From (4.7a), we see that

$$[\nabla(U_0 - U_i), \nabla \Phi] - \frac{1}{2} \{ [C(U_0 - U_i), \Phi] + [U_0 - U_i, C\Phi] \} \rightarrow \Theta \tag{4.9}$$

as $i \rightarrow \infty$. Also, from (4.7b), we have

$$\int_{\Omega} (|U_i|^{4/(n-2)} * U_i - |U_0|^{4/(n-2)} * U_0) * \Phi \rightarrow \Theta \tag{4.10}$$

as $i \rightarrow \infty$. So, from (4.8), we conclude that

$$(\mathbf{J}'(U_0) - \mathbf{J}'(U_i))(\Phi) \rightarrow \Theta \tag{4.11}$$

as $i \rightarrow \infty$. Then, by using the foregoing convergence and our hypothesis $\mathbf{J}'(U_i) \rightarrow \Theta$, we conclude that, for all $\Phi \in \mathbf{H}_0^1(\Omega)$, $\mathbf{J}'(U_0)(\Phi) = \Theta$ and therefore

$$\mathbf{J}'(U_0)(U_0) = [\nabla U_0, \nabla U_0] - [\mathbf{C}U_0, U_0] - \int_{\Omega} |U_0|^{p+1} = 0. \tag{4.12}$$

Let $V_i = U_i - U_0$, then

$$\begin{aligned} \overrightarrow{o(\mathbf{1})} &= \mathbf{J}'(U_i)(V_i) = [\nabla V_i, \nabla V_i] + [\nabla U_0, \nabla V_i] \\ &\quad - \frac{1}{2} \{ [\mathbf{C}U_i, V_i] + [U_i, \mathbf{C}V_i] \} \\ &\quad - \int_{\Omega} |U_i|^{4/(n-2)} * U_i * V_i \end{aligned} \tag{4.13}$$

as $i \rightarrow \infty$.

From (4.7a), (4.7b), (4.13), and identity (3.12), we see that, as $i \rightarrow \infty$,

$$\begin{aligned} [\nabla V_i, \nabla V_i] &= \int_{\Omega} |U_i|^{4/(n-2)} * U_i * V_i + \overrightarrow{o(\mathbf{1})} \\ &= \int_{\Omega} |U_0 + V_i|^{4/(n-2)} * (U_0 + V_i) * (U_0 + V_i - U_0) + \overrightarrow{o(\mathbf{1})} \\ &= \int_{\Omega} |U_0 + V_i|^{p+1} - \int_{\Omega} |U_0 + V_i|^{4/(n-2)} * (U_0 + V_i) * U_0 + \overrightarrow{o(\mathbf{1})} \\ &= \|U_0\|_{p+1}^{p+1} + \|V_i\|_{p+1}^{p+1} \\ &\quad - \int_{\Omega} |U_0 + V_i|^{4/(n-2)} * (U_0 + V_i) * U_0 + \overrightarrow{o(\mathbf{1})}. \end{aligned} \tag{4.14}$$

Since $V_i \rightarrow \Theta$ in $L^r(\Omega)$, for $1 \leq r < 2^*$, we get

$$[\nabla V_i, \nabla V_i] = \|V_i\|_{2^*}^{2^*} + \overrightarrow{o(\mathbf{1})}. \tag{4.15}$$

From (4.2) and (4.5), we deduce that

$$\mathbf{J}(U_i) = \frac{1}{n} \{ [\nabla U_i, \nabla U_i] - [\mathbf{C}U_i, U_i] \} + \overrightarrow{o(\mathbf{1})} \tag{4.16}$$

as $i \rightarrow \infty$.

Now, since $V_i \rightarrow \Theta$ in $L^2(\Omega)$ and $V_i \rightharpoonup \Theta$ in $H_0^1(\Omega)$, from (4.16), we obtain

$$J(U_i) = \frac{1}{n} \{ [\nabla U_0, \nabla U_0] - [CU_0, U_0] \} + \frac{1}{n} [\nabla V_i, \nabla V_i] + \overrightarrow{o(1)} \quad (4.17)$$

as $i \rightarrow \infty$.

Now, from (4.12), we see that $[\nabla U_0, \nabla U_0] - [CU_0, U_0] \geq \Theta$, then, by (4.17), we have

$$[\nabla V_i, \nabla V_i] \leq nJ(U_i) + \overrightarrow{o(1)} \quad (4.18)$$

as $i \rightarrow \infty$.

The foregoing inequality, united with our hypothesis $J(U_i) \rightarrow \vec{c} \in (\Theta, (1/n)\vec{S}^{n/2})$, produces

$$[\nabla V_i, \nabla V_i] \leq \text{const} < \vec{S}^{n/2} \quad (4.19)$$

for all i large enough.

It is important to note that $[\nabla \Phi, \nabla \Phi] = \|\Phi\|_{H_0^1(\Omega)}^2$ and $\Phi \in H_0^1(\Omega)$. Now, since $H_0^1(\Omega) \hookrightarrow L^{2^*}(\Omega)$ and

$$\sqrt{\vec{S}} \|\Phi\|_{L^{2^*}(\Omega)} \leq \|\Phi\|_{H_0^1(\Omega)}, \quad (4.20)$$

we deduce from (4.15) that

$$\|V_i\|_{H_0^1(\Omega)}^2 \left(\vec{S}^{2^*/2} - \|V_i\|_{H_0^1(\Omega)}^{2^*-2} \right) \leq \overrightarrow{o(1)} \quad (4.21)$$

as $i \rightarrow \infty$. Now, from (4.19) and since $n \geq 4$, we deduce that $\vec{S}^{2^*/2} - \|V_i\|_{H_0^1(\Omega)}^{2^*-2}$ is greater than a positive constant for i large enough. Then from (4.21), we conclude that $V_i \rightarrow \Theta$. \square

Now, we are ready for the following theorem.

THEOREM 4.2. *Suppose that*

- (e) $\lambda + \delta, \gamma + \theta < \lambda_1$;
- (f) $\lambda + \delta, \gamma + \theta > 0$.

Then problem (4.1) has at least a nonzero solution.

Proof. First, we conclude, from hypothesis (e), that $\vec{S}_C > \Theta$, then for all $U_0 = (u_0, v_0) \in H_0^1(\Omega)$, $\|U_0\|_{L^{2^*}(\Omega)}^{2^*} = \vec{1}$, we have

$$M = [\nabla U_0, \nabla U_0] - [CU_0, U_0] > \vec{S}_C > \Theta. \quad (4.22)$$

Now, we show that, for all $U_0 = (u_0, v_0) \in \mathbf{H}_0^1(\Omega)$ and $\|U_0\|_{\mathbf{L}^{2^*}(\Omega)}^2 = \vec{\mathbf{I}}$,

$$\frac{1}{n} \vec{\mathbf{S}}_C^{n/2} \leq \sup_{\vec{\mathbf{t}} > \Theta} \mathbf{J}(\vec{\mathbf{t}} * U_0) \leq \frac{1}{n} \vec{\mathbf{S}}^{1/n}. \tag{4.23}$$

In fact, let

$$\varphi(\vec{\mathbf{t}}) = \mathbf{J}(\vec{\mathbf{t}} * U_0) = \frac{1}{2} \mathbf{M} * \vec{\mathbf{t}}^2 - \frac{1}{2^*} \vec{\mathbf{t}}^{2^*}, \quad \vec{\mathbf{t}} \in \mathbb{R}^2. \tag{4.24}$$

It is clear that $\varphi(\vec{\mathbf{t}}) > \Theta$, for $\vec{\mathbf{t}} > \Theta$ small enough and $\lim_{t \rightarrow \infty} \varphi(\vec{\mathbf{t}}) \rightarrow -\infty$. Now, by a direct calculation, we see that, for $\vec{\mathbf{t}} = \mathbf{M}^{(n-2)/4}$, $\varphi(\vec{\mathbf{t}})$ reaches its *maximum* value. Then, from (4.22), we get

$$\sup_{\vec{\mathbf{t}} > \Theta} \varphi(\vec{\mathbf{t}}) = \frac{1}{n} \mathbf{M}^{n/2} > \frac{1}{n} \vec{\mathbf{S}}_C^{n/2}. \tag{4.25}$$

Finally, from our hypothesis (f), we deduce that $\vec{\mathbf{S}}_C < \vec{\mathbf{S}}$, then

$$\sup_{\vec{\mathbf{t}} > \Theta} \varphi(\vec{\mathbf{t}}) < \frac{1}{n} \vec{\mathbf{S}}^{n/2}. \tag{4.26}$$

Therefore, (4.25) and (4.26) produce (4.23). Now, by Theorem 4.1, $\mathbf{J}(\Theta) = \Theta$, $\varphi(\vec{\mathbf{t}}) \rightarrow -\infty$, and (4.23), we apply the mountain pass lemma of Ambrosetti and Rabinowitz [1], adapted in this case to our functional \mathbf{J} , to prove the existence of $\vec{\mathbf{c}} \in \mathbb{R}^2$ such that

$$\vec{\mathbf{c}} = \inf \sup \mathbf{J}(U) \in \left(\Theta, \frac{1}{n} \vec{\mathbf{S}}^{n/2} \right), \tag{4.27}$$

where inf and sup are taken on suitable sets and $\mathbf{J}^{-1}(\vec{\mathbf{c}}) \neq \Phi$. Therefore, $U \in \mathbf{J}^{-1}(\vec{\mathbf{c}})$ is a nonzero weak solution of (4.1). □

The biharmonic equation

The results of foregoing theorem can be applied to the following class of nonlinear biharmonic equation under Navier-Dirichlet boundary conditions:

$$\Delta^2 u = \theta u + |u|^{2^*-2} u \tag{4.28}$$

on Ω , $u = \Delta u = 0$ on $\partial\Omega$, and $\theta > 0$. Indeed, $U = (u, v)$ with $v = -\Delta u$ satisfies the problem

$$-\vec{\Delta} U = \mathbf{A}U + \mathbf{G}(U) \tag{4.29}$$

on Ω and $U = \Theta$ on $\partial\Omega$, where

$$\mathbf{A} = \begin{pmatrix} 0 & 1 \\ \theta & 0 \end{pmatrix}, \quad G(U) = (|u|^{2^*-2}u, 0). \quad (4.30)$$

As a direct application of Theorem 4.2, we obtain the following theorem.

THEOREM 4.3. *Suppose that*

- (g) $\theta, 1 < \lambda_1$;
- (h) $\theta > 0$ (this is a cooperative case).

Then problem (4.28) has a weak solution $u \in H^3(\Omega) \cap H_0^1(\Omega)$ with $\Delta u \in H_0^1(\Omega)$.

References

- [1] A. Ambrosetti and P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, *J. Functional Analysis* **14** (1973), 349–381.
- [2] H. Brézis and T. Kato, *Remarks on the Schrödinger operator with singular complex potentials*, *J. Math. Pures Appl.* (9) **58** (1979), no. 2, 137–151.
- [3] H. Brézis and E. Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, *Proc. Amer. Math. Soc.* **88** (1983), no. 3, 486–490.
- [4] H. Brézis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, *Comm. Pure Appl. Math.* **36** (1983), no. 4, 437–477.
- [5] D. G. Costa and C. A. Magalhães, *A variational approach to subquadratic perturbations of elliptic systems*, *J. Differential Equations* **111** (1994), no. 1, 103–122.
- [6] J. Fleckinger, J. Hernández, and F. de Thélin, *On maximum principles and existence of positive solutions for some cooperative elliptic systems*, *Differential Integral Equations* **8** (1995), no. 1, 69–85.
- [7] J. López-Gómez and M. Molina-Meyer, *The maximum principle for cooperative weakly coupled elliptic systems and some applications*, *Differential Integral Equations* **7** (1994), no. 2, 383–398.
- [8] S. Pohozaev, *Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$* , *Soviet Math. Dokl.* **6** (1965), 1408–1411, translated from Russian *Dokl. Akad. Nauk SSSR*, **165** (1965), 33–36.
- [9] M. Zuluaga, *Nonzero and positive solutions of a superlinear elliptic system*, *Arch. Math. (Brno)* **37** (2001), no. 1, 63–70.

Mario Zuluaga: Departamento de Matemáticas, Universidad Nacional de Colombia, Bogota, Colombia

E-mail address: mzuluaga@matematicas.unal.edu.co

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie