

ON THE GLOBAL SOLVABILITY OF SOLUTIONS TO A QUASILINEAR WAVE EQUATION WITH LOCALIZED DAMPING AND SOURCE TERMS

E. CABANILLAS LAPA, Z. HUARINGA SEGURA, AND F. LEON BARBOZA

Received 18 July 2004 and in revised form 9 March 2005

We prove existence and uniform stability of strong solutions to a quasilinear wave equation with a locally distributed nonlinear dissipation with source term of power nonlinearity of the type $u'' - M(\int_{\Omega} |\nabla u|^2 dx) \Delta u + a(x)g(u') + f(u) = 0$, in $\Omega \times]0, +\infty[$, $u = 0$, on $\Gamma \times]0, +\infty[$, $u(x, 0) = u_0(x)$, $u'(x, 0) = u_1(x)$, in Ω .

1. Introduction

Let Ω be a bounded domain of \mathbb{R}^N with a smooth boundary $\Gamma = \partial\Omega$. We consider the initial-boundary value problem

$$\begin{aligned} u'' - M\left(\int_{\Omega} |\nabla u|^2 dx\right) \Delta u + a(x)g(u') + f(u) &= 0, & \text{in } \Omega \times]0, +\infty[, \\ u &= 0, & \text{on } \Gamma \times]0, +\infty[, \\ u(x, 0) = u_0(x), \quad u'(x, 0) = u_1(x), & \text{in } \Omega, \end{aligned} \tag{1.1}$$

where $M(s)$ is a C^1 -class function on $[0, +\infty[$ satisfying $M(s) \geq m_0 > 0$, for $s \geq 0$, with m_0 constant, a is a smooth nonnegative function but vanishes somewhere in $\overline{\Omega}$, $f(u)$ is a nonlinear term like $f(u) \sim -|u|^\alpha u$, and g is a real-valued function.

The problem (1.1), when $M(s) = 1$ and f is some type of nonlinear function, has been studied by Zuazua [10] and Nakao [9]. Recently, Cabanillas et al. have treated in [2, 3] a more delicate case where M is not a constant function ($f(u) = 0, -h_0 u$). Kouemou-Patcheu [6] investigated the case $M(s) = a_0 + bs$ with $a(x) = 1$ in Ω and $f(u) = 0$. We fix $x^0 \in \mathbb{R}^N$ and we set

$$m(x) = x - x^0, \quad R = \sup \{ |m(x)|; x \in \Omega \}, \quad \Gamma_0 = \{x \in \Gamma; m(x) \cdot \nu(x) > 0\}, \tag{1.2}$$

where $\nu(x)$ denotes the outward unit normal at $x \in \Gamma$. Let $a = a(x)$ be a smooth nonnegative function such that

$$a(x) \geq a_0 > 0, \quad \text{a.e. in } \omega, \tag{1.3}$$

where ω is a neighborhood of Γ_0 and a_0 is a positive constant. By neighborhood of Γ_0 , we actually mean the intersection of Ω and a neighborhood of Γ_0 .

The goal of this work is to obtain global existence and decay estimates of the strong solutions of the quasilinear wave equation (1.1) when M is not a constant function, the function a satisfies (1.3), g is a C^1 , odd, increasing function, and $f(u) \sim -|u|^\alpha u$.

2. Preliminaries and main result

Throughout this paper, the functions considered are all real valued and the notations for their norm are adopted as usual (e.g., Lions [7]).

We consider the following general hypotheses.

(A.1) Assumptions on M :

$$M \in C^1([0, +\infty[), \quad M(s) \geq m_0 > 0, \quad \forall s \geq 0, \tag{2.1}$$

$$|M'(s)| \leq \beta s^{\gamma/2}, \quad \forall s \geq 0 \tag{2.2}$$

for some constants $\beta \geq 0, \gamma \geq 0$.

(A.2) Assumptions on a :

$$a \in C^2(\Omega) \cap C(\overline{\Omega}), \quad |\Delta a(x)| \leq a_1 a(x), \quad a_1 > 0. \tag{2.3}$$

(A.3) Assumptions on f :

f is a C^1 -class function on \mathbb{R} and satisfies

$$|f(u)| \leq h_0 |u|^{\alpha+1}, \quad |f'(u)| \leq h_0 |u|^\alpha, \quad \forall u \in \mathbb{R}, \tag{2.4}$$

with some constant $h_0 > 0$ and

$$0 < \alpha < \frac{2}{(N-4)^+}, \tag{2.5}$$

where $(N-4)^+ = \max\{N-4, 0\}$.

(A.4) g is a C^1 odd increasing function and

$$C_1 |s| \leq |g(s)| \leq C_2 |s|^q \quad \text{if } |s| \geq 1 \text{ with } 1 \leq q \leq \frac{2}{(N-4)^+}, \tag{2.6}$$

$$C_3 |s|^{p+1} \leq g(s)s \quad \text{if } |s| < 1, 1 \leq p < +\infty,$$

where $C_i, i = 1, 2, 3$, are positive constants.

We have the following fundamental inequalities.

LEMMA 2.1 (Sobolev-Poincaré inequality). *Let α be a number with $0 \leq \alpha < \infty$ ($N = 1, 2$) or $0 \leq \alpha \leq 4/(N-2)$ ($N \geq 3$), then there is a constant $C_* > 0$ such that*

$$|u|_{\alpha+2} \leq C_* |\nabla u|_2 \quad \text{for } u \in H_0^1(\Omega). \tag{2.7}$$

LEMMA 2.2 (Gagliardo-Nirenberg inequality). *Let $1 \leq r < q \leq +\infty$ and $p \leq q$. Then, the inequality*

$$|u|_{W^{k,q}} \leq C |u|_{W^{m,p}}^\theta |u|_r^{1-\theta} \quad \text{for } u \in W^{m,p} \cap L^r \tag{2.8}$$

holds with some $C > 0$ and

$$\theta = \left(\frac{k}{N} + \frac{1}{r} - \frac{1}{q}\right) \left(\frac{m}{N} + \frac{1}{r} - \frac{1}{p}\right)^{-1} \tag{2.9}$$

provided that $0 < \theta \leq 1$ (assume that $0 < \theta < 1$ if $q = +\infty$).

LEMMA 2.3. Let $E : [0, +\infty[\rightarrow [0, +\infty[$ be a nonincreasing function and assume that there are two constants $p \geq 1$ and $A > 0$ such that

$$\int_S^{+\infty} E^{(p+1)/2}(t) dt \leq AE(S), \quad 0 \leq S < +\infty. \tag{2.10}$$

Then,

$$E(t) \leq \begin{cases} CE(0)e^{-\lambda t}, & \forall t \geq 0 \text{ if } p = 1, \\ CE(0)(1+t)^{-2/(p-1)}, & \forall t \geq 0 \text{ if } p > 1, \end{cases} \tag{2.11}$$

where C and λ are positive constants independent of the initial energy $E(0)$.

We will construct a stable set in $H_0^1 \cap H^2$. For this, we define the functionals

$$\begin{aligned} J(u) &= \frac{1}{2} \widetilde{M}(|\nabla u|^2) + \int_{\Omega} F(u) dx, \quad \text{for } u \in H_0^1, \\ I(u) &= M(|\nabla u|^2) |\nabla u|^2 + \int_{\Omega} f(u) \cdot u dx, \quad \text{for } u \in H_0^1, \\ E(u, v) &= \frac{1}{2} |v|^2 + J(u), \quad \text{for } (u, v) \in H_0^1 \times L^2, \end{aligned} \tag{2.12}$$

where

$$\widetilde{M}(s) = \int_0^s M(\xi) d\xi, \quad F(\lambda) = \int_0^\lambda f(s) ds. \tag{2.13}$$

LEMMA 2.4. Let $0 < \alpha < 4/(N - 4)^+$. Then, for any $K > 0$, there exists a number $\varepsilon_0 = \varepsilon_0(K)$ such that if $|\Delta u| \leq K$ and $|\nabla u| \leq \varepsilon_0$,

$$J(u) \geq \frac{m_0}{4} |\nabla u|^2, \quad I(u) \geq \frac{m_0}{2} |\nabla u|^2. \tag{2.14}$$

Proof. By the Gagliardo-Nirenberg inequality, we deduce that

$$|u|_{\alpha+2}^{\alpha+2} \leq C |u|_{2N/(N-2)}^{(\alpha+2)(1-\theta)} |\Delta u|^{(\alpha+2)\theta} \leq C |\nabla u|^{(\alpha+2)(1-\theta)} |\Delta u|^{(\alpha+2)\theta} \tag{2.15}$$

with

$$\theta = \left(\frac{N-2}{2N} - \frac{1}{\alpha+2} \right)^+ \left(\frac{2}{N} + \frac{N-2}{2N} - \frac{1}{2} \right)^{-1} = \left(\frac{(N-2)\alpha-4}{2(\alpha+2)} \right)^+ \leq 1. \tag{2.16}$$

Here, we note that

$$(\alpha+2)(1-\theta)-2 = \begin{cases} \alpha & \text{if } 0 < \alpha < \frac{4}{N-2} \\ & (0 < \alpha < +\infty, \text{ for } N = 1, 2), \\ \frac{(4-N)\alpha+4}{2} & \text{if } \frac{4}{N-2} < \alpha < \frac{4}{N-4} \\ & \left(\frac{4}{N-2} < \alpha < +\infty, N = 3, 4 \right). \end{cases} \tag{2.17}$$

Hence, if $|\Delta u| \leq K$, we get

$$\begin{aligned} J(u) &\geq \frac{m_0}{2} |\nabla u|^2 - \frac{h_0}{\alpha+2} |u|^{\alpha+2} \\ &\geq \frac{m_0}{2} |\nabla u|^2 - Ch_0 |\nabla u|^{(\alpha+2)(1-\theta)} |\Delta u|^{(\alpha+2)\theta} \\ &\geq \left\{ \frac{m_0}{2} - Ch_0 K^{(\alpha+2)\theta} |\nabla u|^{(\alpha+2)(1-\theta)-2} \right\} |\nabla u|^2. \end{aligned} \tag{2.18}$$

Using (2.17), we can define $\varepsilon_0 = \varepsilon_0(K)$ by

$$CK^{(\alpha+2)\theta} \varepsilon_0^{(\alpha+2)(1-\theta)-2} \leq \frac{m_0}{4}. \tag{2.19}$$

Thus, we obtain

$$J(u) \geq \frac{m_0}{4} |\nabla u|^2 \tag{2.20}$$

if $|\nabla u| \leq \varepsilon_0$. In a completely analogous way, we can get (2.14) for $I(u)$. □

We define our stable set W_K by

$$W_K = \left\{ (u, v) \in (H_0^1 \cap H^2) \times H_0^1 : |\Delta u| < K, |\nabla v| < K, \sqrt{4m_0^{-1}E(u_0, v_0)} < \varepsilon_0 \right\} \tag{2.21}$$

for $K > 0$.

Remark 2.5. If we consider $f(u) \cdot u \geq 0$, then we need not take $\varepsilon_0(K)$, and W_K is replaced by

$$\widetilde{W}_K = \{ (u, v) \in (H_0^1 \cap H^2) \times H_0^1 : |\Delta u| < K, |\nabla v| < K \}. \tag{2.22}$$

3. Statement of the results

In this section, we will state our main theorem.

THEOREM 3.1 (local existence). *Let initial data $\{u_0, u_1\}$ belong to $(H_0^1 \cap H^2) \times H_0^1$ and let the assumptions (A.1)–(A.4) be fulfilled. Then there exists a unique local solution u of (1.1) belonging to*

$$C_w^0([0, T[; H_0^1 \cap H^2) \cap C_w^1([0, T[; H_0^1) \cap C^0([0, T[; H_0^1) \cap C^1([0, T[; L^2(\Omega)) \quad (3.1)$$

for some $T = T(|\Delta u_0|, |\nabla u_1|) > 0$.

Moreover, at least one of the following statements is valid:

- (i) $T = +\infty$,
- (ii) $|\nabla u'(t)|^2 + |\Delta u(t)|^2 \rightarrow \infty$ as $t \rightarrow T^-$,
- (iii) $M(|\nabla u(t)|^2) \rightarrow 0$ as $t \rightarrow T^-$.

The proof of this theorem is well known.

THEOREM 3.2 (global existence and decay property). *Suppose (A.1)–(A.4) hold. Then there exists an open set S_0 in $(H_0^1 \cap H^2) \times H_0^1$, which contains $(0, 0)$ such that if $(u_0, u_1) \in S_0$, the problem (1.1) admits a unique global solution $u(t)$ on the class*

$$L^\infty([0, +\infty[; H_0^1 \cap H^2) \cap W^{1,\infty}([0, +\infty[; H_0^1) \cap W^{2,\infty}([0, +\infty[; L^2). \quad (3.2)$$

Moreover, the energy determined by the solution u has the decay states

$$\begin{aligned} E(u(t), u'(t)) &\leq C_0 e^{-\lambda t} \quad \text{if } p = 1, \\ E(u(t), u'(t)) &\leq \tilde{C}_0 (1+t)^{-2/(p-1)} \quad \text{if } p > 1, \end{aligned} \quad (3.3)$$

where C_0 , \tilde{C}_0 , and λ are certain positive constants depending on $|\nabla u_0|$, $|u_1|$, and other quantities.

Proof. We divide the proof into several lemmas. For the moment, we denote $E(u(t), u'(t))$ by $E(t)$.

LEMMA 3.3. *Let $u(t)$ be a local solution to the problem (1.1) on $[0, T[$, $T > 0$. Then*

$$\forall 0 \leq S \leq T < +\infty, \quad E(S) - E(T) = \int_S^T \int_\Omega a(x) u' g(u') dx dt. \quad (3.4)$$

Multiplying the equation in (1.1) by $u'(t)$ and integrating on $[S, T[$, we get

$$\begin{aligned} - \int_S^T \int_\Omega a(x) u' g(u') dx dt &= \left[\frac{1}{2} |u'(t)|^2 + \frac{1}{2} \tilde{M}(|\nabla u(t)|^2) + F(u(t)) \right]_S^T \\ &= E(T) - E(S). \end{aligned} \quad (3.5)$$

It is easy to see the identity

$$E'(t) = - \int_\Omega a u' g(u') dx \leq 0. \quad (3.6)$$

In particular, $E(t)$ is nonincreasing and

$$E(t) \leq E(0) \tag{3.7}$$

as long as the local solutions exist. □

LEMMA 3.4. *Let $u(t)$ be a local solution to the problem (1.1) satisfying $(u(t), u'(t)) \in W_K$ on $[0, T[$ for some $K > 0$. Then,*

$$E(t) \leq \begin{cases} CI_0 e^{-\lambda t}, & \text{on } [0, T[\quad \text{if } p = 1, \\ \hat{q}(1+t)^{-2/(p-1)}, & \text{if } p > 1, \end{cases} \tag{3.8}$$

where $I_0^2 = E(0)$, $\lambda = \lambda(K, I_0)$, and $\hat{q} = \hat{q}(K, I_0)$ denote certain positive constants continuously depending on K and I_0 .

The proof of this lemma is based on the following identities given by the multiplier method. We omit to write the differential elements in the integrals, in order to simplify the expressions.

LEMMA 3.5. *Let $q \in [W^{1,\infty}(\Omega)]^N$, $\beta \in \mathbb{R}$, and $\xi \in W^{1,\infty}(\Omega)$. Then*

$$\begin{aligned} & \int_S^T \int_\Gamma M(|\nabla u(t)|^2) q \cdot \nu \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma \\ &= (u', 2q \cdot \nabla u + \beta u) E^\sigma \Big|_S^T + \int_S^T \int_\Omega (\operatorname{div}(q) - \beta) [|u'|^2 - M(|\nabla u(t)|^2) |\nabla u(t)|^2 E^\sigma] \\ &+ 2 \int_S^T \int_\Omega M(|\nabla u(t)|^2) \frac{\partial q_k}{\partial x_i} \frac{\partial u}{\partial x_k} \frac{\partial u}{\partial x_i} E^\sigma - \sigma \int_S^T \int_\Omega u' (2q \cdot \nabla u + \beta u) E^{\sigma-1} E' \\ &+ \int_S^T \int_\Omega \operatorname{ag}(u') (2q \cdot \nabla u + \beta u) E^\sigma - \int_S^T \int_\Omega \operatorname{div}(q) F(u) E^\sigma + \beta \int_S^T \int_\Omega f(u) \cdot u E^\sigma, \end{aligned} \tag{3.9}$$

$$\begin{aligned} & (u', \xi u) E^\sigma \Big|_S^T + \int_S^T \int_\Omega \xi [M(|\nabla u|^2) |\nabla u|^2 - |u'|^2] E^\sigma \\ & - \sigma \int_S^T \int_\Omega u' u \xi E^{\sigma-1} E' + \int_S^T \int_\Omega M(|\nabla u(t)|^2) (\nabla u, u \nabla \xi) E^\sigma \\ & + \int_S^T \int_\Omega \operatorname{ag}(u') \xi u E^\sigma + \int_S^T \int_\Omega \xi f(u) u E^\sigma = 0. \end{aligned} \tag{3.10}$$

For the proof, see Lions [8] or Komornik [5].

Proof of Lemma 3.5. We proceed in several steps.

Step 1. Applying (3.9) with $q(x) = m(x)$, observing that $\operatorname{div} q = N$, we obtain

$$\begin{aligned}
 & (u', 2m \cdot \nabla u + \beta u)E^\sigma \Big|_S^T + (N - \beta) \int_S^T \int_\Omega |u'|^2 E^\sigma \\
 & + (\beta - N + 2) \int_S^T \int_\Omega M(|\nabla u|^2) |\nabla u|^2 E^\sigma - \sigma \int_S^T \int_\Omega u' (2m \cdot \nabla u + \beta u) E^{\sigma-1} E' \\
 & + \int_S^T \int_\Omega ag(u')(2m \cdot \nabla u + \beta u)E^\sigma + \beta \int_S^T \int_\Omega f(u) \cdot uE^\sigma - N \int_S^T \int_\Omega F(u)E^\sigma \\
 & = \int_S^T \int_\Gamma M(|\nabla u|^2) \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma \leq RM_0 \int_S^T \int_{\Gamma_0} \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma,
 \end{aligned} \tag{3.11}$$

where

$$m_0 \leq M(|\nabla u|^2) \leq \max \left\{ M(s), 0 \leq s \leq \frac{4E(0)}{m_0} \right\} \equiv M_0. \tag{3.12}$$

Throughout the remaining part of this work, positive constants will be denoted by C and will change line to line. Here, we observe that under the assumption $(u(t), u'(t)) \in W_K$, the functionals $E(t)$, $e(t) = (1/2)(|u'(t)|^2 + |\nabla u(t)|^2)$ and $|u'(t)|^2 + I(u(t))$ are all equivalent, by Lemma 2.4.

We take $\beta \in]N - 2, N[$ and $\theta_0 = \min\{2(N - \beta), \beta - N + 2\}$, we deduce that

$$\begin{aligned}
 \theta_0 \int_S^T E^{\sigma+1} & \leq RM_0 \int_S^T \int_{\Gamma_0} \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma + \left| (u', 2m \cdot \nabla u + \beta u) \Big|_S^T \right| \\
 & + \left| \int_S^T \int_\Omega ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| \\
 & + \sigma \left| \int_S^T \int_\Omega u' (2m \cdot \nabla u + \beta u) E^{\sigma-1} E' \right| \\
 & + \beta \left| \int_S^T \int_\Omega f(u) \cdot uE^\sigma \right| + N \left| \int_S^T \int_\Omega F(u)E^\sigma \right|.
 \end{aligned} \tag{3.13}$$

Since the energy is nonincreasing, using the result of Komornik [5], we find that

$$\left| (u', 2m \cdot \nabla u + \beta u) \Big|_S^T E^\sigma \right| \leq CE(S), \tag{3.14}$$

$$\left| \sigma \int_S^T \int_\Omega u' (2m \cdot \nabla u + \beta u) E^{\sigma-1} E' \right| \leq CE(S). \tag{3.15}$$

By the Hölder inequality, we have

$$\begin{aligned} \left| \int_S^T \int_{\Omega_1} ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| &\leq \int_S^T E^\sigma \left(\int_{\Omega_1} a^2 g^2(u') \right)^{1/2} \left(\int_{\Omega_1} |2m \cdot \nabla u + \beta u| \right)^{1/2} \\ &\leq \int_S^T E^\sigma \left(\int_{\Omega_1} a^2 (u' g(u'))^{2/(p+1)} \right)^{1/2} E^{1/2} \\ &\leq C \int_S^T E^{\sigma+1/2} |E'|^{1/(p+1)}, \end{aligned} \tag{3.16}$$

$$\begin{aligned} \left| \int_S^T \int_{\Omega_2} ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| &\leq C \int_S^T E^\sigma \left\{ \int_{\Omega_2} |ag(u')| |\nabla u| + \int_{\Omega_2} |ag(u')| |u| \right\} \\ &\leq C \int_S^T E^\sigma |ag(u')|_{1+q^{-1}} \{ |\nabla u|_{q+1} + |u| \}. \end{aligned} \tag{3.17}$$

We observe here, from Lemma 2.2, that

$$|\nabla u|_{q+1} \leq C |\nabla u|^{1/(q+1)} |\Delta u|^{q/(q+1)} \leq CK^{q/(q+1)} E^{1/2(q+1)}, \tag{3.18}$$

$$\begin{aligned} |ag(u')|_{1+q^{-1}} &= \left[\int_{\Omega_2} |ag(u')|^{(q+1)/q} \right]^{q/(q+1)} \\ &\leq C \left[\int_{\Omega_2} au'g(u') \right]^{q/(q+1)} \leq C |E'|^{q/(q+1)}. \end{aligned} \tag{3.19}$$

From (3.17), (3.18), and (3.19), we have

$$\left| \int_S^T \int_{\Omega_2} ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| \leq C \int_S^T |E|^{\sigma+1/2(q+1)} |E'|^{q/(q+1)}, \tag{3.20}$$

where we set for each $t \geq 0$,

$$\Omega_1 = \Omega_1(t) = \{x \in \Omega : |u'(x, t)| \leq 1\}, \quad \Omega_2 = \Omega \setminus \Omega_1. \tag{3.21}$$

Thus, from (3.16) and (3.20), we get

$$\left| \int_S^T \int_{\Omega} ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| \leq C \int_S^T \left[E^{\sigma+1/2} |E'|^{1/(p+1)} + |E|^{\sigma+1/2(q+1)} |E'|^{q/(q+1)} \right]. \tag{3.22}$$

Now, using the Young inequality, we obtain

$$\left| \int_S^T \int_{\Omega} ag(u')(2m \cdot \nabla u + \beta u)E^\sigma \right| \leq \varepsilon \int_S^T E^{\sigma+1} + E(S), \quad \varepsilon > 0. \tag{3.23}$$

It follows from (3.14), (3.15), and (3.23) that

$$\frac{\theta_0}{2} \int_S^T E^{\sigma+1} \leq CE(S) + \int_S^T \int_{\Omega} \left(\frac{\beta^2}{\theta_0} |f(u)|^2 + NF(u) \right) E^{\sigma} + RM_0 \int_S^T \int_{\Gamma_0} \left| \frac{\partial u}{\partial \nu} \right|^2 E^{\sigma}. \quad (3.24)$$

To estimate the last term in (3.24), we utilize (3.10) with $\xi = \eta$, where $\eta \in W^{1,\infty}(\Omega)$ is a function (constructed by Zuazua in [10]) which satisfies

$$\begin{aligned} 0 &\leq \eta \leq 1, \\ \eta &= 1, \quad \text{in } \hat{\omega}, \quad \frac{|\nabla \eta|^2}{\eta} \in L^{\infty}(\omega), \\ \eta &= 0, \quad \text{in } \Omega \setminus \omega, \end{aligned} \quad (3.25)$$

and $\hat{\omega}$ is an open set in Ω , with $\Gamma_0 \subseteq \hat{\omega} \subsetneq \omega$.

First, we have from (3.10)

$$\begin{aligned} \int_S^T \int_{\Omega} \eta M(|\nabla u|^2) |\nabla u|^2 E^{\sigma} &= (-u', \eta u) E^{\sigma} \Big|_S^T - \int_S^T \int_{\Omega} ag(u') u \eta E^{\sigma} \\ &\quad - \int_S^T \int_{\Omega} M(|\nabla u|^2) \nabla u \cdot u \nabla \eta E^{\sigma} \\ &\quad + \int_S^T \int_{\Omega} \eta |u'|^2 E^{\sigma} - \int_S^T \int_{\Omega} \eta f(u) u E^{\sigma} \\ &\quad + \sigma \int_S^T \int_{\Omega} u' u \xi E^{\sigma-1} E'. \end{aligned} \quad (3.26)$$

Simple calculations, using the Young inequalities, show that

$$\left| - (u', \eta u) E^{\sigma} \Big|_S^T + \sigma \int_S^T \int_{\Omega} u' u \xi E^{\sigma-1} E' \right| \leq CE(S), \quad (3.27)$$

$$\left| \int_S^T (ag(u'), \eta u) E^{\sigma} \right| \leq CE(S) + \frac{\varepsilon}{2} \int_S^T E^{\sigma+1}, \quad \varepsilon > 0, \quad (3.28)$$

$$\left| - \int_S^T \int_{\Omega} M(|\nabla u|^2) \nabla u \cdot u \nabla \eta E^{\sigma} \right| \leq C \int_S^T \int_{\omega} |u|^2 E^{\sigma} + \frac{1}{2} \int_S^T \int_{\Omega} \eta M(|\nabla u|^2) |\nabla u|^2 E^{\sigma}. \quad (3.29)$$

From (3.26)–(3.29), we obtain

$$\begin{aligned} \frac{1}{2} \int_S^T \int_{\Omega} \eta M(|\nabla u|^2) |\nabla u|^2 E^{\sigma} &\leq C \left[E(S) + \int_S^T \int_{\omega} (|u'|^2 + |u|^2 E^{\sigma}) \right] \\ &\quad + \frac{C_*}{m_0^2} \int_S^T \int_{\Omega} |f(u)|^2 E^{\sigma} + \varepsilon \int_S^T E^{\sigma+1}. \end{aligned} \quad (3.30)$$

Step 2. We take a vector field $h \in [W^{1,\infty}(\Omega)]^N$ such that

$$h = \nu, \quad \text{on } \Gamma_0, \quad h \cdot \nu \geq 0, \quad \text{on } \Gamma, \quad h = 0, \quad \text{on } \Omega \setminus \hat{\omega}. \quad (3.31)$$

Choosing $\beta = 0$ and $q = h$ in (3.9), we get

$$\begin{aligned} \left| \int_S^T \int_\Omega ag(u')h \cdot |\nabla u|E^\sigma \right| &\leq C \int_S^T \left[E^{\sigma+1/2}|E'|^{1/(p+1)} + E^{\sigma+1/2(q+1)}|E'|^{q/(q+1)} \right], \\ m_0 \int_S^T \int_{\Gamma_0} \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma &\leq CE(S) + 3\alpha_0 \int_S^T \int_{\hat{\omega}} (|u'|^2 + M(|\nabla u|^2)|\nabla u|^2)E^\sigma \\ &\quad + \alpha_0 \int_S^T \int_\Omega |F(u)|E^\sigma + \varepsilon' \int_S^T E^{\sigma+1}, \quad \varepsilon' > 0, \end{aligned} \tag{3.32}$$

where $\sum_{i,j=1}^N |\partial h_j/\partial x_i| \leq \alpha_0$, for all $x \in \bar{\Omega}$.

Combining (3.30) and (3.32), we have

$$\begin{aligned} m_0 \int_S^T \int_{\Gamma_0} \left| \frac{\partial u}{\partial \nu} \right|^2 E^\sigma &\leq C \left[E(S) + \int_S^T \int_\omega (|u'|^2 + |u|^2)E^\sigma \right] \\ &\quad + \frac{\sigma C_*^2 \alpha_0}{m_0^2} \int_S^T \int_\Omega |f(u)|^2 E^\sigma + \alpha_0 \int_S^T \int_\Omega |F(u)|E^\sigma + \varepsilon \int_S^T E^{\sigma+1}. \end{aligned} \tag{3.33}$$

We conclude from (3.24) and (3.33) that

$$\begin{aligned} \frac{\theta_0}{2} \int_S^T E^{\sigma+1} &\leq \left(\frac{\sigma C_*^2 \alpha_0 R M_0}{m_0^3} + \frac{\beta^2}{\theta_0} \right) \int_S^T \int_\Omega |f(u)|^2 \\ &\quad + \left(N + \frac{\alpha_0 R M_0}{m_0} \right) \int_S^T \int_\Omega |F(u)| \\ &\quad + C \left[E(S) + \int_S^T \int_\omega (|u'|^2 + |u|^2) \right]. \end{aligned} \tag{3.34}$$

Now, in order to absorb the last term into the right-hand side of (3.34), we adapt a method introduced in Conrad and Rao [4]. To this end, we consider $z(t) \in H_0^1(\Omega)$, solution of

$$-\Delta z = \chi(\omega)u, \quad \text{in } \Omega, \quad z = 0, \quad \text{on } \Gamma, \tag{3.35}$$

where $\chi(\omega)$ is the characteristic function of ω . It is easy to verify that z' is solution of the problem

$$-\Delta z' = \chi(\omega)u', \quad \text{in } \Omega, \quad z' = 0, \quad \text{on } \Gamma. \tag{3.36}$$

A simple computation gives

$$|z| \leq C|u|_{L^2(\omega)}, \quad |z'| \leq C|u'|_{L^2(\omega)}, \quad (\nabla z, \nabla u) = |u|_{L^2(\omega)}^2. \tag{3.37}$$

Next, we “multiply” the equation in (1.1) by zE^σ , integrate by parts on $\Omega \times]S, T[$, and use (3.37). Thus, we find

$$\int_S^T \int_\omega M(|\nabla u|^2) |\nabla u|^2 E^\sigma = -(u', z)E^\sigma|_S^T + \int_S^T (u', z')E^\sigma + \sigma \int_S^T E^{\sigma-1} E'(u', z') - \int_S^T (ag(u'), z)E^\sigma - \int_S^T (f(u), z)E^\sigma. \tag{3.38}$$

Here, we note that

$$\left| -(u', z)E^\sigma|_S^T + \sigma \int_S^T E^{\sigma-1} E'(u', z') \right| \leq CE(S), \tag{3.39}$$

$$\left| \int_S^T (u', z)E^\sigma \right| \leq C \int_S^T \int_\omega |u'|^2 E^\sigma + \varepsilon \int_S^T E^{\sigma+1}, \quad \varepsilon > 0, \tag{3.40}$$

$$\left| \int_S^T (ag(u'), z)E^\sigma \right| \leq CE(S) + \varepsilon' \int_S^T E^{\sigma+1}, \quad \varepsilon' > 0, \tag{3.41}$$

$$\left| - \int_S^T (f(u), z)E^\sigma \right| \leq \frac{1}{2m_0} \int_S^T \int_\Omega |f(u)|^2 E^\sigma + \frac{m_0}{2} \int_S^T \int_\omega |u|^2 E^\sigma. \tag{3.42}$$

Using (3.39)–(3.41), we have in (3.38)

$$\int_S^T \int_\omega |u|^2 E^\sigma \leq C \left(E(S) + \int_S^T \int_\omega |u'|^2 E^\sigma \right) + \frac{1}{m_0^2} \int_S^T \int_\Omega |f(u)|^2 E^\sigma + \varepsilon \int_S^T E^{\sigma+1}, \quad \varepsilon > 0. \tag{3.43}$$

Then inserting (3.43) into (3.34) gives

$$\int_S^T E^{\sigma+1} \leq C \left(E(S) + \int_S^T \int_\omega |u'|^2 E^\sigma \right) + \delta_0 \int_S^T \int_\Omega |f(u)|^2 E^\sigma + \delta_1 \int_S^T |F(u)| E^\sigma, \tag{3.44}$$

where

$$\delta_0 = \frac{8}{\theta_0} \left[\frac{\sigma C_* \alpha_0 R M_0}{m_0^3} + \frac{\beta^2}{\theta_0} + \frac{1}{m_0^2} \right], \quad \delta_1 = \frac{8}{\theta_0} \left[N + \frac{\alpha_0 R M_0}{m_0} \right]. \tag{3.45}$$

Now, we observe that

$$\begin{aligned} \delta_0 \int_\Omega |f(u)|^2 &\leq \delta_0 h_0 |u|_{2(\alpha+1)}^{2(\alpha+1)} \leq C \delta_0 h_0 |u|_{2N/(N-2)}^{2(1-\theta_1)(\alpha+1)} |\Delta u|^{2\theta_1(\alpha+1)} \\ &\leq C \delta_0 h_0 \varepsilon_0^{2(1-\theta_1)(\alpha+1)-2} K^{2\theta_1(\alpha+1)} |\nabla u|^2 \end{aligned} \tag{3.46}$$

if $N \geq 3$ and $\alpha > 2/(N - 2)$, with $\theta_1 = ((\alpha + 1)(N - 2) - N)/2(\alpha + 1)$.

Further, we have that since $\alpha \leq 2/(N - 4)^+$,

$$(1 - \theta_1)(\alpha + 1) \geq 1. \tag{3.47}$$

Thus,

$$\delta_0 h_0 |u|_{2(\alpha+1)}^{2(\alpha+1)} \leq C \delta_0 h_0 \varepsilon_0^{2(1-\theta_1)(\alpha+1)-2} K^{2\theta_1(\alpha+1)} |\nabla u|^2. \tag{3.48}$$

When $N \leq 2$ or $\alpha \leq 2/(N - 2)^+$, we see that (3.48) holds with $\theta_1 = 0$. Thus, under a little more stronger assumption than (2.19),

$$Ch_0 \left(\varepsilon_0^{(\alpha+2)(1-\theta_1)-2} K^{(\alpha+2)\theta} + \varepsilon_0^{2(1-\theta_1)(\alpha+1)-2} K^{2\theta_1(\alpha+1)} \right) \leq \frac{m_0}{4}, \tag{3.49}$$

we obtain from (3.44) that

$$\int_S^T E^{\sigma+1} dt \leq C \left(E(S) + \int_S^T \int_\omega |u'|^2 E^\sigma \right). \tag{3.50}$$

In order to absorb the second term in (3.50), we consider two cases.

(i) The case $p = 1$. We take $\sigma = 0$, hence by (2.6),

$$\int_S^T E \leq CE(S) + \frac{C}{a_0} \int_S^T \int_\omega a |u'|^2 \leq CE(S) + \int_S^T \int_\omega a |g(u')| |u'| \leq CE(S). \tag{3.51}$$

Applying Lemma 2.3, we obtain

$$E(t) \leq CI_0 e^{-\lambda t}, \quad \lambda = \lambda(K, I_0). \tag{3.52}$$

(ii) When $p > 1$, we take $\sigma = (p - 1)/2$. It follows from (2.6) that

$$\begin{aligned} \int_S^T E^{(p+1)/2} &\leq CE(S) + \frac{C}{a_0} \int_S^T \int_\omega a |u'|^2 E^{(p-1)/2} \\ &\leq CE(S) + C \int_S^T \int_\Omega a |u'|^2 E^{(p-1)/2} \\ &\leq C \left[E(S) + \int_S^T E^{(p-1)/2} \left(\int_{\Omega_1} a |u'|^2 + \int_{\Omega_2} a |u'|^2 \right) \right] \\ &\leq C \left[E(S) + \int_S^T (E^{(p-1)/2} |E'|^{2/(p+1)} + E^{(p-1)/2} |E'|) \right] \\ &\leq C [E(S) + E^{(p+1)/2}(S)] + \varepsilon \int_S^T E^{(p+1)/2}, \quad \varepsilon > 0. \end{aligned} \tag{3.53}$$

Hence

$$\int_S^T E^{(p+1)/2} \leq CE(S). \tag{3.54}$$

Using Lemma 2.3 again, we conclude that

$$E(t) \leq q(K, I_0) (1 + t)^{-2/(p-1)}, \quad t \geq 0, \tag{3.55}$$

as long as the local solutions exist, where $q(K, I_0)$ denotes a certain constant continuously depending on K and I_0 . □

We are now in a position to obtain H^2 a priori bounds. Set

$$E_1(t) = M \left(|\nabla u(t)|^2 \right) |\Delta u(t)|^2 + |\nabla u'(t)|^2. \tag{3.56}$$

Then we have the following estimate, which is the heart of this paper.

LEMMA 3.6. Assume that $u(t)$ is a solution satisfying $(u(t), u'(t)) \in W_K$ on $[0, T[$, for some $K > 0$. Then,

$$|\nabla u'(t)|^2 + |\Delta u(t)|^2 \leq Q^2(I_0, I_1, K), \tag{3.57}$$

with

$$\lim_{I_0 \rightarrow 0} Q^2(I_0, I_1, K) = \frac{I_1^2}{\min\{1, m_0\}}, \tag{3.58}$$

and where

$$I_1^2 = |\nabla u_1|^2 + M(|\nabla u_0|^2) |\Delta u_0|^2. \tag{3.59}$$

Proof. For $E_1(t)$ with respect to t , we get

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} E_1(t) + (ag'(u') \nabla u'(t), \nabla u'(t)) \\ = -(f'(u) \nabla u, \nabla u') + M'(|\nabla u(t)|^2) (\nabla u'(t), \nabla u(t)) |\Delta u(t)|^2 - (g(u') \nabla a, \nabla u'). \end{aligned} \tag{3.60}$$

Using the assumptions on a , g , and f , it follows from (3.60) that

$$\begin{aligned} \frac{d}{dt} E_1(t) &\leq C \left\{ \int_{\Omega} (|u|^\alpha |\nabla u| |\nabla u'|) + |\nabla u|_2^{\gamma+1} |\nabla u'|_2 |\Delta u|_2^2 + \sum_{i=1}^n \int_{\Omega} \frac{\partial a}{\partial x_i} \frac{\partial}{\partial x_i} \int_0^u g(\xi) d\xi \right\} \\ &\leq C \left\{ \int_{\Omega} (|u|^\alpha |\nabla u| |\nabla u'|) + |\nabla u|_2^{\gamma+1} |\nabla u'|_2 |\Delta u|_2^2 + \int_{\Omega} \left| \int_0^u g \right| |\Delta a| \right\} \\ &\leq C \left\{ \left(\int_{\Omega} |u|^{2\alpha} |\nabla u|^2 \right)^{1/2} \left(\int_{\Omega} |\nabla u'|^2 \right)^{1/2} + E(t)^{(\gamma+1)/2} K^3 + \int_{\Omega} ag(u') u' \right\}. \end{aligned} \tag{3.61}$$

Further, we observe that

$$\begin{aligned} \left(\int_{\Omega} |u|^{2\alpha} |\nabla u|^2 \right)^{1/2} &\leq |u|_{N\alpha}^\alpha |\nabla u|_{2N/(N-2)} \\ &\leq C |u|_{2N/(N-2)}^{\alpha(1-\theta_0)} |\Delta u|_2^{\alpha\theta_0} |\Delta u|_2 \quad (\text{by Lemma 2.2}) \\ &\leq C |\nabla u|_2^{\alpha(1-\theta_0)} |\Delta u|_2^{\alpha\theta_0+1} \\ &\leq CE(t)^{\alpha(1-\theta_0)/2} K^{\alpha\theta_0+1}, \end{aligned} \tag{3.62}$$

with

$$\theta_0 = \left(\frac{N-2}{2} - \frac{1}{\alpha} \right)^+ = \frac{((N-2)\alpha - 2)^+}{2\alpha} \leq 1. \tag{3.63}$$

Then, it follows from (3.61) and (3.62) that

$$\frac{d}{dt} E_1(t) \leq C \{ E(t)^{\alpha(1-\theta_0)/2} K^{\alpha\theta_0+2} + E(t)^{(\gamma+1)/2} K^3 - E'(t) \}. \tag{3.64}$$

Thus, integrating (3.64) over $[0, t[$, we obtain $\int_{\Omega} \left| \int_0^t g \right| |\Delta a|$,

$$\begin{aligned} & |\Delta u(t)|^2 + |\nabla u'(t)|^2 \\ & \leq \frac{1}{\min\{1, m_0\}} \left\{ I_1^2 + CK^{\alpha\theta_0+2} \int_0^{+\infty} E(t)^{\alpha(1-\theta_0)/2} dt + CK^3 \int_0^{+\infty} E(t)^{(\gamma+1)/2} dt + CI_0^2 \right\} \\ & \leq \frac{1}{\min\{1, m_0\}} \left\{ I_1^2 + CI_0^2 + CI_0^{(\gamma+1)/2} K^3 + CI_0^{\alpha(1-\theta_0)/2} K^{\alpha\theta_0+2} \right\} \\ & \equiv Q^2(I_0, I_1, K), \quad \text{on } [0, T[. \end{aligned} \tag{3.65}$$

Define

$$\begin{aligned} S_K &= \{(u_0, u_1) \in (H_0^1 \cap H^2) \times H_0^1 : Q^2(I_0, I_1, K) < K\}, \\ S_0 &= \bigcup_{K>0} S_K. \end{aligned} \tag{3.66}$$

Since

$$\lim_{I_0 \rightarrow 0} Q^2(I_0, I_1, K) = \frac{I_1^2}{\min\{1, m_0\}}, \tag{3.67}$$

the S_K is not empty if $I_1 < \min\{1, m_0\}K^2$ and I_0 is sufficiently small.

If $(u_0, u_1) \in S_K$ for some $K > 0$, then the corresponding local solution $u(t)$ exists on some interval $[0, T[$ and satisfies

$$E_1(u(t)) \leq \min\{1, m_0\}K^2 \quad \text{on } 0 \leq t < T. \tag{3.68}$$

From (3.43), we have $(u(t), u'(t)) \in W_K$.

Hence, it follows from Lemma 3.6 that

$$|\nabla u'(t)|^2 + |\Delta u(t)|^2 \leq Q^2(I_0, I_1, K) < K^2, \quad \text{on } [0, T[. \tag{3.69}$$

Next, we affirm that

$$(u(t), u'(t)) \in S_K, \quad \text{on } [0, T[. \tag{3.70}$$

In fact, suppose that there is a number $t^* \in [0, T[$ such that $(u(t), u'(t)) \in S_K$ on $[0, T[$ and $(u(t^*), u'(t^*)) \notin S_K$, then it follows that

$$Q^2(E(u(t^*)), E_1(u(t^*)), K) \geq K^2. \tag{3.71}$$

Note that $Q^2(I_0, I_1, K)$ is the increasing function with respect to I_0 and I_1 . Hence, we see from (3.48), the energy identity (3.4), and Lemma 3.6 that

$$I_1^2 + CI_0^2 + C_1(I_0)K^3 + C_2(I_0)K^{\alpha\theta_0+2} \geq \min\{1, m_0\}K^2. \tag{3.72}$$

But, then, since

$$\lim_{I_0 \rightarrow 0} Q^2(I_0, I_1, K) = \frac{I_1^2}{\min\{1, m_0\}}, \tag{3.73}$$

we may take I_1 so that

$$\frac{2I_1^2}{\min\{1, m_0\}} < K^2 \quad (3.74)$$

with sufficiently small I_0 . This contradicts (3.72).

Now, since we can repeat the continuation procedure indefinitely, we conclude that $u(t)$ can be continued globally on $[0, +\infty[$ and $(u(t), u'(t)) \in S_0$, for all $t \geq 0$.

The uniqueness follows from a similar argument as in [1]. \square

References

- [1] M. Aassila, *Global existence of solutions to a wave equation with damping and source terms*, Differential Integral Equations **14** (2001), no. 11, 1301–1314.
- [2] E. Cabanillas, J. Bernui, and Z. Huaranga, *Existencia Global y Decaimiento, de la Energía de una Ecuación de Kirchoff con Disipación Localizada*, Pesquimat Rev de la F.C.M **6** (2003), no. 1, 21–31.
- [3] ———, *Global Existence and Decay Estimates to a quasilinear wave equation with locally distributed Damping*, Pesquimat, Rev de la F.C.M - UNMSM V N **2** (2003).
- [4] F. Conrad and B. Rao, *Decay of solutions of the wave equation in a star-shaped domain with nonlinear boundary feedback*, Asymptotic Anal. **7** (1993), no. 3, 159–177.
- [5] V. Komornik, *Exact Controllability and Stabilization, the Multiplier Method*, RAM: Research in Applied Mathematics, Masson, Paris, 1994.
- [6] S. Kouémou-Patcheu, *Global existence and exponential decay estimates for a damped quasilinear equation*, Comm. Partial Differential Equations **22** (1997), no. 11-12, 2007–2024.
- [7] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [8] ———, *Contrôlabilité Exacte, Perturbations et Stabilisation de Systèmes Distribués. Tome 1. Contrôlabilité Exacte. [Exact Controllability, Perturbations and Stabilization of Distributed Systems. Vol. 1. Exact Controllability]*, Recherches en Mathématiques Appliquées [Research in Applied Mathematics], vol. 8, Masson, Paris, 1988.
- [9] M. Nakao, *Decay of solutions of the wave equation with a local degenerate dissipation*, Israel J. Math. **95** (1996), 25–42.
- [10] E. Zuazua, *Exponential decay for the semilinear wave equation with locally distributed damping*, Comm. Partial Differential Equations **15** (1990), no. 2, 205–235.

E. Cabanillas Lapa: Instituto de Investigación, Facultad de Ciencias Matemáticas, Universidad Nacional Mayor de San Marcos, Lima, Peru

E-mail address: cleugenio@yahoo.com

Z. Huaranga Segura: Instituto de Investigación, Facultad de Ciencias Matemáticas, Universidad Nacional Mayor de San Marcos, Lima, Peru

E-mail address: zachariashs@yahoo.es

F. Leon Barboza: Instituto de Investigación, Facultad de Ciencias Matemáticas, Universidad Nacional Mayor de San Marcos, Lima, Peru

E-mail address: fleonb2004@yahoo.es

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.otero@usc.es