

A STUDY OF THE WAVES AND BOUNDARY LAYERS DUE TO A SURFACE PRESSURE ON A UNIFORM STREAM OF A SLIGHTLY VISCOUS LIQUID OF FINITE DEPTH

ARGHYA BANDYOPADHYAY

Received 31 May 2006; Revised 25 October 2006; Accepted 22 November 2006

The 2D problem of linear waves generated by an arbitrary pressure distribution $p_0(x, t)$ on a uniform viscous stream of finite depth h is examined. The surface displacement ζ is expressed correct to $O(\nu)$ terms, for small viscosity ν , with a restriction on $p_0(x, t)$. For $p_0(x, t) = p_0(x)e^{i\omega t}$, exact forms of the steady-state propagating waves are next obtained for all x and not merely for $x \gg 0$ which form a wave-quartet or a wave-duo amid local disturbances. The long-distance asymptotic forms are then shown to be uniformly valid for large h . For numerical and other purposes, a result essentially due to Cayley is used successfully to express these asymptotic forms in a series of powers of powers of $\nu^{1/2}$ or $\nu^{1/4}$ with coefficients expressed directly in terms of nonviscous wave frequencies and amplitudes. An approximate thickness of surface boundary layer is obtained and a numerical study is undertaken to bring out the salient features of the exact and asymptotic wave motion in question.

Copyright © 2006 Arghya Bandyopadhyay. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The classical problem of waves due to any initial variable pressure distribution on the surface of an ocean of finite depth has attracted the attention of many researchers mainly because of its connection with several important physical problems such as ship-waves, impact of aerial blast waves on the ocean, storms, tsunamis, and so forth. Although the physical settings of these problems are different, there exists a similarity in the analyses of the wave integrals and also of the wave behavior in many cases. For example, the linear waves generated by initial surface disturbances in an inviscid liquid exhibits the same qualitative behavior as that of the tsunamis (cf. Wehausen and Laitone [11, page 618]). Our objective here, however, is to study the effect of viscosity on the waves in question without any further elaboration of such interrelationships.

2 A study of the waves and boundary layers

It is well known that while the formal solution of the linear problem of waves produced by surface disturbances on a stationary or moving viscous fluid of finite depth is easily obtained, substantial difficulty arises in evaluating or analyzing this solution which can be seen in the work of Bandyopadhyay [1], Nikitin and Potetyunko [6], and Pramanik [7]. Such analyses have so far remained confined to derivation of an asymptotic expression of the surface displacement ζ for small viscosity ν at large times and distances, and there seems to have no attempt at analytical and numerical studies of the wave integrals in its exact form (I). Also the approximation of ζ for smallness of ν is generally carried out in a rather heuristic way without paying attention to the restriction that may have to be placed on the pressure models for the approximation to succeed (II).

In this paper, we examine the two-dimensional form of the problem stated above for a uniformly flowing stream of small viscosity, our main objective being to find the restriction mentioned in (II) for a general pressure model $p_0(x, t)$ and to achieve (I) for a special model $p_0(x)e^{i\omega t}$. For the former, we first express ζ correct to $O(\nu)$ terms by a proven approximation formula of Bandyopadhyay [1], and show incidentally that an $O(\sqrt{\nu})$ analysis of ζ (cf. Pramanik [7] for $p_0(x)e^{i\omega t}$) fails to bring out the effect of viscosity on ζ for large depths, while an $O(\nu)$ analysis succeeds in correcting this shortcoming (cf. (3.9) together with (3.7), (3.8), and (3.10)). Secondly, in Section 4 we initiate the study (I) for the special pressure distribution $p_0(x)e^{i\omega t}$, and verify that a steady-state exists everywhere and not merely for large distances (contrast Pramanik [7]). This steady state as it is shown in Section 4.2 consists of a propagating wave system ζ_p^* compounded by standing local disturbances for all times and distances. After the exact evaluation of the former, we find that this system of propagating waves ζ_p^* is either a damped and constant-velocity wave-duo or a wave-quartet according to the fact that the stream speed u_0 does or does not exceed a certain critical value. At large distances, the standing waves fade out in comparison to the leading term, denoted by ζ^{**} , of the asymptotic expression of ζ_p^* . When ζ^{**} is retained correct up to $O(\sqrt{\nu})$ terms, it agrees with the corresponding result obtained by Pramanik [7]; this expression, however, as is shown here makes the long-distance wave form inaccurate for large h , while in our case the presence of $O(\nu)$ terms in ζ_p^* rectifies such inappropriate result for large depth (Section 5).

We next show (Sections 6.1 and 6.2) that it is advantageous to write ζ^{**} in the following form by a result essentially due to Cayley (Whittaker and Watson [12, page 147]):

$$\zeta^{**} = \begin{cases} \zeta_0 + \sqrt{\nu}\zeta_2 + \nu\zeta_4, & \text{when } u_0 \neq u_0^*, \\ \zeta'_0 + \nu^{1/4}\zeta'_1 + \nu^{1/2}\zeta'_2 + \nu^{3/4}\zeta'_3 + \nu\zeta'_4, & \text{when } u_0 = u_0^*, \end{cases} \quad (1.1)$$

where each ζ_j (or ζ'_j) is wave-quartet or wave-duo, and in either case the wave amplitudes and wave frequencies are all directly expressed in terms of their nonviscous values. Such expansions also facilitate a numerical study of the wave motion. Some interesting features of the wave motion are brought out through a computational study of the exact wave integral ζ_p^* and its asymptotic forms which are discussed in Section 8. Finally, since the motion may be considered to have a high Reynolds number $\sqrt{gh^3}/\nu$, a question naturally arises as to the formation of boundary layers both at the surface and the bottom and the shapes they assume vis-à-vis those of waves. In this regard, in Section 7, we determine the

cross-vertical vorticity $\omega_z(x, y, t)$ and find its approximate value correct to $O(\nu)$ terms. Its Cayley expansion is next derived for $|x| \rightarrow \infty$ by the same procedure as adopted for ζ^{**} . These expressions are utilized for the determination of the approximate thickness of the boundary layer at the surface at large horizontal distances from the source.

In passing, mention may be made of two important papers, one by Beale [2] who is concerned about the existence of solution of the above problem in a general three-dimensional nonlinear setting when the initial surface and velocity fields are prescribed, and the other paper by Ehrenmark and Body [4] who are however interested in the solution of the inviscid problem in the presence of a plane beach under the varying conditions on the surface as well as on the bed.

2. The problem and its integral solution

For the two-dimensional motion of a horizontally unbounded viscous liquid of constant depth h parallel to the xy -plane, a coordinate system is chosen such that the origin is in the undisturbed free surface which is the xy -plane; the y -axis is vertically upwards and the x -axis is positive to the right, while the stream moves in the Ox direction with speed u_0 relative to the frame of reference.

Waves of height $\zeta(x, y)$ above the undisturbed surface are generated by the action of the pressure distribution $p_0(x, t)$ which is suddenly applied on $y = 0$ at $t = 0$. $p_0(x, t)$ is assumed to have a Fourier transform in x and also to have a Laplace transformation in t . If ζ and the disturbance velocity components (u, v) parallel to the axes are small, the equation of continuity and the linearized Stokes-Navier equations are, respectively,

$$\begin{aligned} \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} &= 0, \\ \left(\frac{\partial}{\partial t} + u_0 \frac{\partial}{\partial x} \right) (u, v) &= -\frac{1}{\rho} \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) p + \nu \nabla^2 (u, v), \end{aligned} \quad (2.1)$$

where $p = p_1 + \rho g y$, $\nabla^2 \equiv \partial^2/\partial x^2 + \partial/\partial y^2$, $-\infty < x < \infty$, $-h \leq y \leq \zeta \approx 0$, $t \geq 0$. For a fluid at rest, the initial conditions are

$$u = v = 0, \quad \zeta = 0 \quad \text{for } t = 0. \quad (2.2)$$

The linearized normal stress and the no-shear stress condition on the surface together with the surface boundary condition are, respectively,

$$-p_0(x, t) = -p + \rho g \zeta + 2\mu \nu_y, \quad u_y + v_x = 0, \quad \zeta_t + u_0 \zeta_x = v \quad \text{for } y = 0, t > 0. \quad (2.3)$$

The condition of no-slip at the bottom is

$$u = v = 0 \quad \text{for } y = -h, t > 0. \quad (2.4)$$

Lastly the conditions at infinity are u, v, p and their x -derivatives are zero as $|x| \rightarrow \infty$.

4 A study of the waves and boundary layers

Here subscripts t, x, y refer to partial differentiation. Our main interest is in the expression for ζ .

For convenience, we will suppress arguments of the functions and their transforms in general. Let U, V, H, P, P_0 denote the Fourier transforms of u, v, ζ, p, p_0 , respectively, in x , and let $\tilde{U}, \tilde{V}, \tilde{H}, \tilde{P}, \tilde{P}_0$ denote the Laplace transforms of the former in t, k , and s being the respective transform parameters.

The formal solution of the problem may be written as

$$[u, v, p, \zeta] = \frac{1}{(2\pi)^{3/2}i} \int_{-\infty}^{\infty} e^{-ikx} dk \int_{c-i\infty}^{c+i\infty} [\tilde{U}, \tilde{V}, \tilde{P}, \tilde{H}] e^{ts} ds, \quad (2.5)$$

where the constant c has its usual meaning, and the transform integrals are given by

$$\begin{aligned} \tilde{U} &= \frac{i \operatorname{sgn} k}{\rho(s - iku_0)} [|k| \tilde{A} \{ \cosh |k|(y+h) - \cosh m(y+h) \} \\ &\quad + \tilde{B} \{ |k| \sinh |k|(y+h) - m \sinh m(y+h) \}], \\ \tilde{V} &= \frac{|k|}{\rho(s - iku_0)m} [\tilde{A} \{ |k| \sinh m(y+h) - m \sinh |k|(y+h) \} \\ &\quad + m \tilde{B} \{ \cosh m(y+h) - \cosh |k|(y+h) \}], \\ \tilde{H} &= -|k| \{ \mu D(s - iku_0)m \}^{-1} (m \sinh |k|h \cosh mh - |k| \sinh mh \cosh |k|h), \\ D\tilde{A} &= (k^2 + m^2) \cosh mh - 2k^2 \cosh |k|h, \\ mD\tilde{B} &= 2mk^2 \sinh |k|h - |k|(k^2 + m^2) \sinh mh, \\ -\tilde{P}_0(s - iku_0)Dv^{-1} &= 4k^2(k^2 + m^2) - \cosh mh \cosh |k|h \left\{ (k^2 + m^2)^2 + 4k^4 + \frac{\sigma^2}{\nu^2} \right\} \\ &\quad + |k|m^{-1} \sinh mh \sinh |k|h \left\{ (k^2 + m^2)^2 + 4k^2m^2 + \left(\frac{gk}{\sigma\nu} \right)^2 \right\}, \end{aligned} \quad (2.6)$$

where

$$\sigma^2 = [\sigma(k)]^2 = g|k| \tanh |k|h, \quad m^2 = k^2 + \frac{s}{\nu} - \frac{iku_0}{\nu}. \quad (2.7)$$

By (2.6), we have

$$\tilde{H} = \frac{|k|\tilde{P}_0}{\rho} \frac{F(\nu^{1/2})}{f(\nu^{1/2})}, \quad -\tilde{P}_0 = -P + \rho g \tilde{H} + 2\mu \tilde{V}_y, \quad (2.8)$$

where

$$\begin{aligned}
 F(\nu^{1/2}) &= (s - iku_0 + \nu k^2)^{1/2} \tanh |k|h - \nu^{1/2} |k| \tanh \left(\frac{s - iku_0 + \nu k^2}{\nu} \right)^{1/2} h, \\
 f(\nu^{1/2}) &= 4(s - iku_0 + \nu k^2)^{1/2} (s - iku_0 + 2\nu k^2) \nu k^2 \operatorname{sech} |k|h \operatorname{sech} \left(\frac{s - iku_0 + \nu k^2}{\nu} \right)^{1/2} h \\
 &\quad - (s - iku_0 + \nu k^2)^{1/2} \{ (s - iku_0 + 2\nu k^2)^2 + 4\nu^2 k^4 + \sigma^2 \} + |k| \nu^{1/2} \\
 &\quad \times \{ (s - iku_0 + 2\nu k^2)^2 + 4\nu k^2 (s - iku_0 + \nu k^2) + g|k| \coth |k|h \} \\
 &\quad \times \tanh |k|h \times \tanh \left(\frac{s - iku_0 + \nu k^2}{\nu} \right)^{1/2} h.
 \end{aligned} \tag{2.9}$$

By the inversion theorem for Laplace transform and the convolution theorem, one then gets

$$H = \frac{|k|}{2\pi i \rho} \left[\int_0^t P_0(k, t - \tau) d\tau \int_{c-i\infty}^{c+i\infty} e^{s\tau} \frac{F(\nu^{1/2})}{f(\nu^{1/2})} ds \right]. \tag{2.10}$$

The Fourier inversion theorem now gives the exact integral expression for ζ ,

$$\zeta = \frac{1}{(2\pi)^{3/2} i \rho} \int_{-\infty}^{\infty} \left[\int_0^t P_0(k, t - \tau) d\tau \int_{c-i\infty}^{c+i\infty} e^{s\tau} \frac{F(\nu^{1/2})}{f(\nu^{1/2})} ds \right] |k| e^{-ikx} dk. \tag{2.11}$$

The corresponding integral expressions for u , v , p follow in a similar way.

3. Waves in a slightly viscous fluid

If the kinematic coefficient of viscosity ν of the fluid is small, it may be shown that

$$\begin{aligned}
 \zeta &= -\frac{1}{(2\pi)^{3/2} i \rho} \int_{-\infty}^{\infty} |k| e^{-ikx} dk \left[\int_0^t P_0(k, t - \tau) \times e^{iku_0 \tau} d\tau \int_{c-i\infty}^{c+i\infty} e^{s'\tau} \left(\frac{T_1}{\Delta_1} \right) ds' \right] \\
 &\quad + O(\nu^{3/2}), \quad \text{as } \nu \rightarrow 0,
 \end{aligned} \tag{3.1}$$

provided as $k \rightarrow \infty$, $P_0(k_1, k_2, \tau)$ is of the order $O(k^{-17/4-n})$, $n > 0$, uniformly in ψ , and τ with $k_1 = k \cos \psi$, $k_2 = k \sin \psi$ [11].

Here $s' = s - iku_0$, and

$$T_1 \equiv T_1(s', k) = s' \tanh |k|h - |k|(\nu s')^{1/2} + \frac{1}{2} \nu k^2 \tanh |k|h, \tag{3.2}$$

$$\Delta_1 \equiv \Delta_1(s', k) = s'^3 + s' \sigma^2 - (\nu s')^{1/2} g^{-1} (s'^2 \sigma^2 + g^2 k^2) + \frac{1}{2} \nu k^2 (9s'^2 + \sigma^2).$$

6 A study of the waves and boundary layers

This approximation process leaves the conditions unaffected since we have from (2.5), (2.8), and Sneddon [9, page 184] that

$$\begin{aligned}
 p(x, 0, 0+) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \lim_{s \rightarrow \infty} [s \{ \tilde{P}(k, 0, s) \}_{(F/f) \rightarrow (-T_1/\Delta_1)}] e^{-ikx} dk \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \lim_{s \rightarrow \infty} [s \{ \tilde{P}_0(k, s) \}_{(F/f) \rightarrow (-T_1/\Delta_1)}] e^{-ikx} dk \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} P_0(k, 0) e^{-ikx} dk = p_0(x, 0).
 \end{aligned} \tag{3.3}$$

Evaluation of the s' -integral (3.1). The s' -integrand in (3.1) has a branch point at $s' = 0$ as well as possible poles arising from those zeros of Δ_1 for which $T_1 \neq 0$. We suppose that the s' -plane is cut along the negative real axis from 0 to $-\infty$, and that $s'^{1/2}$ is real and positive when s' is real and positive. Considered as a sixth-degree polynomial in $s'^{1/2}$, Δ_1 has, for $\nu = 0$, a double zero at the point zero and four other simple zeros at $\sqrt{\sigma} e^{\pm i\pi/4}$, $\sqrt{\sigma} e^{\pm 3\pi/4}$.

For the three corresponding values of s' , namely

$$s'_{00} = 0, \quad s'_{10} = i\sigma, \quad s'_{20} = -i\sigma, \tag{3.4}$$

we may assume the following expansion for the possible zeros s'_j , $j = 0, 1, 2$ of Δ_1 in powers of $\nu^{1/2}$:

$$s'_j = s'_{j0} + s'_{j1} \nu^{1/2} + s'_{j2} \nu + s'_{j3} \nu^{3/2} + \dots \tag{3.5}$$

It is then found that

$$s'_{01} = 0, \quad s'_{02} - s'^{1/2}_{02} \left(\frac{k}{\tanh kh} \right) + \frac{1}{2} k^2 = 0. \tag{3.6}$$

To our degree of accuracy, this shows that $T_1(s'_0, k) = T_1(\nu s'_{02}, k-) = 0$.

Leaving aside then the zero s_0 of Δ_1 and determining the other coefficients $s_{j\ell}$ for $\ell = 1, 2, 3$, we find that the s' -integrand of (3.1) has two simple poles at $s' = s'_j$, $j = 1, 2$, given by

$$\begin{aligned}
 s'_j &= \alpha_j + \frac{1}{2} [1 + (-1)^j i] \nu^{3/2} (2\sigma)^{-1/2} |k|^3 \operatorname{cosech}^3 2|k|h \\
 &\times \left(\frac{-13}{4} + \frac{1}{2} \cosh 4kh - \cosh 6kh \right) + \dots,
 \end{aligned} \tag{3.7}$$

where

$$\begin{aligned}
 \alpha_j &= \alpha_j(k) = (-1)^{j-1} i\sigma - \{1 + (-1)^{j-1} i\} |k| \left(\frac{\sigma\nu}{2} \right)^{1/2} \operatorname{cosech} 2|k|h \\
 &- 2\nu k^2 \frac{\cosh 2kh + \cosh 4kh - 1}{\cosh 4kh - 1}, \quad j = 1, 2.
 \end{aligned} \tag{3.8}$$

(cf. Biesel's result cited by Wehausen and Laitone in [11, page 644].)

Using a modified Bromwich contour (Tranter [10]) now completes the s' -integration of (3.1) and the result is

$$\zeta = -\frac{1}{\sqrt{2\pi\rho}} \int_{-\infty}^{\infty} |k| e^{-ikx} \left[\int_{\tau=0}^t P_0(k, t-\tau) \times e^{iku_0\tau} f_2(k, \tau) d\tau \right] dk, \quad (3.9)$$

where

$$f_2(k, t) = \sum_{j=1}^2 \frac{T_1(s'_j, k)}{\Delta'_1(s'_j, k)} e^{s'_j t} - \frac{1}{2\pi i} \int_0^{\infty} e^{-t\chi} \left\{ \left[\frac{T_1}{\Delta_1} \right]_{s'=\chi e^{i\pi}} - \left[\frac{T_1}{\Delta_1} \right]_{s'=\chi e^{-i\pi}} \right\} d\chi. \quad (3.10)$$

Introducing conditions similar to those of evaluating (3.1), we may show that the use of the values s'_j (given by (3.7)) instead of $s' = \alpha_j$ for the roots would add only a term of order $\nu^{3/2}$ to ζ as calculated on the basis of (3.9) and (3.10), correct to the first power of ν in the amplitude. We also have

$$\sum_{j=1}^2 \frac{T_1(s_j, k)}{\Delta'_1(s_j, k)} e^{s_j t} = (g|k|)^{-1} e^{-\sigma_2 t} \left[\left(\sigma - \frac{\sigma_1}{2} \right) \sin(\sigma - \sigma_1)t + \frac{\sigma_2}{2} \cos(\sigma - \sigma_1)t \right], \quad (3.11)$$

where

$$\sigma_1 = |k| \left(\frac{\sigma\nu}{2} \right)^{1/2} \operatorname{cosech} 2|k|h, \quad \sigma_2 = \sigma_1 + 2\nu k^2 \left(1 + \frac{\cosh 2kh}{\cosh 4kh - 1} \right). \quad (3.12)$$

The final expression for ζ results when $f_2(k, t)$ in (3.10) is modified in accordance with (3.11) and (3.12). The surface displacement thus consists of damped gravity waves corresponding to the part of (3.9) containing the integral free term of (3.10) together with a diffusive disturbance pattern corresponding to the remaining part of (3.9). We also note that in the former part, there also appears to be a phase lag as we combine the sine and cosine terms into a single sine or cosine term (cf. Prosperetti [8]).

4. General character of the steady-state surface motion for an oscillatory pressure distribution

When the applied pressure distribution is of the form $p_0(x, t) = p_0(x)e^{i\omega t}H(t)$, $H(t)$ is the Heaviside unit function, we have by (3.11)-(3.12) that

$$\begin{aligned} \zeta = & -\frac{1}{\sqrt{2\pi\rho g}} e^{i\omega t} \int_{-\infty}^{\infty} P_0(k) e^{-ikx} \frac{K(k)}{2} dk \int_0^t \exp [i(\sigma - \sigma_1) - i\omega + iku_0 - \sigma_2] \tau d\tau \\ & - \frac{1}{\sqrt{2\pi\rho g}} e^{i\omega t} \int_{-\infty}^{\infty} P_0(k) e^{-ikx} \frac{\overline{K}(k)}{2} dk \int_0^t \exp [-i(\sigma - \sigma_1) - i\omega + iku_0 - \sigma_2] \tau d\tau \\ & + \frac{e^{i\omega t}}{\rho i (2\pi)^{3/2}} \int_{-\infty}^{\infty} P_0(k) |k| e^{-ikx} dk \int_0^t e^{(-i\omega + iku_0 - \chi)\tau} d\tau \\ & \times \int_0^{\infty} \left\{ \left[\frac{T_1}{\Delta_1} \right]_{s'=\chi e^{i\pi}} - \left[\frac{T_1}{\Delta_1} \right]_{s'=\chi e^{-i\pi}} \right\} d\chi. \end{aligned} \quad (4.1)$$

Here

$$K(k) = \frac{\sigma_2}{2} - i\left(\sigma - \frac{\sigma_1}{2}\right). \quad (4.2)$$

Following the evaluation of the τ -integrals in the first two terms of (4.1), we apply Watson's lemma (see Bleistein and Handelsman [3, page 103]) to the resulting k -integral to obtain its limit as $t \rightarrow \infty$, x remaining fixed. For the third term of (4.1), we have

$$\begin{aligned} C_3 &= \int_{-\infty}^{\infty} P_0(k) |k| dk \int_0^1 t e^{(-i\omega + iku_0)t\tau'} d\tau' \\ &\times \int_0^{\infty} \frac{\chi^{5/2} \nu^{1/2} \operatorname{sech}^2 kh [\chi - (1/2)\nu k^2 (8 \cosh^2 kh + 1)] e^{-\chi t\tau'} d\chi}{\Delta_1(\chi e^{i\pi}, k) \cdot \Delta_1(\chi e^{-i\pi}, k)}. \end{aligned} \quad (4.3)$$

Considering $\Delta_1(s', k)$ as a sixth-degree polynomial in $\sqrt{s'}$, we may approximately express the product $\Delta_1]_{\chi e^{i\pi}} \cdot \Delta_1]_{\chi e^{-i\pi}}$ as

$$(\chi + a_o^2)(\chi + a_o'^2)(\chi + \alpha_1')^2(\chi + \alpha_2')^2, \quad (4.4)$$

where a_o, a_o' are the two values of $\sqrt{\nu}x$, where x is given by $x^2 - x(k/\tanh kh) + (1/2)k^2 = 0$ and α_1', α_2' are the same as α_1, α_2 of (3.8) but correct only up to the term in $\sqrt{\nu}$ therein. It is easy to see that $t \times \chi$ -integral tends to zero as $t \rightarrow \infty$ for $0 < \varepsilon < \tau' \leq 1$, while the continuity of the $\chi - \tau'$ integral in $0 \leq \varepsilon < 1$ ensures that

$$C_3 \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (4.5)$$

The steady-state surface displacement ζ^* thus comes out to be

$$\zeta^* \sim \frac{e^{i\omega t}}{\sqrt{2\pi\rho g}} (C_1 + C_2), \quad (4.6)$$

where

$$C_1 = \int_{-\infty}^{\infty} P_0(k) \frac{(\sigma - \sigma_1/2) + i\sigma_2/2}{2(-\sigma + \sigma_1 + \omega - ku_0 - i\sigma_2)} e^{-ikx} dk, \quad (4.7)$$

$$C_2 = \int_{-\infty}^{\infty} P_0(k) \frac{-(\sigma - \sigma_1/2) + i\sigma_2/2}{2(\sigma - \sigma_1 + \omega - ku_0 - i\sigma_2)} e^{-ikx} dk. \quad (4.8)$$

The picture of the ocean surface presented by ζ^* is, as will be shown below, one of a damped propagating plane wave system proceeding amid local standing disturbances at any distance which need not necessarily be far away from the origin. This view necessitates the identification of the infinities of the integrands, of C_1 and C_2 , or more precisely, the zeros of the denominators D_{\mp} of these integrands, which are pertinent to our assumption

of small ν ,

$$D_- = \omega - ku_0 - 2i\nu k^2 \left(1 + \frac{\cosh 2kh}{\cosh 4kh - 1} \right) - (i-1)\sigma_1 - \sigma = 0, \quad (4.9)$$

$$D_+ = \omega - ku_0 - 2i\nu k^2 \left(1 + \frac{\cosh 2kh}{\cosh 4kh - 1} \right) - (i+1)\sigma_1 + \sigma = 0. \quad (4.10)$$

Such zeros may be obtained in the form of perturbation expression of (4.9), (4.10) in positive powers of ν with the corresponding zeros of $(D_{\mp})_{\nu=0}$ as initial values.

4.1. Zeros of D_{\mp} for small ν . First we note the following. The equation $D_-(k, 0) = 0$ has only real roots; there are either

- (a) three distinct roots $k_{01}(> 0)$ and $k_{0j}(< 0)$ $j = 2, 3$ or;
- (b) one single root $k_{01}(> 0)$ and a double root $K_0(< 0)$, where $k_{02} < K_0 < k_{03}$ or;
- (c) a single root $k_{01}(> 0)$.

These three cases arise accordingly as $u_0 \leq$ or $-(d/dk)(\sqrt{gk \tanh kh})|_{k=K_0} = u_0^*$ (say) (see Figure 4.1).

(I) Correspondingly, under the same condition on u_0 , and for small ν , we may conclude that the equation $D_-(k, \nu) = 0$ possesses either (i) three roots k_j , $j = 1, 2, 3$ or (ii) three roots k_1, K_2, K_3 , the last 2 corresponding to the double root K_0 , or (iii) only one root k_1 ; these roots $(k_j; K_\ell)_{j=1,2,3, \ell=2,3}$ again reduce to the roots k_{0j} or K_0 $j = 1, 2, 3$ of $D_-(k, 0) = 0$ in the respective cases, as $\nu \rightarrow 0+$.

(II) The equation $D_+(k, 0) = 0$ has only one real root $k_{04}(> 0)$ (Figure 4.2), and also complex roots outside the circle $|k| = \omega'/u'_0$ in the second and third quadrants of the k -plane for $-\pi/4 \leq \text{Im}(k) \leq \pi/4$ in directions less than $2 \sin^{-1}(1/2\sqrt{u'_0 \omega'})$. For small ν , and corresponding to these roots, the equation $D_+(k, \nu) = 0$ then possesses, respectively, the root k_4 and a set of distinct complex roots which, however, leads to waves with enhanced damping due to presence of viscosity. Consequently, we ignore these latter roots and confine attention only to the root k_4 which again, as mentioned in (I), tends to k_{04} as $\nu \rightarrow 0+$.

4.2. Exact form of propagating waves. Considering C_1 of (4.7) first for the case $u_0 < u_0^*$, we use the result of Section 4.1(I) to rewrite it as


$$C_1 = C_{10} + C_{11} + C_{12}, \quad (4.11)$$

where


$$[C_{10}, C_{11}, C_{12}] = \int_{-\infty}^{\infty} e^{-ikx} dk \left[N_1(k) \frac{\partial}{\partial k} \left\{ \ell n \frac{D_-(k)}{\Pi_1^3(k-k_j)} \right\}, \sum_1^3 \frac{N_1(k) - N_1(k_j)}{k - k_j}, \sum_1^3 \frac{N_1(k_j)}{k - k_j} \right],$$

$$N_1(k) = \frac{P_0(k)}{2D_-(k)} \cdot \left(\sigma - \frac{\sigma_1}{2} + \frac{i\sigma_2}{2} \right). \quad (4.12)$$


10 A study of the waves and boundary layers



(a) Three distinct real zeros when $\omega' = 0.2$ and $u'_0 = 0.5$.



(b) Two real zeros, one single and the other repeated when $\omega' = 0.42$ and $u'_0 = 0.5$.



(c) Only a single real zero when $\omega' = 3.5$ and $u'_0 = 1.5$.

Figure 4.1. Illustrative graphs of $D_-(k, 0)$ for the three cases of possible real zeros.




Figure 4.2. Illustrative graph of $D_+(k, 0)$ showing the occurrence of its real zero. $\omega' = 0.3$ and $u'_0 = 0.7$.

Observing that C_{10} is independent of k_j , and that the integrands of C_{10} and C_{11} are regular functions in the immediate vicinity (indented at $k = 0$) of the real axis in the lower or upper half (for $x < 0$ or $x > 0$) of the complex k -plane, we conclude that C_{10} and C_{11} contribute only standing waves to ζ^* of the same frequency as that of the applied pressure.

For C_{12} , we use the result of Gradshteyn and Ryzhik [5, page 445, (2), (4), (6), (8)],

$$\int_{-\infty}^{\infty} \frac{e^{-ikx}}{k - k_j} dk = \pi \left(\sqrt{-\frac{1}{k_j^2}} k_j \cosh [x\sqrt{-k_j^2}] - i \cos [x\sqrt{k_j^2}] \text{Sign}[x] + \left\{ \sqrt{-k_j^4} \text{Sign}[x] \sin [x\sqrt{k_j^2}] \right\} \div \sqrt{-\frac{1}{k_j^2}} \cdot k_j^3 + i \sinh [x\sqrt{-k_j^2}] \right). \quad (4.13)$$

It is then easily verified that the three components of ζ^* arising from three terms of C_{12} each satisfies the wave equation

$$\frac{\partial^2 y}{\partial t^2} = c_j^2 \frac{\partial^2 y}{\partial x^2}, \quad \forall x, t, \text{ where } c_j = \frac{\omega}{k_j}, \quad (4.14)$$

and hence each of these components represents a propagating wave with complex velocity ω/k_j . Thus the progressive wave part of ζ^* arising from C_1 is

$$\zeta_{p1}^* = \frac{e^{i\omega t}}{\sqrt{2\pi\rho g}} \sum_1^3 N_1(k_j) \pi \left(\sqrt{-\frac{1}{k_j^2}} k_j \cosh [x\sqrt{-k_j^2}] - i \cos [x\sqrt{k_j^2}] \text{Sign}[x] + \left\{ \sqrt{-k_j^4} \text{Sign}[x] \sin [x\sqrt{k_j^2}] \right\} \div \sqrt{-\frac{1}{k_j^2}} \cdot k_j^3 + i \sinh [x\sqrt{-k_j^2}] \right). \quad (4.15)$$

For $u_0 = u_0^*$, we have, in view of Section 4.1(I), three progressive waves again together with standing disturbances for all x and t , the former being given by an expression of the same form as above with K_2 and K_3 replacing k_2 and k_3 , respectively.

For $u_0 > u_0^*$, Section 4.1(I) shows that $\zeta_{p_1}^*$ consists of only one progressive wave for all x and t given by the first term of $\zeta_{p_1}^*$ above.

C_2 of (4.8) may be treated similarly, and yields only one progressive wave $\zeta_{p_2}^*$ with complex velocity ω/k_4 together with standing disturbances for all x and t , $\zeta_{p_2}^*$ being of the same form as the first term of $\zeta_{p_1}^*$ with $N_1(k_1)$ replaced by $N_2(k_4)$, where $N_2(k) = (P_0(k)/2D'_+(k)) \cdot (\sigma_1/2 - \sigma + i\sigma_2/2)$.

The final expression for the propagating part of ζ^* arising from C_1 and C_2 is then

$$\begin{aligned} \zeta_p^* &= \zeta_{p_1}^* + \zeta_{p_2}^*, \\ &= \frac{e^{i\omega t}}{\sqrt{2\pi\rho g}} \sum_1^3 N_1(k_j) \pi \left(\sqrt{-\frac{1}{k_j^2}} k_j \cosh [x\sqrt{-k_j^2}] - i \cos [x\sqrt{k_j^2}] \text{Sign}[x] \right. \\ &\quad \left. + \{ \sqrt{-k_j^4} \text{Sign}[x] \sin [x\sqrt{k_j^2}] \} \div \sqrt{-\frac{1}{k_j^2}} \cdot k_j^3 + i \sinh [x\sqrt{-k_j^2}] \right) \\ &\quad + N_2(k_4) \pi \left(\sqrt{-\frac{1}{k_4^2}} k_4 \cosh [x\sqrt{-k_4^2}] - i \cos [x\sqrt{k_4^2}] \text{Sign}[x] \right. \\ &\quad \left. + \{ \sqrt{-k_4^4} \text{Sign}[x] \sin [x\sqrt{k_4^2}] \} \div \sqrt{-\frac{1}{k_4^2}} \cdot k_4^3 + i \sinh [x\sqrt{-k_4^2}] \right). \end{aligned} \quad (4.16)$$

It may be noted that the progressive wave part of ζ^* forms a wave-quartet when $u_0 \leq u_0^*$, and a wave-duo when $u_0 > u_0^*$, for all x and t .

5. Asymptotic expression of ζ^* for large $|x|$

We first observe that the contributions of the standing waves associated with $\zeta_{p_1}^*$ and $\zeta_{p_2}^*$, as referred to in Section 4.2, to ζ^* are insignificant compared to the dominant terms of $\zeta_{p_1}^*$ and $\zeta_{p_2}^*$ for large $|x|$. The dominant term of the asymptotic expression of ζ^* , for large $|x|$, denoted by ζ^{**} , can then be obtained from the asymptotic expansion of ζ_p^* , given by (4.16). We have then

$$\begin{aligned} \zeta^{**} &\cong \frac{e^{i\omega t}}{\rho g} (2\pi)^{1/2} i \left[\{ N_1(k_1) e^{-ik_1 x} + H(u_0^* - u_0) N_1(k_3) e^{-ik_3 x} + N_2(k_4) e^{-ik_4 x} \} H(x) \right. \\ &\quad \left. + H(u_0^* - u_0) N_1(k_2) e^{-ik_2 x} H(-x) \right], \\ \zeta^{**} &\cong \frac{e^{i\omega t}}{\rho g} (2\pi)^{1/2} i \left[\{ N_1(k_1) e^{-ik_1 x} + N_1(K_3) e^{-iK_3 x} + N_2(k_4) e^{-ik_4 x} \} H(x) \right. \\ &\quad \left. + N_1(K_2) e^{-iK_2 x} H(-x) \right], \quad u_0 = u_0^*. \end{aligned} \quad (5.1)$$

Although these two asymptotic expressions ζ^{**} of ζ^* are in agreement in form with those of Pramanik [7], who performed an $O(\sqrt{\nu})$ analysis, there are sharp distinctions with regard to the content which arises mainly due to our $O(\nu)$ analysis of the problem. It will be seen later (Section 6) that a restriction to an $O(\sqrt{\nu})$ analysis makes the long-distance wave form inaccurate for large h .

6. Cayley expansions of k_j , $N_\ell(k_j)$, and of ζ^{**}

6.1. Expansions of k_j and $N_\ell(k_j)$ in terms of their nonviscous values in a power series of $\nu^{1/2}$ or $\nu^{1/4}$ for small ν . For actual as well as numerical calculations of the roots ($k_j; K_\ell$) of $D_\mp(k, \nu) = 0$ and also of the wave amplitudes $N_\ell(k_j)$ for small ν , we may use a result essentially due to Cayley (see Whittaker and Watson [12, page 147]). This enables us to express the above quantities in terms of the corresponding roots in the nonviscous problem in a power series of $\nu^{1/2}$ or $\nu^{1/4}$, respectively, for the two types of roots. To this purpose, we rewrite (4.9) and (4.10) in their equivalent forms ((a) and (b)) for the (first, second types) of roots, respectively,

(a) for

$$u_0 \neq u_0^*, \quad (6.1)$$

$$[(4.9), (4.10)] \longrightarrow \varphi_\mp(k) - \sqrt{\nu}f(k) = 0, \quad f(k) \neq 0, \quad (6.2)$$

where

$$\varphi_-(k) = e^{-3i\pi/4} \sinh 2|k|h \left\{ \sigma^{1/2} - \sqrt{-4(2 \sinh^2 2kh + \cosh 2kh)(\omega - ku_0 - \sigma) + \sigma} \right\}, \quad (6.3)$$

$$f(k) = 2|k|(2 \sinh^2 2kh + \cosh 2kh),$$

and $\varphi_+(k)$ is obtained from $\varphi_-(k)$ on replacing $e^{-(3i\pi)/4}$ and $(\omega - ku_0 - \sigma)$ by $e^{(3i\pi)/4}$ and $(\omega - ku_0 + \sigma)$, respectively. We have (Cayley), for the three roots of (4.9) corresponding to the nonrepeated roots of $D_-(k, 0) = 0$, and for the single root of (4.10) corresponding to the root of $D_+(k, 0) = 0$, the result

$$k_j = k_{0j} + \sqrt{\nu}k_{1j} + \nu k_{2j} + \dots, \quad (6.4)$$

where

$$k_{1j} = \frac{f(k_{0j})}{\varphi'_\mp(k_{0j})}, \quad k_{2j} = \frac{1}{2\{\varphi'_\mp(k_{0j})\}^3} \left| \begin{array}{cc} \varphi'_\mp & f^2 \\ \varphi''_\mp & (f^2)' \end{array} \right|_{k=k_{0j}}, \quad (6.5)$$

upper sign for $u_0 < u_0^*$, $j = 1, 2, 3$ the lower sign when $j = 4$;

(b) for $u_0 = u_0^*$, we replace (4.9) by the two equations

$$\varphi_1(k) - (-1)^\ell \nu^{1/4} f_1(k) = 0, \quad \ell = 2, 3, \quad (6.6)$$

where

$$\varphi_1(k) = \sqrt{\varphi_-(k)}, \quad f_1(k) = \sqrt{f(k)} \quad (6.7)$$

so that by the same result cited above, the roots K_j corresponding to a double root $K_0 \equiv K_{02} \equiv K_{03}$ may be expressed in a positive integral powers of $\nu^{1/4}$ as

$$K_j = K_{0j} + (-1)^j \nu^{1/4} K_{1j} + \nu^{1/2} K_{2j} + (-1)^j \nu^{3/4} K_{3j} + \nu K_{4j} \pm \dots, \quad j = 2, 3, \quad (6.8)$$

here $K_{02} = K_{03} = K_0$ and the coefficients K_{ij} , $i = 1, 2, 3, 4$ are given by the same formula as in the case (6.2) with φ_1 and f_1 replacing φ_- and f .

For $N_\ell(k_j)$, series expansions similar to the above follow from Cayley's general expansion formula [12, page 147]. The corresponding coefficients are the same as those given later (Section 6.2) in consideration with the asymptotic expression of ζ^{**} .

An interesting conclusion from the above expansions is that ζ_p^* in (4.16) is obtained correctly for large depth only when an $O(\nu)$ analysis of the problem is carried out instead of restricting it to an $O(\sqrt{\nu})$ (or, $O(\nu^{3/4})$) one. For instance, in the case $u_0 < u_0^*$, if $kh \gg 1$ $\varphi_\mp \sim (1/2)e^{\mp 3i\pi/4} e^{4kh} \sqrt{\omega - ku_0 \mp \sigma}$ and $f \sim |k|e^{4kh}$, $k_{1j} \rightarrow 0$ as $h \rightarrow \infty$ but k_{2j} does not tend to zero in the same limit. Hence if we keep ζ_p^* in (4.16) correct up to $O(\sqrt{\nu})$ terms, we get the paradoxical result that ζ_p^* is independent of ν as $h \rightarrow \infty$. The presence of $O(\nu)$ term in ζ_p^* shows that there is no such paradox, and leads to a correct value of ζ_p^* for large depths. When $u_0 = u_0^*$, since $\varphi'_1 = \sqrt{\varphi''_1/2}$, we may show similarly that $K_{\ell j} \rightarrow 0$, $\ell = 1, 2, 3$, as $h \rightarrow \infty$ but K_{4j} does not tend to zero as $h \rightarrow \infty$; it follows as before that ζ_p^* for large depths is given correctly only when terms at least of the order $O(\nu)$ are retained. This justifies our assertion in Section 1 regarding the uniform validity of ζ_p^* for large h .

6.2. Cayley expansion of ζ^{} .** Use of Cayley expansion of k_j and $N_\ell(k_j)$ as mentioned above helps us to visualize ζ^{**} in a novel way. For then ζ^{**} assumes the following form (a) or (b):

(a) when $u_0 \neq u_0^*$,

$$\zeta^{**} = \frac{\sqrt{2\pi}i}{\rho g} [\zeta_0 + \sqrt{\nu}\zeta_2 + \nu\zeta_4], \quad (6.9)$$

where

$$\begin{aligned} \zeta_\ell &= H(x)\{M_\ell \times E_1\} + H(-x)H(u_0^* - u_0)N_{\ell 1}(k_{02})e^{-i(k_2 x - \omega t)}, \quad \ell = 0, 2, 4, \\ M_\ell &= \begin{bmatrix} N_{\ell 1}(k_{01}) & H(u_0^* - u_0) & N_{\ell 1}(k_{03}) & N_{\ell 2}(k_{04}) \end{bmatrix}_{1 \times 3}, \quad \ell = 0, 2, 4, \\ E_1 &= \begin{bmatrix} e^{-i(k_1 x - \omega t)} & e^{-i(k_3 x - \omega t)} & e^{-i(k_4 x - \omega t)} \end{bmatrix}^T, \\ N_{0j}(\alpha) &= N_j(\alpha), \quad N_{2j} = \frac{f(\alpha)}{\varphi'_\mp(\alpha)} N'_j(\alpha), \end{aligned} \quad (6.10)$$

$$N_{4j} = \frac{1}{2\{\varphi'_\mp(\alpha)\}^3} \left| \begin{array}{c} \varphi'_\mp \\ (f^2 N'_j) \\ \varphi''_\mp \\ (f^2 N'_j)' \end{array} \right|_{k=\alpha}, \quad j = 1, 2,$$

where f and φ_\mp are given by (6.4), φ_- is to be taken when $\alpha = k_{01}, k_{02}, k_{03}$, while φ_+ of (6.4) is to be modified to φ_+ as mentioned in Section 4.2 when $\alpha = k_{04}$;

(b) when $u_0 = u_0^*$,

$$\zeta^{**} = \frac{\sqrt{2\pi i}}{\rho g} [\zeta'_0 + \nu^{1/4} \zeta'_1 + \nu^{1/2} \zeta'_2 + \nu^{3/4} \zeta'_3 + \nu \zeta'_4], \quad (6.11)$$

where

$$\begin{aligned} \zeta'_\ell &= H(x)\{J_\ell \times E_2\} + H(-x)J_{\ell 1}(K_{02})e^{-i(k_2 x - \omega t)}, \quad \ell = 0, 1, 2, 3, 4, \\ J_\ell &= \begin{cases} \begin{bmatrix} N_{\ell 1}(k_{01}) & N_{\ell 1}(K_{03}) & N_{\ell 2}(k_{04}) \end{bmatrix}_{1 \times 3}, & \ell = 0, 2, 4, \\ \begin{bmatrix} 0 & -N_{\ell 1}(K_{03}) & 0 \end{bmatrix}_{1 \times 3}, & \ell = 1, 3, \end{cases} \\ E_2 &= \begin{bmatrix} e^{-i(k_1 x - \omega t)} & e^{-i(K_3 x - \omega t)} & e^{-i(k_4 x - \omega t)} \end{bmatrix}^T, \\ N_{0j}(\alpha) &= N_j(\alpha), \\ N_{2j} &= \frac{1}{2\{\varphi'_\mp(\alpha)\}^3} \left| \begin{array}{c} \varphi'_\mp \\ \varphi''_\mp \end{array} \right|_{k=\alpha} \begin{array}{c} (f_1^2 N'_j) \\ (f_1^2 N'_j)' \end{array} \quad \text{when } \alpha = k_{01}, k_{04}, j = 1, 2, \\ N_{11} &= \frac{f_1(K_{03})}{\varphi'_1(K_{03})} N'_j(K_{03}), \text{ and so forth.} \end{aligned} \quad (6.12)$$

We note that each ζ_j is a wave-quartet or a wave-duo and the terms of (6.9) and (6.11) are scaled down in powers of $\nu^{1/2}$ or $\nu^{1/4}$, respectively. Further the exponential parts of the four or two components of ζ_j remain invariant for all j while the nonexponential parts are independent of ν , these being functions of k_{0j} only. The strength of these expansions becomes more evident as we proceed to make a numerical study of the wave motion and also compare these solutions in exact form with those of their asymptotic counterpart.

7. Cross-vertical vorticity and boundary layers and their asymptotic expansion

In what follows, we show how to obtain approximate expressions of the thickness of the boundary layers which are formed at the surface and at the bottom because of the motion having a high Reynolds number ($\sqrt{gh^3}/\nu$). We note that viscosity generates only the cross-vertical vorticity ω_z which is obtained from (2.5) with the help of (2.6) and (2.9) as

$$\begin{aligned} \omega_z &= \frac{1}{(2\pi)2^{3/2}i\rho} \int_{-\infty}^{\infty} \text{sgn } k e^{-ikx} dk \int_{c-i\infty}^{c+i\infty} \tilde{P}_0 e^{st} ds \\ &\times \left([2k^2(m - ik^2)\{|k| \sinh m(y+h) \text{sech } mh - m \cosh m(y+h) \text{sech } mh \tanh |k|h\} \right. \\ &\quad + k^2(k^2 + m^2)(1-i)\{m \sinh |k|(y+h) \text{sech } |k|h \\ &\quad \quad \quad \left. - |k| \cosh |k|(y+h) \text{sech } |k|h \tanh mh\} \\ &\quad - |k|(k^2 + m^2)(m^2 - ik^2) \sinh my \text{sech } mh \text{sech } |k|h \\ &\quad \left. - 2k^4(1-i)m \sinh |k|y \text{sech } mh \text{sech } |k|h\right] \div (\nu^{-3/2} f(\nu^{1/2})). \end{aligned} \quad (7.1)$$

If δ_s , δ_b are the thicknesses of the surface and bottom boundary layers, $\omega_z = 0$ for $\delta_s \leq |y| \leq h - \delta_b$, because the motion is irrotational just outside the boundary layers. An approximate value of δ_s may thus be obtained as

$$\delta_s = -\frac{\omega_z)_{y=0}}{\partial\omega_z/\partial y]_{y=0}}, \quad (7.2)$$

with a similar expression for δ_b . To obtain the expression for thickness δ_s^* of the layer below the leading steady-state wave, we analyze the expressions for $\omega_z^*)_{y=0}$ and $\partial\omega_z^*/\partial y]_{y=0}$ in the same way as done for the determination of ζ correct up to $O(\nu)$ terms. This gives, possibly for further restrictions on the order of $P_0(k)$ as $|k| \rightarrow \infty$, the following:

$$\omega_z^*)_{y=0} \cong \frac{e^{i\omega t}}{\sqrt{2\pi\rho g}}(C_4 + C_5), \quad \frac{\partial\omega_z^*}{\partial y}]_{y=0} \cong \frac{e^{i\omega t}}{\sqrt{2\pi\rho g}}(C_6 + C_7), \quad (7.3)$$

$$\delta_s^* = -\frac{\omega_z^*)_{y=0}}{\partial\omega_z^*/\partial y]_{y=0}},$$

here

$$C_4 = \int_{-\infty}^{\infty} \frac{P_0(k)S_1(k)}{-\sigma + \sigma_1 + \omega - ku_0 - i\sigma_2} e^{-ikx} dk,$$

$$C_5 = \int_{-\infty}^{\infty} \frac{-P_0(k)\bar{S}_1(k)}{\sigma - \sigma_1 + \omega - ku_0 - i\sigma_2} e^{-ikx} dk, \quad (7.4)$$

$$C_6 = \int_{-\infty}^{\infty} \frac{P_0(k)S_2(k)}{-\sigma + \sigma_1 + \omega - ku_0 - i\sigma_2} e^{-ikx} dk,$$

$$C_7 = \int_{-\infty}^{\infty} \frac{-P_0(k)\bar{S}_2(k)}{\sigma - \sigma_1 + \omega - ku_0 - i\sigma_2} e^{-ikx} dk,$$

where $S_1(k)$ and $S_2(k)$ are given by

$$S_1(k) = k^2 \left[\left(\frac{3\sigma\sigma_2}{2} + \sigma^2 - \sigma_1^2 - \frac{3\sigma\sigma_1}{2} \right) + i \left(\frac{3\sigma\sigma_2}{2} - \sigma^2 - \sigma_1^2 + \frac{3\sigma\sigma_1}{2} \right) \right],$$

$$S_2(k) = \frac{k^2\sigma^{3/2}}{2^{1/2}} \left[\left\{ 4(\sigma_2 - \sigma_1) + 2\sigma + \frac{3\sigma_1^2}{4} + k^2 \tanh |k|h + 5|k|\sigma_1\nu^{1/2} \right\} \right. \\ \left. + i \left\{ 4(\sigma_2 + \sigma_1) - 2\sigma - \frac{3\sigma_1^2}{4} - 7k^2 \tanh |k|h + |k|(5\sigma_1 - 2\sigma) \right\} \right]. \quad (7.5)$$

Denoting the asymptotic value of ω_z^* $_{y=0}$ and $\partial\omega_z^*/\partial y$ $_{y=0}$ at large distances by ω_z^{**} and $\partial\omega_z^{**}/\partial y$, we have

$$\omega_z^{**} \cong \frac{e^{i\omega t}}{\rho g} (2\pi)^{1/2} i \left[\{G_1(k_1)e^{-ik_1x} + H(u_0^* - u_0)G_1(k_3)e^{-ik_3x} + G_2(k_4)e^{-ik_4x}\}H(x) \right. \\ \left. + H(u_0^* - u_0)G_1(k_2)e^{-ik_2x}H(-x) \right], \quad (7.6)$$

where

$$G_1(\alpha) = \left[\frac{P_0(k)S_1(k)}{\{-\sigma + \sigma_1 + \omega - ku_0 - i\sigma_2\}'} \right]_{k=\alpha}, \\ G_2(k_4) = \left[\frac{-P_0(k)\bar{S}_1(k)}{\{\sigma - \sigma_1 + \omega - ku_0 - i\sigma_2\}'} \right]_{k=k_4},$$

$$\frac{\partial\omega_z^{**}}{\partial y} \cong \frac{e^{i\omega t}}{\rho g} (2\pi)^{1/2} i \left[\{H_1(k_1)e^{-ik_1x} + H(u_0^* - u_0)H_1(k_3)e^{-ik_3x} + H_2(k_4)e^{-ik_4x}\}H(x) \right. \\ \left. + H(u_0^* - u_0)H_1(k_2)e^{-ik_2x}H(-x) \right], \quad (7.7)$$

where

$$H_1(\alpha) = \left[\frac{P_0(k)S_2(k)}{\{-\sigma + \sigma_1 + \omega - ku_0 - i\sigma_2\}'} \right]_{k=\alpha}, \\ H_2(k_4) = \left[\frac{-P_0(k)\bar{S}_2(k)}{\{\sigma - \sigma_1 + \omega - ku_0 - i\sigma_2\}'} \right]_{k=k_4}, \quad (7.8)$$

here accent (') denotes d/dk .

By applying Cayley's expansions again as in the case of (6.9), we can find the asymptotic expansion of ω_z^{**} and $\partial\omega_z^{**}/\partial y$ at large distances in powers of $\nu^{1/2}$ as when $u_0 \neq u_0^*$,

$$\omega_z^{**} = \frac{\sqrt{2\pi}i}{\rho g} [\omega_0 + \sqrt{\nu}\omega_2 + \nu\omega_4], \quad (7.9)$$

where

$$\omega_\ell = H(x)\{M_\ell \times E_1\} + H(-x)H(u_0^* - u_0)N_{\ell 1}(k_{02})e^{-i(k_2x - \omega t)}, \quad \ell = 0, 2, 4, \\ M_\ell = \left[G_{\ell 1}(k_{01}) \quad H(u_0^* - u_0)G_{\ell 1}(k_{03}) \quad G_{\ell 2}(k_{04}) \right]_{1 \times 3}, \quad \ell = 0, 2, 4, \\ E_1 = \left[e^{-i(k_1x - \omega t)} \quad e^{-i(k_3x - \omega t)} \quad e^{-i(k_4x - \omega t)} \right]^T, \\ G_0(\alpha) = G_j(\alpha), \quad G_{2j} = \frac{f(\alpha)}{\varphi_{\mp}^{\prime}(\alpha)} G_j^{\prime}(\alpha), \\ G_{4j} = \frac{1}{2\{\varphi_{\mp}^{\prime}(\alpha)\}^3} \left| \begin{array}{c} \varphi_{\mp}^{\prime} \\ \varphi_{\mp}^{\prime\prime} \end{array} \right| \left| \begin{array}{c} (f^2 G_j^{\prime}) \\ (f^2 G_j^{\prime})' \end{array} \right|_{k=\alpha}, \quad j = 1, 2, \quad (7.10)$$

when $u_0 \neq u_0^*$,

$$\frac{\partial \omega_z^{**}}{\partial z} = \frac{\sqrt{2\pi}i}{\sqrt{\gamma\rho g}} [\omega_0 + \sqrt{\gamma}\omega_2 + \gamma\omega_4], \quad (7.11)$$

where

$$\begin{aligned} \tilde{\omega}_\ell &= H(x)\{M_\ell \times E_1\} + H(-x)H(u_0^* - u_0)N_{\ell 1}(k_{02})e^{-i(k_2x - \omega t)}, \quad \ell = 0, 2, 4, \\ M_\ell &= \left[H_{\ell 1}(k_{01}) \quad H(u_0^* - u_0)H_{\ell 1}(k_{03}) \quad H_{\ell 2}(k_{04}) \right]_{1 \times 3}, \quad \ell = 0, 2, 4, \\ E_1 &= \left[e^{-i(k_1x - \omega t)} \quad e^{-i(k_3x - \omega t)} \quad e^{-i(k_4x - \omega t)} \right]^T, \\ H_0(\alpha) &= H_j(\alpha), \quad H_{2j} = \frac{f(\alpha)}{\varphi_\mp'(\alpha)} H_j'(\alpha), \\ H_{4j} &= \frac{1}{2\{\varphi_\mp'(\alpha)\}^3} \left| \begin{array}{cc} \varphi_\mp' & (f^2 H_j') \\ \varphi_\mp' & (f^2 H_j')' \end{array} \right|_{k=\alpha}, \quad j = 1, 2, \end{aligned} \quad (7.12)$$

where f and φ_\mp are given by (6.4), φ_- is to be taken when $\alpha = k_{01}, k_{02}, k_{03}$, while φ_- of (6.4) is to be modified to φ_+ as mentioned in Section 4.2 when $\alpha = k_{04}$.

From (7.2), we have by the help of (7.9) the approximate thickness of the surface boundary layer in the asymptotic case as

$$\delta_s^{**} = \frac{\omega_0}{\tilde{\omega}_0} \sqrt{\gamma} + \frac{\omega_2 \tilde{\omega}_0 - \omega_0 \tilde{\omega}_2}{\tilde{\omega}_0^2} \gamma. \quad (7.13)$$

We note that δ_s^* and δ_s^{**} are both independent of t , in other words, the shape of the steady-state boundary layer is not affected by the periodicity of the surface forcing.

8. Physical conclusions with numerical illustrations of the exact and asymptotic forms of steady-state progressive waves and boundary layer thickness

Here we mention some physical features of the exact steady-state progressive waves exact and asymptotic as expressed by the result derived in Sections 4.2 and 6.2. We also undertake a numerical evaluation of (4.16), (6.9), and (6.11) to illustrate these features along with that of the surface boundary layer thicknesses in the exact and asymptotic forms as shown in (7.3) and (7.13). For this, we assume a general parabolic distribution of applied surface pressure of the form

$$p_0(x) = \begin{cases} A(1-x^2)^{\lambda-1} H(t) & \text{when } |x| \leq 1, \lambda > \frac{15}{4}, \\ 0 & \text{when } |x| > 1, \end{cases} \quad H(t) \text{ is the Heaviside unit function,} \quad (8.1)$$

which gradually falls from a maximum value at the origin to zero on a finite strip of the free surface. Clearly then, we have $P_0(k) = \{8\sqrt{2}A\Gamma(1/2)J_4(k)\}/k^4$, taking $\lambda = 9/2$.




Figure 8.1




Figure 8.2

We first provide a numerical evaluation of the progressive part of the exact wave form represented by ζ_p^* (4.16) when $u_0 > u_0^*$ for moderate distances. If the asymptotic wave form (6.9) is also computed under the same situation, we notice a close resemblance between the two wave forms albeit with a lateral shift (Figure 8.1). This indicates the




Figure 8.3




Figure 8.4

role of the damping factor in obliterating quickly the differences between the two with increasing distances.

Next we show that at large distances, the leading term ζ_0 of the Cayley asymptotic expansion (6.9) is alone a sufficient approximation of ζ_p^* , which is the exact progressive wave component of ζ^* . Another observation of interest is that for any specified range,




Figure 8.5




Figure 8.6. When the depth is ten times, and as a result we have taken $\omega' = 6.3 \times \sqrt{10}$, $u'_0 = (1/\sqrt{10}) \times 10$, $\nu' = (1/10^{3/2}) \times 0.0002$.

when the distances are doubled, the wave amplitudes are on the average approximately halved. These are shown in Figures 8.2 to 8.4.

For the long-distance asymptotics (6.9), we note that the other two components $\sqrt{\nu}\zeta_2$ and $\nu\zeta_4$ are insignificant compared to ζ_0 . However ζ_2 and ζ_4 are illustrated separately in Figure 8.5 to emphasize that ultimately ζ_4 dominates ζ_2 which diminishes to zero with




Figure 8.7. The form of dominant ζ_{vis} , near the critical value of u'_0 . $\omega' = 0.1$, $u'_0 = 0.8$, ($u'^*_0 = 0.787$), $\nu' = 0.0002$.




Figure 8.8. The situation of nonviscous ζ , near the critical value of u'_0 . $\omega' = 0.1$, $u'_0 = 0.8$, ($u'^*_0 = 0.787$), $\nu' = 0.0002$.

increasing ocean depth (Figure 8.6). Indeed the orders of magnitude of ζ_2 and ζ_4 are reversed as the ocean depth increases or viscosity diminishes, this also signifies the importance of an $O(\nu)$ analysis of the problem. In Figures 8.7 and 8.8, we compare the wave-heights in the vicinity of a certain critical speed u'^*_0 of the running stream in the viscous and nonviscous fluids, respectively. The latter demonstrates the exceedingly large wave heights which are known to occur in this case; the former however illustrates the finiteness of the wave-amplitude in the same. In Figure 8.9, a comparison sketch of the waves is shown in the viscous and nonviscous case, and the viscous wave damping is well observed here.

We next determine the steady-state surface boundary layer thickness δ^*_s from (7.3) by the same numerical process as used for evaluating the exact progressive wave forms in Section 4.2. The corresponding variation of thickness of the surface boundary layer with distance, for fixed depth and time, is shown in Figure 8.10 along with that of the exact steady-state progressive wave form ζ^*_p evaluated earlier. In conclusion, we would like to




Figure 8.9




Figure 8.10

mention the following dimensionless parameters:

$$\omega' = \omega \sqrt{\frac{h}{g}}, \quad u'_0 = \frac{u_0}{\sqrt{gh}}, \quad \nu' = \frac{\nu}{\sqrt{gh^3}}, \quad x' = \frac{x}{h}, \quad k' = kh, \quad (8.2)$$

which are being used while performing the computational study of the problem for Figures 4.1, 4.2, and 8.1–8.10.

In Figures 8.1–8.5 and 8.9–8.10, the values taken for the nondimensional parameters are $\omega' = 6.3$, $u'_0 = 10$, $\nu' = 0.0002$.

Acknowledgments

The author is deeply indebted to Professor A. R. Sen of the Department of Mathematics, Jadavpur University, Calcutta, for his help and suggestions during the preparation of this paper. The author is also grateful to the referees for suggesting some improvements, which have since been incorporated in the paper.

References

- [1] A. Bandyopadhyay, *Effects of viscosity on linear gravity waves due to surface disturbances in water of finite depth*, ZAMM. Zeitschrift für Angewandte Mathematik und Mechanik **83** (2003), no. 3, 205–213.
- [2] J. T. Beale, *Large-time regularity of viscous surface waves*, Archive for Rational Mechanics and Analysis **84** (1984), no. 4, 307–352.
- [3] N. Bleistein and R. A. Handelsman, *Asymptotic Expansions of Integrals*, Holt, Rinehart & Winston, New York, 1975.
- [4] U. T. Ehrenmark and G. L. Body, *The initial-value problem on a plane beach*, The Quarterly Journal of Mechanics and Applied Mathematics **54** (2001), no. 3, 389–411.
- [5] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, 4th ed., Academic Press, New York, 1965.
- [6] A. K. Nikitin and E. N. Potetyunko, *The three-dimensional Cauchy-Poisson problem for waves on the surface of a viscous liquid of finite depth*, Soviet Physics. Doklady **12** (1967), no. 5, 422–424.
- [7] A. K. Pramanik, *Waves due to an oscillatory pressure on a shallow viscous stream*, The Quarterly Journal of Mechanics and Applied Mathematics **26** (1973), no. 2, 193–204.
- [8] A. Prosperetti, *Viscous effects on small-amplitude surface waves*, Physics of Fluids **19** (1976), no. 2, 195–203.
- [9] I. N. Sneddon, *The Use of Integral Transforms*, McGraw-Hill, New York, 1972.
- [10] C. J. Tranter, *Integral Transforms in Mathematical Physics*, Methuen, London, 1951.
- [11] J. V. Wehausen and E. V. Laitone, *Surface waves*, Handbuch der Physik, Vol. 9, Part 3, Springer, Berlin, 1960, pp. 446–778.
- [12] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, Cambridge University Press, Cambridge, 1920.

Arghya Bandyopadhyay: Department of Mathematics, Khalisani College, Hooghly 712 138, India
E-mail address: b.arghya@gmail.com

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk