

*Research Article*

## **Lie Group Analysis of a Flow with Contaminant-Modified Viscosity**

Raseelo J. Moitsheki

Received 1 November 2006; Revised 19 January 2007; Accepted 8 May 2007

Recommended by Meir Shillor

A class of coupled system of diffusion equations is considered. Lie group techniques resulted in a rich array of admitted point symmetries for special cases of the source term. We also employ potential symmetry methods for chosen cases of concentration and a zero source term. Some invariant solutions are constructed using both classical Lie point and potential symmetries.

Copyright © 2007 Raseelo J. Moitsheki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### **1. Introduction**

There has been a great interest in studying the fluid dynamics of pollutants in rivers, due to increasingly important environmental and engineering problems, and more and more refined models have been proposed [1–6]. Shulka [4] obtained analytical solutions by the Fourier transform method for the case of unsteady transport dispersion of nonconservative pollutant/biochemical oxygen demand with first-order decay under each of the sine and cosine variations of waste discharge concentration at upstream boundary and nonzero initial condition through the river.

In this paper, we focus on a system of partial differential equations derived from both Navier-Stokes and concentration equations for fluid flow. We employ the methods of group of transformations for differential equations to analyze the system. In particular, we make use of Lie point and potential symmetry techniques. Over the past 120 years, the use of groups based on local symmetries originally due to Lie [7] has played an important role in obtaining invariant/similarity solutions of differential equations (see, e.g., [8–13]). Among other types, classical Lie point symmetries are classified as local symmetries and potential symmetries as nonlocal symmetries.

Calculations of symmetries are usually very long and tedious. However, we use the free available program Dimsym [14], which is written as a subprogram for the computer algebra package Reduce [15] to construct the admitted symmetries. This paper is structured as follows; In Section 2, we discuss the derivation of the governing equation. In Section 3, we use classical Lie point symmetry techniques to analyze the system of equations for a river pollution. Some exotic cases for the source term led to extra symmetries being admitted. In Section 4, we discuss the potential symmetries admitted by the system and construct some invariant solutions. Lastly we have conclusion in Section 5.

## 2. Governing equations

The equations governing the water pollution problem is derived from both navier-stokes and concentration equations for fluid flow. The basic assumption in the derivation of such equations is initially the river is incompressible, fully developed with constant viscosity, then a given pollutant is injected into the river and the fluid viscosity then changes due to the concentration of the pollutant, that is, the fluid dynamic viscosity is now pollutant-concentration-dependent. The problem now is to determine the diffusion of pollutant with time and space in the river and the effect of pollutant on the river velocity profiles.

The basic one-dimensional equations in original variables are

$$\rho \frac{\partial u}{\partial t} = -\frac{\partial P}{\partial x} + \frac{\partial}{\partial y} \left( \mu \frac{\partial u}{\partial y} \right) \quad (\text{equation for fluid motion}), \quad (2.1)$$

$$\rho \frac{\partial c}{\partial t} = \frac{\partial}{\partial y} \left( D \frac{\partial c}{\partial y} \right) + S(y, t) \quad (\text{pollutant concentration equation}), \quad (2.2)$$

where

- (i)  $\rho$  = constant river density,
- (ii)  $D$  = mass diffusion,
- (iii)  $S(y, t)$  = external source of pollutant,
- (iv)  $\mu$  = river dynamic viscosity,
- (v)  $u$  = river velocity,
- (vi)  $c$  = pollutant concentration,
- (vii)  $P$  = river pressure,
- (viii)  $t$  = time,
- (ix)  $(x, y)$  = the axial and transverse coordinate, respectively.

At time  $t > 0$ , the fully developed stream is disturbed by injection of pollutant from an external source. The river viscosity and the mass diffusivity is assumed to vary as follows:

$$\frac{\mu}{\mu_0} = \left( \frac{c}{c_0} \right)^\lambda, \quad \frac{D}{D_0} = \left( \frac{c}{c_0} \right)^\lambda, \quad (2.3)$$

where  $\mu_0$ ,  $c_0$ ,  $D_0$ , and  $\lambda$  are the river viscosity coefficient, characteristic concentration, mass diffusivity coefficient, and constant exponent, respectively. The following dimensionless variables are introduced:

$$\begin{aligned} u' &= \frac{uL}{v_0}, & y' &= \frac{y}{L}, & x' &= \frac{x}{L}, & t' &= \frac{tv_0}{L^2}, & P' &= \frac{L^2 P}{\rho v_0^2}, & v_0 &= \frac{\mu_0}{\rho}, \\ R &= \frac{v_0}{D_0}, & K &= -\frac{\partial P'}{\partial x'}, & S' &= \frac{SL^2}{v_0 c_0}, & c' &= \frac{c}{c_0}. \end{aligned} \quad (2.4)$$

Neglecting the prime symbol for clarity, we obtain a dimensionless system of partial differential equations (PDEs):

$$\frac{\partial u}{\partial t} = K + \frac{\partial}{\partial y} \left( c^\lambda \frac{\partial u}{\partial y} \right), \quad (2.5)$$

$$\frac{\partial c}{\partial t} = \frac{1}{R} \frac{\partial}{\partial y} \left( c^\lambda \frac{\partial c}{\partial y} \right) + S(y, t), \quad (2.6)$$

where  $R$  is the Schmidt number and  $K$  is the imposed constant pressure axial gradient.

### 3. Point symmetry reductions of the system of (2.5) and (2.6)

In the initial Lie point symmetry analysis of the system of PDEs (2.5) and (2.6), with  $S$  being an arbitrary function of  $y$  and  $t$ , the admitted generic point symmetries or the principal Lie algebra are two dimensional and spanned by the base vectors

$$\Gamma_1 = \frac{\partial}{\partial u}, \quad \Gamma_2 = (u - Kt) \frac{\partial}{\partial u}. \quad (3.1)$$

The principal Lie algebra extends for the cases listed in Table 3.1. Wherever they appear  $A$ ,  $\alpha$ ,  $\beta$ ,  $m$ ,  $p$ , and  $w$  are arbitrary constants. Note that, following multiplication by the constant  $R$  and then letting it vanish, the resulting equation (2.6) becomes an ordinary differential equation (ODE). This case may not be physically realistic but leads to extra admitted symmetries. The exercise of searching for the forms of arbitrary functions that extend the principal Lie algebra is called *group classification*. The problem of group classification was introduced by Ovsiannikov [16] and recent accounts on this topic may be found for example in [9, 17–19]. In this section we adopt methods in [9] to perform group classification of the system of (2.5) and (2.6).

*Invariant solutions.* Here we consider only two examples for the cases which are more realistic. Note that reduction by other symmetries leads to reduced nonlinear ordinary differential equations.

*Example 3.1.* If we consider from Table 3.1 the case  $S = 0$  with arbitrary  $R$  and  $\lambda$ , then  $\Gamma_3$ -invariant solution is given by

$$u = Kt + c_1, \quad c = y^{2/\lambda} \left\{ -\frac{1}{R} \left( 2 + \frac{4}{\lambda} \right) t + k_1 \right\}^{-1/\lambda}, \quad (3.2)$$

where  $c_1$ , and  $k_1$  are arbitrary constants of integration.

Table 3.1. Special cases of  $S(y, t)$  and extension of the principal Lie algebra.

$S(y, t)$	Constants	Symmetries
Arbitrary	$R = 0, \lambda$ arbitrary	$\Gamma_3 = \partial_y; \Gamma_4 = \frac{2c}{\lambda} \partial_c + y \partial_y$ $\Gamma_5 = \frac{H'(t)}{\lambda} c \partial_c + KH(t) \partial_u + H(t) \partial_t$
0	$R, \lambda$ arbitrary $R = 1, \lambda$ arbitrary $R = 1, \lambda = -2$	$\Gamma_3 = 2Kt \partial_u + y \partial_y + 2t \partial_t; \Gamma_4 = \partial_t;$ $\Gamma_5 = 2c \partial_c + \lambda y \partial_y; \Gamma_6 = \partial_y$ $\Gamma_7 = c \partial_u$ $\Gamma_8 = -c \partial_c + y \partial_y$
$A(\alpha y + \beta)^m e^{wt}$	$R, \lambda$ arbitrary	$\Gamma_3 = c \partial_c + \frac{\lambda}{2} \left( y + \frac{\beta}{\alpha} \right) \partial_y + \frac{1}{2w} (2 - \lambda m) \partial_t$
$e^{ny} t^m$	$R, \lambda$ arbitrary	$\Gamma_3 = \frac{1}{\lambda(m+1)+1} \{ -K \lambda n t \partial_u + n c \partial_c$ $+ [\lambda(m+1)+1] \partial_y - \lambda n t \partial_t \}$
$y^n t^m$	$R, \lambda$ arbitrary $\lambda = \frac{-1}{m+1}, m = \frac{n+2}{-2}$	$\Gamma_3 = \frac{1}{\lambda(m+1)+1} \{ -K(\lambda n + 2)t \partial_u + [2(m+1)+n] c \partial_c$ $+ [\lambda(m+1)+1] y \partial_y - \lambda n t \partial_t \}$ $\Gamma_3 = 2Kt \partial_u - n c \partial_c + 2t \partial_t$ $\Gamma_4 = n c \partial_c + y \partial_y$
$e^{my+nt}$	$\lambda, R$ arbitrary	$\Gamma_3 = \frac{-Km}{n} \partial_u + \partial_y - \frac{m}{n} \partial_t$
$Ae^{wt}$	$R, \lambda$ arbitrary	$\Gamma_3 = 2Kt \partial_u + 2wc \partial_c + \lambda w y \partial_y + 2 \partial_t;$ $\Gamma_4 = \partial_y$
$y^m f(t)$	$\lambda = 2/m$	$\Gamma_3 = m c \partial_c + y \partial_y$
$f(t)$	$R, \lambda$ arbitrary	$\Gamma_3 = \partial_y$
$t^m$	$R, \lambda$ arbitrary	$\Gamma_4 = 2(m+1)c \partial_c + 2Kt \partial_u + 2t \partial_t + [1 + (m+1)\lambda] y \partial_y$
$f(y)$	$R, \lambda$ arbitrary	$\Gamma_3 = K \partial_u$
$e^{my}$	$R, \lambda$ arbitrary	$\Gamma_4 = c \partial_c - K \lambda t \partial_u + \frac{\lambda+1}{m} \partial_y - \lambda t \partial_t$
$y^m$	$R, \lambda$ arbitrary	$\Gamma_4 = c \partial_c + \frac{2-\lambda m}{m+2} K t \partial_u + \frac{\lambda+1}{m+2} y \partial_y + \frac{2-\lambda m}{m+2} t \partial_t$
$m(ny - pt)$	$R, \lambda$ arbitrary	$\Gamma_3 = \frac{p}{n} \partial_y + \partial_t$

Example 3.2. For the case  $S = A(\alpha y + \beta)^m e^{wt}$ , we have  $\Gamma_3$  leads to the invariant solution in functional form

$$c = \exp\left(\frac{2wt}{2-\lambda m}\right) G(\eta), \quad u = H(\eta), \quad \eta = \left(y + \frac{\beta}{\alpha}\right) \exp\left(\frac{-\lambda wt}{2-\lambda m}\right), \quad (3.3)$$

with  $G$  and  $H$  satisfying the system of coupled ODEs

$$\begin{aligned} (2 - \lambda m)G^\lambda G'' + \lambda(2 - \lambda m)G^{\lambda-1}(G')^2 + \lambda R w \eta G' + \lambda^m A R(2 - \lambda m)\eta^m - 2Rw &= 0, \\ \lambda(2 - \lambda m)G^{\lambda-1}G'H' + (2 - \lambda m)G^\lambda H'' + w\lambda\eta H' + K(2 - \lambda m) &= 0, \end{aligned} \quad (3.4)$$

where the prime indicate differentiation with respect to  $\eta$ . This system of ODE has no exact solution for arbitrary values of  $\lambda$ . If  $\lambda = 0$ , then we obtain invariant solutions

$$c = \exp\left(\frac{2wt}{2 - \lambda m}\right)\left(\frac{Rw}{2}y^2 + c_1y + c_2\right), \quad u = -\frac{Ky^2}{2} + c_3y + c_4, \quad (3.5)$$

wherein  $c_i$ 's are constants. However, in this case the system is now linear and uncoupled. Symmetry reductions by other admitted symmetries lead to highly nonlinear ODEs. We herein omit those reductions.

In the next section, we analyze (2.5) and (2.6) using potential symmetry. It is possible when using potential symmetries to construct invariant solutions which cannot be obtained by point symmetries.

#### 4. Potential symmetry reductions and invariant solutions

**4.1. Trivial case.** The most trivial case is given when  $S = 0$  and the solution of (2.2) is a constant, say  $\sqrt[3]{y}$ , that is, given a constant river pollutant concentration. This trivial case implies that (2.1) becomes a linear diffusion equation with a constant diffusion coefficient and a constant  $K$ . In other words, the system of (2.1) and (2.2) reduces to a single PDE written in conserved form

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial y}\left(Ky + y\frac{\partial u}{\partial y}\right). \quad (4.1)$$

Equation (4.1) may be embedded in a system of first order partial differential equation called auxiliary system [9].  $K$  may be equated to zero by the transformation  $\bar{u} = u + Kt$ . However, here we allow  $K$  to be nonzero. Pucci and Saccomandi [20] provided the necessary condition for a PDE to admit potential symmetries. A PDE may be expressed in an auxiliary system in more than one inequivalent way (see, e.g., [20]). For example [21], the inhomogeneous nonlinear diffusion equation

$$u_t = \left[\left(\frac{y}{u}\right)^2 u_y\right]_y \quad (4.2)$$

may be embedded into the auxiliary systems

$$v_y = u, \quad v_t = \left(\frac{y}{u}\right)^2 u_y, \quad (4.3)$$

$$v_y = \frac{u}{y}, \quad v_t = \frac{y}{u^2} u_y - \frac{1}{u}. \quad (4.4)$$

Equation (4.4) admits potential symmetries whereas (4.3) does not. Methods are being developed to find the most general conservation laws or auxiliary systems which admit nonlocal symmetries for the given governing equation (see, e.g., [22, 23]). We observe that (4.1) may be expressed in two distinct systems of first-order PDEs, namely,

$$v_y = u, \quad v_t = Ky + \gamma u_y, \tag{4.5}$$

$$v_y = u - Kt, \quad v_t = \gamma u_y, \tag{4.6}$$

where  $v$  is the potential or nonlocal variable. The first system arises from a simple and natural split of (4.1) whilst the second was, relatively, not quite obvious to construct. We note that system (4.6) possesses point symmetries which induce potential symmetries for (4.1), whereas system (4.5) does not. One may employ the “direct method of finding auxiliary systems” [23]. If we assume that (4.1) can be embedded in the auxiliary system

$$v_y = f(y, t, u, u_t), \tag{4.7}$$

$$v_t = g(y, t, u, u_y). \tag{4.8}$$

For the compatibility of (4.7) and (4.8) we require

$$f_t + u_t f_u + u_{tt} f_{u_t} = g_y + u_y g_u + u_{yy} g_{u_y}, \tag{4.9}$$

on solutions of (4.1). Note that the system of (4.7) and (4.8) is the definition of the conservation law. Hence, by simple calculation, from (4.9) it follows that

$$f = c_1 u + J(y, t), \quad g = \gamma c_1 u_y + H(y, t), \tag{4.10}$$

with  $c_1$  being a constant,  $J$  and  $H$  being functions of  $y$ , and  $t$  satisfying the equation

$$J_t + Kc_1 = H_y. \tag{4.11}$$

Without loss of generality we let  $c_1 = 1$ . The auxiliary system (4.5) may be obtained by letting  $J = 0$ , that way  $H = Ky$ , wherein the integration constant vanishes. Also, auxiliary system (4.6) may be obtained by letting  $H = 0$ , that way  $J = -Kt$  with the integration constant being zero. Note that if the constant of integration is nonzero, then beyond translations, both the auxiliary systems admit infinite point symmetries. The System (4.6) admits a finite six-dimensional Lie algebra spanned by

$$\begin{aligned} \Gamma_1 &= \frac{1}{4y} (-y^2 u - 6y t u - 2y v + k y^2 t + 10K \gamma t^2) \frac{\partial}{\partial u} + t^2 \frac{\partial}{\partial t} - \frac{1}{4y} (y^2 v + 2\gamma v t) \frac{\partial}{\partial v} + y t \frac{\partial}{\partial y}, \\ \Gamma_2 &= 2K t \frac{\partial}{\partial u} + v \frac{\partial}{\partial v} + y \frac{\partial}{\partial y} + 2t \frac{\partial}{\partial t}, \quad \Gamma_3 = K \frac{\partial}{\partial u} + \frac{\partial}{\partial t}, \\ \Gamma_4 &= (u - K t) \frac{\partial}{\partial u} + v \frac{\partial}{\partial v}, \quad \Gamma_5 = \frac{\partial}{\partial y}, \\ \Gamma_6 &= \frac{1}{2y} (-y u - v + K y t) \frac{\partial}{\partial u} - \frac{\gamma v}{2y} \frac{\partial}{\partial v} + t \frac{\partial}{\partial y}, \quad \Gamma_7 = \frac{\partial G}{\partial y} \frac{\partial}{\partial u} + G \frac{\partial}{\partial v}, \end{aligned} \tag{4.12}$$

where  $G = G(y, t)$  is a solution of the linear diffusion equation

$$G_t = \gamma G_{yy}. \quad (4.13)$$

Clearly,  $\Gamma_1$  and  $\Gamma_6$  induce nonlocal symmetries for the governing equation (4.1).

We construct the invariant solution for the system (4.8) using the potential symmetry operator  $\Gamma_1$  and  $\Gamma_6$ . The  $\Gamma_1$ -invariant solution is given in the functional form by

$$\begin{aligned} u &= t^{-3/2} \exp\left(\frac{y^2}{4\gamma t}\right) \int \frac{1}{4\gamma t} \exp\left(-\frac{y^2}{4\gamma t}\right) \left[10\gamma K t^{5/2} - 2y \exp\left(-\frac{y^2}{4\gamma t}\right) G(\rho) + K y^2 t^{3/2}\right] dt; \\ v &= \frac{1}{\sqrt{t}} \exp\left(-\frac{y^2}{4\gamma t}\right) G(\rho), \end{aligned} \quad (4.14)$$

where the similarity variable  $\rho = y/t$ , and the  $\Gamma_6$ -invariant solutions are

$$u = -\frac{A y}{2\gamma \sqrt{t^3}} \exp\left(-\frac{y^2}{4\gamma t}\right) + K t; \quad v = \frac{A}{\sqrt{t}} \exp\left(-\frac{y^2}{4\gamma t}\right). \quad (4.15)$$

Here,  $A$  is an arbitrary constant.

**4.2. Nontrivial case.** One may note that the solution of (2.6) appear as the coefficient in (2.5). Unfortunately, as expected, not all the solutions to (2.6) lead to extra symmetries for (2.5). However, we observe that the invariant solution  $c = k_1/y$  to (2.6) with  $S = 0$  (see, e.g., [9]) leads to a rich array of symmetries, particularly potential symmetries, being admitted by (2.5). In the classical Lie point symmetry analysis we observe that (2.5) admits beside the infinite symmetry generator, the translation in  $t$  and scaling of  $y$ . The linear combination of these finite point symmetries leads to a highly nonlinear ordinary differential equation which was not analytically easy to solve.

On the other hand, (2.5) with  $c = k_1/y$  and  $\lambda = -2$ , may be embedded in an auxiliary system

$$v_y = u - K t, \quad v_t = \left(\frac{k_1}{y}\right)^{-2} u_y, \quad (4.16)$$

where  $v$  is a potential variable. Note that we choose the auxiliary system that gives rise to potential symmetries. The case  $\lambda = -2$  is not realistic, however it is interesting from the mathematical point of view. The integrated form which is equivalent to the auxiliary system (4.16) and arising from letting the potential variable be given by

$$v = \int (u(y, t) - K t) dy + J(t), \quad (4.17)$$

where  $J(t)$  is the integration constant, is given by the nonconstant coefficient linear diffusion equation,

$$v_t = \left(\frac{y}{k_1}\right)^2 v_{yy}. \quad (4.18)$$

Note that (4.18) does not admit potential symmetries, but the auxiliary system (4.16) admits the symmetries

$$\Gamma_1 = \left[ \left( -\frac{k_1^2}{4}u + \frac{k_1^2 K}{4}t - \frac{1}{2}ut - \frac{k_1^2 K}{4y}v + \frac{1}{2}Kt^2 \right) \log y - \frac{1}{4k_1^2}ut^2 - \frac{3}{2}ut + \frac{1}{2y}vt + \frac{Kt^3}{4k_1^2} - \frac{5K}{2}t^2 \right] \frac{\partial}{\partial u} + \left[ \left( -k_1^2 y^2 v + 2vt \right) \log y + t^2 v - 2vt \right] \frac{\partial}{\partial v} + 4ty \log y \frac{\partial}{\partial y} + t^2 \frac{\partial}{\partial t}; \tag{4.19}$$

$$\Gamma_2 = \left[ \left( \frac{u}{4} + \frac{Kt}{4} \right) \log y - \frac{ut}{4k_1^2} - \frac{u}{2} + \frac{v}{4y} + \frac{Kt^2}{4k_1^2} + \frac{3K}{2}t \right] \frac{\partial}{\partial u} + t \frac{\partial}{\partial t} + \frac{y}{2} \log y \frac{\partial}{\partial y} + \left[ \frac{v}{4} \log y - \frac{vt}{4k_1^2} \right] \frac{\partial}{\partial v}; \tag{4.20}$$

$$\Gamma_3 = \frac{\partial}{\partial u} + K \frac{\partial}{\partial t}; \quad \Gamma_4 = (u - Kt) \frac{\partial}{\partial u} + v \frac{\partial}{\partial v}; \quad \Gamma_5 = (u - Kt) \frac{\partial}{\partial u} + y \frac{\partial}{\partial y}; \tag{4.21}$$

$$\Gamma_6 = \left[ \frac{k_1^2}{2}(u - Kt) \log y + \frac{ut}{2} + \frac{k_1^2 v}{2y} - \frac{Kt^2}{2} \right] \frac{\partial}{\partial u} - yt \frac{\partial}{\partial y} + \frac{v}{2}(k_1^2 \log y - t) \frac{\partial}{\partial v};$$

and the infinite symmetry generator

$$\Gamma_\infty = \frac{\partial H(y,t)}{\partial y} \frac{\partial}{\partial u} + H(y,t) \frac{\partial}{\partial v}, \tag{4.22}$$

where  $H$  is an arbitrary solution of (4.18).  $\Gamma_1, \Gamma_2,$  and  $\Gamma_6$  are potential symmetries. Equation (4.18) may naturally split into another auxiliary system using a different potential variable, say  $w$ . However, the potential symmetries that may rise in terms of the variable  $w$  are not transformable to the symmetries in given (4.20). The genuine potential symmetries lead to the functional form of the invariant solution. As an example, we construct the solution using the  $\Gamma_6$  generator.  $\Gamma_6$ -invariant solution is given by

$$v = \sqrt{y} \exp \left( -\frac{k_1^2}{4t}(\log y)^2 \right) G(t), \tag{4.23}$$

$u =$

$$\left\{ \int \frac{\exp \left( -k_1^2(\log y)^2/4t \right) (k_1^2 K t y \log y - k_1^2 \sqrt{y} \exp \left( -k_1^2(\log y)^2/4t \right) G(t) + K y t^2}{2y^{5/2}t} dy + c_1 \right\} \cdot y^{(k_1^2/4t) \log y + 1/2}, \tag{4.24}$$

where  $G(t)$  is an arbitrary function of  $t$  and  $c_1$  is a constant.

### 5. Some discussions and concluding remarks

The analysis in this paper refers to flow along a single channel. Even if water viscosity does not vary much, the analysis applies equally well to industrial fluid mixtures, for example water-alcohol mixtures, wherein this effect may be more important.



The problem has been mathematically simplified by assuming both viscosity and diffusivity to vary with concentration in the same way, that is, the power law with the same exponent. Further analysis of the system of (2.5) and (2.6) with distinct exponents for viscosity and diffusivity may be an interesting topic for future study.

Lie group analysis resulted in some exotic admitted symmetries and hence some invariant solution. We observed that the problem is difficult when the source term is nonzero. However, we made use of nonconstant concentration that appeared as coefficient in the river velocity equation. A rich symmetry structure indicated in Table 3.1 may lead to reductions and extra invariant solutions.

## Acknowledgments

The author wishes to thank the National Research Foundation of South Africa under Thuthuka program, for the generous financial support, and Prof. O. D. Makinde for the useful discussions. The author is grateful to the reviewers for the most valuable comments and improvements to this manuscript.

## References

- [1] H. A. Basha and F. S. El-Habel, "Analytical solution of the one-dimensional time-dependent transport equation," *Water Resources Research*, vol. 29, no. 9, pp. 3209–3214, 1993.
- [2] R. B. Bird, W. E. Stewart, and E. N. Lightfoot, *Transport Phenomena*, John Wiley & Sons, New York, NY, USA, 1960.
- [3] M. Munafò, G. Cecchi, F. Baiocco, and L. Mancini, "River pollution from non-point sources: a new simplified method of assessment," *Journal of Environmental Management*, vol. 77, no. 2, pp. 93–98, 2005.
- [4] P. Shulka, "Analytical solutions for unsteady transport dispersion of nonconservative pollutant with time-dependent periodic waste discharge concentration," *Journal of Hydraulic Engineering*, vol. 128, no. 9, pp. 866–869, 2002.
- [5] G. I. Taylor, "The dispersion of matter in turbulent flow through pipe," *Proceedings of the Royal Society of London*, vol. 233, no. 1155, pp. 446–448, 1954.
- [6] C. Zheng and G. D. Bennet, *Applied Contaminant Transport Modeling: Theory and Practice*, Van Nostrand Reinhold, New York, NY, USA, 1995.
- [7] S. Lie, "Über integration durch bestimmte integrals von einer klasse linearer partieller differentialgleichungen," *Archivum Mathematicum*, vol. 6, pp. 328–368, 1881.
- [8] G. W. Bluman and S. C. Anco, *Symmetry and Integration Methods for Differential Equations*, vol. 154 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 2002.
- [9] G. W. Bluman and S. Kumei, *Symmetries and Differential Equations*, vol. 81 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1989.
- [10] N. H. Ibragimov, Ed., *CRC Handbook of Lie Group Analysis of Differential Equations. Vol. 1: Symmetries, Exact Solutions and Conservation Laws*, CRC Press, Boca Raton, Fla, USA, 1994.
- [11] N. H. Ibragimov, Ed., *CRC Handbook of Lie Group Analysis of Differential Equations. Vol. 2: Applications in Engineering and Physical Sciences*, CRC Press, Boca Raton, Fla, USA, 1995.
- [12] N. H. Ibragimov, Ed., *CRC Handbook of Lie Group Analysis of Differential Equations. Vol. 3: New Trends in Theoretical Development and Computational Methods*, CRC Press, Boca Raton, Fla, USA, 1996.
- [13] P. J. Olver, *Applications of Lie Groups to Differential Equations*, vol. 107 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 1986.

- [14] J. Sherring, “DIMSYM symmetry determination and linear differential equation package,” La-trobe University, 1993.
- [15] A. C. Hearn, *REDUCE*, Rand Corporation Publication CP78, Santa Monica, Calif, USA, 1985.
- [16] L. V. Ovsiannikov, “Group relations of the equation of non-linear heat conductivity,” *Doklady Akademii Nauk SSSR*, vol. 125, pp. 492–495, 1959.
- [17] N. M. Ivanova and C. Sophocleous, “On the group classification of variable-coefficient nonlinear diffusion-convection equations,” *Journal of Computational and Applied Mathematics*, vol. 197, no. 2, pp. 322–344, 2006.
- [18] C. Sophocleous, “Symmetries and form-preserving transformations of generalised inhomogeneous nonlinear diffusion equations,” *Physica A*, vol. 324, no. 3-4, pp. 509–529, 2003.
- [19] O. O. Vaneeva, A. G. Johnpillai, R. O. Popovych, and C. Sophocleous, “Enhanced group analysis and conservation laws of variable coefficient reaction-diffusion equations with power nonlinearities,” *Journal of Mathematical Analysis and Applications*, vol. 330, no. 2, pp. 1363–1386, 2007.
- [20] E. Pucci and G. Saccomandi, “Potential symmetries and solutions by reduction of partial differential equations,” *Journal of Physics A: Mathematical and General*, vol. 26, no. 3, pp. 681–690, 1993.
- [21] R. J. Moitsheki, P. Broadbridge, and M. P. Edwards, “Systematic construction of hidden non-local symmetries for the inhomogeneous nonlinear diffusion equation,” *Journal of Physics A: Mathematical and General*, vol. 37, no. 34, pp. 8279–8286, 2004.
- [22] S. C. Anco and G. W. Bluman, “Direct construction method for conservation laws of partial differential equations—part II: general treatment,” *European Journal of Applied Mathematics*, vol. 13, no. 5, pp. 567–585, 2002.
- [23] J. Goard, personal correspondence, 2004.

Raseelo J. Moitsheki: Department of Mathematics, Vaal University of Technology, Private bag X021, Vanderbijlpark 1900, South Africa  
Email address: joelm@vut.ac.za

## Special Issue on Space Dynamics

### Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

### Lead Guest Editor

**Antonio F. Bertachini A. Prado**, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; [prado@dem.inpe.br](mailto:prado@dem.inpe.br)

### Guest Editors

**Maria Cecilia Zanardi**, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; [cecilia@feg.unesp.br](mailto:cecilia@feg.unesp.br)

**Tadashi Yokoyama**, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; [tadashi@rc.unesp.br](mailto:tadashi@rc.unesp.br)

**Silvia Maria Giuliatti Winter**, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; [silvia@feg.unesp.br](mailto:silvia@feg.unesp.br)