

Research Article

Asymptotic Behavior of a Competition-Diffusion System with Variable Coefficients and Time Delays

Miguel Uh Zapata, Eric Avila Vales, and Angel G. Estrella

*Facultad de Matemáticas, Universidad Autónoma de Yucatán, Periférico Norte, Tablaje 13615,
C.P. 97119, Mérida, Yucatán, Mexico*

Correspondence should be addressed to Eric Avila Vales, avila@uady.mx

Received 26 June 2007; Accepted 15 January 2008

Recommended by Malgorzata Peszynska

A class of time-delay reaction-diffusion systems with variable coefficients which arise from the model of two competing ecological species is discussed. An asymptotic global attractor is established in terms of the variable coefficients, independent of the time delays and the effect of diffusion by the upper-lower solutions and iteration method.

Copyright © 2008 Miguel Uh Zapata et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and main result

The Lotka-Volterra competition model with diffusion and time delays has been the object of analysis by numerous authors under different approaches. For the case of two species, Ruan and Zhao [1] considered uniform persistence and global extinction; Lu [2] studied global attractivity, and Gourley and Ruan [3] analyzed stability and traveling fronts. The periodic case has also been considered, Feng and Wang [4] studied asymptotic stability and Zhou et al. [5] investigated the Hopf bifurcation. The cases of three and N -species have also been analyzed in [6–8].

In this paper, we consider the asymptotic behavior of solutions for the competition-diffusion system with time delays of the following two species:

$$\frac{\partial u_1}{\partial t} = A\Delta u_1(t, x) + u_1(t, x) \left[a_1(t, x) - b_1(t, x)u_1(t, x) - c_1(t, x) \int_0^\infty u_1(t-\tau, x) d\mu_1(\tau) - d_1u_2(t-r_2, x) \right],$$

$t > 0, \quad x \in \Omega,$

$$\frac{\partial u_2}{\partial t} = A\Delta u_2(t, x) + u_2(t, x) \left[a_2(t, x) - b_2(t, x)u_2(t, x) - c_2(t, x) \int_0^\infty u_2(t-\tau, x) d\mu_2(\tau) - d_2u_1(t-r_1, x) \right],$$

$t > 0, \quad x \in \Omega,$

$$\begin{aligned} \frac{\partial u_1}{\partial \eta} = \frac{\partial u_2}{\partial \eta} = 0, \quad t > 0, \quad x \in \partial\Omega, \\ u_i(t, x) = \phi_i(t, x) \quad (i = 1, 2) \quad t \leq 0, \quad x \in \overline{\Omega}, \end{aligned} \quad (1.1)$$

where $\Omega \subseteq \mathbb{R}^N$ ($N \geq 1$) is a bounded domain with smooth boundary $\partial\Omega$, $\partial/\partial\eta$ denotes the differentiation outward normal on $\partial\Omega$, $A \geq 0$, $d_i > 0$, $r_i > 0$, $0 < a_i \leq a_i(t, x) \leq A_i$, $0 < b_i \leq b_i(t, x) \leq B_i$, $0 \leq c_i \leq c_i(t, x) \leq C_i$, $\mu_i(\cdot)$ is of bounded variation with $\mu_i(0) = 0$, $\phi_i \in C^1((-\infty, 0]) \times \overline{\Omega}$ is bounded and nonnegative and $0 \neq \phi_i(0, \cdot) \in C^1(\overline{\Omega})$ ($i = 1, 2$), and u_1 and u_2 are the density functions of two species competing for a shared limited resource.

The functions $a_1(x, t)$ and $a_2(x, t)$ denote the intrinsic growth rate of the species, $b_1(x, t)$ and $b_2(x, t)$ represent self-limitation rates, and $c_1(x, t)$ and $c_2(x, t)$ represent the coefficients of the infinite continuous delay. The constants d_1, d_2 represent the competition rates. The distributed time delay should be viewed as the effects of past history.

Let $M_i(t)$ denote the total variation of $\mu_i(\cdot)$ on $[0, t]$ and let $M_i^\pm(t) = (M_i(t) \pm \mu_i(t))/2$, for all $t \in \mathbb{R}^+$. Then, $M_i(t)$ and $M_i^\pm(t)$ are nonnegative and nondecreasing on \mathbb{R}^+ . It easy to see that

$$M_i^+(t) + M_i^-(t) = M_i(t), \quad M_i^+(t) - M_i^-(t) = \mu_i(t). \quad (1.2)$$

Denote $M_{0i} = \lim_{t \rightarrow \infty} M_i(t)$, $M_{0i}^\pm = \lim_{t \rightarrow \infty} M_i^\pm(t)$, and $\mu_{0i} = \lim_{t \rightarrow \infty} \mu_i(t)$. Then,

$$M_{0i}^+ + M_{0i}^- = M_{0i}, \quad M_{0i}^+ - M_{0i}^- = \mu_{0i}. \quad (1.3)$$

Our result can be stated as follows.

Theorem 1.1. Assume that $b_i > c_i M_{0i}^-$, $a_i - (C_i M_{0i}^+ A_i) / (b_i - c_i M_{0i}^-) > 0$, and

$$\frac{d_2 A_2 (b_2 - c_2 M_{02}^-)}{(b_1 - c_1 M_{01}^-) [a_2 (b_2 - c_2 M_{02}^-) - C_2 M_{02}^+ A_2]} < \frac{A_2}{A_1} < \frac{(b_2 - c_2 M_{02}^-) [a_1 (b_1 - c_1 M_{01}^-) - C_1 M_{01}^+ A_1]}{d_1 A_1 (b_1 - c_1 M_{01}^-)}. \quad (1.4)$$

Then, for any $\phi_i \in C^1((-\infty, 0]) \times \overline{\Omega}$ with $\phi(0, x) \neq 0$, the solution of (1.1) satisfies

$$\begin{aligned} 0 < \alpha_1 \leq \liminf_{t \rightarrow \infty} \min_{x \in \overline{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \overline{\Omega}} u_1(t, x) \leq \beta_1, \\ 0 < \alpha_2 \leq \liminf_{t \rightarrow \infty} \min_{x \in \overline{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \overline{\Omega}} u_2(t, x) \leq \beta_2, \end{aligned} \quad (1.5)$$

where $\alpha_1, \alpha_2, \beta_1, \beta_2$ are constants given by the linear system

$$\begin{aligned} a_1 - C_1 M_{01}^+ \beta_1 - d_1 \beta_2 - \alpha_1 (B_1 - c_1 M_{01}^-) &= 0, \\ a_2 - C_2 M_{02}^+ \beta_2 - d_2 \beta_1 - \alpha_2 (B_2 - c_2 M_{02}^-) &= 0, \\ A_1 - c_1 M_{01}^+ \alpha_1 - d_1 \alpha_2 - \beta_1 (b_1 - c_1 M_{01}^-) &= 0, \\ A_2 - c_2 M_{02}^+ \alpha_2 - d_2 \alpha_1 - \beta_2 (b_2 - c_2 M_{02}^-) &= 0. \end{aligned} \quad (1.6)$$

Remark 1.2. If $\mu_i(\tau) = \int_0^\tau f_i(t)dt$ with $f_i \in C(\mathbb{R}^+) \cap L^1(\mathbb{R}^+)$, $A = 1$, and $0 < a_i = a_i(t, x) = A_i$, $0 < b_i = b_i(t, x) = B_i$, $c_i = c_i(t, x) = C_i = 1$, then by Theorem 1.1, we get

$$\begin{aligned} 0 < \alpha_1 &\leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_1(t, x) \leq \beta_1, \\ 0 < \alpha_2 &\leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_2(t, x) \leq \beta_2, \end{aligned} \quad (1.7)$$

where $\alpha_1, \alpha_2, \beta_1, \beta_2$ are constants given by the linear system

$$\begin{aligned} a_1 - M_{01}^+ \beta_1 - d_1 \beta_2 - \alpha_1 (b_1 - M_{01}^-) &= 0, \\ a_2 - M_{02}^+ \beta_2 - d_2 \beta_1 - \alpha_2 (b_2 - M_{02}^-) &= 0, \\ a_1 - M_{01}^+ \alpha_1 - d_1 \alpha_2 - \beta_1 (b_1 - M_{01}^-) &= 0, \\ a_2 - M_{02}^+ \alpha_2 - d_2 \alpha_1 - \beta_2 (b_2 - M_{02}^-) &= 0. \end{aligned} \quad (1.8)$$

Solving this system, we have the following result:

$$\begin{aligned} \alpha_1 = \beta_1 &= \frac{a_1(b_2 + M_{02}^+ - M_{02}^-) - a_2 d_1}{(b_1 + M_{01}^+ - M_{01}^-)(b_2 + M_{02}^+ - M_{02}^-) - d_1 d_2}, \\ \alpha_2 = \beta_2 &= \frac{a_2(b_1 + M_{01}^+ - M_{01}^-) - a_1 d_2}{(b_1 + M_{01}^+ - M_{01}^-)(b_2 + M_{02}^+ - M_{02}^-) - d_1 d_2}, \end{aligned} \quad (1.9)$$

which coincides with the result of [9], where the authors considered a system like (1.1) with constant coefficients.

Reaction-diffusion systems with delay have been treated by many authors. There are two ways to approach them. The first one is in the framework of semigroup theory of dynamical systems [1, 10]. The second one is a method of upper and lower solutions, using associated monotone iterations; several authors have studied their dynamic properties [2, 9, 11]. Sometimes the birth and death rates depend on both space and time, so when we consider instantaneous and delayed interference within the species and the diffusive effects of the species, system (1.1) will be the appropriate model.

The way we organize the paper is as follows: we first introduce several results which play an important role in the proof of Theorem 1.1 which we will prove in Section 2. We will provide some numerical simulations in Section 3 in order to illustrate our theory.

The following results are developed in [11]. They considered the Volterra reaction-diffusion equations with variable coefficients:

$$\begin{aligned} \frac{\partial u}{\partial t} &= A \Delta u(t, x) + u(t, x) \left[a(t, x) - b(t, x)u(t, x) - c(t, x) \int_0^\infty u(t - \tau, x) d\mu(\tau) \right], \quad t > 0, \quad x \in \Omega, \\ \frac{\partial u}{\partial \eta} &= 0, \quad t > 0, \quad x \in \partial\Omega, \\ u(t, x) &= \phi(t, x), \quad t \leq 0, \quad x \in \bar{\Omega}, \end{aligned} \quad (1.10)$$

where $0 < a_1 \leq a(t, x) \leq A_1$, $0 < b_1 \leq b(t, x) \leq B_1$, $0 \leq c_1 \leq c(t, x) \leq C_1$, and $\phi \in C^1((-\infty, 0]) \times \bar{\Omega}$ is bounded and nonnegative and $0 \neq \phi(0, \cdot) \in C^1(\bar{\Omega})$. Let $M(t)$ denote the total variation of $\mu(\cdot)$ and define M^\pm, M_{0i}^\pm of the same form as that of M_i^\pm, M_{0i}^\pm , respectively.

Lemma 1.3. Assume that $b_1 > c_1 M_0^-$ and $a_1 - (C_1 M_0^+ A_1)/(b_1 - c_1 M_0^-) > 0$, then the solution of (1.10) satisfies

$$\alpha \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u(t, x) \leq \beta, \quad (1.11)$$

where

$$\begin{aligned} \alpha &= \frac{(b_1 - c_1 M_0^-) a_1 - C_1 M_0^+ A_1}{(b_1 - c_1 M_0^-)(B_1 - c_1 M_0^-) - C_1^2 (M_0^+)^2}, \\ \beta &= \frac{(B_1 - c_1 M_0^-) A_1 - C_1 M_0^+ a_1}{(b_1 - c_1 M_0^-)(B_1 - c_1 M_0^-) - C_1^2 (M_0^+)^2}. \end{aligned} \quad (1.12)$$

If $M^+(t) = 0$, then $\mu(t) = -M^-(t)$. Thus,

$$\begin{aligned} \frac{\partial u}{\partial t} &= A \Delta u(t, x) + u(t, x) \left[a(t, x) - b(t, x) u(t, x) + c(t, x) \int_0^\infty u(t - \tau, x) dM^-(\tau) \right], \quad t > 0, \quad x \in \Omega, \\ \frac{\partial u}{\partial \eta} &= 0, \quad t > 0, \quad x \in \partial \Omega, \\ u(t, x) &= \phi(t, x), \quad t \leq 0, \quad x \in \bar{\Omega}. \end{aligned} \quad (1.13)$$

Lemma 1.4. Assume that $b_1 > c_1 M_0^-$ and $a_1 > 0$, then the solution of (1.13) satisfies

$$0 < \alpha \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u(t, x) \leq \beta, \quad (1.14)$$

where

$$\alpha = \frac{a_1}{B_1 - c_1 M_0^-}, \quad \beta = \frac{A_1}{b_1 - c_1 M_0^-}. \quad (1.15)$$

Lemma 1.5. If $u(t, x)$ is the solution of (1.10), then $0 < u(t, x)$.

Now, we introduce the existence-comparison result for the competition-diffusion system (1.1) which is a particular case of Theorem 2.2 in [12].

Definition 1.6. A pair of smooth functions $\tilde{u} = (\tilde{u}_1, \tilde{u}_2)$ and $\hat{u} = (\hat{u}_1, \hat{u}_2)$ are called upper-lower solutions of (1.1) if $\tilde{u}_i \geq \hat{u}_i$ ($i = 1, 2$) in $\mathbb{R} \times \bar{\Omega}$ and the following differential inequalities hold:

$$\begin{aligned} \frac{\partial \tilde{u}_i}{\partial t} - A \Delta \tilde{u}_i &\geq \tilde{u}_i(t, x) \left[a_i(t, x) - b_i(t, x) \tilde{u}_i(t, x) - c_i(t, x) \int_0^\infty \tilde{u}_i(t - \tau, x) d\mu_i(\tau) - d_i \hat{u}_j(t - r_j, x) \right], \\ & \quad j \neq i, \quad t > 0, \quad x \in \Omega, \\ \frac{\partial \hat{u}_i}{\partial t} - A \Delta \hat{u}_i &\leq \hat{u}_i(t, x) \left[a_i(t, x) - b_i(t, x) \hat{u}_i(t, x) - c_i(t, x) \int_0^\infty \hat{u}_i(t - \tau, x) d\mu_i(\tau) - d_i \tilde{u}_j(t - r_j, x) \right], \\ & \quad j \neq i, \quad t > 0, \quad x \in \Omega, \end{aligned}$$

$$\frac{\partial \hat{u}_i}{\partial \eta} \leq 0 \leq \frac{\partial \tilde{u}_i}{\partial \eta}, \quad t > 0, \quad x \in \partial \Omega,$$

$$\hat{u}_i(t, x) \leq \phi_i(t, x) \leq \tilde{u}_i(t, x), \quad t \leq 0, \quad x \in \bar{\Omega} \quad (i = 1, 2). \quad (1.16)$$

Lemma 1.7. *If there exists a pair of upper-lower solutions $\tilde{u} = (\tilde{u}_1, \tilde{u}_2)$ and $\hat{u} = (\hat{u}_1, \hat{u}_2)$ of (1.1), then the problem (1.1) has a unique solution $u^* = (u_1^*, u_2^*)$ and $\hat{u}_i \leq u_i^* \leq \tilde{u}_i$, $i = 1, 2$.*

2. Proof of the main result

The method of proof is via successive improvements of upper-lower solutions of suitable systems.

For given $\phi = (\phi_1, \phi_2)$ as initial conditions for the system (1.1), let K_1, K_2 be constants such that

$$K_1 \geq \max \left\{ \|\phi_1\|, \frac{A_1}{b_1 - c_1 M_{01}^-} \right\}, \quad K_2 \geq \max \left\{ \|\phi_2\|, \frac{A_2}{b_2 - c_2 M_{02}^-} \right\}, \quad (2.1)$$

where $\|\phi_i\| = \sup\{|\phi_i(t, x)| : (t, x) \in (-\infty, 0] \times \bar{\Omega}\}$, $i = 1, 2$.

Then, $(0, 0)$ and (K_1, K_2) are a pair of lower-upper solutions of (1.1). By Lemma 1.7, there exists a unique global nonnegative solution (u_1, u_2) of (1.1) and it satisfies $0 \leq u_1(t, x) \leq K_1$, $0 \leq u_2(t, x) \leq K_2$.

Define $\bar{u}_1^{(1)}(x, t)$ and $\bar{u}_2^{(1)}(x, t)$ by

$$\begin{aligned} \frac{\partial \bar{u}_1^{(1)}}{\partial t} &= A \Delta \bar{u}_1^{(1)} + \bar{u}_1^{(1)} \left[a_1(t, x) - b_1(t, x) \bar{u}_1^{(1)} + c_1(t, x) \int_0^\infty \bar{u}_1^{(1)}(t - \tau, x) dM_1^-(\tau) \right] \quad t > 0, \quad x \in \Omega, \\ \frac{\partial \bar{u}_2^{(1)}}{\partial t} &= A \Delta \bar{u}_2^{(1)} + \bar{u}_2^{(1)} \left[a_2(t, x) - b_2(t, x) \bar{u}_2^{(1)} + c_2(t, x) \int_0^\infty \bar{u}_2^{(1)}(t - \tau, x) dM_2^-(\tau) \right] \quad t > 0, \quad x \in \Omega, \\ \frac{\partial \bar{u}_1^{(1)}}{\partial \eta} &= \frac{\partial \bar{u}_2^{(1)}}{\partial \eta} = 0, \quad t > 0, \quad x \in \partial\Omega, \\ \bar{u}_i^{(1)}(t, x) &= K_i, \quad t \leq 0, \quad x \in \bar{\Omega} \quad (i = 1, 2). \end{aligned} \quad (2.2)$$

Then, $(0, 0)$ and $(\bar{u}_1^{(1)}, \bar{u}_2^{(1)})$ are lower and upper solutions, and by Lemma 1.7,

$$0 \leq u_1 \leq \bar{u}_1^{(1)}, \quad 0 \leq u_2 \leq \bar{u}_2^{(1)}. \quad (2.3)$$

By Lemma 1.4, we can get

$$0 < \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} \bar{u}_i^{(1)}(t, x) \leq \beta_i^{(0)}, \quad (2.4)$$

where

$$\beta_i^{(0)} = \frac{A_i}{b_i - c_i M_{0i}^-}, \quad i = 1, 2. \quad (2.5)$$

Then from (2.3) and (2.4), we get

$$0 \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_i(t, x) \leq \beta_i^{(0)}. \quad (2.6)$$

From (2.6) and the definition of $M_{0i}^+ = \lim_{t \rightarrow \infty} M_i^+(t)$, we have that for any sufficiently small $\varepsilon > 0$, there exist $t'_1 > 0$ and $t_1 > t'_1$ such that

$$\begin{aligned} \max_{x \in \bar{\Omega}} \bar{u}_i^{(1)}(t, x) &< \beta_i^{(0)} + \varepsilon, \quad \text{for } t \geq t'_1, \\ \bar{u}_i^{(1)}(t - r_i, x) &< \beta_i^{(0)} + \varepsilon, \quad \text{for } t \geq t_1, \\ C_i(M_{0i}^+ - M_i^+(t - t'_1)) &< \varepsilon, \quad \text{for } t \geq t_1. \end{aligned} \quad (2.7)$$

Define $\underline{u}_1^{(1)}(t, x)$ and $\underline{u}_2^{(1)}(t, x)$ by

$$\begin{aligned} \frac{\partial \underline{u}_1^{(1)}}{\partial t} &= A \Delta \underline{u}_1^{(1)} + \underline{u}_1^{(1)} \left[a_1(t, x) - b_1(t, x) \underline{u}_1^{(1)} + c_1(t, x) \int_0^\infty \underline{u}_1^{(1)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - c_1(t, x) \int_0^\infty \bar{u}_1^{(1)}(t - \tau, x) dM_1^+(\tau) - d_1 \bar{u}_2^{(1)}(t - r_2, x) \right], \quad t > t_1, \quad x \in \Omega, \\ \frac{\partial \underline{u}_2^{(1)}}{\partial t} &= A \Delta \underline{u}_2^{(1)} + \underline{u}_2^{(1)} \left[a_2(t, x) - b_2(t, x) \underline{u}_2^{(1)} + c_2(t, x) \int_0^\infty \underline{u}_2^{(1)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - c_2(t, x) \int_0^\infty \bar{u}_2^{(1)}(t - \tau, x) dM_2^+(\tau) - d_2 \bar{u}_1^{(1)}(t - r_1, x) \right], \quad t > t_1, \quad x \in \Omega, \\ \frac{\partial \underline{u}_1^{(1)}}{\partial \eta} &= \frac{\partial \underline{u}_2^{(1)}}{\partial \eta} = 0, \quad t > t_1, \quad x \in \partial \Omega, \\ \underline{u}_i^{(1)}(t, x) &= \frac{1}{2} u_i(t, x), \quad (t, x) \in (-\infty, t_1] \times \bar{\Omega}. \end{aligned} \quad (2.8)$$

Then, $(\underline{u}_1^{(1)}, \underline{u}_2^{(1)})$ and $(\bar{u}_1^{(1)}, \bar{u}_2^{(1)})$ are a pair of lower-upper solutions of (1.1), and by Lemma 1.7,

$$\underline{u}_1^{(1)} \leq u_1 \leq \bar{u}_1^{(1)}, \quad \underline{u}_2^{(1)} \leq u_2 \leq \bar{u}_2^{(1)}. \quad (2.9)$$

From $\bar{u}_i^{(1)}(t, x) \leq K_i$, for all $t \in \mathbb{R}$, and (2.8), for $t \geq t_1$, we get

$$\begin{aligned} &c_i(t, x) \int_0^\infty \bar{u}_i^{(1)}(t - \tau, x) dM_i^+(\tau) + d_i \bar{u}_j^{(1)}(t - r_j, x) \\ &= c_i(t, x) \int_0^{t-t'_1} \bar{u}_i^{(1)}(t - \tau, x) dM_i^+(\tau) + c_i(t, x) \int_{t-t'_1}^\infty \bar{u}_i^{(1)}(t - \tau, x) dM_i^+(\tau) + d_i \bar{u}_j^{(1)}(t - r_j, x) \\ &\leq c_i(t, x) \int_0^{t-t'_1} (\beta_i^{(0)} + \varepsilon) dM_i^+(\tau) + c_i(t, x) \int_{t-t'_1}^\infty K_i dM_i^+(\tau) + d_i (\beta_j^{(0)} + \varepsilon) \\ &\leq C_i (M_i^+(t - t_1) - M(0)) (\beta_i^{(0)} + \varepsilon) + K_i C_i (M_{0i}^+ - M_i^+(t - t_1)) + d_i (\beta_j^{(0)} + \varepsilon) \\ &\leq C_i M_{0i}^+ (\beta_i^{(0)} + \varepsilon) + d_i (\beta_j^{(0)} + \varepsilon) + K_i \varepsilon. \end{aligned} \quad (2.10)$$

It follows from (2.9) that, for $t > t_1$, $x \in \Omega$,

$$\begin{aligned} \frac{\partial \underline{u}_1^{(1)}}{\partial t} &\geq A\Delta \underline{u}_1^{(1)} + \underline{u}_1^{(1)} \left[a_1(t, x) - b_1(t, x)\underline{u}_1^{(1)} + c_1(t, x) \int_0^\infty \underline{u}_1^{(1)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - C_1 M_{01}^+(\beta_1^{(0)} + \varepsilon) - d_1(\beta_2^{(0)} + \varepsilon) - K_1 \varepsilon \right], \\ \frac{\partial \underline{u}_2^{(1)}}{\partial t} &\geq A\Delta \underline{u}_2^{(1)} + \underline{u}_2^{(1)} \left[a_2(t, x) - b_2(t, x)\underline{u}_2^{(1)} + c_2(t, x) \int_0^\infty \underline{u}_2^{(1)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - C_2 M_{02}^+(\beta_2^{(0)} + \varepsilon) - d_2(\beta_1^{(0)} + \varepsilon) - K_2 \varepsilon \right]. \end{aligned} \quad (2.11)$$

By the comparison principle, we get, for $t > t_1$, $x \in \Omega$,

$$\underline{u}_1^{(1)} \geq v_1^{(1)}, \quad \underline{u}_2^{(1)} \geq v_2^{(1)}, \quad (2.12)$$

where $v_1^{(1)}$ and $v_2^{(1)}$ are the solutions of the following problem, respectively:

$$\begin{aligned} \frac{\partial v_1^{(1)}}{\partial t} &= A\Delta v_1^{(1)} + v_1^{(1)} \left[a_1(t, x) - b_1(t, x)v_1^{(1)} + c_1(t, x) \int_0^\infty v_1^{(1)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - C_1 M_{01}^+(\beta_1^{(0)} + \varepsilon) - d_1(\beta_2^{(0)} + \varepsilon) - K_1 \varepsilon \right], \\ \frac{\partial v_1^{(1)}}{\partial \eta} &= 0, \quad t > t_1, \quad x \in \partial\Omega, \\ v_1^{(1)}(t, x) &= \frac{1}{2}u_1(t, x), \quad (t, x) \in (-\infty, t_1] \times \overline{\Omega}, \\ \frac{\partial v_2^{(1)}}{\partial t} &= A\Delta v_2^{(1)} + v_2^{(1)} \left[a_2(t, x) - b_2(t, x)v_2^{(1)} + c_2(t, x) \int_0^\infty v_2^{(1)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - C_2 M_{02}^+(\beta_2^{(0)} + \varepsilon) - d_2(\beta_1^{(0)} + \varepsilon) - K_2 \varepsilon \right], \\ \frac{\partial v_2^{(1)}}{\partial \eta} &= 0, \quad t > t_1, \quad x \in \partial\Omega, \\ v_2^{(1)}(t, x) &= \frac{1}{2}u_2(t, x), \quad (t, x) \in (-\infty, t_1] \times \overline{\Omega}. \end{aligned} \quad (2.13)$$

Using the three initial conditions and ε sufficiently small, we have

$$\begin{aligned} a_1 - C_1 M_{01}^+(\beta_1^{(0)} + \varepsilon) - d_1(\beta_2^{(0)} + \varepsilon) - K_1 \varepsilon &> 0, \\ a_2 - C_2 M_{02}^+(\beta_2^{(0)} + \varepsilon) - d_2(\beta_1^{(0)} + \varepsilon) - K_2 \varepsilon &> 0. \end{aligned} \quad (2.14)$$

By Lemma 1.4, we get

$$\begin{aligned}
0 &< \frac{a_1 - C_1 M_{01}^+ \beta_1^{(0)} - d_1 \beta_2^{(0)}}{B_1 - c_1 M_{01}^-} - \varepsilon \frac{C_1 M_{01}^+ + d_1 + K_1}{B_1 - c_1 M_{01}^-} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} v_1^{(1)}(t, x), \\
0 &< \frac{a_2 - C_2 M_{02}^+ \beta_2^{(0)} - d_2 \beta_1^{(0)}}{B_2 - c_2 M_{02}^-} - \varepsilon \frac{C_2 M_{02}^+ + d_2 + K_2}{B_2 - c_2 M_{02}^-} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} v_2^{(1)}(t, x).
\end{aligned} \tag{2.15}$$

Then from (2.12), (2.14), and ε sufficiently small, we can conclude that

$$\begin{aligned}
0 &< \alpha_1^{(0)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_1(t, x) \leq \beta_1^{(0)}, \\
0 &< \alpha_2^{(0)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_2(t, x) \leq \beta_2^{(0)},
\end{aligned} \tag{2.16}$$

where

$$\alpha_1^{(0)} = \frac{a_1 - C_1 M_{01}^+ \beta_1^{(0)} - d_1 \beta_2^{(0)}}{B_1 - c_1 M_{01}^-}, \quad \alpha_2^{(0)} = \frac{a_2 - C_2 M_{02}^+ \beta_2^{(0)} - d_2 \beta_1^{(0)}}{B_2 - c_2 M_{02}^-}. \tag{2.17}$$

For any sufficiently small $\varepsilon > 0$, there exist $t'_2 > t_1$ and $t_2 > t'_2$ such that

$$\begin{aligned}
\min_{x \in \bar{\Omega}} \underline{u}_i^{(1)}(t, x) &> \alpha_i^{(0)} - \varepsilon, \quad \text{for } t \geq t'_2, \\
\underline{u}_i^{(1)}(t - r_i, x) &> \alpha_i^{(0)} + \varepsilon, \quad \text{for } t \geq t_2, \\
M_{0i}^+ - \varepsilon &< M_i^+(t - t'_2), \quad \text{for } t \geq t_2,
\end{aligned} \tag{2.18}$$

Define $\bar{u}_1^{(2)}$ and $\bar{u}_2^{(2)}$ by

$$\begin{aligned}
\frac{\partial \bar{u}_1^{(2)}}{\partial t} &= A \Delta \bar{u}_1^{(2)} + \bar{u}_1^{(2)} \left[a_1(t, x) - b_1(t, x) \bar{u}_1^{(2)} + c_1(t, x) \int_0^\infty \bar{u}_1^{(2)}(t - \tau, x) dM_1^-(\tau) \right. \\
&\quad \left. - c_1(t, x) \int_0^\infty \underline{u}_1^{(1)}(t - \tau, x) dM_1^+(\tau) - d_1 \underline{u}_2^{(1)}(t - r_2, x) \right], \quad t > t_2, \quad x \in \Omega, \\
\frac{\partial \bar{u}_2^{(2)}}{\partial t} &= A \Delta \bar{u}_2^{(2)} + \bar{u}_2^{(2)} \left[a_2(t, x) - b_2(t, x) \bar{u}_2^{(2)} + c_2(t, x) \int_0^\infty \bar{u}_2^{(2)}(t - \tau, x) dM_2^-(\tau) \right. \\
&\quad \left. - c_2(t, x) \int_0^\infty \underline{u}_2^{(1)}(t - \tau, x) dM_2^+(\tau) - d_2 \underline{u}_1^{(1)}(t - r_1, x) \right], \quad t > t_2, \quad x \in \Omega, \\
\frac{\partial \bar{u}_1^{(2)}}{\partial \eta} &= \frac{\partial \bar{u}_2^{(2)}}{\partial \eta} = 0, \quad t > t_2, \quad x \in \partial \Omega, \\
\bar{u}_i^{(2)}(t, x) &= K_i, \quad (t, x) \in (-\infty, t_2] \times \bar{\Omega} \quad (i = 1, 2).
\end{aligned} \tag{2.19}$$

Then $(\underline{u}_1^{(1)}, \underline{u}_2^{(1)})$ and $(\bar{u}_1^{(2)}, \bar{u}_2^{(2)})$ are a pair of lower-upper solutions of (1.1). By Lemma 1.7,

$$\underline{u}_1^{(1)} \leq u_1 \leq \bar{u}_1^{(2)}, \quad \underline{u}_2^{(1)} \leq u_2 \leq \bar{u}_2^{(2)}. \quad (2.20)$$

From (2.19) and (2.20), for $t > t_2$, $x \in \Omega$, we get

$$\begin{aligned} \frac{\partial \bar{u}_1^{(2)}}{\partial t} &\leq A\Delta \bar{u}_1^{(2)} + \bar{u}_1^{(2)} \left[a_1(t, x) - b_1(t, x)\bar{u}_1^{(2)} + c_1(t, x) \int_0^\infty \bar{u}_1^{(2)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - c_1 M_{01}^+(\alpha_1^{(0)} - \varepsilon) - d_1(\alpha_2^{(0)} - \varepsilon) + \varepsilon c_1 \alpha_1^{(0)} \right], \\ \frac{\partial \bar{u}_2^{(2)}}{\partial t} &\leq A\Delta \bar{u}_2^{(2)} + \bar{u}_2^{(2)} \left[a_2(t, x) - b_2(t, x)\bar{u}_2^{(2)} + c_2(t, x) \int_0^\infty \bar{u}_2^{(2)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - c_2 M_{02}^+(\alpha_2^{(0)} - \varepsilon) - d_2(\alpha_1^{(0)} - \varepsilon) + \varepsilon c_2 \alpha_2^{(0)} \right]. \end{aligned} \quad (2.21)$$

By the comparison principle, for $t > t_2$, $x \in \Omega$, we get

$$\bar{u}_1^{(2)} \leq w_1^{(1)}, \quad \bar{u}_2^{(2)} \leq w_2^{(1)}, \quad (2.22)$$

where $w_1^{(1)}$ and $w_2^{(1)}$ are the solutions of the following problem, respectively:

$$\begin{aligned} \frac{\partial w_1^{(1)}}{\partial t} &= A\Delta w_1^{(1)} + w_1^{(1)} \left[a_1(t, x) - b_1(t, x)w_1^{(1)} + c_1(t, x) \int_0^\infty w_1^{(1)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - c_1 M_{01}^+(\alpha_1^{(0)} - \varepsilon) - d_1(\alpha_2^{(0)} - \varepsilon) + \varepsilon c_1 \alpha_1^{(0)} \right], \quad t > t_2, \quad x \in \Omega, \\ \frac{\partial w_1^{(1)}}{\partial \eta} &= 0, \quad t > t_2, \quad x \in \partial\Omega, \\ w_1^{(1)}(t, x) &= K_1, \quad (t, x) \in (-\infty, t_2] \times \bar{\Omega}, \\ \frac{\partial w_2^{(1)}}{\partial t} &= A\Delta w_2^{(1)} + w_2^{(1)} \left[a_2(t, x) - b_2(t, x)w_2^{(1)} + c_2(t, x) \int_0^\infty w_2^{(1)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - c_2 M_{02}^+(\alpha_2^{(0)} - \varepsilon) - d_2(\alpha_1^{(0)} - \varepsilon) + \varepsilon c_2 \alpha_2^{(0)} \right], \quad t > t_2, \quad x \in \Omega, \\ \frac{\partial w_2^{(1)}}{\partial \eta} &= 0, \quad t > t_2, \quad x \in \partial\Omega, \\ w_2^{(1)}(t, x) &= K_2, \quad (t, x) \in (-\infty, t_2] \times \bar{\Omega}. \end{aligned} \quad (2.23)$$

From (2.16), (2.18), and sufficiently small ε , we get

$$\begin{aligned} a_1 - c_1 M_{01}^+(\alpha_1^{(0)} - \varepsilon) - d_1(\alpha_2^{(0)} - \varepsilon) &> 0, \\ a_2 - c_2 M_{02}^+(\alpha_2^{(0)} - \varepsilon) - d_2(\alpha_1^{(0)} - \varepsilon) &> 0. \end{aligned} \quad (2.24)$$

By Lemma 1.4, we get

$$0 < \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} w_1^{(1)}(t, x) \leq \frac{A_1 - c_1 M_{01}^+ \alpha_1^{(0)} - d_1 \alpha_2^{(0)}}{b_1 - c_1 M_{01}^-} - \varepsilon \frac{c_1 M_{01}^+ + d_1 - c_1 \alpha_1^{(0)}}{b_1 - c_1 M_{01}^-},$$

$$0 < \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} w_2^{(1)}(t, x) \leq \frac{A_2 - c_2 M_{02}^+ \alpha_2^{(0)} - d_2 \alpha_1^{(0)}}{b_2 - c_2 M_{02}^-} - \varepsilon \frac{c_2 M_{02}^+ + d_2 - c_2 \alpha_2^{(0)}}{b_2 - c_2 M_{02}^-}.$$
(2.25)

Then from (2.22), (2.24), and sufficiently small ε ,

$$0 < \alpha_1^{(0)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_1(t, x) \leq \beta_1^{(1)},$$

$$0 < \alpha_2^{(0)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_2(t, x) \leq \beta_2^{(1)},$$
(2.26)

where

$$\beta_1^{(1)} = \frac{A_1 - c_1 M_{01}^+ \alpha_1^{(0)} - d_1 \alpha_2^{(0)}}{b_1 - c_1 M_{01}^-}, \quad \beta_2^{(1)} = \frac{A_2 - c_2 M_{02}^+ \alpha_2^{(0)} - d_2 \alpha_1^{(0)}}{b_2 - c_2 M_{02}^-}.$$
(2.27)

Then,

$$0 < \alpha_1^{(0)} \leq \beta_1^{(1)} \leq \beta_1^{(0)},$$

$$0 < \alpha_2^{(0)} \leq \beta_2^{(1)} \leq \beta_2^{(0)}.$$
(2.28)

For any sufficiently small $\varepsilon > 0$, there exist $t'_3 > t_2$ and $t_3 > t'_3$ such that

$$\max_{x \in \bar{\Omega}} \bar{u}_i^{(2)}(t, x) < \beta_i^{(1)} + \varepsilon, \quad \text{for } t \geq t'_3,$$

$$\bar{u}_i^{(1)}(t - r_i, x) < \beta_i^{(1)} + \varepsilon, \quad \text{for } t \geq t_3,$$
(2.29)

$$C_i(M_{0i}^+ - M_i^+(t - t'_3)) < \varepsilon, \quad \text{for } t \geq t_3.$$

Define $\underline{u}_1^{(2)}(t, x)$ and $\underline{u}_2^{(2)}(t, x)$ by

$$\frac{\partial \underline{u}_1^{(2)}}{\partial t} = A \Delta \underline{u}_1^{(2)} + \underline{u}_1^{(2)} \left[a_1(t, x) - b_1(t, x) \underline{u}_1^{(2)} + c_1(t, x) \int_0^\infty \underline{u}_1^{(2)}(t - \tau, x) dM_1^-(\tau) \right. \\ \left. - c_1(t, x) \int_0^\infty \bar{u}_1^{(2)}(t - \tau, x) dM_1^+(\tau) - d_1 \bar{u}_2^{(2)}(t - r_2, x) \right], \quad t > t_3, \quad x \in \Omega,$$

$$\frac{\partial \underline{u}_2^{(2)}}{\partial t} = A \Delta \underline{u}_2^{(2)} + \underline{u}_2^{(2)} \left[a_2(t, x) - b_2(t, x) \underline{u}_2^{(2)} + c_2(t, x) \int_0^\infty \underline{u}_2^{(2)}(t - \tau, x) dM_2^-(\tau) \right. \\ \left. - c_2(t, x) \int_0^\infty \bar{u}_2^{(2)}(t - \tau, x) dM_2^+(\tau) - d_2 \bar{u}_1^{(2)}(t - r_1, x) \right], \quad t > t_3, \quad x \in \Omega,$$

$$\frac{\partial \underline{u}_1^{(2)}}{\partial \eta} = \frac{\partial \underline{u}_2^{(2)}}{\partial \eta} = 0, \quad t > t_3, \quad x \in \partial \Omega,$$

$$\underline{u}_i^{(2)}(t, x) = \frac{1}{2} u_i(t, x), \quad (t, x) \in (-\infty, t_3] \times \bar{\Omega}.$$

(2.30)

Then, $(\underline{u}_1^{(2)}, \underline{u}_2^{(2)})$ and $(\bar{u}_1^{(2)}, \bar{u}_2^{(2)})$ are a pair of lower-upper solutions of (1.1). By Lemma 1.7,

$$\underline{u}_1^{(2)} \leq u_1 \leq \bar{u}_1^{(2)}, \quad \underline{u}_2^{(2)} \leq u_2 \leq \bar{u}_2^{(2)}. \quad (2.31)$$

From (2.30) and (2.31), for $t > t_3$, $x \in \Omega$, we get

$$\begin{aligned} \frac{\partial \underline{u}_1^{(2)}}{\partial t} &\geq A\Delta \underline{u}_1^{(2)} + \underline{u}_1^{(2)} \left[a_1(t, x) - b_1(t, x)\underline{u}_1^{(2)} + c_1(t, x) \int_0^\infty \underline{u}_1^{(2)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - C_1 M_{01}^+(\beta_1^{(1)} + \varepsilon) - d_1(\beta_2^{(1)} + \varepsilon) - K_1 \varepsilon \right], \\ \frac{\partial \underline{u}_2^{(2)}}{\partial t} &\geq A\Delta \underline{u}_2^{(2)} + \underline{u}_2^{(2)} \left[a_2(t, x) - b_2(t, x)\underline{u}_2^{(2)} + c_2(t, x) \int_0^\infty \underline{u}_2^{(2)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - C_2 M_{02}^+(\beta_2^{(1)} + \varepsilon) - d_2(\beta_1^{(1)} + \varepsilon) - K_2 \varepsilon \right]. \end{aligned} \quad (2.32)$$

By the comparison principle, for $t > t_3$, $x \in \Omega$, we get

$$\underline{u}_1^{(2)} \geq v_1^{(2)}, \quad \underline{u}_2^{(2)} \geq v_2^{(2)}, \quad (2.33)$$

where $v_1^{(2)}$ and $v_2^{(2)}$ are the solutions of the following problem, respectively:

$$\begin{aligned} \frac{\partial v_1^{(2)}}{\partial t} &= A\Delta v_1^{(2)} + v_1^{(2)} \left[a_1(t, x) - b_1(t, x)v_1^{(2)} + c_1(t, x) \int_0^\infty v_1^{(2)}(t - \tau, x) dM_1^-(\tau) \right. \\ &\quad \left. - C_1 M_{01}^+(\beta_1^{(1)} + \varepsilon) - d_1(\beta_2^{(1)} + \varepsilon) - K_1 \varepsilon \right], \quad t > t_3, \quad x \in \Omega, \\ \frac{\partial v_1^{(2)}}{\partial \eta} &= 0, \quad t > t_3, \quad x \in \partial\Omega, \\ v_1^{(2)}(t, x) &= \frac{1}{2}u_1(t, x), \quad (t, x) \in (-\infty, t_3] \times \bar{\Omega}, \\ \frac{\partial v_2^{(2)}}{\partial t} &= A\Delta v_2^{(2)} + v_2^{(2)} \left[a_2(t, x) - b_2(t, x)v_2^{(2)} + c_2(t, x) \int_0^\infty v_2^{(2)}(t - \tau, x) dM_2^-(\tau) \right. \\ &\quad \left. - C_2 M_{02}^+(\beta_2^{(1)} + \varepsilon) - d_2(\beta_1^{(1)} + \varepsilon) - K_2 \varepsilon \right], \quad t > t_3, \quad x \in \Omega, \\ \frac{\partial v_2^{(2)}}{\partial \eta} &= 0, \quad t > t_3, \quad x \in \partial\Omega, \\ v_2^{(2)}(t, x) &= \frac{1}{2}u_2(t, x), \quad (t, x) \in (-\infty, t_3] \times \bar{\Omega}. \end{aligned} \quad (2.34)$$

From (2.29), (2.30), and ε sufficiently small, we get

$$\begin{aligned} a_1 - C_1 M_{01}^+ (\beta_1^{(1)} + \varepsilon) - d_1 (\beta_2^{(1)} + \varepsilon) - K_1 \varepsilon &> 0, \\ a_2 - C_2 M_{02}^+ (\beta_2^{(1)} + \varepsilon) - d_2 (\beta_1^{(1)} + \varepsilon) - K_2 \varepsilon &> 0. \end{aligned} \quad (2.35)$$

By Lemma 1.4 we get

$$\begin{aligned} 0 &< \frac{a_1 - C_1 M_{01}^+ \beta_1^{(1)} - d_1 \beta_2^{(1)}}{B_1 - c_1 M_{01}^-} - \varepsilon \frac{C_1 M_{01}^+ + d_1 + K_1}{B_1 - c_1 M_{01}^-} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} v_1^{(1)}(t, x), \\ 0 &< \frac{a_2 - C_2 M_{02}^+ \beta_2^{(1)} - d_2 \beta_1^{(1)}}{B_2 - c_2 M_{02}^-} - \varepsilon \frac{C_2 M_{02}^+ + d_2 + K_2}{B_2 - c_2 M_{02}^-} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} v_2^{(1)}(t, x). \end{aligned} \quad (2.36)$$

From (2.33), (2.35), and for ε , we conclude that

$$\begin{aligned} 0 &< \alpha_1^{(1)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_1(t, x) \leq \beta_1^{(1)}, \\ 0 &< \alpha_2^{(1)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_2(t, x) \leq \beta_2^{(1)}, \end{aligned} \quad (2.37)$$

where

$$\alpha_1^{(1)} = \frac{a_1 - C_1 M_{01}^+ \beta_1^{(1)} - d_1 \beta_2^{(1)}}{B_1 - c_1 M_{01}^-}, \quad \alpha_2^{(1)} = \frac{a_2 - C_2 M_{02}^+ \beta_2^{(1)} - d_2 \beta_1^{(1)}}{B_2 - c_2 M_{02}^-}. \quad (2.38)$$

Then,

$$\begin{aligned} 0 &< \alpha_1^{(0)} \leq \alpha_1^{(1)} \leq \beta_1^{(1)} \leq \beta_1^{(0)}, \\ 0 &< \alpha_2^{(0)} \leq \alpha_2^{(1)} \leq \beta_2^{(1)} \leq \beta_2^{(0)}. \end{aligned} \quad (2.39)$$

Define the sequences $\alpha_1^{(k)}$, $\alpha_2^{(k)}$, $\beta_1^{(k)}$, and $\beta_2^{(k)}$ as follows:

$$\begin{aligned} \alpha_1^{(k)} &= \frac{a_1 - C_1 M_{01}^+ \beta_1^{(k)} - d_1 \beta_2^{(k)}}{B_1 - c_1 M_{01}^-}, & \alpha_2^{(k)} &= \frac{a_2 - C_2 M_{02}^+ \beta_2^{(k)} - d_2 \beta_1^{(k)}}{B_2 - c_2 M_{02}^-}, \\ \beta_1^{(k+1)} &= \frac{A_1 - c_1 M_{01}^+ \alpha_1^{(k)} - d_1 \alpha_2^{(k)}}{b_1 - c_1 M_{01}^-}, & \beta_2^{(k+1)} &= \frac{A_2 - c_2 M_{02}^+ \alpha_2^{(k)} - d_2 \alpha_1^{(k)}}{b_2 - c_2 M_{02}^-}, \\ \beta_1^{(0)} &= \frac{A_1}{b_1 - c_1 M_{01}^-}, & \beta_2^{(0)} &= \frac{A_2}{b_2 - c_2 M_{02}^-}. \end{aligned} \quad (2.40)$$

Lemma 2.1. *For the above-defined sequences, one has*

$$\left[\alpha_1^{(k+1)}, \beta_1^{(k+1)} \right] \subseteq \left[\alpha_1^{(k)}, \beta_1^{(k)} \right], \quad \left[\alpha_2^{(k+1)}, \beta_2^{(k+1)} \right] \subseteq \left[\alpha_2^{(k)}, \beta_2^{(k)} \right], \quad k \geq 0. \quad (2.41)$$

Proof. For $k = 0$, it has been shown that $[\alpha_1^{(1)}, \beta_1^{(1)}] \subseteq [\alpha_1^{(0)}, \beta_1^{(0)}]$ and $[\alpha_2^{(1)}, \beta_2^{(1)}] \subseteq [\alpha_2^{(0)}, \beta_2^{(0)}]$. Using induction, we can complete the proof.

Lemma 2.1 implies that

$$\lim_{k \rightarrow \infty} \alpha_1^{(k)}, \quad \lim_{k \rightarrow \infty} \alpha_2^{(k)}, \quad \lim_{k \rightarrow \infty} \beta_1^{(k)}, \quad \lim_{k \rightarrow \infty} \beta_2^{(k)} \quad (2.42)$$

exist, denoted as α_1 , α_2 , β_1 , and β_2 , respectively. From (2.41), we have the following linear system by which we can obtain the numbers α_1 , α_2 , β_1 , and β_2 :

$$\begin{aligned} \alpha_1 &= \frac{a_1 - C_1 M_{01}^+ \beta_1 - d_1 \beta_2}{B_1 - c_1 M_{01}^-}, & \alpha_2 &= \frac{a_2 - C_2 M_{02}^+ \beta_2 - d_2 \beta_1}{B_2 - c_2 M_{02}^-}, \\ \beta_1 &= \frac{A_1 - c_1 M_{01}^+ \alpha_1 - d_1 \alpha_2}{b_1 - c_1 M_{01}^-}, & \beta_2 &= \frac{A_2 - c_2 M_{02}^+ \alpha_2 - d_2 \alpha_1}{b_2 - c_2 M_{02}^-}. \end{aligned} \quad (2.43)$$

□

Lemma 2.2. *For the solutions of (1.1), one has*

$$0 < \alpha_1^{(k)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_1(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_1(t, x) \leq \beta_1^{(k)}, \quad k \geq 0, \quad (2.44)$$

$$0 < \alpha_2^{(k)} \leq \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} u_2(t, x) \leq \limsup_{t \rightarrow \infty} \max_{x \in \bar{\Omega}} u_2(t, x) \leq \beta_2^{(k)}, \quad k \geq 0. \quad (2.45)$$

Proof. We have shown that (2.45) and (3.1) are valid for $k = 0, 1$. Using induction and repeating the above process, we can complete the proof. □

Combining the above lemmas, we can complete the proof of Theorem 1.1

Remark 2.3. Following the same kind of proof for Theorem 1.1, it can be shown that the same conclusions hold if instead of system (1.1) we work with solutions of

$$\frac{\partial v_1}{\partial t} = A \Delta v_1(t, x) + v_1(t, x) \left[a_1(t, x) - b_1(t, x) v_1(t, x) - c_1(t, x) \int_0^\infty v_1(t - \tau, x) d\mu_1(\tau) - d_1 v_2(t - r_2, x) \right],$$

$$t > 0, \quad x \in \Omega,$$

$$\frac{\partial v_2}{\partial t} = B \Delta v_2(t, x) + v_2(t, x) \left[a_2(t, x) - b_2(t, x) v_2(t, x) - c_2(t, x) \int_0^\infty v_2(t - \tau, x) d\mu_2(\tau) - d_2 v_1(t - r_1, x) \right],$$

$$t > 0, \quad x \in \Omega,$$

$$\frac{\partial v_1}{\partial \eta} = \frac{\partial v_2}{\partial \eta} = 0, \quad t > 0, \quad x \in \partial \Omega,$$

$$v_i(t, x) = \phi_i(t, x) \quad (i = 1, 2) \quad t \leq 0, \quad x \in \bar{\Omega}, \quad (2.46)$$

as expected since they do not include the coefficients of diffusion. We thank one of the referees for his comments regarding this matter.

3. Numerical simulations

In this section, we present some numerical results that agree with Theorem 1.1 proved above. We used the method of upper and lower solutions as developed by Pao [12, 13] discretizing the systems into finite difference systems. On both examples, the domain used is $\Omega = (0, 12)$.

Example 3.1. In this example, we work with coefficients that depend on t and initial values that depend on (t, x) .


Consider the system

$$\begin{aligned} \frac{\partial u_1}{\partial t} &= \frac{\partial^2 u_1}{\partial x^2} + u_1 \left[(\sin(t) + 2) - \left(\frac{1}{1+x^2} + 10 \right) u_1 \right. \\ &\quad \left. - (\cos(t) + 1.01) \int_0^\infty \frac{1}{1+\tau^2} u_1(t-\tau, x) d\tau - \frac{1}{10} u_2(t-2, x) \right], \quad t \in [0, T], x \in [0, 12], \\ \frac{\partial u_2}{\partial t} &= \frac{\partial^2 u_2}{\partial x^2} + u_2 \left[\left(\sqrt{\sin^2(t) + 1} \right) - \left(\frac{x}{1+x^2} + 7.5 \right) u_2 \right. \\ &\quad \left. - (\sin(t^2) + 1.01) \int_0^\infty \frac{1}{1+\tau^2} u_2(t-\tau, x) d\tau - \frac{1}{2} u_1(t-1.5, x) \right], \quad t \in [0, T], x \in [0, 12], \\ \frac{\partial u_i}{\partial x}(0, t) &= \frac{\partial u_i}{\partial x}(12, t) = 0 \quad (i = 1, 2), t > 0, \\ u_1(t, x) &= \frac{1}{2} \sin\left(\frac{1}{2}t\right) + 1.01, \quad t \leq 0, x \in [0, 12], \\ u_2(t, x) &= \frac{1}{2} \left(\sin\left(\frac{1}{2}t\right) + \sin\left(\frac{1}{2}x\right) \right) + 1.01, \quad t \leq 0, x \in [0, 12]. \end{aligned} \tag{3.1}$$


It is easy to see that this system satisfies the conditions of the main theorem (Theorem 1.1), therefore the global attractors are defined by the solutions of the linear system

$$\begin{aligned} \alpha_1(11 - 0) &= 1 - 2.01 \frac{\pi}{2} \beta_1 - \frac{1}{10} \beta_2, \\ \alpha_2(1.5 - 0) &= 1 - 2.01 \frac{\pi}{2} \beta_2 - \beta_1, \\ \beta_1(10 - 0) &= 3 - 0.01 \frac{\pi}{2} \alpha_1 - \frac{1}{10} \alpha_2, \\ \beta_2(0.5 - 0) &= 2 - 0.01 \frac{\pi}{2} \alpha_2 - \alpha_1, \end{aligned} \tag{3.2}$$

that is, the global attractors for u_1 and u_2 are given by $[0.002245, 0.299884]$ and $[0.011192, 0.285680]$, respectively. Figure 1 shows the numerical simulation of the solution of this system and on it we can notice the global attractors obtained before.



(a)




(b)


Figure 1: Numerical simulation of Example 3.1.

Example 3.2. According to the main theorem (Theorem 1.1), the global attractors depend only on the coefficients and not on the initial values. In this example, we only change the initial functions of the above example:

$$\begin{aligned}
 u_1(t, x) &= \sin(x) + 1.01, \quad t \leq 0, \quad x \in [0, 12], \\
 u_2(t, x) &= \frac{1}{200}(1 + t^2) \sin\left(\frac{\pi x}{12}\right) + 0.01, \quad t \leq 0, \quad x \in [0, 12],
 \end{aligned} \tag{3.3}$$



(a)



(b)

Figure 2: Numerical simulation of Example 3.2.

to obtain the same attractors as in Example 3.1. Figure 2 shows the numerical simulation of the solution of this system and we observe that they have the attractors expected.

Acknowledgments

The research of the first and second authors was supported in part by CONACYT under Grant SEP-2003-C02-44029. The research of the second author was supported in part by SNI under Grant 15284. The research of the third author was supported in part by UADY under Grant PRIORI-FMAT-04-002.

References

- [1] S. G. Ruan and X.-Q. Zhao, "Persistence and extinction in two species reaction-diffusion systems with delays," *Journal of Differential Equations*, vol. 156, no. 1, pp. 71–92, 1999.
- [2] X. Lu, "Persistence and extinction in a competition-diffusion system with time delays," *The Canadian Applied Mathematics Quarterly*, vol. 2, no. 2, pp. 231–246, 1994.
- [3] S. A. Gourley and S. Ruan, "Convergence and travelling fronts in functional differential equations with nonlocal terms: a competition model," *SIAM Journal on Mathematical Analysis*, vol. 35, no. 3, pp. 806–822, 2003.
- [4] W. Feng and F. Wang, "Asymptotic periodicity and permanence in a competition-diffusion system with discrete delays," *Applied Mathematics and Computation*, vol. 89, no. 1–3, pp. 99–110, 1998.
- [5] L. Zhou, Y. Tang, and S. Hussein, "Periodic bifurcation solution for a delay competition diffusion system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 9, pp. 6073–6084, 2001.
- [6] W. Feng and X. Lu, "Harmless delays for permanence in a class of population models with diffusion effects," *Journal of Mathematical Analysis and Applications*, vol. 206, no. 2, pp. 547–566, 1997.
- [7] C. V. Pao, "Global asymptotic stability of Lotka-Volterra 3-species reaction-diffusion systems with time delays," *Journal of Mathematical Analysis and Applications*, vol. 281, no. 1, pp. 186–204, 2003.
- [8] C. V. Pao, "Global asymptotic stability of Lotka-Volterra competition systems with diffusion and time delays," *Nonlinear Analysis: Real World Applications*, vol. 5, no. 1, pp. 91–104, 2004.
- [9] Y. Wang and Y. Meng, "Asymptotic behavior of a competition-diffusion system with time delays," *Mathematical and Computer Modelling*, vol. 38, no. 5-6, pp. 509–517, 2003.
- [10] J. Wu and X.-Q. Zhao, "Diffusive monotonicity and threshold dynamics of delayed reaction diffusion equations," *Journal of Differential Equations*, vol. 186, no. 2, pp. 470–484, 2002.
- [11] B. Shi and Y. Chen, "A priori bounds and stability of solutions for a Volterra reaction-diffusion equation with infinite delay," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 44, no. 1, pp. 97–121, 2001.
- [12] C. V. Pao, "Systems of parabolic equations with continuous and discrete delays," *Journal of Mathematical Analysis and Applications*, vol. 205, no. 1, pp. 157–185, 1997.
- [13] C. V. Pao, *Nonlinear Parabolic and Elliptic Equations*, New York, NY, USA, Plenum Press, 1992.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br