

BLOWUP OF SOLUTIONS OF A NONLINEAR WAVE EQUATION

ABBES BENAÏSSA AND SALIM A. MESSAOUDI

Received 4 April 2001 and in revised form 20 October 2001

We establish a blowup result to an initial boundary value problem for the nonlinear wave equation $u_{tt} - M(\|B^{1/2}u\|^2)Bu + ku_t = |u|^{p-2}u$, $x \in \Omega$, $t > 0$.

1. Introduction

We consider the initial boundary value problem (IBVP) for the nonlinear wave equation

$$\begin{aligned} u_{tt} - Au + ku_t &= |u|^{p-2}u, & x \in \Omega, t > 0, \\ u(x, t) &= 0, & x \in \partial\Omega, t \geq 0, \\ u(x, 0) &= u_0(x), \quad u_t(x, 0) = u_1(x), & x \in \Omega, \end{aligned} \quad (1.1)$$

where

$$\begin{aligned} Au &= M\left(\|B^{1/2}u\|^2\right)e^{-\Phi(x)} \operatorname{div}\left(e^{\Phi(x)}\nabla u\right), \\ \|B^{1/2}u\|^2 &= \int_{\Omega} e^{\Phi(x)}|\nabla u|^2 dx, \end{aligned} \quad (1.2)$$

$p > 2$ is a constant, k is a positive constant, $M : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function, $\Phi \in L^\infty(\Omega)$, and $\Omega \subset \mathbb{R}^n$ is a bounded domain with a smooth boundary Γ so that the divergence theorem can be applied.

When $M \equiv 1$ and $\Phi \equiv 0$, for the case $k = 0$, it is well known that the source term $|u|^{p-2}u$ is responsible for finite blowup (global nonexistence) of solutions with negative initial energy (see [1, 9]). The interaction

between the damping term and the source has been first considered by Levine [11, 12]. For $k > 0$, the author showed that solutions, with negative initial energy, blow up in finite time. In [5], Georgiev and Todorova extended Levine’s result to the case of nonlinear damping of the form $|u_t|^m u_t$. This result was generalized to an abstract setup by Levine and Serrin [14], Levine et al. [13], and Vitillaro [18]. In [16], Messaoudi extended the result of Levine to the situation where $\Phi \neq 0$.

When $\Phi \equiv 0$ and M is not a constant function, the equation without the damping and source terms is often called the wave equation of Kirchhoff type which has been introduced by Kirchhoff [10] in order to study the nonlinear vibrations of an elastic string. The existence of local and global solutions in Sobolev and Gevrey classes was investigated by many authors (see [2, 3, 4, 6, 7, 8, 15, 17]).

In the present paper, we investigate the blowup of solutions of the initial boundary value problem (1.1). We show that, for suitably chosen initial data, any strong solution blows up in finite time. Our work is based on the results of [14].

2. Main result

In order to state our main result, we introduce the weighted space

$$L^s(\Omega, \Phi) := \left\{ v : \Omega \rightarrow \mathbb{R} / \int_{\Omega} e^{\Phi(x)} |u_0|^s dx < \infty \right\}, \tag{2.1}$$

$$E(0) = \frac{1}{2} \int_{\Omega} e^{\Phi(x)} u_1^2 dx + \frac{1}{2} \bar{M} \left(\|B^{1/2} u_0\|^2 \right) - \frac{1}{p} \int_{\Omega} e^{\Phi(x)} |u_0|^p dx.$$

We also make the following hypothesis:

$$M \in C(\mathbb{R}_+, \mathbb{R}_+), \quad \bar{M}(s) = \int_0^s M(k) dk, \tag{2.2}$$

such that

$$r \bar{M}(s) \geq s M(s), \quad \forall s \geq 0, \quad 1 < r < \frac{p}{2}. \tag{2.3}$$

THEOREM 2.1. *Let $p > 2$ and assume that (2.2) and (2.3) hold. Then, for any initial data satisfying $E(0) < 0$, the solution of (1.1) blows up in finite time.*

Proof. Except for the operator Au , this problem is similar to [14, problem (4.1)–(4.3)] for $l = 2$. So the proof goes exactly like the one of [14, Theorem 5]. It remains only to show that Au and $F(u) = |u|^{p-2}u$ satisfy

conditions (1s) and (2s) in [14, page 346]. To do this, we set

$$\begin{aligned} V &= Y = L^2(\Omega, \Phi), & W &= L^p(\Omega, \Phi), \\ D &= H_0^1(\Omega, \Phi) = \{u \in H_0^1(\Omega, \Phi) / u, \nabla u \in Y\}. \end{aligned} \tag{2.4}$$

It is clear that A and F are Frechet derivatives of the C^1 real-valued potentials given by

$$\mathcal{A}u = \frac{1}{2} \bar{M}(\|B^{1/2}u\|^2), \quad \mathcal{F}(u) = \frac{1}{p} \|u\|_W^p. \tag{2.5}$$

Now we have, by virtue of (2.3),

$$\begin{aligned} \langle Au, u \rangle_V &= \int_{\Omega} e^{\Phi(x)} u M(\|B^{1/2}u\|^2) e^{-\Phi(x)} \operatorname{div}(e^{\Phi(x)} \nabla u) \\ &= M(\|B^{1/2}u\|^2) \int_{\Omega} u \operatorname{div}(e^{\Phi(x)} \nabla u) \\ &= (\|B^{1/2}u\|^2) \|B^{1/2}u\|^2 \leq r \bar{M}(\|B^{1/2}u\|^2) \leq 2r \mathcal{A}u, \\ \langle F(u), u \rangle_V - 2r \mathcal{F}(u) &= \left(1 - \frac{2r}{p}\right) \|u\|_W^p = (p - 2r) \mathcal{F}(u). \end{aligned} \tag{2.6}$$

Therefore, conditions (1s) and (2s) in [14, page 346] are satisfied. This completes the proof. □

Remark 2.2. Conditions (3s) and (1d)–(3d) of [14] are automatically satisfied since $Pu_t = u_t$ and $Q(t, u_t) = ku_t$ are linear. See the proof of [14, Theorem 5].

Acknowledgments

The authors would like to thank the referee for his valuable remarks. Special thanks from the second author goes to King Fahd University of Petroleum and Minerals Library for its continuous support.

References

- [1] J. M. Ball, *Remarks on blow-up and nonexistence theorems for nonlinear evolution equations*, Quart. J. Math. Oxford Ser. (2) **28** (1977), no. 112, 473–486.
- [2] P. D’Ancona and S. Spagnolo, *Global solvability for the degenerate Kirchhoff equation with real analytic data*, Invent. Math. **108** (1992), no. 2, 247–262.
- [3] R. W. Dickey, *Infinite systems of nonlinear oscillation equations related to the string*, Proc. Amer. Math. Soc. **23** (1969), 459–468.

- [4] ———, *Infinite systems of nonlinear oscillation equations*, J. Differential Equations **8** (1970), 16–26.
- [5] V. Georgiev and G. Todorova, *Existence of a solution of the wave equation with nonlinear damping and source terms*, J. Differential Equations **109** (1994), no. 2, 295–308.
- [6] D. Gourdin and M. Mechab, *Problème de Goursat non linéaire dans les espaces de Gevrey pour les équations de Kirchhoff généralisées*, J. Math. Pures Appl. (9) **75** (1996), no. 6, 569–593 (French).
- [7] F. Hirose, *Global solvability for the generalized degenerate Kirchhoff equation with real-analytic data in \mathbf{R}^n* , Tsukuba J. Math. **21** (1997), no. 2, 483–503.
- [8] K. Kajitani and K. Yamaguti, *On global real analytic solutions of the degenerate Kirchhoff equation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **21** (1994), no. 2, 279–297.
- [9] V. K. Kalantarov and O. A. Ladyzhenskaya, *The occurrence of collapse for quasilinear equations of parabolic and hyperbolic type*, J. Soviet Math. **10** (1978), 53–70.
- [10] G. Kirchhoff, *Vorlesungen über Mechanik*, Teubner, Leipzig, 1883 (German).
- [11] H. A. Levine, *Instability and nonexistence of global solutions to nonlinear wave equations of the form $Pu_{tt} = -Au + \mathcal{F}(u)$* , Trans. Amer. Math. Soc. **192** (1974), 1–21.
- [12] ———, *Some additional remarks on the nonexistence of global solutions to nonlinear wave equations*, SIAM J. Math. Anal. **5** (1974), 138–146.
- [13] H. A. Levine, S. Ro Park, and J. Serrin, *Global existence and global nonexistence of solutions of the Cauchy problem for a nonlinearly damped wave equation*, J. Math. Anal. Appl. **228** (1998), no. 1, 181–205.
- [14] H. A. Levine and J. Serrin, *Global nonexistence theorems for quasilinear evolution equations with dissipation*, Arch. Rational Mech. Anal. **137** (1997), no. 4, 341–361.
- [15] L. A. Medeiros and M. M. Miranda, *Solutions for the equation of nonlinear vibrations in Sobolev spaces of fractionary order*, Mat. Apl. Comput. **6** (1987), no. 3, 257–276.
- [16] S. A. Messaoudi, *Blow up in solutions of a semilinear wave equation*, Int. J. Appl. Math. **1** (1999), no. 6, 621–626.
- [17] S. Spagnolo, *The Cauchy problem for Kirchhoff equations*, Rend. Sem. Mat. Fis. Milano **62** (1992), 17–51 (1994).
- [18] E. Vittilaro, *Global nonexistence theorems for a class of evolution equation with dissipation*, Arch. Rational Mech. Anal. **137** (1997), 341–361.

Abbes Benaissa: Department of Mathematics, Djillali Liabes University, P.O. Box 89, Sidi Bel Abbes 22000, Algeria

Salim A. Messaoudi: Mathematical Sciences Department, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

E-mail address: messaoud@kfupm.edu.sa

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru