

ON THE CONTROL OF A TRUNCATED GENERAL IMMIGRATION PROCESS THROUGH THE INTRODUCTION OF A PREDATOR

E. G. KYRIAKIDIS

Received 17 December 2003; Accepted 15 February 2005

This paper is concerned with the problem of controlling a truncated general immigration process, which represents a population of harmful individuals, by the introduction of a predator. If the parameters of the model satisfy some mild conditions, the existence of a control-limit policy that is average-cost optimal is proved. The proof is based on the uniformization technique and on the variation of a fictitious parameter over the entire real line. Furthermore, an efficient Markov decision algorithm is developed that generates a sequence of improving control-limit policies converging to the optimal policy.

Copyright © 2006 E. G. Kyriakidis. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In many problems dealing with the optimal control of a stochastic process under the criterion of minimizing the expected long-run average cost per unit time it is possible to prove that the optimal policy initiates the controlling action if and only if the state of the process exceeds a critical level. Such a policy is usually called control-limit policy. A method that in some problems leads to the proof of the optimality of a control-limit policy is a parametric analysis introduced by Federgruen and So [2] in a queueing model. According to this method first it is shown that an optimal control-limit policy exists when a parameter (possibly fictitious) takes sufficiently small values. This assertion is then extended inductively from interval to interval of the parameter values. An important advantage of the Federgruen-So method is that in many cases, as a corollary, it can be proved that any local minimum within the set of the average costs of control-limit policies is a global minimum within this set. This result enables us to compute very quickly the optimal policy using the usual bisection procedure or a special-purpose policy iteration algorithm that creates a sequence of strictly improving control-limit policies.

The present paper is concerned with the problem of controlling a pest population, which grows stochastically according to a general immigration process in a habitat with finite capacity, through the introduction of a predator. It is assumed that the predator

2 Control of a truncated general immigration process

captures the pests one at a time and then emigrates from the habitat. The capture rate of the predator depends on the number of pests. A finite-state continuous time Markov decision model is constructed and it is proved that there exists an average- cost optimal control-limit policy, if the parameters of the model satisfy some mild conditions. The proof is based on Federgruen-So method.

Note that the Federgruen-So method has been applied to two other Markov decision models for pest control. These models differ from the present one in the way the pest population grows or in the way the pest population is controlled. Specifically, in the first of these models (see [7]) it was assumed that the pest population grows according to a general immigration process in a habitat with finite capacity and it is controlled through total catastrophes, which annihilate instantaneously the pest population size. In the second model (see [8]) it was assumed that the pest population grows according to a Poisson process in a habitat with unlimited capacity and it is controlled through the introduction of a predator. The capture rate of the predator was assumed to be constant.

The structure of the rest of the paper is as follows. In Section 2 we give a detailed description of the Markov decision process. In Section 3, firstly, a necessary and sufficient condition is found under which the condition of never controlling is optimal. When this condition fails, the optimality of control-limit policies is shown by applying the Federgruen-So technique. In Section 4 a tailor-made policy iteration algorithm is developed that generates a sequence of improving control-limit policies and converges to the optimal policy.

2. The model

Consider a population of individuals that cause some kind of damage (e.g., pests) which grow stochastically according to a general immigration process in a habitat with carrying capacity N , where N is a positive integer. Assume that the immigration rate that corresponds to each state i , $0 \leq i \leq N - 1$ is equal to $\nu_i > 0$. The immigration rate ν_N that corresponds to the state N is necessarily equal to zero, since N is the carrying capacity of the habitat. It is assumed that the damage done by the pests is represented by a cost c_i , $0 \leq i \leq N$, for each unit of time during which the population size is i . We impose the natural assumptions that the sequence $\{c_i\}$ is non decreasing and $c_0 = 0$.

We suppose that there is a controller who observes the evolution of the population continuously and may take an action that introduces a predator in the habitat, whenever a new state is entered. That is, the controller takes actions on a discrete-time mode. More specifically, we assume that there exists a controlling mechanism which can be in one of two modes: on or off. Whenever the mechanism is turned off the pest population evolves without being influenced. When it is turned on, a predator is introduced in the habitat after some random time that is exponentially distributed. The presence of the predator immediately stops the immigrations of the pests, that is, the rates ν_i , $0 \leq i \leq N - 1$, take immediately the value 0. As soon as the predator is introduced in the habitat, it captures the pests one at a time until their population size is reduced to zero and then it emigrates with rate $\vartheta > 0$. It is assumed that the predator captures the pests with rate $\sigma_i > 0$ when their population size is i , $1 \leq i \leq N$. The unit of time has been chosen in such a way that

the rate at which the predator is introduced in the habitat is equal to one. Thus, when the controlling mechanism is on, the length of time until the introduction of the predator is exponentially distributed with unit mean. Whenever the controlling mechanism is on, it incurs a cost of $K > 0$ per time unit.

Let i and i' be the states of the process at which the population size of the pests is i , $0 \leq i \leq N$, and the predator is absent from their habitat or present, respectively. A stationary policy f is defined by a sequence $\{f_i : 0 \leq i \leq N\}$ where f_i is the action taken when the process is at state i . It is assumed that $f_i = 1$, when the controlling mechanism is on, and $f_i = 0$ when the controlling mechanism is off. If the stationary policy $f \equiv \{f_i : 0 \leq i \leq N\}$ is used, our assumptions imply that we have a continuous time Markov chain model for the population growth of the pests with state space $S = \{0, 0', 1, 1', \dots, N, N'\}$.

Our goal is to find a policy that minimizes the expected long-run average cost per unit time for every initial state among all stationary policies. The decision epochs include the epochs at which an immigration of a pest occurs and the epochs at which the predator emigrates. An intuitively appealing class of policies is the class of control-limit policies $\{P_n : n = 0, 1, \dots, N\}$, where P_n is the stationary policy under which the controlling action is taken if and only if the population size of the pests is equal to or exceeds n . It seems reasonable that the optimal policy will be of control-limit type if K is sufficiently small. In an earlier paper (see [6]) a similar model was introduced, in which the pest population grows in a habitat with unlimited capacity according to a simple immigration process. The cost rates c_i and the captures rates were taken as $c_i = i$ and $\sigma_i = \sigma$, $i \geq 0$. In that work the optimality of a particular control-limit policy within the wider class of all stationary policies was established by proving that it satisfies the optimality equation and certain conditions given by [1].

In the present model, it seems difficult to repeat the same proof since the expression for the average cost under a control-limit policy is too complicated. However, if we impose some mild conditions on the parameters of the model, we can prove the existence of an optimal control-limit policy by applying the Federgruen-So technique, which, as it was mentioned in the previous section, is based on a variation of a parameter over the entire real line. The same technique has been applied in some other queueing and maintenance models (see Federgruen & So [3, 4], So [13], So & Tang [14, 15]) and in two pest control models (see Kyriakidis [7, 8]).

The conditions that we impose on the parameters of the model are given below.

Condition 1.

$$\sum_{j=i}^N \left(\prod_{k=i+1}^j \frac{\nu_{k-1}}{1 + \nu_k} \right) \left(\sum_{k=1}^j \sigma_k^{-1} \right) \leq (1 + \nu_i) \left(\nu_{i-1} \sigma_i^{-1} + \sum_{j=1}^i \sigma_j^{-1} \right), \quad (2.1)$$

with $1 \leq i < N$ and $\prod_{k=i+1}^i \equiv 1$.

Condition 2.

$$\sum_{j=i}^N \left((1 + \nu_i)^{-1} \prod_{k=i+1}^j \frac{\nu_{k-1}}{1 + \nu_k} \right) \left(c_j + \sum_{k=1}^j \frac{c_k}{\sigma_k} \right) \geq \frac{\nu_{i-1} c_i}{\sigma_i} + c_{i-1} + \sum_{j=1}^i \frac{c_j}{\sigma_j}, \quad (2.2)$$

with $1 \leq i < N$ and $\prod_{k=i+1}^i \equiv 1$.

4 Control of a truncated general immigration process

The proposition below, which can be proved by induction on i , gives a sufficient condition for the validity of Condition 1.

PROPOSITION 2.1. *If $\nu_0 \geq \nu_1 \geq \dots \geq \nu_{N-1}$ and $\sigma_1 \leq \sigma_2 \leq \dots \leq \sigma_N$ then Condition 1 holds.*

3. The optimality of control-limit policies

If the process is never controlled the long-run average cost per unit time is c_N since N is an absorbing state in this case. In the proposition below a necessary and sufficient condition is given under which the policy of never controlling is optimal. Its proof is presented in the appendix.

PROPOSITION 3.1. *The policy that never introduces the predator in the habitat is optimal if and only if*

$$c_N \left(\frac{1}{\nu_0} + \frac{1}{\vartheta} \right) + \sum_{i=1}^{N-1} (c_N - c_i) \left(\frac{1}{\nu_i} + \frac{1}{\sigma_i} \right) \leq K. \quad (3.1)$$

Assume now that the relation (3.1) is not valid. In this case the policy that never introduces the predator is not optimal. The average cost of a stationary policy which prescribes action 0 at state N is equal to the average cost of the policy that never introduces the predator since N is an absorbing state under such a stationary policy. Consequently, we can restrict ourselves only to the stationary policies that prescribe action 1 at state N . All the results that we will present in the rest of this section are concerned with the optimal policy among these stationary policies. Let r be a real number (possibly negative) that represents a fictitious cost incurred each unit of time the process is occupying the state $0'$. In Theorem 3.5 it will be shown that a control-limit policy is optimal for any fixed value of r , in particular for $r = 0$.

Let $T_{i0'}^{(n)}$ and $T_{i'0'}^{(n)}$, $0 \leq i \leq N$, be the expected time until the process under the policy P_n , $0 \leq n \leq N$, reaches the state $0'$, given that the initial state is i or i' , respectively. Let also $C_{i0'}^{(n)}$ and $C_{i'0'}^{(n)}$, $0 \leq i \leq N$, be the expected cost until the process under the policy P_n , $0 \leq n \leq N$, reaches the state $0'$, given that the initial state is i or i' , respectively. Conditioning on the first transition from the state i , we obtain:

$$T_{i0'}^{(n)} = \frac{1 + \nu_i T_{i+1,0'}^{(n)} + \sum_{j=1}^i \sigma_j^{-1}}{\nu_i + 1}, \quad n \leq i \leq N-1, \quad (3.2)$$

$$C_{i0'}^{(n)} = \frac{c_i + K + \nu_i C_{i+1,0'}^{(n)} + \sum_{j=1}^i c_j / \sigma_j}{\nu_i + 1}, \quad n \leq i \leq N-1. \quad (3.3)$$

Note also that

$$T_{N0'}^{(n)} = 1 + \sum_{j=1}^N \frac{1}{\sigma_j}, \quad C_{N0'}^{(n)} = c_N + K + \sum_{j=1}^N \frac{c_j}{\sigma_j}. \quad (3.4)$$

Given the above values of $T_{N0'}^{(n)}$ and $C_{N0'}^{(n)}$, the quantities $T_{i0'}^{(n)}$ and $C_{i0'}^{(n)}$, $i = N-1, \dots, n$ can be found from (3.2) and (3.3), recursively.

Let g_n denote the expected long-run average cost per unit time under the policy P_n , $0 \leq n \leq N$. The process under the policy P_n is a regenerative process, where the successive entries into state $0'$ can be taken as regenerative epochs between successive cycles. From a well-known regenerative argument (see [11, Proposition 5.9]) it follows that g_n is equal to the expected cost of a cycle divided by the expected time of the cycle. Hence,

$$g_n = \frac{\sum_{i=0}^{n-1} c_i/\nu_i + C_{n0'}^{(n)} + r/\vartheta}{\sum_{i=0}^{n-1} 1/\nu_i + T_{n0'}^{(n)} + 1/\vartheta}, \quad 0 \leq n \leq N. \quad (3.5)$$

Let $h_i^{(n)}$, $0 \leq i \leq N$, be the relative value associated with the policy P_n , $0 \leq n \leq N$, that corresponds to the state i and let $w_i^{(n)}$, $0 \leq i \leq N$, be the relative value associated with the policy P_n , $0 \leq n \leq N$, that corresponds to the state i' . These quantities are defined by (see relation (3.1.7) in Tijms [16])

$$h_i^{(n)} = C_{i0'}^{(n)} - g_n T_{i0'}^{(n)}, \quad (3.6)$$

$$w_i^{(n)} = C_{i'0'}^{(n)} - g_n T_{i'0'}^{(n)}. \quad (3.7)$$

Clearly,

$$w_0^{(n)} = 0, \quad (3.8)$$

since $g_n = C_{0'0'}^{(n)}/T_{0'0'}^{(n)}$, by the usual regenerative argument. According to the semi-Markov version of Theorem 3.1.1 in Tijms [16] (see [16, page 220]) the numbers $h_i^{(n)}$, $w_i^{(n)}$, $0 \leq i \leq N$ and g_n satisfy the system of equations:

$$h_i^{(n)} = \frac{c_i - g_n}{\nu_i} + h_{i+1}^{(n)}, \quad 0 \leq i \leq n-1, \quad (3.9)$$

$$h_i^{(n)} = \frac{c_i + K - g_n + \nu_i h_{i+1}^{(n)} + w_i^{(n)}}{\nu_i + 1}, \quad n \leq i \leq N-1, \quad (3.10)$$

$$w_i^{(n)} = \frac{c_i - g_n}{\sigma_i} + w_{i-1}^{(n)}, \quad 1 \leq i \leq n, \quad (3.11)$$

$$w_0^{(n)} = \frac{r - g_n}{\vartheta} + h_0^{(n)}. \quad (3.12)$$

Let $A_i^{(n)} = T_{i0'}^{(n)} - \sum_{j=1}^i \sigma_j^{-1}$ and $B_i^{(n)} = C_{i0'}^{(n)} - \sum_{j=1}^i (c_j/\sigma_j)$.

The results of Lemmas 3.2, 3.3 and 3.4 will be used in the proof of Theorem 3.5. The proof of Lemma 3.2 is similar to the proof of Proposition 3 in [8] and the proof of Lemma 3.3 is similar to the proof of Lemma 2 in [7].

LEMMA 3.2. *The policy P_n is optimal if and only if*

$$\begin{aligned} \frac{c_i - g_n}{\nu_i} + h_{i+1}^{(n)} &\leq \frac{c_i + K - g_n + \nu_i h_{i+1}^{(n)} + w_i^{(n)}}{\nu_i + 1}, \quad 0 \leq i \leq n-1, \\ \frac{c_i + K - g_n + \nu_i h_{i+1}^{(n)} + w_i^{(n)}}{\nu_i + 1} &\leq \frac{c_i - g_n}{\nu_i} + h_{i+1}^{(n)}, \quad n \leq i \leq N-1. \end{aligned} \quad (3.13)$$

6 Control of a truncated general immigration process

LEMMA 3.3. Assume that the policy P_n , $n < N$, is optimal for some fixed value R of the parameter r . Then, it is impossible for the policy P_n to be optimal for all $r \geq R$ (simultaneously).

LEMMA 3.4. (i) Condition 1 implies that the sequence $\{A_i^{(n)}\}$ is non-increasing in i , $n \leq i < N$, for each $n = 0, 1, \dots$

(ii) Condition 2 implies that the sequence $\{B_i^{(n)}\}$ is non-decreasing in i , $n \leq i < N$, for each $n = 0, 1, \dots$

THEOREM 3.5. There exists a sequence $R_0 < R_1 \leq R_2 \leq \dots \leq R_N < R_{N+1}$ with $R_0 = -\infty$ and $R_{N+1} = +\infty$ such that the policy P_n , $0 \leq n \leq N$ is optimal for all $r \in [R_n, R_{n+1}]$, where $R_{n+1} = \sup\{w : w \geq R_n, \text{ the policy } P_n \text{ is optimal for all } r \in [R_n, w]\}$.

Proof. The proof is by induction on n . We first establish that a number $R > -\infty$ exists such that the policy P_0 is optimal for all $r \leq R$. In view of Lemma 3.2, it suffices to show that the numbers $h_i^{(0)}$ and $w_i^{(0)}$, $0 \leq i \leq N$, and g_0 satisfy the inequalities:

$$\frac{c_i + K - g_0 + \nu_i h_{i+1}^{(0)} + w_i^{(0)}}{\nu_i + 1} \leq \frac{c_i - g_0}{\nu_i} + h_{i+1}^{(0)}, \quad 0 \leq i \leq N - 1. \quad (3.14)$$

Using (3.6) and (3.7), with $n = 0$ the above inequalities reduce to

$$g_0 \left[\nu_i (T_{i+1,0'}^{(0)} - T_{i',0'}^{(0)}) + 1 \right] \leq c_i + \nu_i C_{i+1,0'}^{(0)} - \nu_i C_{i',0'}^{(0)} - \nu_i K, \quad (3.15)$$

with $0 \leq i \leq N - 1$.

Note that the process under P_0 must pass through the state i' before it enters the state $0'$, if the initial state is $i + 1$. Hence, $T_{i+1,0'}^{(0)} - T_{i',0'}^{(0)} > 0$. From (3.5) we have that $g_0 \rightarrow -\infty$ as $r \rightarrow -\infty$. Thus, there exists a number $R > -\infty$ such that (3.15) hold simultaneously for all $r \leq R$. From Lemma 3.3 it follows that $R_1 < +\infty$, where $R_1 = \sup\{w : w \geq R \text{ and the policy } P_0 \text{ is optimal for all } r \leq w\}$.

Suppose that there exists a sequence $R_0 < R_1 \leq R_2 \leq \dots \leq R_n$, where $n < N$, such that the policy P_s , $0 \leq s \leq n$, is optimal for all $r \in [R_s, R_{s+1}]$ with $R_{s+1} = \sup\{w : w \geq R_s \text{ and the policy } P_s \text{ is optimal for all } r \in [R_s, w]\} < +\infty$. We will show that the policy P_{n+1} is optimal for $r = R_{n+1}$. To achieve this, we use the standard uniformization technique (see Serfozo [12]) to transform the original Markov decision process into an equivalent one in which the times between transitions have the same exponential parameter $\nu = \max_{1 \leq i \leq N} \{\nu_0, 1 + \nu_i, \sigma_i, \vartheta\}$ whatever the state and the action are. The reformulated Markov decision process has the same average cost as the original one under any stationary policy. Thus both models have the same optimal policy. Let \tilde{g}_n and $\tilde{h}_i^{(n)}$, $\tilde{w}_i^{(n)}$, $0 \leq i \leq N$, denote the average cost and the relative values under the policy P_n in the new model. Let also $\tilde{T}_{i0'}^{(n)}$, $\tilde{T}_{i',0'}^{(n)}$, and $\tilde{C}_{i0'}^{(n)}$, $\tilde{C}_{i',0'}^{(n)}$, $0 \leq i \leq N$, be the expected times and costs, respectively, until the new process under the policy P_n , $1 \leq n \leq N$, reaches the state $0'$, given that the initial state is i or i' .

Consider now some $\varepsilon > 0$. If $r = R_{n+1} + \varepsilon$, the policy P_n is not optimal for the original and, consequently, for the new model. Hence, according to the corresponding result of Lemma 3.2 for the equivalent model one of the following two cases occurs:

Case 1. For some i with $0 \leq i \leq n-1$:

$$\frac{c_i + K - \tilde{g}_n + \nu_i \tilde{h}_{i+1}^{(n)} + [\nu - (\nu_i + 1)] \tilde{h}_i^{(n)} + \tilde{w}_i^{(n)}}{\nu} < \frac{c_i - \tilde{g}_n + \nu_i \tilde{h}_{i+1}^{(n)} + (\nu - \nu_i) \tilde{h}_i^{(n)}}{\nu}. \quad (3.16)$$

The above inequality is equivalent to $\psi_i(R_{n+1} + \varepsilon) > 0$, with

$$\begin{aligned} \psi_i(r) &= \tilde{h}_i^{(n)} - \tilde{w}_i^{(n)} - K \\ &= \tilde{C}_{i0'}^{(n)} - \tilde{C}_{i'0'}^{(n)} - \tilde{g}_n (\tilde{T}_{i0'}^{(n)} - \tilde{T}_{i'0'}^{(n)}) - K \\ &= C_{i0'}^{(n)} - C_{i'0'}^{(n)} - g_n (T_{i0'}^{(n)} - T_{i'0'}^{(n)}) - K, \end{aligned} \quad (3.17)$$

where the last equality follows from the fact that the original and the reformulated process have the same generator (see Serfozo [12]). Since $\tilde{T}_{i0'}^{(n)} - \tilde{T}_{i'0'}^{(n)} > 0$ and g_n as given in (3.5) is increasing in r we deduce that $\psi_i(r)$ is decreasing in r . Thus,

$$0 < \psi_i(R_{n+1} + \varepsilon) < \psi_i(R_{n+1}) \leq 0, \quad (3.18)$$

where the last inequality follows from the optimality of P_n for $r = R_{n+1}$. Clearly, this is a contradiction and the following Case 2 must arise.

Case 2. For some i with $n \leq i \leq N$:

$$\frac{c_i - \tilde{g}_n + \nu_i \tilde{h}_{i+1}^{(n)} + (\nu - \nu_i) \tilde{h}_i^{(n)}}{\nu} < \frac{c_i + K - \tilde{g}_n + \nu_i \tilde{h}_{i+1}^{(n)} + [\nu - (\nu_i + 1)] \tilde{h}_i^{(n)} + \tilde{w}_i^{(n)}}{\nu}. \quad (3.19)$$

The above inequality is equivalent to $\psi_i(R_{n+1} + \varepsilon) < 0$, with

$$\begin{aligned} \psi_i(r) &= \tilde{h}_i^{(n)} - \tilde{w}_i^{(n)} - K \\ &= C_{i0'}^{(n)} - C_{i'0'}^{(n)} - g_n (T_{i0'}^{(n)} - T_{i'0'}^{(n)}) - K \\ &= B_i^{(n)} - g_n A_i^{(n)} - K. \end{aligned} \quad (3.20)$$

From Lemma 3.4 we deduce that $\psi_i(r)$, $n \leq i \leq N$, is non-decreasing in i . Thus,

$$\psi_n(R_{n+1} + \varepsilon) \leq \psi_i(R_{n+1} + \varepsilon) < 0. \quad (3.21)$$

Consider a sequence $\{\varepsilon_\ell\} \downarrow 0$. In view of the above inequality we have that for all ℓ , $\psi_n(R_{n+1} + \varepsilon_\ell) < 0$. From the continuity of $\psi_n(r)$ in r it follows that $\psi_n(R_{n+1}) \leq 0$. However, $\psi_n(R_{n+1}) \geq 0$ since the policy P_n is optimal for $r = R_{n+1}$. Thus, $\psi_n(R_{n+1}) = 0$. The last equality means that in the new model for $r = R_{n+1}$ the actions prescribed, for each state i , by the policy P_{n+1} minimizes the right-side of the optimality equation (see [16, equation (3.5.4)]), which is satisfied by the numbers $\tilde{h}_i^{(n)}$, $\tilde{w}_i^{(n)}$, $0 \leq i \leq N$, and \tilde{g}_n . Thus the policy P_{n+1} is optimal for $r = R_{n+1}$ in the new and, consequently, in the original model.

8 Control of a truncated general immigration process

Consider again the original model. If $n + 1 < N$, we define $R_{n+2} = \sup\{w : w \geq R_{n+1} \text{ and the policy } P_{n+1} \text{ is optimal for all } r \in [R_{n+1}, w]\}$. From Lemma 3.3 it follows that $R_{n+2} < \infty$. If $n + 1 = N$, it can be shown that the policy P_N is optimal for all $r \geq R_N$, using a similar analysis as in Case 1 without transforming the model. \square

LEMMA 3.6. $\{T_{n0'}^{(n)}\}$, $0 \leq n \leq N$, is non-decreasing in n .

Proof. Conditioning on the first transition from state n we have that

$$T_{n0'}^{(n)} = \frac{1}{\nu_n + 1} + \frac{\nu_n}{\nu_n + 1} T_{n+1,0'}^{(n+1)} + \frac{1}{\nu_n + 1} \sum_{j=1}^n \frac{1}{\sigma_j}. \quad (3.22)$$

Hence,

$$T_{n+1,0'}^{(n+1)} - T_{n0'}^{(n)} = \frac{1}{\nu_n + 1} \left(T_{n+1,0'}^{(n+1)} - 1 - \sum_{j=1}^n \frac{1}{\sigma_j} \right). \quad (3.23)$$

Conditioning on the time until the introduction of the predator we obtain

$$\begin{aligned} T_{n+1,0'}^{(n+1)} &= \int_0^\infty \left[t + \sum_{j=n+1}^N p_{n+1,j}(t) \left(\sum_{k=1}^j \frac{1}{\sigma_k} \right) \right] e^{-t} dt \\ &\geq 1 + \int_0^\infty \left[\sum_{j=n+1}^N p_{n+1,j}(t) \left(\sum_{k=1}^j \frac{1}{\sigma_k} \right) \right] e^{-t} dt \\ &= 1 + \sum_{k=1}^n \frac{1}{\sigma_j}, \end{aligned} \quad (3.24)$$

where $p_{n+1,j}(t)$ is the probability that the state of the (uncontrolled) general immigration process at time t will be j , given that the state at time 0 is $n + 1$. The relations (3.23) and (3.24) give the result of the lemma. \square

From (3.5) and Lemma 3.6 we deduce that g_n can be written as

$$g_n = r\pi_n + \tilde{g}_n, \quad (3.25)$$

where \tilde{g}_n is independent of r and π_n is decreasing in n . Using this result and Theorem 3.5, the following proposition, which will be useful in the computation of the optimal control-limit policy, can be proved in the same way as the Lemma 5.2 in Federgruen and So [2].

PROPOSITION 3.7. For any fixed r any local minimum within the set $\{g_n : 0 \leq n \leq N\}$ is a global minimum within this set.

4. The computation of the optimal policy

In this section we assume that $r = 0$. So, we consider again the model introduced in Section 2. In view of Theorem 3.5, if condition (3.1) fails, there exists an optimal control-limit policy P_{n^*} . From Proposition 3.7 it follows that the optimal critical point n^* can be

found by the standard bisection procedure or by a tailor-made policy iteration algorithm. The tailor-made policy iteration algorithm, which is based on Tijms's embedding technique (see Tijms [16, page 234]), generates a sequence of strictly improving control-limit policies that converges to P_{n^*} . Similar algorithms have developed in queueing, inventory and maintenance models (see [10] and [16, Section 3.6]) and in other pest control models (see [5, 7]). From a great number of examples we have tested it seems that the tailor-made policy iteration algorithm is more efficient than the bisection procedure.

Tailor-made policy iteration algorithm

Step 1. Check (3.1). If it is true then the policy of never controlling is optimal. Otherwise go to Step 2.

Step 2 (Initialization). Choose an initial critical integer n , $0 \leq n \leq N$.

Step 3 (Value-determination step). For the current policy P_n , compute its average cost g_n , using (3.2), (3.3), (3.4), (3.5) and the associated relative values $h_i^{(n)}$, $0 \leq i \leq n$, using (3.8), (3.10), (3.12).

Step 4 (Policy-improvement step). (a) Find, if it exists, the smallest integer \tilde{n} such that $1 \leq \tilde{n} < n$ and

$$\frac{c_i + K - g_n + \nu_i h_{i+1}^{(n)} + w_i^{(n)}}{\nu_i + 1} \leq h_i^{(n)}, \quad \tilde{n} \leq i < n, \quad (4.1)$$

where, $w_i^{(n)}$ is computed from the relations (3.8) and (3.11), and go to Step 3 with n replaced by \tilde{n} . Else go to (b).

(b) Find, if it exists, the largest integer \tilde{n} such that $n < \tilde{n} \leq N$ and

$$\frac{c_i - g_n}{\nu_i} + h_{i+1}^{(n)} \leq h_i^{(n)}, \quad n \leq i \leq \tilde{n} - 1. \quad (4.2)$$

The numbers $h_i^{(n)}$, $n + 1 \leq i \leq N$, can be found, if it is necessary, by (3.2), (3.3), (3.4), (3.6).

Step 5 (Convergence test). If it is not possible to find an integer \tilde{n} such that Steps 4(a) or 4(b) are satisfied, then the algorithm is stopped. The optimal policy is P_n and its average cost is g_n .

We give as illustration a numerical example in which $N = 160$, $\nu_i = 20(1 - i/N)$, $\sigma_i = 40$, $c_i = i$, $1 \leq i \leq N$, $\vartheta = 30$, $K = 80$. This example clearly satisfies the condition of Proposition 2.1 and, therefore, Condition 1 holds. It can be also verified numerically that Condition 2 holds. If the initial policy is the policy P_{160} the successive policies that are generated by the algorithm are the policies P_{160} , P_8 , P_{45} , P_{31} , P_{33} with average costs 129.7, 56.87, 47.58, 46.47, 46.44, respectively.

Appendix

Proof of Proposition 3.1. Suppose that the policy of never controlling is optimal. Its average cost is equal to c_N . Assume that (3.1) is not true. From the well-known regenerative

argument (see Ross [11, Proposition 5.9]) it follows that the average cost g_N under the policy P_N is given by

$$g_N = \frac{\sum_{i=0}^{N-1} (c_i/\nu_i) + c_N + K + \sum_{i=1}^N (c_i/\sigma_i)}{\sum_{i=0}^{N-1} (1/\nu_i) + 1 + \sum_{i=1}^N (1/\sigma_i) + (1/\vartheta)}. \quad (\text{A.1})$$

It can be seen that $g_N < c_N$. This is a contradiction.

Suppose that (3.1) holds. From Miller [9, Theorem 10] it follows that the policy of never controlling is optimal if there exist two sequences $\{h_i\}$ and $\{w_i\}$ that correspond to the states i and i' , $0 \leq i \leq N$, respectively, such that

$$\begin{aligned} c_N &= c_i + \nu_i h_{i+1} - \nu_i h_i, & 0 \leq i \leq N-1 \\ c_N &\leq c_i + K + \nu_i h_{i+1} + w_i - (\nu_i + 1)h_i, & 1 \leq i \leq N \\ c_N &= c_i + \sigma_i w_{i-1} - \sigma_i w_i, & 1 \leq i \leq N \\ c_N &= c_0 + \vartheta h_0 - \vartheta w_0. \end{aligned} \quad (\text{A.2})$$

It can be readily checked that the expressions:

$$\begin{aligned} h_i &= \sum_{j=0}^{i-1} \frac{c_N - c_j}{\nu_j} + \frac{c_N}{\vartheta} + w_0, & 0 \leq i \leq N \\ w_i &= \sum_{j=1}^i \frac{c_j - c_N}{\sigma_j} + w_0, & 0 \leq i \leq N \end{aligned} \quad (\text{A.3})$$

satisfy the above four relations for any value of w_0 . Hence the policy of never controlling is optimal. \square

Proof of Lemma 3.4. (i) To prove that for each $n = 0, 1, \dots, N-1$ the sequence $\{A_i^{(n)}\}$ is non-increasing in i , $n \leq i < N$, it suffices to show that

$$T_{i+1,0'}^{(0)} - T_{i0'}^{(0)} \leq \frac{1}{\sigma_{i+1}}, \quad 0 \leq i < N-1. \quad (\text{A.4})$$

Using (3.2) we see that the above relation is equivalent to the following one:

$$T_{i+1,0'}^{(0)} \leq 1 + \frac{\nu_i + 1}{\sigma_{i+1}} + \sum_{j=1}^i \frac{1}{\sigma_j}, \quad 1 \leq i+1 < N. \quad (\text{A.5})$$

Conditioning on the time until the introduction of the predator we obtain that

$$\begin{aligned} T_{i+1,0'}^{(n+1)} &= \int_0^\infty \left[t + \sum_{j=i+1}^N p_{i+1,j}(t) \left(\sum_{k=1}^j \frac{1}{\sigma_k} \right) \right] e^{-t} dt \\ &= 1 + \sum_{j=i+1}^N p_{i+1,j}^* \left(\sum_{k=1}^j \frac{1}{\sigma_k} \right), \end{aligned} \quad (\text{A.6})$$

where, $p_{i+1,j}(t)$ is the probability that the state of the (uncontrolled) general truncated immigration process will be j at time t given that the state at time 0 is $i+1$, and

$$p_{i+1,j}^* = \int_0^\infty p_{i+1,j}(t)e^{-t} dt. \quad (\text{A.7})$$

Using (A.6) we see that (A.5) is equivalent to

$$\sum_{j=i}^N p_{ij}^* \left(\sum_{k=i}^j \sigma_k^{-1} \right) \leq \nu_{i-1} \sigma_i^{-1} + \sum_{j=i}^i \sigma_j^{-1}, \quad 1 \leq i < N. \quad (\text{A.8})$$

Taking Laplace transforms with respect to t in the Kolmogorov forward equation for the probabilities $p_{ij}(t)$ we obtain the following expression for $p_{i,j}^*$, $i \leq j \leq N$.

$$p_{ij}^* = \left[\prod_{k=i+1}^j \nu_{k-1} (1 + \nu_k)^{-1} \right] (1 + \nu_i)^{-1}, \quad i \leq j \leq N, \quad \prod_{k=i+1}^i \equiv 1. \quad (\text{A.9})$$

Using (A.9) it can be seen that the relation (A.8) is equivalent to Condition 1.

(ii) To prove that, for each $n = 0, 1, \dots, N-1$ the sequence $\{B_i^{(n)}\}$ is non-decreasing in i , $n \leq i < N$ it suffices to show that

$$C_{i+1,0'}^{(0)} + C_{i0'}^{(0)} \geq \frac{c_{i+1}}{\sigma_{i+1}}, \quad 0 \leq i < N-1. \quad (\text{A.10})$$

Using (3.3) we see that the above relation is equivalent to the following one:

$$C_{i+1,0'}^{(0)} \geq c_i + K + \sum_{j=1}^{i+1} \frac{c_j}{\sigma_j} + \frac{\nu_i c_{i+1}}{\sigma_{i+1}}, \quad 1 \leq i+1 < N. \quad (\text{A.11})$$

Conditioning on the time until the introduction of the predator we obtain that

$$C_{i+1,0'}^{(0)} = \int_0^\infty \left[\int_0^t \{E[c_{X(s)} | X(0) = i+1] + K\} ds + \sum_{j=i+1}^N p_{i+1,j}(t) \left(\sum_{k=1}^j \frac{c_k}{\sigma_k} \right) \right] e^{-t} dt, \quad (\text{A.12})$$

where $X(s)$ is the population size of the (uncontrolled) truncated general immigration process at time s . Applying a well-known property of Laplace transforms (e.g., see Tijms [16, page 362]) the above expression reduces to

$$\begin{aligned} C_{i+1,0'}^{(0)} &= \int_0^\infty E[c_{X(t)} | X(0) = i+1] e^{-t} dt \\ &+ \int_0^\infty \left[\sum_{j=i+1}^N p_{i+1,j}(t) \left(\sum_{k=1}^j \frac{c_k}{\sigma_k} \right) \right] e^{-t} dt + K \\ &= \sum_{j=i+1}^N p_{i+1,j}^* c_j + \sum_{j=i+1}^N p_{i+1,j}^* \left(\sum_{k=1}^j \frac{c_k}{\sigma_k} \right) + K. \end{aligned} \quad (\text{A.13})$$

Using (A.13) it can be seen that (A.11) is equivalent to

$$\sum_{j=1}^N p_{ij}^* \left(c_j + \sum_{k=1}^j \frac{c_k}{\sigma_k} \right) \geq c_{i-1} + \sum_{j=1}^i \frac{c_j}{\sigma_j} + \frac{\nu_{i-1} c_i}{\sigma_i}, \quad 1 \leq i < N. \quad (\text{A.14})$$

In view of (A.9) the last relation is equivalent to Condition 2.

References

- [1] J. Bather, *Optimal stationary policies for denumerable Markov chains in continuous time*, Advances in Applied Probability **8** (1976), no. 1, 144–158.
- [2] A. Federgruen and K. C. So, *Optimal time to repair a broken server*, Advances in Applied Probability **21** (1989), no. 2, 376–397.
- [3] ———, *Optimal maintenance policies for single-server queueing systems subject to breakdowns*, Operations Research **38** (1990), no. 2, 330–343.
- [4] ———, *Optimality of threshold policies in single-server queueing systems with server vacations*, Advances in Applied Probability **23** (1991), no. 2, 388–405.
- [5] E. G. Kyriakidis, *A Markov decision algorithm for optimal pest control through uniform catastrophes*, European Journal of Operational Research **64** (1993), no. 1, 38–44.
- [6] ———, *Optimal pest control through the introduction of a predator*, European Journal of Operational Research **81** (1995), no. 2, 357–363.
- [7] ———, *Optimal control of a truncated general immigration process through total catastrophes*, Journal of Applied Probability **36** (1999), no. 2, 461–472.
- [8] ———, *Optimal control of a simple immigration process through the introduction of a predator*, Probability in the Engineering and Informational Sciences **17** (2003), no. 1, 119–135.
- [9] B. L. Miller, *Finite state continuous time Markov decision processes with an infinite planning horizon*, Journal of Mathematical Analysis and Applications **22** (1968), no. 3, 552–569.
- [10] R. D. Nobel and H. C. Tijms, *Optimal control for an $M^X/G/1$ queue with two service modes*, European Journal of Operational Research **113** (1999), no. 3, 610–619.
- [11] S. M. Ross, *Applied Probability Models With Optimization Applications*, Dover, New York, 1992.
- [12] R. F. Serfozo, *An equivalence between continuous and discrete time Markov decision processes*, Operations Research **27** (1979), no. 3, 616–620.
- [13] K. C. So, *Optimality of control limit policies in replacement models*, Naval Research Logistics **39** (1992), no. 5, 685–697.
- [14] K. C. So and C. S. Tang, *Optimal batch sizing and repair strategies for operations with repairable jobs*, Management Science **41** (1995), no. 5, 894–908.
- [15] ———, *Optimal operating policy for a bottleneck with random rework*, Management Science **41** (1995), no. 4, 620–636.
- [16] H. C. Tijms, *Stochastic Models. An Algorithmic Approach*, Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics, John Wiley & Sons, Chichester, 1994.

E. G. Kyriakidis: Department of Financial and Management Engineering, University of the Aegean, 31 Fostini Str., 82100 Chios, Greece

E-mail address: kyriak@aegean.gr

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br